

$$\Phi = \chi^{-1} \Psi \chi.$$

In such a case the iterates of the two maps will also be conjugate and, if they are numerical methods, they will have similar stability properties and performance (e.g. the same effective order). It is difficult to separate the relevance of the two properties in cases where symplectic and reversible maps are conjugate. This is however rarely the case and certainly does not hold generically for discrete maps in many dimensions [209]. There is no direct correspondence between reversible and symplectic maps, however each class of maps admits certain theorems of dynamical systems which are in many ways analogous (for example, the Kolmogorov-Arnold-Moser, or KAM, theory for symplectic maps near elliptic fixed points [386] has an analogue for reversible maps [97]).

Of particular importance for molecular dynamics are the following properties: a symplectic map will preserve volume, whereas a time-reversible map need not do so, and a symplectic integrator will approximately conserve energy due to the existence of the perturbed Hamiltonian, whereas a time-reversible integrator may give rise to a drift in energy [166].

3.7 Molecular Dynamics with Hard-Sphere Collisions

We close this chapter with a discussion of molecular dynamics with hard-core repulsion. This is a system which is not, properly speaking, Hamiltonian, but which has dynamics that are largely defined by a Hamiltonian system. The added complication is the nonsmooth character of the motion as impulses are applied at points of collision. The discussion below is a summary of work in [40, 184].

We assume, as usual a Hamiltonian $H = \mathbf{p}^T \mathbf{M}^{-1} \mathbf{p} / 2 + U(\mathbf{q})$ but we add the inequality constraint $\|\mathbf{q}_i - \mathbf{q}_j\| \geq \sigma_i + \sigma_j$ where $\sigma_i, i = 1, \dots, N$ is a core radius. The condition $\|\mathbf{q}_i - \mathbf{q}_j\| = \sigma_i + \sigma_j$, some i, j defines the constraint surface. When the particles are not touching, they move along Newtonian paths defined by the standard equations of motion. At impact, they exchange momentum and energy according to the rules of elastic collision. Specifically, at the point of contact, the momentum vectors of the two spheres are adjusted according to the rule:

$$\mathbf{p} := \mathbf{p} + \alpha \mathbf{u}_\perp$$

where $\mathbf{u}_\perp$ is normal to the constraint surface and α is a parameter chosen to maintain the conservation of energy. Let $\mathcal{R}_c$ denote the action of the collision operator on the vector of positions and momenta. Then we can write the evolution formally as

$$\mathcal{F}_\tau^{\text{h.s.}}(\mathbf{q}, \mathbf{p}) = \mathcal{G}_{\Delta\tau_r} \circ \mathcal{R}_c \circ \mathcal{G}_{\Delta\tau_{r-1}} \dots \circ \mathcal{R}_c \circ \mathcal{G}_{\Delta\tau_0}$$