

Chapter 3 Describing, Exploring, and Comparing Data

Measures of Center (3.1)

Objectives

In this lesson, we will learn how to

- describe the meaning of a measure of center;
- compute and interpret the mean, median, mode, and midrange;
- decide when a measure of center is resistant;
- apply round-off rules for measures of center;
- compute an approximate mean from a frequency distribution;
- compute a weighted mean, such as a grade-point average.

1. Measures of Center

A **measure of center** is a value at the center or middle of a data set.

Common measures of center include

mean, median, mode, midrange.

Big Idea

A measure of center gives us a single value that represents a typical or central value in a data set. However, different measures of center can give different answers.

2. Mean

The **mean**, or arithmetic mean, is found by adding all data values and dividing by the number of data values.

Terminology Caution

The word *average* is sometimes used for the mean, but it can also refer loosely to other measures of center. In statistics, be precise: say mean, median, mode, or midrange.

Mean

For a sample,

$$\bar{x} = \frac{\sum x}{n}.$$

For a population,

$$\mu = \frac{\sum x}{N}.$$

Notation

- $\sum$ means "the sum of."
- x represents an individual data value.
- n is the number of values in a sample.
- N is the number of values in a population.
- $\bar{x}$, pronounced "x-bar," is the sample mean.
- μ , pronounced "mu," is the population mean.

Important Properties of the Mean

- The mean uses every data value.
- Sample means from the same population tend to vary less than other measures of center.
- The mean can be greatly affected by an extreme value, or outlier.

A statistic is **resistant** if extreme values do not cause it to change very much. The mean is not resistant.

Example 1: Mean: Find the mean of the first eleven wait times, in minutes, for Space Mountain at 10 AM:

50, 25, 75, 35, 50, 25, 30, 50, 45, 25, 20.

3. Median

The **median** is the middle value when the original data values are arranged in order.

Finding the Median

First sort the data values.

- If the number of values is odd, the median is the exact middle value.
- If the number of values is even, the median is the mean of the two middle values.

Important Properties of the Median

- The median is resistant because it is not affected much by a few extreme values.
- The median does not directly use every data value.

Example 2: Median with an Odd Number of Values: Find the median of the first eleven wait times, in minutes, for Space Mountain at 10 AM:

50, 25, 75, 35, 50, 25, 30, 50, 45, 25, 20.

Compare the Mean and Median

For these same wait times, the mean is 39.1 minutes while the median is 35.0 minutes. The relatively high value of 75 pulls the mean upward, but it has much less effect on the median.

Example 3: Median with an Even Number of Values: Find the median after including the twelfth wait time:

50, 25, 75, 35, 50, 25, 30, 50, 45, 25, 20, 50.

3A. Choosing Between the Mean and Median

Which Measure Should We Use?

- Use the mean when the distribution is approximately symmetric and has no influential outliers.
- Prefer the median when the distribution is strongly skewed or contains extreme values.
- Always examine the distribution before choosing a measure of center.

Outlier Experiment: In the original eleven wait times, replace the largest value, 75, with 750. What happens to the mean and median?

4. Mode

The **mode** is the value or values that occur with the greatest frequency.

Finding the Mode

- A data set can have one mode.
- A data set can have more than one mode.
- A data set can have no mode.
- The mode can be used with qualitative data.

When two values occur with the same greatest frequency, the data set is called **bimodal**. When more than two values occur with the same greatest frequency, the data set is called **multimodal**.

Example 4: Mode: Find the mode of the first eleven wait times, in minutes, for Tower of Terror at 10 AM:

35, 35, 20, 50, 95, 75, 45, 50, 30, 35, 30.

Other Mode Examples:

- The wait times 30, 30, 50, 50, 75 have two modes: 30 and 50.

- The wait times 20, 30, 35, 50, 75 have no mode because no value is repeated.

5. Midrange

The **midrange** is the value halfway between the maximum and minimum values.

Midrange

$$\text{midrange} = \frac{\text{maximum value} + \text{minimum value}}{2}.$$

The midrange is very easy to compute, but it uses only the maximum and minimum values, so it is not resistant. It is rarely used in practice. Do not confuse the midrange with the median: the midrange uses only the two extremes, while the median depends on the ordered position of the data.

Example 5: Midrange: Find the midrange of the first eleven wait times, in minutes, for Space Mountain at 10 AM:

50, 25, 75, 35, 50, 25, 30, 50, 45, 25, 20.

6. Round-Off Rules

Round-Off Rules for Measures of Center

- For the mean, median, and midrange, round to one more decimal place than the original data.
- For the mode, leave the value as is, because the mode is one of the original data values.

7. Critical Thinking with Measures of Center

We can calculate measures of center from many sets of numbers, but we should ask whether the calculation is meaningful.

Be Careful

Mean and median may not be meaningful when the numbers are really labels, ranks, jersey numbers, or values from a biased list.

Even when a measure of center is mathematically valid, it can be misleading if the data come from a biased or unrepresentative sample. Always consider how the data were collected and whether the sample represents the population of interest.

Example 6: When Measures of Center Are Not Meaningful: For each situation, decide why the mean or median may not be meaningful.

- (a) Zip codes of several famous locations.
- (b) Ranks of selected medical schools.
- (c) Jersey numbers of football players.
- (d) Top 5 CEO compensation values.
- (e) The 50 state mean ages, one from each state.

8. Mean from a Frequency Distribution

When data are summarized in a frequency distribution, we can approximate the mean by using class midpoints. Because the original observations are unavailable, we temporarily treat every value in a class as if it were equal to that class midpoint. This simplifying assumption is why the resulting mean is an approximation rather than an exact value.

Mean from a Frequency Distribution

$$\bar{x} = \frac{\sum(f \cdot x)}{\sum f},$$

where f is the frequency and x is the class midpoint.

Example 7: Mean from a Frequency Distribution: Use the frequency distribution below to approximate the mean.

Time (seconds)	f	x	$f \cdot x$
0 – 14	6	7.0	42.0
15 – 29	18	22.0	396.0
30 – 44	14	37.0	518.0
45 – 59	5	52.0	260.0
60 – 74	5	67.0	335.0
75 – 89	1	82.0	82.0
90 – 104	1	97.0	97.0
Totals	50		1730.0

Connection to a Weighted Mean

The grouped-data formula is a weighted mean: each class midpoint x is weighted by its frequency f .

9. Weighted Mean

A **weighted mean** is used when different data values have different weights.

Weighted Mean

$$\bar{x} = \frac{\sum(w \cdot x)}{\sum w},$$

where w represents the weight assigned to each data value.

Example 8: Computing Grade-Point Average: A student takes five courses with grades and credits as follows:

Grade	Quality Points x	Credits w
A	4	3
A	4	4
B	3	3
C	2	3
F	0	1

Compute the student's GPA.

Summary

Key Ideas

- The mean uses every data value, but it is not resistant.
- The median is the middle value and is resistant.
- Use the mean for roughly symmetric data without influential outliers; prefer the median for skewed data or data with extreme values.
- The mode is the most frequent value and can be used with qualitative data.
- The midrange is easy to compute, but it is not resistant and is rarely used in practice.
- Always ask whether a measure of center is meaningful and whether the sample is representative.
- For grouped data, the mean is approximate because each class is represented by its midpoint.
- A weighted mean is used when some values count more than others.