

Problem 1

Calculate the Lagrangian $\mathcal{L}(x, \alpha)$.

$$\mathcal{L}(x, \alpha) = -(x_1 + x_2) + \alpha_1(x_1^2 + x_2^2 - 1)$$

Obtain the Lagrangian dual function $g(\alpha)$.

$$\begin{aligned} \frac{\nabla \mathcal{L}(x, \alpha)}{\partial x_1} &= 0 \Leftrightarrow \\ 2\alpha_1 x_1 - 1 &= 0 \Leftrightarrow \\ x_1 &= \frac{1}{2\alpha_1} \end{aligned}$$

$$\begin{aligned} \frac{\nabla \mathcal{L}(x, \alpha)}{\partial x_2} &= 0 \Leftrightarrow \\ 2\alpha_1 x_2 - 1 &= 0 \Leftrightarrow \\ x_2 &= \frac{1}{2\alpha_1} \end{aligned}$$

Solve the dual problem (Plug in $g(\alpha)$ the derived x).

$$\begin{aligned} \frac{\nabla g(\alpha)}{\partial \alpha} &= 0 \Leftrightarrow \\ \frac{1}{\alpha_1^2} - \frac{1}{2\alpha_1^2} - 1 &= 0 \Leftrightarrow \\ \alpha_1^2 &= \frac{1}{2} && \text{(by constraint } \alpha_i \geq 0) \\ \alpha_1 &= \frac{1}{\sqrt{2}} \end{aligned}$$

Problem 2**Problem 3**

By the formulation of the SVM problem we have

$$\begin{aligned} \text{minimize} \quad & f_0(w, b) = \frac{1}{2} w^\top w \\ \text{subject to} \quad & f_i(w, b) = y_i(w^\top x_i + b) - 1 \geq 0, \quad \text{for } i = 1, \dots, N \end{aligned} \quad (1)$$

Clearly we can rewrite (1) to

$$f_i(w, b) = -y_i(w^\top x_i + b) + 1 \leq 0, \quad \text{for } i = 1, \dots, N$$

By Slater's constraint qualification we have that the duality gap of the SVM problem is zero if $f_0(x)$, $f_1(x)$, ... $f_N(x)$ are convex and the constraints $f_1(x)$, ... $f_N(x)$ are affine. Clearly both assumptions are met, because the first is simply the L_2 -norm, which is convex and the constraints are linear functions in w shifted by an offset b , which makes them affine. Thus the duality gap is zero.

Problem 4

a). Let $\mathbf{X} \in \mathbb{R}^{n \times d}$, where each row is a datapoint $\mathbf{x} \in \mathbb{R}^d$, $\mathbf{Y} \in \mathbb{R}^{n \times n}$, where each column is the label y_i of the i -th datapoint replicated n times and $\boldsymbol{\alpha} \in \mathbb{R}^n$, where each column is the i -th coefficient α_i .

By the Hadamard product we have

$$\mathbf{Q} = \mathbf{X}\mathbf{X}^\top \odot (\mathbf{Y} \odot \mathbf{Y}^\top)$$

$$g(\boldsymbol{\alpha}) = \boldsymbol{\alpha}^\top \mathbf{Q} \boldsymbol{\alpha}$$

b). We recognize $\mathbf{X}\mathbf{X}^\top$ as the empirical covariance matrix, which is by construction positive semi-definite.