DPG Spring Meeting 2026

Electronic coarse-graining for accurate and scalable modeling of organic semiconductors

•Andriy Zhugayevych[1], Denis Andrienko[1], Sergei Tretiak[2]

[1] *Max Planck Institute for Polymer Research*
[2] *Los Alamos National Laboratory, USA*

March 9, 2026

Acknowledgments

Naomi Kinaret, Kun-Han Lin (noncrystalline morphologies)
Andriy Kadashchuk (U Bayreuth) (experiment)
MPIP Polymer Theory Department (polymer theory)
MPIP Molecular Electronics Department (experiment)

Funding

MPIP/MPG fellowship

MAX PLANCK INSTITUTE FOR POLYMER RESEARCH

Outline

1) Motivation, 2) Methodology, 3) Applications

Motivation: Accurate modeling of organic semiconductors

$\Longrightarrow$ Coarse-graining of molecular and electronic degrees of freedom

Electronic coarse-graining — reducing size of electronic basis w/o loss of accuracy

class	basis	examples
✓ Broadband semiconductors ("free" e/h)	valence orbitals	 Si, graphene  , black P, GaAs
✓ Small-molecule organic semiconductors (intermolecular hopping)	HOMO /LUMO	 naphthalene, rubrene, TCNQ, C8-BTBT
?? Other organic semiconductors (intra- and intermolecular dynamics)	fragment HOMO /LUMO	 polymers, large molecules, MOF, OLED, OPV

Goal is to model complex organic semiconductors at level of small-molecule ones: 1) black-box, 2) scalable, 3) compatible technology, 4) controllable accuracy

Methodology of electronic coarse-graining: Main ideas

- We need only frontier molecular orbitals
- Large enough clusters approximate infinite solid
- See technical details at https://cmsos.github.io/escp



TBH = Tight Binding Hamiltonian

... will be illustrated with two examples: polythiophene and graphene

Electronic coarse-graining: example of polythiophene

(valence band of single planar polythiophene chain, min basis = 2 LMOs per monomer)

bands in all-electron basis





bands in coarse-grained basis





all-electron density



CG density (2 MOs)

The proposed approach solves electronic structure problem for any π-conjugated system which can be fragmentized into π-closed-shell monomers

Electronic coarse-graining: example of graphene





The same approach can be applied to other semiconductors

e.g. to calculate electronic structure of infinite systems with methods available only for finite systems

Applications

(where conventional approaches produce substantial systematic errors)

- Electronic structure of crystalline polymers



- Density of electronic states in organic semiconductors



- Guest-to-guest tunneling in host-guest systems



- Electronic transport through A-D-A molecules for solar cells

  J Chem Phys 159, 024107 (2023)

- Charge storage mechanism in metal-organic polymers

  Chem Sci 13, 8161 (2022)

Electronic structure of crystalline polymers

Challenges

- Combination of strong intrachain and weak interchain couplings
- Unit cell and supercell are too large for brute force methods
- No experimental crystal structure JPCC 122, 9141 (2018); JCTC 19, 8481 (2023)

$\Longrightarrow$ Little is known about electronic structure of crystalline polymers





Electronic coarse-graining allows for calculation of electronic structure of polymers at the level of accurate density functionals

Why PEDOT is one of best polymers for charge transport?

(Here we compare only thiophene-based polymers: PolyThiophene, PEDOT, P3HT)



- How large intermolecular electronic couplings can be?
  250 meV in perfectly aligned π-stacks (max-symmetry π-stack of PPP)
- What prevents perfect alignment?
  Flat potential energy surface wrt to tilts and shifts
  (0 to max coupling upon shift by 1 bondlength due to BLA pattern)
- What is the "best" thiophene-based polymer?
  2D PEDOT – (1) well-aligned to maximize couplings and (2) has small thermal fluctuations of intermolecular couplings

---

- What other guidelines we can give for thiophenes?
  Localize HOMO on sulfur to increase couplings
  (TT,TBT fusing is good, TTT,TTTT is not)

How to correctly compare simulated electronic DOS?

(Critically important for heterogeneous systems such as host-guest systems)



Right panel: in red – HOMO-DOS (biased), in gray – full DOS (correct)

Compare complete electronic DOS for electronically different systems

(e.g. UGH vs other OLED host and guest molecules)

Is guest-to-guest tunneling efficient in host-guest systems?

How to accurately calculate tunneling couplings ("superexchange")?

Challenges leading to loss of accuracy

- Effects of environment (correct SCF)
- Multiple pathways
- Electronic localization



Contact distance is 12 Å,
coupling is 0.13 meV,
3 bridging molecules

Couplings are too small, see talk CPP57.2 by Naomi Kinaret

Summary

Electronic Coarse-Graining is a powerful tool (sometimes the only one) for calculation and analysis of electronic structure of complex systems:

- Anything that can be monomerized (polymers, COFs, MOFs)
- Solids and clusters of large molecules
- Defects in solids
- Infinite systems from finite-size calculations

Project web-page

- Electronic Coarse-Graining (including copy of this presentation)
  https://zhugayevych.me/research/ECG
- Computational Materials Science of Organic Semiconductors
  https://cmsos.github.io