

# PC3233 AM1 Assignment 1

Hor Zhu Ming

January 2024

## 1 Angular Momenta

### 1.1 Prove $[J^2, J_i] = 0$

*Proof.* Since  $J^2 = J_x^2 + J_y^2 + J_z^2 = \sum_{j=x,y,z} J_j^2$ .

$$\begin{aligned}[J^2, J_i] &= \left[ \sum_j J_j^2, J_i \right] \\ &= \sum_j [J_j^2, J_i] \\ &= \sum_j [J_j, J_i] J_j + J_j [J_j, J_i]\end{aligned}$$

Since  $[J_j, J_i] = i\hbar\epsilon_{ijk}J_k$ , where  $\hbar$  is set to be 1. Then,

$$[J_j, J_i] = \begin{cases} 0 & \text{if } i = j \\ i\epsilon_{jik}J_k & \text{if } i \neq j \end{cases}$$

Therefore,

$$\begin{aligned}\sum_j [J_j, J_i] J_j + J_j [J_j, J_i] &= 0 + i\epsilon_{yxz}J_zJ_y + i\epsilon_{yxz}J_yJ_z + i\epsilon_{zxy}J_yJ_z + i\epsilon_{zxy}J_zJ_y \\ &= i(-1)J_zJ_y + i(-1)J_yJ_z + i(1)J_yJ_z + i(1)J_zJ_y \\ &= 0\end{aligned}$$

Hence,  $[J^2, J_i] = 0$ . □

### 1.2 Prove $[J^2, J_{\pm}] = 0$

*Proof.* Since  $J_{\pm} = J_x \pm iJ_y$ ,

$$\begin{aligned}[J^2, J_{\pm}] &= [J^2, J_x \pm iJ_y] \\ &= [J^2, J_x] \pm i[J^2, J_y] \\ &= 0 \pm 0 \quad \because \text{From the Solution 1.1} \\ &= 0\end{aligned}$$

Hence,  $[J^2, J_{\pm}] = 0$ . □

### 1.3 Prove $[J_z, J_{\pm}] = \pm J_{\pm}$

*Proof.*

$$\begin{aligned}
 [J_z, J_{\pm}] &= [J_z, J_x \pm iJ_y] \\
 &= [J_z, J_x] \pm i[J_z, J_y] \\
 &= iJ_y \pm i(-iJ_x) \quad \because [J_x, J_y] = i\epsilon_{xyz}J_z \\
 &= iJ_y \pm J_x \\
 &= \pm(\pm iJ_y + J_x) \\
 &= \pm(J_x \pm iJ_y) \\
 &= \pm J_{\pm}
 \end{aligned}$$

Hence,  $[J_z, J_{\pm}] = \pm J_{\pm}$ . □

### 1.4 Prove $J^2 = J_{\pm}J_{\mp} + J_z^2 \mp J_z$

Goal: Simplify the RHS of the equation to get  $J^2$ .

*Proof.*

$$\begin{aligned}
 J_{\pm}J_{\mp} + J_z^2 \mp J_z &= (J_x \pm iJ_y)(J_x \mp iJ_y) + J_z^2 \mp J_z \\
 &= J_x^2 \mp iJ_xJ_y \pm iJ_yJ_x + J_y^2 + J_z^2 \mp J_z \\
 &= J_x^2 + J_y^2 + J_z^2 \mp iJ_xJ_y \pm iJ_yJ_x \mp J_z \\
 &= J^2 \mp iJ_xJ_y \pm iJ_yJ_x \mp J_z \\
 &= \begin{cases} J^2 - iJ_xJ_y + iJ_yJ_x - J_z \\ J^2 + iJ_xJ_y - iJ_yJ_x + J_z \end{cases} \\
 &= \begin{cases} J^2 + i(J_yJ_x - J_xJ_y) - J_z \\ J^2 + i(J_xJ_y - J_yJ_x) + J_z \end{cases} \\
 &= \begin{cases} J^2 + i[J_y, J_x] - J_z \\ J^2 + i[J_x, J_y] + J_z \end{cases} \\
 &= \begin{cases} J^2 + i(-iJ_z) - J_z \\ J^2 + i(iJ_z) + J_z \end{cases} \\
 &= \begin{cases} J^2 + J_z - J_z \\ J^2 - J_z + J_z \end{cases} \\
 &= \begin{cases} J^2 \\ J^2 \end{cases} \\
 &= J^2
 \end{aligned}$$

Hence,  $J^2 = J_{\pm}J_{\mp} + J_z^2 \mp J_z$ . □

```
[1]: # import library
import sympy as sym
from IPython.display import display, Latex
```

## 2 Legendre polynomials

### 2.1

Write a program in your favorite math software to obtain the **first ten** Legendre Polynomials  $P_l(x)$  using Rodrigues' formula

$$P_l(x) = \frac{1}{2^l l!} \frac{d^l}{dx^l} [(x^2 - 1)^l]$$

```
[2]: x, l = sym.symbols("x l")
factorial = sym.factorial
diff = sym.diff

def legendre_polynomials(l, x):
    coefficient = 1 / (2 ** l * factorial(l))
    polynomial_expression = diff((x ** 2 - 1) ** l, x, l)
    res = sym.factor(coefficient * polynomial_expression)
    display(Latex(f"$P_{l}(x) = {sym.latex(res)}$"))
    return res

for i in range(10):
    legendre_polynomials(i, x)
```

$$\begin{aligned}
P_0(x) &= 1 \\
P_1(x) &= x \\
P_2(x) &= \frac{3x^2-1}{2} \\
P_3(x) &= \frac{x(5x^2-3)}{2} \\
P_4(x) &= \frac{35x^4-30x^2+3}{8} \\
P_5(x) &= \frac{x(63x^4-70x^2+15)}{8} \\
P_6(x) &= \frac{231x^6-315x^4+105x^2-5}{16} \\
P_7(x) &= \frac{x(429x^6-693x^4+315x^2-35)}{16} \\
P_8(x) &= \frac{6435x^8-12012x^6+6930x^4-1260x^2+35}{128} \\
P_9(x) &= \frac{x(12155x^8-25740x^6+18018x^4-4620x^2+315)}{128}
\end{aligned}$$

### 2.2

The Legendre polynomials satisfy the recursion relation:

$$lP_l(x) = (2l-1)xP_{l-1}(x) - (l-1)P_{l-2}(x)$$

Write a program in your favorite math software to determine  $P_2(x)$  through  $P_{10}(x)$  (starting with  $P_0(x) = 1$  and  $P_1(x) = x$ ) using the above recursion relation.

```
[3]: term = 10

P0 = 1
display(Latex(f"$P_{0}(x) = {P0}$"))
P1 = x
display(Latex(f"$P_{1}(x) = {P1}$"))
for l in range(2, term):
    tem = (2 * l - 1) * x * P1 - (l - 1) * P0
    result = sym.factor(tem / l)
    display(Latex(f"$P_{l}(x) = {sym.latex(result)}$"))
    P0 = P1
    P1 = result
```

$$P_0(x) = 1$$

$$P_1(x) = x$$

$$P_2(x) = \frac{3x^2-1}{2}$$

$$P_3(x) = \frac{x(5x^2-3)}{2}$$

$$P_4(x) = \frac{35x^4-30x^2+3}{8}$$

$$P_5(x) = \frac{x(63x^4-70x^2+15)}{8}$$

$$P_6(x) = \frac{231x^6-315x^4+105x^2-5}{16}$$

$$P_7(x) = \frac{x(429x^6-693x^4+315x^2-35)}{16}$$

$$P_8(x) = \frac{6435x^8-12012x^6+6930x^4-1260x^2+35}{128}$$

$$P_9(x) = \frac{x(12155x^8-25740x^6+18018x^4-4620x^2+315)}{128}$$