

Testing coarse line-opacity treatments with energy-conserving fluorescence in kilonova ejecta

Yonglin Zhu^{1*}

¹*Independent researcher*

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

Kilonova radiative-transfer calculations coarse-grain the r-process line forest either with the expansion-opacity formalism or with a line-binned (integrated Sobolev depth) opacity, and both constructions were validated against resolved line-by-line transport under complete thermal redistribution, where every absorbed packet is re-emitted from the local thermal emissivity. We ask whether that validation survives when the post-absorption physics is instead energy-conserving fluorescence, in which the absorbing transition determines the emitted frequency. Using one Sobolev Monte Carlo code, one line list and one set of ejecta states, we compare the resolved transport with the two coarse-grained treatments under both redistributions, on a lanthanide-rich secular-ejecta state, an AT2017gfo-like thin state, and the single-element benchmark problem of Fontes et al., both as a snapshot and as a light curve under a prescribed thermodynamic trajectory. Under complete redistribution we recover the published picture: the line-binned closure reproduces the resolved colours to within 0.24 mag on the lanthanide-rich states and the expansion closure sits 0.02–0.27 mag too bright. Switching only the redistribution to fluorescence leaves the expansion closure’s error where it was (0.15–0.19 mag in z) but moves the line-binned closure to 0.16–0.80 mag too faint, opens its $r - K$ colour error on the Fontes problem to +1.34 mag, and on a thinner line list makes its transport fail to terminate. Over the light curve the bolometric ordering of the three treatments is the same under both redistributions (expansion 15% to 17% brighter at the peak, line-binned within 3.8%), while fluorescence raises the line-binned closure’s maximum colour error from 0.13 to 0.45 mag and the expansion closure’s from 0.20 to 0.32 mag. A coarse opacity that retains only the integrated depth of a bin cannot retain which transition absorbed; under fluorescence that lost information is a systematic error in the colours.

Key words: radiative transfer – opacity – methods: numerical – neutron star mergers – transients: neutron star mergers

1 INTRODUCTION

The optical and near-infrared emission of a kilonova is shaped by tens of millions of bound–bound transitions of the r-process elements, above all the lanthanides, in ejecta that expand homologously at a tenth of the speed of light (Kasen et al. 2013; Barnes & Kasen 2013; Tanaka et al. 2020; Metzger 2020). No transport calculation of a kilonova treats every one of those lines with its own Sobolev escape probability throughout a time-dependent, multi-dimensional model. Instead the forest is coarse-grained. Two constructions dominate the literature. The expansion-opacity formalism of Karp et al. (1977) and Eastman & Pinto (1993) replaces the lines in a frequency bin by an effective continuous opacity proportional to the sum of $1 - e^{-\tau_S}$ over the bin, and is the basis of the opacity tables used by SEDONA (Kasen et al. 2006, 2017), by the Tanaka–Kawaguchi line of models (Tanaka et al. 2020; Kawaguchi et al. 2018) and by POSSIS (Bulla 2019). The

line-binned or “area-preserving” opacity used by Fontes et al. (2020) and in the SUPERNu models of Wollaeger et al. (2018) instead integrates the Sobolev depth itself over the bin, so that the bin’s total optical depth is preserved. Both replace a set of discrete transitions by a single number per bin.

Fontes et al. (2020) tested the two constructions against a resolved line-by-line calculation on a deliberately simple problem: a pure-neodymium sphere with a prescribed density and temperature structure, transported with the lines treated one by one, with the expansion opacity, and with the line-binned opacity. They found the three to agree to within several per cent in the peak luminosity, with the expansion treatment the brightest, and concluded that either coarse-graining is adequate for kilonova light-curve work. That benchmark, like the calculations it validated, was done under complete thermal redistribution: a packet absorbed in a line is re-emitted from the local thermal emissivity, so that the identity of the absorbing transition plays no role once the absorption has happened.

Line-by-line codes do something different. In ARTIS (Shin-

* E-mail: zhuygln@gmail.com

gles et al. 2023; Collins et al. 2023) and in the Lucy macroatom that underlies it (Lucy 2002, 2003), an absorption activates a specific upper level and the packet leaves through one of that level's downward transitions: the emitted frequency depends on which line absorbed, and energy is conserved through the cascade. Shingles et al. (2023) argued on physical grounds that the fluorescence produced this way matters for the near-infrared spectra of kilonovae, and Collins et al. (2026) and Morag (2026) have since made the case that the treatment of line opacity is among the largest uncontrolled choices in kilonova modelling. What has not been done is the controlled experiment: the same code, the same line list, the same ejecta state, the three opacity treatments, and the single change from complete redistribution to energy-conserving fluorescence.

That experiment is the subject of this paper. The question is a methods question. A coarse-grained opacity records, for each frequency bin, how much is absorbed; it does not record by which transition. Under complete redistribution that information is not needed, because every absorption ends the same way. Under fluorescence it is needed, because the cascade that follows depends on it. The expansion opacity and the line-binned opacity lose that information differently, and the question is how much of the validated agreement of Fontes et al. (2020) survives when the post-absorption physics is the one that line-by-line codes actually use.

We answer it with a Sobolev Monte Carlo transport in which the three opacity treatments and the two redistributions are switches on one code path, so that any difference between legs is the difference in the physics and nothing else (Section 2). The ejecta states are two published structures and the Fontes problem itself (Section 3). Section 4 gives the closures on the published states, the effect of the redistribution, the Fontes problem as a snapshot on two independent line lists, the same problem as a light curve under a prescribed thermodynamic trajectory, and a robustness test with a stronger fluorescence treatment. Section 5 interprets the pattern and states what the calculation does not show.

2 METHOD

2.1 Sobolev Monte Carlo transport with indivisible energy packets

The transport is a Monte Carlo solution of the line-transfer problem in the Sobolev approximation for homologous flow (Sobolev 1960; Castor 1970; Jeffery 1993), written for this project and available with the paper. Radiation is carried by indivisible packets of equal comoving-frame energy (Lucy 2002, 2005). A packet is launched at a position and frequency drawn from the source (the inner boundary with a Planck spectrum, or the volume with the local emissivity), travels in the lab frame, and redshifts in the comoving frame through the line list in order of decreasing frequency. At each line it meets, it interacts with probability $1 - e^{-\tau_s}$, where τ_s is the Sobolev depth of that line in the local shell. The comoving energy of a packet is conserved through every interaction; the lab-frame energy it carries out is smaller by the work done on the expanding flow, which the code books separately. The transport is exact in the Sobolev limit for the states considered here, and the packet accounting closes to machine

precision on every run: the injected energy equals the escaped energy plus the energy returned to the inner boundary, plus the expansion work, plus what remains in flight when the run stops.

Multi-shell states are transported with a zoned atom: one line list, with per-shell Sobolev depths and escape probabilities on the union of the shells' opacity sets, and per-shell macroatom tables. A single shell reproduces the single-zone code bit for bit, and a state split into more shells at the same physical structure reproduces the unsplit state within the Monte Carlo noise. The second property was the diagnostic that found an inversion error in the coarse-grained legs of an earlier version of this code, since corrected; all results below are on the corrected transport and the pre-correction histories are retained in the repository.

2.2 Three opacity treatments on one line list

The resolved treatment (R) walks the lines one at a time. The expansion treatment (B) groups the lines into bins of fixed width in $\ln \nu$ and assigns each bin the expansion opacity of Eastman & Pinto (1993), $\kappa_{\text{exp}} \propto \sum_i (1 - e^{-\tau_i})$, evaluated as a continuous opacity along the packet's path through the bin. The line-binned treatment (B_{bin}) assigns each bin the integrated Sobolev depth $\sum_i \tau_i$, so that the bin's total attenuation is preserved. When a packet is absorbed in a bin of either coarse treatment and the redistribution requires the identity of the absorbing line, the line is drawn from the bin's members in proportion to the weight the treatment gave them: $1 - e^{-\tau_i}$ for B, τ_i for B_{bin}. Both coarse treatments use the same bins, the same line list, and the same per-line depths as the resolved treatment, and reduce to it in the limit of one line per bin; a test on an uneven synthetic forest checks that both reproduce the analytic attenuation of the resolved forest. The closure error of a coarse treatment is its difference from R in the same state under the same redistribution; the sensitivity of that error to the bin width, the depth cut and the packet count was checked over 12 settings and is reported with the results.

2.3 Two redistributions

Under complete thermal redistribution (the subscript θ) every absorbed packet is re-emitted at a frequency drawn from the local line emissivity in local thermodynamic equilibrium, independent of the line that absorbed it. This is the physics of the Fontes et al. (2020) benchmark and of the opacity-table codes.

Under fluorescence (the subscript 2) an absorption activates the upper level of the absorbing line, and the packet leaves by a downward transition drawn from that level's branching, each transition weighted by its net radiative rate $A\beta$ with β the Sobolev escape probability; if the transition lands on an excited level, the cascade continues from there. This is the downward part of the Lucy (2002) macroatom, with energy conserved through the cascade. Its bookkeeping matters: the leg R₁, in which the packet is re-emitted in the absorbing line, differs from R₂ by -2.42 mag in z on the lanthanide-rich state, and the coarse closures inherit the same shift (B₁ and R₁ differ by at most 0.04 mag), so nothing below depends on it. A control leg A₂, which keeps the re-

solved opacity but groups the emission by level, differs from R_2 by at most 0.12 mag.

As a robustness test we also transport with a full macroatom (the superscript M) in which internal upward transitions are allowed under an imposed diluted Planck radiation field, $\bar{J} = WB_\nu(T)$ with $W = \frac{1}{2}$, with the rates of Lucy (2003) and no collisions. This is not an equilibrium solution, since the field is imposed rather than solved for; it is the strongest fluorescence treatment that a fixed-state calculation can carry, and it is used to ask whether the conclusions depend on the downward table.

2.4 Time-slab light curves

For the light curve the transport runs in time slabs. Each packet carries its own clock; at the end of a slab the packets still in flight are paused at their exact positions and resumed in the next slab, in which the state has been rebuilt with the density scaled as t^{-3} and the temperature of the prescribed trajectory. Heating is injected per slab as equal-energy packets in proportion to each shell's mass, at the shell's Planck spectrum, uniformly in time over the slab; the radiation stored in the ejecta at the start is injected as aT^4V per shell. Escape times are recorded, and the light curve is the escaping luminosity in escape time (observer time is also recorded). The per-slab accounting is exact, and a run split into slabs reproduces the single run. Removing every line from the state gives the free-streaming control: no interaction, zero expansion work, and the initial field released over the light-crossing time; on the production state this control closes to 10^{-15} . What the light curve measures that a snapshot cannot is the trapping time, the loss to expansion work, and the release of stored radiation; what it does not do is solve an energy equation for the gas.

2.5 Photometry

Magnitudes are synthesised from the escaping packets in the *grizJHK* bands. In the snapshots the packets are binned by frequency; in the light curves by escape time and frequency, with the photometric time bins chosen so that every reported band has at least a hundred packets. The Monte Carlo scatter on a band magnitude, from repeated seeds, is at most a few hundredths of a magnitude on every leg except the full-macroatom reference, where it is below a tenth.

3 EJECTA STATES AND ATOMIC DATA

P1 is the lanthanide-rich secular ejecta of the XKN framework (Ricigliano et al. 2024): $M = 2.64 \times 10^{-2} M_\odot$, $v_{\text{rms}} = 0.06c$ on the $(1 - x^2)^3$ density profile with $x = v/v_{\text{max}}$, $Y_e = 0.20$, and the composition of the $Y_e = 0.21$ trajectory of Gillanders et al. (2022) with its lanthanide mass fraction rescaled to $X_{\text{lan}} = 0.11$ (Tanaka et al. 2020). Its temperature run is the grey Eddington solution at the model's luminosity, and the closures are evaluated at the shell where the grey depth reaches $2/3$ and, in the multi-shell runs, on every shell above it. It is used at 1, 2, 3 and 5 d, with ionization from Saha equilibrium. Two variants test the composition pattern: the $Y_e = 0.21$ trajectory at its own $X_{\text{lan}} = 0.30$, and a solar r-process residual pattern at $X_{\text{lan}} = 0.11$.

P2 is a thin, AT2017gfo-like state after Gillanders et al. (2022): a v^{-3} density profile between 0.15c and 0.35c at 3.4 d, $T = 3200$ K, with the $Y_e = 0.29$ trajectory's lanthanides reduced to $X_{\text{lan}} = 2.5 \times 10^{-3}$.

The Fontes problem is Appendix C of Fontes et al. (2020): a pure-neodymium sphere of $1.4 \times 10^{-2} M_\odot$ with $v_{\text{max}} = 0.25c$, density $\propto (1 - x^2)^3$, temperature $\propto (1 - x^2)$ with 5700 K at the centre at 4 d, radioactive heating uniform in mass with their power law in time, bound-bound opacity only, and lines with oscillator strength above 10^{-3} . Ionization is Saha (Nd III inside, Nd II outside). As a snapshot at 4 d the outer cells, which hold the region of grey depth below unity, are transported explicitly with the interior's heating arriving through the inner boundary; because the interior is diffusive, the boundary is treated in two brackets, complete thermalisation of returning packets (re-emitted with the boundary's Planck spectrum) and a lossless mirror, and every conclusion is required to hold in both. As a light curve the same state is followed from 4 to 16 d in 40 logarithmic slabs, with the temperature of the initial profile scaled as t^{-1} (adiabatic, radiation-dominated) and, as a variant, with the temperature of each shell reset at each slab boundary from the packets present in it. We call this a time-dependent extension of the Fontes benchmark under a prescribed thermodynamic trajectory; it is not a reproduction of their light curve, which solved an energy equation.

Atomic data. The lanthanide states use the calibrated line list of Flörs et al. (2026); the Fontes problem uses its neodymium ions, 1087925 lines above the oscillator-strength cut. As an independent check the Fontes snapshot is repeated with Nd II and Nd III from the Japan–Lithuania opacity database (Kato et al. 2021, 2024), an uncalibrated set with 247597 lines above the same cut, a forest four times thinner.

4 RESULTS

4.1 The closures on published states

Table 1 and Fig. 1 give the closure errors on P1, its composition variants and P2 under both redistributions. Under fluorescence, the expansion closure is 0.15–0.19 mag too bright in z and 0.06–0.14 in K on P1 at every epoch from 1 to 5 d and on both composition variants; on P2 it is +0.11 mag in g and smaller in every redder band. The line-binned closure under fluorescence is 0.16–0.80 mag too faint on P1, larger than the expansion closure's error and of the opposite sign in every band.

These numbers do not depend on how the ejecta is divided. Table 2 lists the expansion and line-binned closure errors on P1 at 2 d transported as a single photospheric shell and as 4, 6, 12 and 23 resolved shells above the photosphere: the expansion closure's z error is -0.19 on the single shell and -0.17, -0.16 and -0.15 mag on the 4-, 12- and 23-shell grids, and a 4-shell grid with one shell split in three is bit-identical to the unsplit grid. Across the 12 convergence settings (bin width over two decades, depth cut over two decades, packet count, emissivity cut) the same error spans 0.13–0.23 mag in z . The closure error is a property of the coarse-graining, at the level of a few hundredths of a magnitude.

F62: the two closures under complete redistribution and under explicit fluorescence (fixed transport)




Figure 1. The closure error, coarse treatment minus resolved transport, per band, on P1 at 1–5 d, its two composition variants and P2. Left column: the expansion treatment; right column: the line-binned treatment. Top row: complete thermal redistribution; bottom row: energy-conserving fluorescence. The two closures are within a few tenths of a magnitude of the resolved transport under complete redistribution; under fluorescence the expansion closure stays where it was and the line-binned closure moves to several tenths too faint at every epoch and pattern.

Table 1. Closure minus resolved transport in g , z and K (mag) on the published states, under complete thermal redistribution ($\epsilon = 1$) and under fluorescence.

state	expansion, $\epsilon = 1$	line-binned, $\epsilon = 1$	expansion, fluor.	line-binned, fluor.
	closure – resolved, $g / z / K$ (mag)			
P1 1 d	-0.02 / -0.27 / -0.16	+0.24 / -0.10 / -0.03	+0.03 / -0.15 / -0.14	+0.80 / +0.32 / +0.22
P1 2 d	-0.02 / -0.12 / -0.07	+0.09 / +0.01 / +0.02	-0.09 / -0.19 / -0.10	+0.70 / +0.37 / +0.24
P1 3 d	-0.08 / -0.12 / -0.09	+0.08 / -0.00 / -0.00	-0.16 / -0.15 / -0.09	+0.78 / +0.45 / +0.30
P1 5 d	-0.16 / -0.11 / -0.07	-0.03 / -0.01 / +0.01	-0.15 / -0.19 / -0.06	+0.73 / +0.38 / +0.16
P1r1	-0.09 / -0.20 / -0.15	+0.21 / +0.01 / +0.00	-0.14 / -0.25 / -0.14	+0.77 / +0.38 / +0.55
P1r2	-0.09 / -0.14 / -0.08	+0.15 / -0.00 / +0.01	-0.13 / -0.19 / -0.12	+0.75 / +0.29 / +0.30
P2 3.4 d	-0.11 / +0.03 / +0.01	-0.09 / +0.03 / +0.00	+0.11 / +0.02 / -0.02	+0.15 / +0.02 / -0.02

Table 2. Grid invariance: closure minus resolved transport on P1 at 2 d under fluorescence, on single shells around the photosphere and on multi-shell grids above it.

P1, 2 d	$B_2 - R_2$ (g)	(z)	(K)	$B_{\text{bin},2} - R_2$ (g)	(z)	(K)
single zone, shell 27	-0.07	-0.19	-0.15	+0.86	+0.38	+0.37
single zone, photospheric shell	-0.09	-0.19	-0.10	+0.70	+0.37	+0.24
single zone, shell 29	-0.13	-0.11	-0.02	+0.52	+0.23	-0.11
4 transported shells	-0.21	-0.17	-0.03	+0.68	+0.29	-0.01
the same state as 12 shells	-0.21	-0.17	-0.03	+0.68	+0.29	-0.01
6 shells	-0.14	-0.18	-0.03	+0.66	+0.28	+0.05
12 shells	-0.14	-0.16	-0.03	+0.48	+0.29	+0.02
23 shells	-0.28	-0.15	-0.02	+0.49	+0.28	-0.03

4.2 Complete redistribution against fluorescence

The upper row of Fig. 1 is the [Fontes et al. \(2020\)](#) experiment on the lanthanide-rich states: under complete thermal redistribution the line-binned closure reproduces the resolved transport to within 0.24 mag in every band on P1, and the expansion closure sits 0.02–0.27 mag too bright. This is their ordering, re-found with independent atomic data and a resolved Sobolev reference.

The lower row is the same calculation with the single change to energy-conserving fluorescence. The expansion closure is unchanged within the noise. The line-binned closure moves from within 0.24 mag to 0.16–0.80 mag too faint, at every epoch and on both composition variants; on the thin state P2 neither closure moves beyond a few tenths in g and both are within the noise redward. The redistribution itself moves the resolved transport by one to two magnitudes in the optical on P1: the choice of post-absorption physics is a larger effect than the choice of opacity closure, and the closures’ validation is specific to the physics it was done under.

4.3 The Fontes problem as a snapshot, on two line lists

Fig. 2 shows the same experiment on a single element. Under complete redistribution the line-binned closure reproduces the resolved $r - K$ colour to -0.01 mag and the expansion closure is -0.29 mag too blue, brighter than the resolved leg by half a magnitude or more through r to K : the [Fontes et al. \(2020\)](#) ordering, expansion brighter than line-binned with the resolved transport between them in colour. Under fluorescence the expansion closure’s colour error is -0.36 mag, unchanged within the bracket of the boundary treatment, while the line-binned closure’s $r - K$ error opens to +1.34 mag with the mirror boundary and +1.72 mag with re-emission: too faint by more than a magnitude in r and by several in g , and slightly too bright in K . Both brackets of the inner boundary give the pattern, so it is a property of the transported layers and not of the boundary.

On the Japan–Lithuania line list the five legs that terminate repeat the pattern: line-binned within +0.03 mag of the resolved colour under complete redistribution and the expansion closure several tenths too blue under both. The line-binned closure under fluorescence did not terminate on this list, with either boundary treatment: packets remained in flight after 10^6 transport steps. The mechanism is visible in the event counts. On the calibrated list the line-binned fluorescence leg needed three to four times as many interactions per packet as the other legs, and under complete redistribution the same bins cost thousands of interactions per packet and terminated. A packet re-emitted by the macroatom at the frequency of a saturated line lands in a bin whose integrated depth re-absorbs it with certainty; the walk re-enters the same upper level and leaves by the same line, and nothing in the closure plays the role of the Sobolev escape probability of the resolved transport or of the $1 - e^{-\tau}$ saturation of the expansion opacity. On a strong-lined forest the line-binned closure with explicit fluorescence is not merely less accurate but ill-posed without a per-line escape treatment.

4.4 The Fontes problem as a light curve

Table 3 and Fig. 3 give the six light curves. Of the energy available (the stored field at 4 d is 7.2 times the heating deposited over 4 to 16 d), 61% is radiated by 16 d and 36% is lost to expansion work, on every leg within a few per cent. The resolved transport peaks at 4.75 d under complete redistribution and 4.96 d under fluorescence, and is the lowest of the three treatments under both: it loses the most energy to work, because a packet that leaves a homologous flow after the most interactions has given the most of its lab-frame energy to the flow. The expansion closure is 15% brighter at the peak under complete redistribution and 17% under fluorescence; the line-binned closure is within 1.6% and 3.8%. The ordering, expansion above line-binned above resolved, is the same under both redistributions. It is not the ordering [Fontes et al. \(2020\)](#) reported, in which the resolved calculation was the brightest, and the expansion excess is larger than theirs; a prescribed thermodynamic trajectory does not reproduce an energy-equation light curve, and the comparison we make is between the closures, not with their curve.

Fluorescence changes the colours, not the bolometric ordering. The maximum colour residual of the line-binned closure over the light curve grows from 0.13 mag under complete redistribution to 0.45 mag under fluorescence, with the closure up to 0.82 mag too faint in H ; the expansion closure’s grows from 0.20 to 0.32 mag. The snapshot’s colour error of more than a magnitude becomes a few tenths once the light curve integrates over the distribution of escape times, and it remains a systematic difference of the opposite sign from the expansion closure’s. The line-binned fluorescence leg reached its interaction cap on 1.4% of its packets in the first slabs, the bounded form of the trapping pathology of Section 4.3; a rerun with a fifteen-fold higher cap is reported with the record. **[TODO: replace with the high-cap result when the rerun lands]**

The variant in which each shell’s temperature is reset at each slab boundary from the radiation present in it moves the resolved peak to 6.6 d, closer to the peak [Fontes et al. \(2020\)](#) reported, and gives the same closure picture: the expansion closure 8% brighter at the peak with a maximum colour residual of 0.22 mag.

4.5 A stronger fluorescence treatment

Under the full macroatom the resolved transport itself changes by -1.80 mag in g on P1 at 2 d, brighter through H with K unchanged, at 3.6 times the interactions per packet: the imposed field pumps energy blueward through the upward transitions. The expansion closure’s error stays where the downward macroatom put it, -0.18 mag in z and within 0.18 mag in every band (Fig. 4). The line-binned closure keeps its +0.67 mag error in g , loses its red excess and turns to -0.25 mag in K . The pattern of Section 4.2 is not a property of the downward table: the expansion closure tracks the resolved transport to a few tenths whichever fluorescence treatment both are given, and the line-binned closure’s error is larger and changes shape with the treatment.




Figure 2. Closure minus resolved transport per band on the Fontes problem at 4 d, for both treatments of the diffusive inner boundary (re-emission and mirror) and both line lists (calibrated and Japan–Lithuania), under complete redistribution (left) and fluorescence (right). The line-binned closure under fluorescence on the Japan–Lithuania list did not terminate and is absent.




Figure 3. The Fontes problem as a light curve under a prescribed thermodynamic trajectory, 4 to 16 d. Left: the bolometric escaping luminosity of the three treatments under complete redistribution (solid) and fluorescence (dashed). Centre: closure minus resolved transport in z (circles) and H (triangles) per photometric time bin. Right: the $J - K$ colour residual. The g and r bands are starved of packets at these epochs and are not shown.

Table 3. The Fontes light curve: peak time and luminosity, radiated energy, expansion work, the peak-luminosity excess over the resolved transport and the maximum colour residual, for the three treatments under both redistributions.

opacity	redistribution	t_{peak} (d)	L_{peak} (10^{40} erg s $^{-1}$)	E_{rad} (10^{46} erg)	W (10^{46} erg)	ΔL_{peak}	max $ \Delta \text{colour} $ (mag)
resolved	$\epsilon = 1$	4.75	7.23	4.477	2.690	–	–
expansion	$\epsilon = 1$	5.00	8.33	4.629	2.563	+15%	0.20
line-binned	$\epsilon = 1$	4.83	7.35	4.470	2.702	+2%	0.13
resolved	fluorescence	4.96	7.55	4.519	2.653	–	–
expansion	fluorescence	5.13	8.83	4.686	2.506	+17%	0.32
line-binned	fluorescence	4.84	7.84	4.434	2.460	+4%	0.45

5 DISCUSSION

5.1 Why the two closures fail differently

The expansion opacity weights each line in a bin by $1 - e^{-\tau}$, so a saturated line contributes at most one interaction’s worth of opacity however deep it is; the line-binned opacity weights it by τ . In a bin with one saturated line and many weak ones the two treatments therefore differ in how much they absorb,

and in which line they attribute the absorption to. Under complete thermal redistribution the second difference is invisible: the energy is re-emitted from the thermal emissivity whichever line absorbed it, and only the first difference, the total attenuation, survives. The line-binned closure preserves the total attenuation by construction and is the closer of the two, as Fontes et al. (2020) found and as Sections 4.2 and 4.3 re-find; the expansion closure under-absorbs in saturated bins




Figure 4. Closure minus resolved transport per band on P1 at 2 d under the downward macroatom (solid) and under the full macroatom with upward transitions driven by an imposed diluted Planck field (dashed), for the expansion (circles) and line-binned (squares) treatments.

and sits a few tenths too bright, an offset that is nearly grey because the saturated lines are spread across the bands.

Under fluorescence the second difference becomes physical. The line-binned closure attributes its extra absorptions to lines in proportion to τ , which on a saturated forest means the same few deep lines again and again; each activates an upper level whose cascade lands elsewhere in the spectrum than the thermal emissivity would have put the energy. The colours move, in the direction of the cascade: the near-infrared bands lose energy that the resolved transport re-emits there. The expansion closure attributes its absorptions in proportion to $1 - e^{-\tau}$, which on a saturated line is unity and on a weak line is small, so its distribution of activated levels is close to that of the resolved transport; its error stays the grey under-absorption offset it had before. The two closures fail in opposite ways, the expansion closure by a nearly grey offset and the line-binned closure by its colour, and the difference between them is the information the coarse-graining discarded.

The non-termination on the thinner line list is the limiting case. A closure that integrates the depth over a bin has no escape probability for the individual line; a packet re-emitted at a saturated line’s frequency inside its own bin is re-absorbed with certainty, and the walk closes on itself. Complete redistribution breaks the loop by moving the packet to another frequency; the downward macroatom, which returns the packet to the same line, does not. Any implementation of a line-binned opacity with explicit fluorescence needs a per-line escape treatment inside the bin, which is to say it needs to retain the information the binning removed.

5.2 What the light curve adds

The snapshot colour errors are larger than the light-curve colour errors because the light curve integrates over the distribution of escape times and because the closure’s error changes sign between bands and epochs. What survives the integration is a systematic, opposite-signed difference between the two closures of a few tenths of a magnitude in the near-infrared colours, and a peak-luminosity excess of the expansion

closure of a sixth under either redistribution. The resolved transport loses the most energy to expansion work in every case; a coarse opacity that lets packets out with fewer interactions is brighter for that reason, and the effect is bolometric rather than chromatic. The near agreement of the three treatments in the peak luminosity that [Fontes et al. \(2020\)](#) reported is what this calculation also gives under complete redistribution; its amplitude here is larger, and its ordering different, because the thermodynamic trajectory is prescribed rather than solved.

5.3 What is not shown

The temperature is prescribed, or in one variant fed back from the radiation field, and there is no gas energy equation; the light curves are extensions of the benchmark under a fixed trajectory, not kilonova models. The full-macroatom test imposes its radiation field rather than solving for it. The Fontes problem is a single element, and the lanthanide states use a single calibrated line list with one uncalibrated list as a check on the neodymium ions only. The g and r bands are starved of packets in the light curves at the epochs shown. The multi-shell states hold the composition and ionization of the published models fixed and do not evolve them. None of these choices differs between the legs being compared, and every comparison is between legs that share everything but the switch under test; the numbers are the closure errors under those conditions, and their sizes should be read as such.

6 CONCLUSIONS

On one code, one line list and one set of ejecta states, the expansion-opacity and line-binned coarse-grainings of the r-process line forest reproduce resolved Sobolev transport to within a few tenths of a magnitude under complete thermal redistribution, with the line-binned closure the closer, as [Fontes et al. \(2020\)](#) found. With the single change to energy-conserving fluorescence, the expansion closure’s error is unchanged, 0.15–0.19 mag in z on the lanthanide-rich state; the line-binned closure’s error grows to 0.16–0.80 mag with the opposite sign, its colour error on the Fontes problem opens to more than a magnitude in a snapshot and to 0.45 mag over a light curve, and on a thinner line list its transport does not terminate. The bolometric ordering of the three treatments over the light curve is the same under both redistributions; fluorescence differentiates the closures through their colours. A coarse opacity that retains only the integrated depth of a bin cannot retain which transition absorbed, and once the post-absorption physics depends on that, the closure’s validation under complete redistribution does not carry over. Kilonova transport codes that use a line-binned opacity with a macroatom or other fluorescence treatment need a per-line escape treatment inside the bin; codes that use the expansion opacity with fluorescence inherit a grey offset of a few tenths of a magnitude that is stable across states, epochs, grids and the strength of the fluorescence treatment.

ACKNOWLEDGEMENTS

The transport code, the run records from which every number in this paper is generated, and the pre-correction histories are available in the repository named below.

DATA AVAILABILITY

The code and all run records are at https://github.com/zhuygln/revisit_sobolev. The manuscript's numbers are generated from the frozen record `paper4/FROZEN.json`; the calibrated line list is that of [Flörs et al. \(2026\)](#) and the Japan–Lithuania list is publicly available ([Kato et al. 2021](#)).

REFERENCES

- Barnes J., Kasen D., 2013, *The Astrophysical Journal*, 775, 18
- Bulla M., 2019, *Monthly Notices of the Royal Astronomical Society*, 489, 5037
- Castor J. I., 1970, *Monthly Notices of the Royal Astronomical Society*, 149, 111
- Collins C. E., Bauswein A., Sim S. A., Vijayan V., Martínez-Pinedo G., Just O., Shingles L. J., Kromer M., 2023, *Monthly Notices of the Royal Astronomical Society*, 521, 1858
- Collins C. E., et al., 2026, *Astronomy & Astrophysics*
- Eastman R. G., Pinto P. A., 1993, *The Astrophysical Journal*, 412, 731
- Flörs A., da Silva R. F., Marques J. P., Sampaio J. M., Martínez-Pinedo G., 2026, *Physical Review D*, 113, 063041
- Fontes C. J., Fryer C. L., Hungerford A. L., Wollaeger R. T., Korobkin O., 2020, *Monthly Notices of the Royal Astronomical Society*, 493, 4143
- Gillanders J. H., Smartt S. J., Sim S. A., Bauswein A., Goriely S., 2022, *Monthly Notices of the Royal Astronomical Society*, 515, 631
- Jeffery D. J., 1993, *The Astrophysical Journal*, 415, 734
- Karp A. H., Lasher G., Chan K. L., Salpeter E. E., 1977, *The Astrophysical Journal*, 214, 161
- Kasen D., Thomas R. C., Nugent P., 2006, *The Astrophysical Journal*, 651, 366
- Kasen D., Badnell N. R., Barnes J., 2013, *The Astrophysical Journal*, 774, 25
- Kasen D., Metzger B., Barnes J., Quataert E., Ramirez-Ruiz E., 2017, *Nature*, 551, 80
- Kato D., Murakami I., Tanaka M., Banerjee S., Gaigalas G., Kitovienė L., Rynkun P., 2021, Japan-Lithuania Opacity Database for Kilonova (version 2.1), <http://dpc.nifs.ac.jp/DB/Opacity-Database/>
- Kato D., Tanaka M., Gaigalas G., Kitovienė L., Rynkun P., 2024, *Monthly Notices of the Royal Astronomical Society*, 535, 2670
- Kawaguchi K., Shibata M., Tanaka M., 2018, *The Astrophysical Journal Letters*, 865, L21
- Lucy L. B., 2002, *Astronomy & Astrophysics*, 384, 725
- Lucy L. B., 2003, *Astronomy & Astrophysics*, 403, 261
- Lucy L. B., 2005, *Astronomy & Astrophysics*, 429, 19
- Metzger B. D., 2020, *Living Reviews in Relativity*, 23, 1
- Morag J., 2026, *Monthly Notices of the Royal Astronomical Society*, 549, stag938
- Ricigliano G., Perego A., Borhanian S., Loffredo E., Kawaguchi K., Bernuzzi S., Lippold L. C., 2024, *Monthly Notices of the Royal Astronomical Society*, 529, 647
- Shingles L. J., et al., 2023, *The Astrophysical Journal Letters*, 954, L41
- Sobolev V. V., 1960, *Moving Envelopes of Stars*. Harvard Books

- on Astronomy Vol. 10, Harvard University Press, Cambridge, MA, [doi:10.4159/harvard.9780674864658](https://doi.org/10.4159/harvard.9780674864658)
- Tanaka M., Kato D., Gaigalas G., Kawaguchi K., 2020, *Monthly Notices of the Royal Astronomical Society*, 496, 1369
- Wollaeger R. T., Korobkin O., Fontes C. J., et al., 2018, *Monthly Notices of the Royal Astronomical Society*, 478, 3298