

Causal graph analysis of COVID-19 observational data in German districts reveals effects of determining factors on reported case numbers

Edgar Steiger^{1*}, Tobias Mußnug¹, Lars Eric Kroll¹

¹ Central Research Institute of Ambulatory Health Care in Germany (Zi), Salzufer 8, D-10587 Berlin, Germany

* esteiger@zi.de

Abstract

~~Mobility, awareness, and weather~~ Several determinants are suspected to be causal drivers for new cases of COVID-19 infection. Correcting for possible confounders, we estimated ~~their causal effects~~ the effects of the most prominent determining factors on reported case numbers. To this end, we used a directed acyclic graph (DAG) as a graphical representation of the hypothesized causal effects of the ~~aforementioned~~ determinants on new reported cases of COVID-19. Based on this, we computed valid adjustment sets of the possible confounding factors. We collected data for Germany from publicly available sources (e.g. Robert Koch Institute, Germany’s National Meteorological Service, Google) for 401 German districts over the period of 15 February to 8 July 2020, and estimated total causal effects based on our DAG analysis by negative binomial regression. Our analysis revealed favorable ~~causal~~ effects of increasing temperature, increased public mobility for essential shopping (grocery and pharmacy) or within residential areas, and awareness measured by COVID-19 burden, all of them reducing the outcome of newly reported COVID-19 cases. Conversely, we saw adverse effects ~~of leading to an increase in new COVID-19 cases for~~ public mobility in retail and recreational areas or workplaces, awareness measured by searches for “corona” in Google, ~~and higher rainfall, leading to an increase in new COVID-19 cases and some~~ socio-demographic factors. Non-pharmaceutical interventions were found to be effective in reducing case numbers. This comprehensive causal graph analysis of a variety of determinants affecting COVID-19 progression gives strong evidence for the driving forces of mobility, public awareness, and temperature, whose implications need to be taken into account for future decisions regarding pandemic management.

Introduction

As the COVID-19 pandemic progresses, research on mechanisms behind the transmission of SARS-CoV-2 shows conflicting evidence [1–3]. While effects of mobility have been extensively discussed, less is known on other factors such as changing awareness in the population [4–6] or the effects of temperature [7–9]. A limiting factor in many studies is the lack of a causal approach to assess the causal contributions of various factors [10]. This can lead to distorted estimates of the causal factors with observational data [10–12].

With COVID-19, we find ourselves in a situation in which information on the causal contribution of various influencing factors in the population is urgently needed to

inform politicians and health authorities. On the other hand, trials cannot be carried out for obvious ethical and legal reasons. Therefore, when assessing the effects of determinants of SARS-CoV-2 spread, special attention must be paid to strategies for the selection of confounding factors.

Another problem with assessing the effects of various determinants of SARS-CoV-2 spread is the heterogeneity of the countries and regions examined for example in the Johns Hopkins University (JHU) COVID-19 database [13]. The comparison of time series of case numbers from different countries and observational periods can be strongly distorted by different factors like testing capacities and regional variations.

Our objective is to provide valid estimates of the ~~causal~~ effects of the main drivers of the pandemic with ~~reduced-bias~~ causal graph approach. We conducted a scoping review of the available studies regarding signaling pathways and determinants of the spread of SARS-CoV-2 infections and the reported new COVID-19 cases. Then we integrated the current findings into a directed acyclic graph for the progress of the pandemic at the regional level. Using the resulting model and the do-calculus we found identifiable effects without blocked causal paths whose effects can be analyzed with observational data [14]. We used regional time series data of all ~~401-German-districts~~ German districts (401) from various publicly available sources to analyze these questions on a regional level. Germany is a good choice in this regard, because it has ample data on contributing factors on the regional level and has had high testing and treatment capacities from early on in the pandemic.

Causal Model

We used a directed acyclic graph (DAG) [11, 12] as a tool to analyze the causal relationships between several exposures and SARS-CoV-2 spread. To get an overview on published associations, a scoping review was conducted from 20th to 22nd of May 2020 within Pubmed and Google scholar. Restrictions were applied to English and German language and the publication date in the last one year. The following search terms were applied to abstracts and title in Pubmed (“COVID-19” OR “COVID19” OR “Corona” OR “Coronavirus” OR “SARS-CoV-2”) and connected separately in each case with the exposure variables (“mobility”, “public awareness”, “awareness”, “google trends”, “ambient temperature”, “temperature”). For “mobility”, we analyzed $n = 8$ studies, $N = 103$ were scanned in Pubmed, together with the first ten pages (100 results) in Google scholar (“awareness”/“public awareness”/“google trends” $n = 9$, $N = 215$; “temperature”/“ambient temperature” $n = 16$, $N = 235$). We integrated these findings where possible into the construction of our DAG, which can be seen in Fig 1.

Fig 1. DAG of determinants of reported COVID-19 cases on the district level. Unobserved variables are light gray, variables marked with an asterisk (*) are confounded by weekday/holiday.

A number of studies report a strong association of **mobility** restrictions on the number of new COVID-19 cases: Restrictive measures (e.g. “stay-at-home” orders, travel bans, or school closures) are shown to possibly reduce the COVID-19 incidence [2, 15–21]. However, some studies point out the combination of various non-pharmaceutical interventions (NPIs) is decisive to prevent new infections [22, 23].

Google Trends [24] data can be used as a tool to get insights into public interest (**awareness**) in the coronavirus disease. Several recent studies imply a connection of relative search volumes (RSV) indices and reported new COVID-19 cases [4–6, 25–30]. Some search terms e.g. “COVID-19” or “coronavirus” predated newly infected cases/total number of cases by roughly 7 to 14 days for different countries [4–6, 26].

Additionally, we acknowledged that individual risk-aware behavior might be a reaction to the current COVID-19 burden (measured as reported cases at the day of exposure).

Mixed evidence is available regarding the effect of **temperature**: On the one hand several papers report an association between increase in temperature and decrease in newly infected COVID-19 cases [7–9, 31–36]. On the other hand, also the opposite has been found [37, 38]. Some studies found no association at all [22, 39–42]. It should be noted that few studies considered other confounding variables than meteorological ones (especially age and population density among others [22, 36, 39]). In addition, the transferability of results between different climate zones is questionable. To avoid possible bias caused by weather variables other than temperature, we included rain, wind, and humidity in our model.

When investigating causal determinants of SARS-CoV-2 infections, a number of confounders have to be considered. Well-known risk factors for SARS-CoV-2 as well as for other infections are demographic factors such as age, gender, socio-economic status (SES), population density, and foreign citizenship/ethnicity [13, 43, 44]. In Germany along with other countries (i.e. Brazil, USA, or the UK), populist parties or politicians and their electorate tend to be more sceptical about effects of containment measures than the other part of the electorate [45, 46]. Therefore we considered both “right-wing populist party votes” and “voter turnout” as possible confounders. Public health interventions were also taken into account (contact restrictions, school closures etc.), as their implementation showed strong correlations with controlling the spread of SARS-CoV-2 [22, 23, 47]. To avoid bias due to reporting delay of case numbers we had to include weekday and German holidays. We included some unobserved variables in our DAG (e.g. “Herd immunity”), too. Please note that “Exposure to SARS-CoV-2” is itself an unobserved variable: German case numbers are reported with delay after date of exposure and symptom onset. *Exposure to the virus* should not be confused with the formal *exposure variables* of the DAG (~~mobility, awareness, temperature~~).

Materials and methods

Data

We collected and aggregated data on reported COVID-19 cases, regional socio-demographic factors, weather, and general mobility on district and state level in Germany for the period of 15 February 2020 to 8 July 2020. Our observation period for the outcome consisted of all dates from ~~23~~ 20 February 2020 to 8 July 2020 (~~$T = 137$~~ $T = 140$), since we used a lag of ~~8~~ 5 days for all confounders. We did not exclude any states or districts ($K = 401$). We analyzed the daily reported number of new cases as outcome (~~$K \cdot T = 54\,937$~~ $K \cdot T = 56\,140$ observations). The set of possible predictors was derived from our causal DAG (see Table 1 and Fig 1). Due to modelling and data limitations, some of the predictors were unobserved or were modelled as a construct consisting of several variables. For our causal graph analysis, we computed adjustment sets ~~in three different scenarios for separate~~ separately for all observed exposures within the DAG : i) ~~mobility of population~~, ii) ~~awareness of COVID-19 (i.e. Google searches for “corona”)~~, iii) ~~weather (i.e. temperature~~ (if the respective exposure was identifiable within the DAG causal analysis framework).

Variables

We downloaded German daily case numbers on district level reported by Robert Koch Institute (RKI, [48]) and aggregated them by date. The number of daily active cases for day d was derived by subtracting the total number of reported cases on day d and day

$d - 14$ (14 days as a conservative estimate for the infectious period, which corresponds here to the required quarantine time in Germany).

To assess the mobility of the German population, we used data publicly available on German state level from Google [49]. Measurements are daily relative changes of mobility in percent compared to the period of 3 January 2020 to 6 February 2020. Missing values (25 out of 13 488) were imputed with value 0 and the state level measurements were passed onto districts within the corresponding state. Google mobility data was available for six different sectors of daily life (“retail and recreation”, “grocery and pharmacy”, “parks”, “transit stations”, “workplaces”, “residential”) which means that “mobility” is a construct consisting of several variables. All variables but “residential” mobility are relative changes of daily visitor numbers to the corresponding sectors compared to the reference period. “Residential” mobility is the relative change of daily time spent at residential areas. The six mobility variables showed high correlations among each other and with other variables. To reduce multicollinearity, we transformed them by principal component analysis (PCA) into six uncorrelated principal components which were used in place of the original variables.

The notion of awareness in the population of COVID-19 describes the general state of alertness about the new infectious disease. As such, it was hard to measure directly. As a proxy, we used the relative interest in the topic term “corona” as indicated by Google searches. The daily data was available on state level [24] and passed onto district level. As a second proxy for awareness, we used the daily reported number of COVID-19 cases on the day of the exposure: Since media reported case numbers prominently, we assumed that this could reflect individual awareness, too.

We constructed daily weather from four variables (“temperature”, “rainfall”, “humidity”, “wind”). Weather data was downloaded from Deutscher Wetterdienst (DWD, [50]) for all weather stations in Germany below 1000 meters altitude with daily records for our observation period. District level daily weather data was aggregated per district by averaging the data from the three nearest weather stations (which includes weather stations inside the district). Missing values were imputed with mean values ($n = 59$ for wind).

The reported number of COVID-19 cases varied strongly by day of the week. Thus, we included “weekday” as a categorical variable. Similarly, the reported cases and the exposure to the virus were affected by official holidays. Within the observation period, this included among others Good Friday, Easter Monday, and Labor Day. To correct for effects of these days, we included two variables in the model, “Holiday (report)” (indicates if the day of the report was a holiday, because governmental health departments were less likely to be on full duty) and “Holiday (exposure)” (indicates if the day of exposure to the virus was a holiday, because the population behaves differently on holidays).

For different official and political ~~measures~~ interventions on a daily basis and the district level we used one-hot encoded daily variables, i.e. ban of mass gatherings, school and kindergarten closures and their gradual reopening, contact restrictions, and mandatory face masks for shopping and public transport.

We included several social, economic, and demographic factors on the district level with direct or indirect influence on the risk of exposure to SARS-CoV-2 in our analysis. All are readily available from INKAR database [51]. We used the share of population that is 65 years or older and the share of population that is younger than 18 years (Age), the share of females in population (Gender), the population density, the share of foreign citizenships and the share of the population seeking refuge (Foreign citizenship), the share of low-income households (Socio-economic status), voter turnout, share of right-wing populist party votes, and the number of nursing (retirement) homes.

All continuous variables but the outcome “Reported new cases of COVID-19” and

Table 1. Observed model variables.

variable	dynamics	level	type	unit/comment	source
Weekday	daily	national	categorical	Sat through Thu as six binary variables, Fri as baseline	-
Holiday (report)	daily	national	binary	-	-
Holiday (exposure)	daily	national	binary	-	-
Mobility					
retail and recreation	daily	state	numeric	percent change compared to reference period	Google [49]
grocery and pharmacy	daily	state	numeric	percent change compared to reference period	Google [49]
parks	daily	state	numeric	percent change compared to reference period	Google [49]
workplaces	daily	state	numeric	percent change compared to reference period	Google [49]
residential	daily	state	numeric	percent change compared to reference period	Google [49]
transit stations	daily	state	numeric	percent change compared to reference period	Google [49]
Awareness					
Searches corona	daily	state	numeric	percent relative to other states and observation period	Google [24]
COVID-19 burden	daily	district	numeric	reported cases on day of exposure	RKI [48]
Weather					
Rainfall	daily	district	numeric	mm (l/sqm)	DWD [50]
Temperature	daily	district	numeric	°C	DWD [50]
Humidity	daily	district	numeric	relative humidity (%)	DWD [50]
Wind	daily	district	numeric	m/s	DWD [50]
Interventions					
Ban of mass gatherings	daily	national	binary	-	-
School and kindergarten closures	daily	state	numeric	0 for no closure, 1 for full closure, 0.5 for partial reopening	-
Contact restrictions	daily	national	binary	-	-
Mandatory face masks	daily	district	binary	-	IZA [52]
Socio-demographic					
Age	constant	district	numeric	2 variables: share of population ≥ 65 years & ≤ 18 years	INKAR [51]
Gender	constant	district	numeric	share of female population	INKAR [51]
Population density	constant	district	numeric	population per sqkm	INKAR [51]
Foreign citizens	constant	district	numeric	2 variables: share of foreign citizens & of population seeking refuge	INKAR [51]
Socio-economic status	constant	district	numeric	share of households with low income	INKAR [51]
Turnout	constant	district	numeric	voter turnout in last election	INKAR [51]
Right-wing populist party votes	constant	district	numeric	share of votes for AfD in last election	INKAR [51]
Nursing homes	constant	district	numeric	number of nursing (retirement) homes	INKAR [51]
Case numbers					
Reported new cases of COVID-19	daily	district	numeric	-	RKI [48]
Active cases	daily	district	numeric	active cases on day of report	RKI [48]

the offset “Active cases” were centered and scaled by one standard deviation for numerical stability. We did not scale variables to unit variance to maintain interpretability of effects on the original scale of variables. Additionally, while we left binary variables as-is. After estimating the effects of variables, we re-scaled continuous variables’ effects to their original scale. Additionally for mobility variables, we re-transformed the effects of the principal components to the original mobility variables. Furthermore, we lagged the effect of all variables (but outcome, offset, and the non-dynamic socio-demographic variables) by 8 days (see Section Results 5 days (optimal lag found by cross-validation) which means that we assumed that their effects on the outcome will be visible after 8-5 days.

Methods

Causal analysis with DAG and adjustment sets

We used a directed acyclic graph as a graphical representation of the hypothesized causal reasoning that leads to exposure to the SARS-CoV-2 virus, onset of COVID-19, and finally reports of COVID-19 cases. We use the terms “causal effect” or “causal relationship” for effect estimates that are based on this causal graph framework. Every node v_i in the graph is the graphical representation of an observed or unobserved variable x_i , a directed edge e_{ij} is an arrow from node v_i to v_j that implies a direct causal relationship from variable x_i onto variable x_j . The set of all nodes is denoted by V , the set of all edges by E , as such, the complete DAG is the tuple $G = (V, E)$. The seminal works of Spirtes and Pearl [53,54] introduce the theory of causal analysis, do-calculus, and how to analyze a DAG to estimate the total or direct causal effect from a variable x_i onto a variable x_j . The direct effect is the effect associated with the edge e_{ij} only (if it exists), while the total effect takes indirect effects via other paths from v_i to v_j into account, too. Here we estimated total effects only, since most of our variables were not hypothesized to have a direct effect on the *reported* number of new COVID-19 cases. In contrast to prediction tasks, where one would include all variables available, it is actually ill-advised to use all available variables to estimate causal effects, due to introducing bias by adjusting for unnecessary variables within the causal DAG. This is why we need to identify a valid set of necessary variables (an adjustment set) to estimate the proper causal effect [54]. The “minimal adjustment set” [55] is a valid adjustment set of variables that does not contain another valid adjustment set as a subset. However, identifying a minimal adjustment set might not be enough to reliably estimate the causal effect. Thus, we identified the “optimal adjustment set” [56] as the set of variables which is a valid adjustment set while having the lowest ~~asymptotic variance in the resulting causal effect estimates~~ Akaike information criterion (AIC).

We analyzed the DAG from Fig 1 with the R Software [57] and the R packages `dagitty` (formal representation of the graph and minimal adjustment sets [12]) and `pcalg` (for finding an optimal adjustment set [58]). For the defined exposures and the outcome “Reported new cases of COVID-19”, we computed the minimal and optimal adjustment sets. Since it was possible that these sets contained unobserved variables that needed to be left out of the regression model, we chose the valid set with the ~~highest pseudo- R^2~~ lowest AIC (see next section) to estimate the final total causal effect from exposure to outcome.

Regression with negative binomial model

We can estimate the causal effect from exposure to outcome by regression [54]. Since the outcome “Reported new cases of COVID-19” is a count variable, one should not employ a linear regression model with Gaussian errors, but instead we assumed a log-linear relationship between the expected value of the outcome Y (new cases) and regressors x , as well as a Poisson or negative binomial distribution for Y :

$$\log(\mathbb{E}[Y|x]) = \alpha + \sum_{i \in S} \beta_i \cdot x_i, \quad (1)$$

where α is the regression intercept, S is the set of adjustment variables for the exposure i^* including the exposure variable itself, β_i are the regression coefficients corresponding to the variables x_i . As such β_{i^*} is the total causal effect from exposure variable x_{i^*} on the outcome Y .

The Poisson regression assumes equality of mean and variance. If this is not the case one observes so-called overdispersion (the variance is higher than the mean), this

indicates one should use regression with a negative binomial distribution instead to estimate the variance parameter separately from the mean. 211 212

We needed to account for the fact that our outcome is not counted per time unit (one day) only, but depends on the number of active COVID-19 cases: Holding all other variables fixed, the number of new cases Y is a constant proportion of the number of active cases A . This was modeled by including an offset $\log(A + 1)$ in the regression model Eq (1): 213 214 215 216 217

$$\begin{aligned} \log(\mathbb{E}[Y|x]) &= \alpha + \log(A + 1) + \sum_{i \in S} \beta_i \cdot x_i \\ \Leftrightarrow \log\left(\frac{\mathbb{E}[Y|x]}{A + 1}\right) &= \alpha + \sum \beta_i \cdot x_i \end{aligned} \quad (2)$$

$$\Leftrightarrow \frac{\mathbb{E}[Y|x]}{A + 1} = \exp(\alpha) \cdot \prod \exp(\beta_i)^{x_i}. \quad (3)$$

Here we added a pseudocount “+1” to ensure a finite logarithm and avoid division by 0. 218 219

One can interpret the model as approximating the log-ratio of new cases and active cases by a linear combination of the regressor variables in Eq (2). If all variables x_i are centered in Eq (3), we have for the baseline $\forall i \ x_i = 0 \Rightarrow \mathbb{E}[Y|x = 0] = \exp(\alpha) \cdot (A + 1)$. In other words, the exponentiated intercept is the baseline daily infection rate (how many people does one infected individual infect in one day). If we hold all variables x_i fixed (e.g. at baseline 0) in Eq (3) but now increase the exposure variable $x_{i^*} = 0$ by one unit to $x_{i^*} + 1 = 0 + 1$, we have

$$\begin{aligned} \mathbb{E}[Y|x'] &= \exp(\alpha) \cdot (A + 1) \cdot \exp(\beta_{i^*}^{x_{i^*}+1}) \prod_{i \neq i^*} \exp(\beta_i)^0 \\ &= \exp(\alpha) \cdot (A + 1) \cdot \exp(\beta_{i^*}), \end{aligned}$$

which means the exponentiated coefficient β_{i^*} describes the rate change of the outcome by one unit increase of the exposure. 220 221

In practice, given observations of Y and x we estimate the regression coefficients α and β_i by maximum likelihood [59]. Our observational measurements are y_{kt} and x_{ikt} , where k indicates the corresponding district and t the date of measurement. 222 223 224

~~When we analyzed different~~ We conducted a log-linear regression (function `glm` with `family=poisson()` for Poisson regression, and `glm.nb` from the MASS package for the negative binomial regression [60]) for the full data set to assess general model adequacy and to estimate the θ parameter of the negative binomial. The proper lag between exposures and outcome was found by 10-fold cross-validation on different lags between 1 and 20 days. Model diagnostics on the final full model did not show severe problems with model assumptions (linearity, distribution of residuals, independence of observations). Analysis of variance inflation factors revealed some problems with multicollinearity. To reduce the effects of multicollinearity, first we transformed the highly correlated mobility variables by PCA as described above. Second, we used a ridge regression approach [61], which is a regularization method that shrinks regression coefficients and alleviates the effect of correlation between variables on their respective regression coefficients. Furthermore, regularized regression allows for better fits on unseen data, thus preventing overfitting the data, too. The hyper-parameter λ of the ridge regression was chosen by 10-fold cross-validation, where the folds were constructed from random subsets of the 401 districts. We used this hyper-parameter with the `cv.glmnet` function from the R package `glmnet` [62] with `family=negative.binomial(theta)` and chose the λ value within one standard 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242

deviation from the minimal λ as regularization hyper-parameter. Afterwards, we calculated the effects of separate exposures on the outcome. For every exposure, we analyzed the different valid adjustment sets given by analysis of the causal DAG (i.e. the minimal and optimal adjustment sets). Then, we first checked if the respective set included unobserved variables. If this was the case for the optimal adjustment set, we discarded the unobserved variables from the set and checked if it was still a valid adjustment set (function `gac` in package `pcalg` [63]). If a minimal adjustment set contained unobserved variables, we discarded the whole set. We conducted a log-linear regression (function `glm` with `family=poisson()` for Poisson regression, If no valid adjustment set for a given exposure was available, we concluded that the effect of this exposure was unidentifiable within our causal graph. We used the function `glmnet` with the parameters θ and `glm.nb` from the `MASS` package for the negative binomial regression [60]) for λ as above on every remaining valid adjustment set as regressors and calculated a Pseudo- R^2 given by $1 - V_m/V_0$, where V_m is the sum of squared prediction errors of the current model and V_0 is the sum of squared prediction errors of the null model (intercept and offset only). That is, our Pseudo- R^2 is 1 minus the fraction of variance unexplained (that is, we applied ridge regression) and calculated the Akaike information criterion (AIC) for this model/set of regressors. Finally, for every exposure, we decided for the model/adjustment set with the highest pseudo- R^2 (if available) with the lowest AIC. We report the exponentiated estimated coefficients along with 99 percent confidence intervals of the estimates for the separate exposures on their original scale.

Results

Descriptive statistics for the included variables are presented in Table 2.

In the observational period, the number of daily reported COVID-19 cases increased till the end of March/beginning of April and continually decreased afterwards till the beginning of June 2020 with a slight increase and decrease afterwards (Fig 2A). On the other hand, the (log-)ratio of reported cases over active cases decreased steeply till the mid of April and increased steadily afterwards with a slight decrease close to the end of the observation period (Fig 2B). Both figures exemplify a considerable variation among the districts (light blue points are individual district's data).

Fig 2. Temporal and district level variation of outcome (log-scale).

In Germany, we observed a rebound in mobility after the initial political measures, reductions in incident cases were associated with a diminishing public interest in COVID-19, and temperatures were overall increasing (cf. Fig 3); with correlations between temporal progression and mobility in retail and recreation $r_{A,D} = -0.02$, awareness ("Searches corona") $r_{A,C} = -0.29$, and temperature $r_{A,B} = 0.79$, $r_{A,D} = 0.8$.

Fig 3. Temporal variation of outcome and main determinants.

0.1 Main results

Main results

We list the results of our causal analysis for the effects of our different exposure variables in Table 3. The estimates are multiplicative rates of increase/decrease for a

Table 2. Descriptive statistics for observed variables.

Variable	mean (SD)
n	56140
Mobility	
retail and recreation	-26.62 (24.60)
grocery and pharmacy	-3.94 (22.77)
parks	47.26 (58.20)
workplaces	-22.96 (20.35)
residential	8.13 (6.49)
transit stations	-29.58 (21.11)
Awareness	
Searches corona	26.94 (18.23)
COVID-19 burden	3.50 (10.28)
Weather	
Rainfall	1.89 (4.01)
Temperature	10.90 (5.33)
Humidity	67.81 (13.03)
Wind	3.63 (1.66)
Interventions	
Ban of mass gatherings	0.83 (0.38)
School and kindergarten closures	0.54 (0.36)
Contact restrictions	0.74 (0.44)
Mandatory face masks	0.49 (0.50)
Socio-demographic	
Age (pop. 65 and older)	22.09 (2.74)
Age (pop. younger 18)	16.17 (1.25)
Gender	50.59 (0.64)
Population density	533.75 (701.84)
Foreign citizens	10.03 (5.14)
Foreign citizens (refugees)	1.88 (1.14)
Socio-economic status	30.64 (6.02)
Turnout	75.08 (3.79)
Right-wing populist party votes	13.39 (5.32)
Nursing homes	36.11 (30.69)
Case numbers (Outcome and offset)	
Reported new cases COVID-19	3.53 (10.29)
Active cases	48.76 (120.86)

one unit increase of the respective variable: Values above 1 lead to an increase, below 1 to a decrease of the infection rate. To put these estimates into perspective, Fig 4 shows the relative causal effect of the different exposure variables on the number of reported COVID-19 cases on a range of sensible values of the exposure variables (95 percent quantiles of data points).

Fig 4. Relative causal effects of exposures.

Within our framework, we saw ~~significant effects for mobility in retail/recreational areas and essential shopping (grocery and pharmacy).~~ For very different effects for individual mobility variables. For mobility in retail/recreation, an increase of 1 percent

Table 3. Effect estimates from causal graph analysis.

cause	effect estimate
Mobility	
retail and recreation	1.0011
grocery and pharmacy	0.9977
parks	0.9997
transit stations	1.0026
workplaces	1.0033
residential	0.9903
Awareness	
Searches corona	1.0089
COVID-19 burden	0.9980
Weather	
Temperature	0.9905
Rainfall	1.0121
Humidity	1.0057
Wind	1.0329
Interventions	
Interventions (ban of mass gatherings)	0.9729
Interventions (school and kindergarten closures)	0.9277
Interventions (contact restrictions)	0.8314
Interventions (mandatory face masks)	0.9064
Demographic	
Age (pop. 65 and older)	0.9953
Age (pop. younger 18)	1.0120
Foreign citizens	1.0048
Foreign citizens (refugees)	0.9985
Gender	0.9925
Nursing homes	1.0011
Population density	1.0000
Socio-economic status	0.9982

point mobility compared to the reference period (03 January to 06 February 2020) leads to an increase of the daily reported case number by about ~~0.8 percent~~ 0.11 percent. Similarly, mobility on workplaces showed an effect of 0.33 increase in case numbers for every 1 percent point increase in mobility, while mobility on transit stations showed an effect of 0.26 increase in case numbers for every 1 percent point increase. Contrarily, ~~a corresponding~~ the remaining three mobility variables showed negative effects on the number of reported COVID-19 cases. An increase of 1 percent point mobility for the areas of grocery/pharmacy leads to a decrease in the reported case number by approximately ~~0.5 percent~~. Mobility on workplaces showed a small effect of 0.3 increase in case numbers for every ~~0.23 percent~~, while increased mobility of 1 percent point within parks leads to a decrease in the reported case number by approximately 0.03 percent, and finally an increase of 1 percent point ~~increase in mobility~~. Other causal effects of mobility were insubstantial and not consistent in their direction (99 percent confidence intervals of estimates include 1) in residential mobility leads to a decrease by approximately 0.97 percent. Fig 4 shows the effects of mobility on a range of possible values. Thus, we expect an increase of daily cases by approximately ~~23-7.8~~ percent if mobility in ~~retail/recreation workplaces~~ reaches baseline levels of 0 percent difference to the reference period. On the other hand, an increase of mobility for

~~grocery/pharmacy-residential areas~~ by 10 percent points compared to the reference period leads to a reduction of the infection rate by approximately ~~7-1.8~~ percent.

“Awareness” had two opposite effects on the outcome in our DAG. Awareness measured by Google searches for *corona* had a positive effect on the number of reported cases. An one percent point increase of the state’s Google searches (relative to other states and the observation period) leads to an increase of approximately ~~1-0.89~~ percent. For example, if a district shows 10 percent points more relative searches for *corona* than another one, we expect approximately ~~11-9.3~~ percent more infections for this district after ~~8-5~~ days. *COVID-19 burden* (reported number of cases on day of exposure) affected the outcome negatively, where every additional daily case in the district leads to a ~~0.6-0.2~~ percent decrease in newly reported case numbers. The ~~last-corresponding~~ plot in Fig 4 visualizes this relationship: For a local outbreak with 20 daily cases as COVID-19 burden, we estimate as total causal effect a subsequent reduction of infection rate by ~~11-3.9~~ percent.

Within our model, we observed ~~a-causal-effect-effects~~ of temperature and all other weather variables. Every increase of 1 degree Celsius in temperature leads to a reduction of the daily reported case numbers by approximately ~~0.8-0.95~~ percent. On the other hand, we found an increasing effect of rainfall: One millimeter (=1 liter per square meter) more rainfall leads to an increase of reported case numbers by approximately ~~1-1~~ ~~1.21~~ percent. We observe effects for humidity and wind as well (higher humidity ~~leading to more cases, and~~ stronger wind leading to ~~less-more~~ cases). In perspective (Fig 4), with temperature we expect an increase by approximately ~~17.8-21~~ percent at a daily average temperature of 0°C compared to a day with 20°C. For rainfall, we expect on a rainy day with 10 mm rainfall a corresponding increase of the infection rate by approximately ~~11-4~~ ~~12.8~~ percent compared to a day with no precipitation.

~~In all cases we~~ The different intervention variables showed the strongest effects in our analysis, see Table 3. While the first intervention (ban of mass gatherings) reduced subsequent daily case numbers by 2.7 percent, the closure of schools/kindergartens reduced infections by an additional 7.2 percent and mandatory face masks reduced this by another 9.4 percent. The effect of contact restrictions was the strongest in our observation period, with an reduction of the case rates by 16.9.

The effects of the different socio-demographic factors are quite small in comparison to the effects described above. We see an increasing effect on case numbers by additional nursing homes between districts. Districts with a younger population, more foreign citizens, higher population density and a lower average social-economic status showed higher case numbers, too.

For all exposures, our analysis pipeline opted to use the ~~reduced~~ (reduced) optimal adjustment set over the minimal adjustment sets because of ~~higher pseudo-R² values (mostly above 0.3)~~ lower AICs, except for ~~mobility, where exposure variable “nursing homes”, for which~~ the minimal adjustment set had ~~a higher pseudo-R²~~. Notably, these sets always include most of our socio-demographic variables as confounders as well as the policy variables (cf. ~~the lowest AIC~~. For an overview of all final adjustments sets, see Table 4, with the exception being COVID-19 burden). We also decided for the lag of 8. We found that there were no valid adjustment sets for the non-identifiable variables turnout and right-wing populist party votes.

We decided for a lag of 5 days based on ~~the highest pseudo-R² values compared to other lags on the chosen adjustment sets~~ cross-validation. Similarly, negative binomial regression was chosen over Poisson regression, because the latter showed overdispersion and ~~overall lower pseudo-R² values~~ ~~an higher AIC value~~.

Table 4. Final adjustment sets for causal analysis.

	Mobility	Searches corona	COVID-19 burden	Temperature	Rainfall	Humidity	Wind	Interventions	Age	Foreign citizens	Gender	Nursing homes	Population density	Socio-economic status
Weekday	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Holiday (report)	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Holiday (exposure)	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Mobility														
Mobility						x								
Awareness														
Searches corona	x			x	x	x	x					x		
COVID-19 burden	x	x		x	x	x	x	x	x	x	x	x	x	x
Weather														
Temperature	x	x	x			x		x	x	x	x		x	x
Rainfall	x	x	x	x		x	x	x	x	x	x		x	x
Humidity	x	x	x					x	x	x	x		x	x
Wind	x	x	x	x	x	x		x	x	x	x		x	x
Interventions														
Interventions	x	x		x	x	x	x		x	x	x	x	x	x
Socio-demographic														
Age	x	x	x	x	x	x	x	x				x	x	x
Gender	x	x	x	x	x	x	x	x	x	x		x	x	x
Population density	x	x	x	x	x	x	x	x	x	x	x	x		x
Foreign citizens		x	x					x	x		x		x	x
Socio-economic status	x	x	x	x	x	x	x	x				x	x	
Turnout			x					x						
Right-wing populist party votes	x	x	x	x	x	x	x	x				x		
Nursing homes	x	x	x	x	x	x	x	x					x	

Discussion

Main findings

Our objective was to identify ~~causal effects~~ effects of determining factors for COVID-19 cases within a causal framework. We found that weather affects the reported number of infections, especially temperature (which has a reducing effect on case numbers) and rainfall (which increases case numbers). We saw that reports of high case numbers in districts led to a reduction in new infection numbers, which indicates risk-averse awareness in the population and/or effective public health measures to suppress a local outbreak. ~~The overall effect of mobility showed no consistent effect, however, in specific areas significant causal effects could be measured~~ Mobility showed distinct effects: Increasing activity in retail and recreational areas, as well as transit stations and workplaces increased reported case numbers, while increased movement for essential shopping (grocery and pharmacy) and in parks or residential areas led to

reduced case numbers. All interventions considered (ban of mass gatherings, school/kindergarten closures, contact restrictions, mandatory face masks) reduced case numbers considerably. Socio-demographic variables had small effects individually, but in conjunction they explained larger case numbers in (urban) areas with younger population, lower socio-economic status, and higher population density.

Furthermore, we made a strong case for the use of causal DAGs in epidemiology and a pandemic like COVID-19: DAGs allow to choose confounders for the analysis in a principled and statistically correct way while reducing possible causes for bias. Also, the DAG formalization allows for discussion about the underlying causal assumptions.

Comparison with previous research

Most research on determinants affecting case numbers of COVID-19 is restricted to single aspects [5, 16, 32, 35]. To reliably identify ~~causal~~-(causal) drivers, one must adjust for confounders. To this end, we used an integrated model with variables from different aspects like mobility, awareness, weather, or socio-demographics and identified confounders by causal analysis with a directed acyclic graph. A causal approach is used in another current COVID-19 analysis [64]. There, however, they identify the causal relationships (reconstruct a DAG), while we estimated ~~causal~~-effects for a given hypothesized causal DAG.

Several studies assessing the impact of public health measures on mobility have each observed a downward trend accompanied by a decrease in the number of newly reported cases [15–17, 19, 23, 47].

Our findings regarding awareness/Google Trends analysis are in good agreement with the correlations found by others [4, 6, 26], who conclude that alertness to COVID-19 rises several days before the highest number of cases are reported. At this point it should be noted, that awareness is substantially influenced by public media coverage, which should be considered, if possible, in future studies [4]. As such, awareness is difficult to measure and here the number of Google searches for “corona” could only be a proxy for this concept.

In addition, in alignment with other recent published studies, our results confirm evidence which associated a negative effect of temperature on new COVID-19 cases [7–9, 31–36]. It is however controversial to other scientific literature describing no effects [22, 39–42] or even converse correlations [37, 38]. The conflicting results might be explained by different climates and characteristics of the populations under study. While we are confident that our strict causal analysis resulted in effect estimates as undistorted as possible, there might be unconsidered bias in those other studies. Further research needs to be done to elucidate the biological characteristics of the novel virus SARS-CoV-2 regarding its ambient temperature survival and transmission. Finally, we found a positive ~~causal~~-effect of increased precipitation (~~rainfall~~) and a raise in COVID-19 cases, which supports previous observations [33].

Limitations and strengths

While use of a causal DAG is itself a strong tool to identify ~~causal~~-(causal) effects (and not just statistical associations), it introduces two limitations: causal assumptions within the graph (depicted by edges) need to be well justified, and the statistical regression model that calculates total causal effects needs to be appropriate for the task at hand. We endorse our graph as a basis for discussion on residual confounding. We did not try to construct the DAG from the available data (cf. [64]). As such, our proposed DAG is not entirely consistent with the data and there are conditional dependencies between variables that cannot be dissolved by adding edges to the DAG (e.g. between the ~~polices~~-(interventions) like contact restrictions and mandatory face

masks). Another way to identify potential problems in the proposed DAG is to perform a sensitivity analysis of its structure by inspecting its maximal ancestral graph (MAG) or its Markov equivalence class represented by a complete partially DAG (CPDAG) and the existence of valid adjustment sets for these generalized graphs [65]. For the MAG derived from our DAG, only the effects for exposures mobility and searches for corona can be estimated with valid adjustment sets, while for the Markov equivalence class all exposures but COVID-19 burden lead to valid adjustments sets. A further analysis of these implications is out of the scope of this paper.

We observed overdispersion and a substantial increase in model performance with a negative binomial regression compared to Poisson regression, which is in line with the results on COVID-19 daily case counts of ~~others~~ [7,9,17,66] [17] and ~~others~~ [7,9,66]. We did not model case counts with a differential equation model like the classic SIR-model [67] and its successors, since these are more suited to prediction ,e.g. [68], while our choice of a negative binomial regression framework allowed us to estimate the effects of confounders more reliably. There are more advanced statistical methods for count data, e.g. zero-inflated models and mixed models. We tested both approaches as extensions to the negative binomial regression and experienced numerical problems and increased computing time, along with an insubstantial increase in model performance. Furthermore, our model assumed that all variables have effects proportional to the size of their measurements. It is possible that some variables show saturation effects or opposite effects for low, medium, or high values. This could be ~~modelled~~ modeled with polynomial or other transformations of the variables, which we did not employ due to limited temporal and spatial data availability. Interaction effects of variables and confounding effects or mediating variables are explicitly taken care of by deriving the valid adjustment sets for a given exposure based on the causal DAG. Use of a fixed DAG with effect estimation via regression assumes that data was generated by the same underlying process for the observation period. By inclusion of the successive mitigation ~~policies~~ interventions as binary variables we were able to explain some of the variance caused by the changing dynamics of case numbers (similar to [66]). While multicollinearity of variables poses less of a problem for a proper causal graph analysis [69], we addressed the problem of multicollinearity in our predictors by two approaches: principal component analysis for the highly collinear mobility variables as well as a regularized regression approach (ridge regression). The latter (in conjunction with cross-validation) also reduced the problem of overfitting.

We stress the point that our effects were deduced on an aggregate (district) level in the absence of available data on an individual level. As such, conclusions about effects cannot be transferred on individuals without the possibility for an ecological fallacy. Furthermore, as we were using administrative data for our analysis, the results are susceptible to the Modifiable Area Unit Problem (MAUP) [70]. The MAUP postulates that different regional aggregations of the units of observation may lead to different results and conclusions. Due to limited available data for the different variables, there is currently no way to overcome these problems that are inherent to all analyses on aggregated data level.

Our observation period was restricted to succession from late winter to spring and summer (February to July). Nevertheless, this transition with increasing temperature was a natural experiment that allowed clues on weather effects.

We could not include data on health care utilization during the pandemic into our models due to the lack of available resources. This is planned for a later follow up to this paper since we rank health care utilization and mobility within health care facilities among the strong factors for COVID-19 progression: personnel in hospitals and private practices is particularly exposed to infection, while the lack of adequate care for other diseases has severe effects on general health of the population. At the same time, health

care facilities are key for testing and surveillance of COVID-19 patients.

While our analysis focused on Germany and its districts, we assume that results may be transferred to other countries by adjusting for their respective weather conditions, mobility habits, socio-demographic characteristics, and other determining factors.

The code and resources for our analysis are available on Github, we invite other researchers to replicate our analysis with different assumptions using the files provided in the repository¹ of the article.

Discussion of causal effects

In our analysis, the adverse ~~effect-effects~~ of mobility in retail/recreation and workplaces and the favorable effect of mobility in grocery/pharmacy ~~indicate that policies and residential areas indicate that interventions~~ like contact restrictions which limit the number of individual interactions can lead to reduced infection numbers. This is due to retail/recreational and workplace areas encompassing mostly places of ~~social-gatherings like restaurants and bars, malls, sports and music venues, among others~~(social gatherings), while if people are doing more of their essential shopping at supermarkets ; ~~they will most likely-and~~ stay at home with less contact to other people, they are less likely to come in contact with infected individuals.

The ~~causal~~-effects of awareness measured via searches for “corona” and the COVID-19 burden are harder to interpret. We assume that within our model, the searches for “corona” are an insufficient proxy for awareness, while the decreasing effect for future case numbers of high daily COVID-19 burden indicates it affects individual risk-behavior and entails effective non-pharmaceutical interventions.

Similarly, the effects of temperature and rainfall can be interpreted as causal effects for indoor and outdoor activities, such that higher temperatures and low rainfall indicate more people spending time outdoor while lower temperatures and high rainfall result in indoor activities, which lead to more infections. Current research suggests this to be due to the prevalent airborne and respiratory droplets and aerosol transmission of the SARS-CoV-2 virus [71]. In this light, we advocate for precautionary measures like increased hygiene, face masks, and air ventilation for unavoidable indoor activities.

Furthermore, our analyses strongly support the effectiveness of non-pharmaceutical interventions. To a lesser extent, the adverse effects of some socio-demographic factors might help to identify areas that are at higher risk of local COVID-19 outbreaks.

Conclusion

To the best of our knowledge, this is the most comprehensive analysis of causes for COVID-19 infections which integrates different data sources (all publicly available). Causal reasoning with a DAG allows us to estimate the possible causal effects more reliably.

Our findings suggest that the ~~causal-infection-driving~~ effects of mobility, awareness, and weather (and to some extent socio-demographic factors) need to be taken into account when deciding for mitigation and suppression ~~measures~~interventions, depending on the recent and future COVID-19 pandemic development.

Acknowledgments

We are thankful for feedback from Thomas Czihal, Johannes Textor, ~~and~~ Ralph Brinks, and an anonymous reviewer who gave helpful suggestions on ~~an earlier draft earlier~~

¹The repository is located here: <https://github.com/zidatalab/causalcovid19>

References

1. WHO Team. Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19); 2020. Available from: [https://www.who.int/publications-detail/report-of-the-who-china-joint-mission-on-coronavirus-disease-2019-\(covid-19\)](https://www.who.int/publications-detail/report-of-the-who-china-joint-mission-on-coronavirus-disease-2019-(covid-19)) [cited 2020-06-25].
2. Chinazzi M, Davis JT, Ajelli M, Gioannini C, Litvinova M, Merler S, et al. The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. *Science*. 2020;368(6489):395–400. doi:10.1126/science.aba9757.
3. Guan W, Ni Z, Hu Y, Liang W, Ou C, He J, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *New England Journal of Medicine*. 2020;doi:10.1056/NEJMoa2002032.
4. Higgins TS, Wu AW, Sharma D, Illing EA, Rubel K, Ting JY. Correlations of Online Search Engine Trends With Coronavirus Disease (COVID-19) Incidence: Infodemiology Study. *JMIR Public Health and Surveillance*. 2020;6(2):e19702. doi:10.2196/19702.
5. Li C, Chen LJ, Chen X, Zhang M, Pang CP, Chen H. Retrospective analysis of the possibility of predicting the COVID-19 outbreak from Internet searches and social media data, China, 2020. *Eurosurveillance*. 2020;25(10). doi:10.2807/1560-7917.ES.2020.25.10.2000199.
6. Yuan X, Xu J, Hussain S, Wang H, Gao N, Zhang L. Trends and Prediction in Daily New Cases and Deaths of COVID-19 in the United States: An Internet Search-Interest Based Model. *Exploratory research and hypothesis in medicine*. 2020;5(2):1–6. doi:10.14218/ERHM.2020.00023.
7. Bannister-Tyrrell M, Meyer A, Faverjon C, Cameron A. Preliminary evidence that higher temperatures are associated with lower incidence of COVID-19, for cases reported globally up to 29th February 2020. *medRxiv*. 2020; p. 2020.03.18.20036731. doi:10.1101/2020.03.18.20036731.
8. Demongeot J, Flet-Berliac Y, Seligmann H. Temperature Decreases Spread Parameters of the New Covid-19 Case Dynamics. *Biology*. 2020;9(5). doi:10.3390/biology9050094.
9. Liu J, Zhou J, Yao J, Zhang X, Li L, Xu X, et al. Impact of meteorological factors on the COVID-19 transmission: A multi-city study in China. *Science of the Total Environment*. 2020;726:138513. doi:10.1016/j.scitotenv.2020.138513.
10. Greenland S, Robins JM, Pearl J. Confounding and Collapsibility in Causal Inference. *Statistical Science*. 1999;14(1):29–46.
11. Schipf S, Knüppel S, Hardt J, Stang A. Directed Acyclic Graphs (DAGs) – Die Anwendung kausaler Graphen in der Epidemiologie. *Gesundheitswesen*. 2011;73(12):888–892. doi:10.1055/s-0031-1291192.

12. Textor J, van der Zander B, Gilthorpe MS, Liškiewicz M, Ellison GT. Robust causal inference using directed acyclic graphs: the R package ‘dagitty’. *International Journal of Epidemiology*. 2017;45(6):1887–1894. doi:10.1093/ije/dyw341.
13. Center for Systems Science and Engineering (CSSE). COVID-19 Data Repository by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University; 2020. Available from: <https://github.com/CSSEGISandData/COVID-19> [cited 2020-07-12].
14. Pearl J, Bareinboim E. External Validity: From Do-Calculus to Transportability Across Populations. *Statistical Science*. 2014;29(4):579–595. doi:10.1214/14-sts486.
15. Chang MC, Kahn R, Li YA, Lee CS, Buckee CO, Chang HH. Modeling the impact of human mobility and travel restrictions on the potential spread of SARS-CoV-2 in Taiwan. *medRxiv*. 2020; p. 2020.04.07.20053439. doi:10.1101/2020.04.07.20053439.
16. Fowler JH, Hill SJ, Obradovich N, Levin R. The Effect of Stay-at-Home Orders on COVID-19 Cases and Fatalities in the United States. *medRxiv*. 2020;doi:10.1101/2020.04.13.20063628.
17. Kraemer MUG, Yang CH, Gutierrez B, Wu CH, Klein B, Pigott DM, et al. The effect of human mobility and control measures on the COVID-19 epidemic in China. *Science (New York, NY)*. 2020;368(6490):493–497. doi:10.1126/science.abb4218.
18. Lasry A, Kidder D, Hast M, Poovey J, Sunshine G, Winglee K, et al. Timing of Community Mitigation and Changes in Reported COVID-19 and Community Mobility - Four U.S. Metropolitan Areas, February 26–April 1, 2020. *MMWR Morbidity and mortality weekly report*. 2020;69(15):451–457. doi:10.15585/mmwr.mm6915e2.
19. Linka K, Peirlinck M, Sahli Costabal F, Kuhl E. Outbreak dynamics of COVID-19 in Europe and the effect of travel restrictions. *Computer Methods in Biomechanics and Biomedical Engineering*. 2020; p. 1–8. doi:10.1080/10255842.2020.1759560.
20. Mazzoli M, Mateo D, Hernando A, Meloni S, Ramasco JJ. Effects of mobility and multi-seeding on the propagation of the COVID-19 in Spain. *medRxiv*. 2020; p. 2020.05.09.20096339. doi:10.1101/2020.05.09.20096339.
21. Xiong C, Hu S, Yang M, Younes HN, Luo W, Ghader S, et al. Data-Driven Modeling Reveals the Impact of Stay-at-Home Orders on Human Mobility during the COVID-19 Pandemic in the U.S. *arXiv e-prints*. 2020; p. arXiv:2005.00667.
22. Jüni P, Rothenbühler M, Bobos P, Thorpe KE, da Costa BR, Fisman DN, et al. Impact of climate and public health interventions on the COVID-19 pandemic: A prospective cohort study. *Canadian Medical Association Journal*. 2020;doi:10.1503/cmaj.200920.
23. Lai S, Ruktanonchai NW, Zhou L, Prosper O, Luo W, Floyd JR, et al. Effect of non-pharmaceutical interventions to contain COVID-19 in China. *Nature*. 2020;doi:10.1038/s41586-020-2293-x.

24. Google LLC. Google Trends, search term “corona”; 2020. Available from: <https://www.google.com/trends> [cited 2020-07-12].
25. Ayyoubzadeh SM, Ayyoubzadeh SM, Zahedi H, Ahmadi M, R Niakan Kalhori S. Predicting COVID-19 Incidence Through Analysis of Google Trends Data in Iran: Data Mining and Deep Learning Pilot Study. *JMIR Public Health and Surveillance*. 2020;6(2):e18828. doi:10.2196/18828.
26. Effenberger M, Kronbichler A, Shin JI, Mayer G, Tilg H, Perco P. Association of the COVID-19 pandemic with Internet Search Volumes: A Google Trends(TM) Analysis. *International Journal of Infectious Diseases*. 2020;95:192–197. doi:10.1016/j.ijid.2020.04.033.
27. Lin YH, Liu CH, Chiu YC. Google searches for the keywords of “wash hands” predict the speed of national spread of COVID-19 outbreak among 21 countries. *Brain, Behavior, and Immunity*. 2020;doi:10.1016/j.bbi.2020.04.020.
28. Mavragani A. Tracking COVID-19 in Europe: Infodemiology Approach. *JMIR public health and surveillance*. 2020;6(2):e18941. doi:10.2196/18941.
29. Walker A, Hopkins C, Surda P. Use of Google Trends to investigate loss-of-smell-related searches during the COVID-19 outbreak. *International Forum of Allergy & Rhinology*. 2020;10(7):839–847. doi:10.1002/alr.22580.
30. Zhou WK, Wang AL, Xia F, Xiao YN, Tang SY. Effects of media reporting on mitigating spread of COVID-19 in the early phase of the outbreak. *Mathematical Biosciences and Engineering*. 2020;17(3):2693–2707. doi:10.3934/mbe.2020147.
31. Qi H, Xiao S, Shi R, Ward MP, Chen Y, Tu W, et al. COVID-19 transmission in Mainland China is associated with temperature and humidity: A time-series analysis. *Science of the Total Environment*. 2020;728:138778. doi:10.1016/j.scitotenv.2020.138778.
32. Shi P, Dong Y, Yan H, Zhao C, Li X, Liu W, et al. Impact of temperature on the dynamics of the COVID-19 outbreak in China. *Science of the Total Environment*. 2020;728:138890. doi:10.1016/j.scitotenv.2020.138890.
33. Sobral MFF, Duarte GB, da Penha Sobral AIG, Marinho MLM, de Souza Melo A. Association between climate variables and global transmission of SARS-CoV-2. *Science of the Total Environment*. 2020;729:138997. doi:10.1016/j.scitotenv.2020.138997.
34. Tosepu R, Gunawan J, Effendy DS, Ahmad LOAI, Lestari H, Bahar H, et al. Correlation between weather and Covid-19 pandemic in Jakarta, Indonesia. *Science of the Total Environment*. 2020;725:138436. doi:10.1016/j.scitotenv.2020.138436.
35. Wang M, Jiang A, Gong L, Luo L, Guo W, Li C, et al. Temperature significant change COVID-19 Transmission in 429 cities. *medRxiv*. 2020;doi:10.1101/2020.02.22.20025791.
36. Wu Y, Jing W, Liu J, Ma Q, Yuan J, Wang Y, et al. Effects of temperature and humidity on the daily new cases and new deaths of COVID-19 in 166 countries. *Science of the Total Environment*. 2020;729:139051. doi:https://doi.org/10.1016/j.scitotenv.2020.139051.

37. Auler AC, Cássaro FAM, da Silva VO, Pires LF. Evidence that high temperatures and intermediate relative humidity might favor the spread of COVID-19 in tropical climate: A case study for the most affected Brazilian cities. *Science of the Total Environment*. 2020;729:139090. doi:10.1016/j.scitotenv.2020.139090.
38. Xie J, Zhu Y. Association between ambient temperature and COVID-19 infection in 122 cities from China. *Science of the Total Environment*. 2020;724:138201. doi:10.1016/j.scitotenv.2020.138201.
39. Briz-Redón A, Serrano-Aroca A. A spatio-temporal analysis for exploring the effect of temperature on COVID-19 early evolution in Spain. *Science of the Total Environment*. 2020;728:138811. doi:10.1016/j.scitotenv.2020.138811.
40. Iqbal N, Fareed Z, Shahzad F, He X, Shahzad U, Lina M. The nexus between COVID-19, temperature and exchange rate in Wuhan city: New findings from partial and multiple wavelet coherence. *Science of the Total Environment*. 2020;729:138916. doi:10.1016/j.scitotenv.2020.138916.
41. Jahangiri M, Jahangiri M, Najafgholipour M. The sensitivity and specificity analyses of ambient temperature and population size on the transmission rate of the novel coronavirus (COVID-19) in different provinces of Iran. *Science of the Total Environment*. 2020;728:138872. doi:10.1016/j.scitotenv.2020.138872.
42. Yao Y, Pan J, Liu Z, Meng X, Wang W, Kan H, et al. No association of COVID-19 transmission with temperature or UV radiation in Chinese cities. *The European Respiratory Journal*. 2020;55(5). doi:10.1183/13993003.00517-2020.
43. de Lusignan S, Dorward J, Correa A, Jones N, Akinyemi O, Amirthalingam G, et al. Risk factors for SARS-CoV-2 among patients in the Oxford Royal College of General Practitioners Research and Surveillance Centre primary care network: a cross-sectional study. *The Lancet Infectious Diseases*. 2020;doi:10.1016/S1473-3099(20)30371-6.
44. Dragano N, Rupprecht CJ, Dortmann O, Scheider M, Wahrendorf M. Higher risk of COVID-19 hospitalization for unemployed: an analysis of 1,298,416 health insured individuals in Germany. *medRxiv*. 2020;doi:10.1101/2020.06.17.20133918.
45. Dohle S, Wingen T, Schreiber M. Acceptance and Adoption of Protective Measures During the COVID-19 Pandemic: The Role of Trust in Politics and Trust in Science; 2020. Available from: osf.io/w52nv.
46. Engle S, Stromme J, Zhou A. Staying at home: mobility effects of COVID-19. Available at SSRN. 2020;doi:10.2139/ssrn.3565703.
47. Cowling BJ, Ali ST, Ng TWY, Tsang TK, Li JCM, Fong MW, et al. Impact assessment of non-pharmaceutical interventions against coronavirus disease 2019 and influenza in Hong Kong: an observational study. *The Lancet Public Health*. 2020;5(5):e279–e288. doi:10.1016/S2468-2667(20)30090-6.
48. Robert Koch-Institut (RKI). Fallzahlen in Deutschland (COVID-19); 2020. Available from: https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Fallzahlen.html [cited 2020-07-12].
49. Google LLC. Google COVID-19 Community Mobility Reports; 2020. Available from: <https://www.google.com/covid19/mobility/> [cited 2020-07-12].

50. Deutscher Wetterdienst (DWD) Climate Data Center (CDC). Recent daily station observations (temperature, pressure, precipitation, sunshine duration, etc.) for Germany, quality control not completed yet, version recent; 2020. Available from: https://opendata.dwd.de/climate_environment/CDC/observations_germany/climate/daily/kl/recent/ [cited 2020-07-12].
51. Bundesinstitut für Bau-, Stadt- und Raumforschung (BBSR). INKAR – Indikatoren und Karten zur Raum- und Stadtentwicklung; 2020. Available from: <https://www.inkar.de/> [cited 2020-06-25].
52. Mitze T, Kosfeld R, Rode J, Wälde K. Face Masks Considerably Reduce COVID-19 Cases in Germany: A Synthetic Control Method Approach. Institute of Labor Economics (IZA); 2020. 13319. Available from: <https://EconPapers.repec.org/RePEc:iza:izadps:dp13319>.
53. Spirtes P, Glymour CN, Scheines R, Heckerman D. Causation, prediction, and search. MIT Press; 2000.
54. Pearl J. Causality. Cambridge: Cambridge University Press; 2009. Available from: <https://www.cambridge.org/core/books/causality/B0046844FAE10CBF274D4ACBDAEB5F5B>.
55. Greenland S, Pearl J, Robins JM. Causal Diagrams for Epidemiologic Research. *Epidemiology*. 1999;10(1):37–48.
56. Henckel L, Perković E, Maathuis MH. Graphical Criteria for Efficient Total Effect Estimation via Adjustment in Causal Linear Models. *arXiv e-prints*. 2019; p. arXiv:1907.02435.
57. R Core Team. R: A Language and Environment for Statistical Computing; 2019. Available from: <https://www.R-project.org/>.
58. Kalisch M, Mächler M, Colombo D, Maathuis MH, Bühlmann P. Causal Inference Using Graphical Models with the R Package pcalg. *Journal of Statistical Software*. 2012;47(11):1–26. doi:10.18637/jss.v047.i11.
59. Hilbe JM, Greene WH. 4 - Count Response Regression Models. In: Rao CR, Miller JP, Rao DC, editors. *Essential Statistical Methods for Medical Statistics*. Boston: North-Holland; 2011. p. 104–145. Available from: <http://www.sciencedirect.com/science/article/pii/B9780444537379500074>.
60. [Venables WN, Ripley BD. Modern Applied Statistics with S. 4th ed. New York: Springer; 2002. Available from: http://www.stats.ox.ac.uk/pub/MASS4.](#)
61. [Hoerl AE, Kennard RW. Ridge Regression: Biased Estimation for Nonorthogonal Problems. Technometrics. 1970;12\(1\):55–67. doi:10.1080/00401706.1970.10488634.](#)
62. [Friedman J, Hastie T, Tibshirani R. Regularization Paths for Generalized Linear Models via Coordinate Descent. Journal of Statistical Software. 2010;33\(1\):1–22.](#)
63. Perković E, Textor J, Kalisch M, Maathuis MH. A Complete Generalized Adjustment Criterion. *arXiv e-prints*. 2015; p. arXiv:1507.01524.
~~Venables WN, Ripley BD. Modern Applied Statistics with S. 4th ed. New York: Springer; 2002. Available from: —~~

64. Gencoglu O, Gruber M. Causal Modeling of Twitter Activity During COVID-19. medRxiv. 2020;doi:10.1101/2020.05.16.20103903.
65. Perković E, Textor J, Kalisch M, Maathuis MH. Complete graphical characterization and construction of adjustment sets in Markov equivalence classes of ancestral graphs. *The Journal of Machine Learning Research*. 2017;18(1):8132–8193.
66. Islam N, Sharp SJ, Chowell G, Shabnam S, Kawachi I, Lacey B, et al. Physical distancing interventions and incidence of coronavirus disease 2019: natural experiment in 149 countries. *BMJ*. 2020;370. doi:10.1136/bmj.m2743.
67. Kermack WO, McKendrick AG. Contributions to the mathematical theory of epidemics–I. 1927. *Bulletin of mathematical biology*. 1991;53(1-2):33–55. doi:10.1007/bf02464423.
68. an der Heiden M, Buchholz U. Modellierung von Beispielszenarien der SARS-CoV-2-Epidemie 2020 in Deutschland. DOI 10.25646/6571.2; 2020. Available from: https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Modellierung_Deutschland.html.
69. [Schisterman EF, Perkins NJ, Mumford SL, Ahrens KA, Mitchell EM. Collinearity and Causal Diagrams: A Lesson on the Importance of Model Specification. *Epidemiology*. 2017;28\(1\).](#)
70. Openshaw S. Ecological Fallacies and the Analysis of Areal Census Data. *Environment and Planning A: Economy and Space*. 1984;16(1):17–31. doi:10.1068/a160017.
71. World Health Organization, et al. Transmission of SARS-CoV-2: implications for infection prevention precautions: Scientific Brief, 09 July 2020. World Health Organization; 2020.