

PARTIAL DIFFERENTIAL EQUATIONS

International Mathematics Masters
National Higher School of Mathematics (NHSM)



INTERNATIONAL
MATHEMATICS MASTER



NATIONAL HIGHER SCHOOL
OF MATHEMATICS
المدرسة الوطنية
العليا في الرياضيات

CONTENTS

THEORY OF (UNBOUNDED) OPERATORS

1.1 Preliminaries on Operators

Let $(X, \|\cdot\|)$ be a Banach space and $\mathcal{L}(X)$ be the Banach space of bounded linear operators.

Definition 1.1.1. An operator $\mathcal{A} : D(\mathcal{A}) \longrightarrow X$ is called an unbounded linear operator (UBLO) if $D(\mathcal{A})$ is a subspace of X and $\sup_{x \neq 0} \frac{\|\mathcal{A}x\|}{\|x\|} = +\infty$

Exercise 1: L

Let $\mathcal{A} : H^1 \longrightarrow L^2$, such that $f \mapsto f'$ and $D(\mathcal{A}) = \{f : \mathbb{R} \rightarrow \mathbb{R} : f, f' \in L^2\}$. Show \mathcal{A} is an unbounded linear operator.

Notation: If \mathcal{A} and \mathcal{B} are unbounded linear operators, then $\mathcal{A} \supset \mathcal{B}$ if and only if $D(\mathcal{A}) \supset D(\mathcal{B})$. That is, for all $x \in D(\mathcal{B})$, $\mathcal{A}x = \mathcal{B}x$.

1.1.1 Resolvent Operator

Definition 1.1.2. Let $\mathcal{A} : D(\mathcal{A}) \longrightarrow X$ be a UBLO.

$$\rho(\mathcal{A}) = \text{Resolvent of } \mathcal{A} = \left\{ \lambda \in \mathbb{C} \mid \begin{array}{l} (\lambda I - \mathcal{A}) : D(\mathcal{A}) \rightarrow X \text{ is bijective, and} \\ (\lambda I - \mathcal{A})^{-1} \in \mathcal{L}(X) \end{array} \right\}$$

$$\sigma(\mathcal{A}) = \text{Spectrum of } \mathcal{A} = \mathbb{C} \setminus \rho(\mathcal{A}).$$

Definition 1.1.3. \mathcal{A} is closed if and only if the graph of \mathcal{A} , denoted as $G(\mathcal{A})$ is closed. Also \mathcal{A} is closable if and only if there exists $\tilde{\mathcal{A}} \supset \mathcal{A}$ such that $G(\tilde{\mathcal{A}}) = \overline{G(\mathcal{A})}$.

Note that $G(\mathcal{A}) = \{(x, \mathcal{A}x) \mid x \in D(\mathcal{A})\}$.

Exercise 2: 3

1. Prove that if it exists, $\tilde{\mathcal{A}}$ is unique, it then denoted by $\overline{\mathcal{A}}$ called closure of \mathcal{A} .
2. Let $\mathcal{A}_\ell = \frac{d}{dx}$ with $(X = C^0([a, b], \mathbb{R}), \|\cdot\| = \sup |f(x)|)$ and $D(\mathcal{A}_\ell) = C^\ell([a, b], \mathbb{R})$. Prove $\overline{\mathcal{A}_\ell} = \mathcal{A}_1$.

Lemma 1.1.1. If \mathcal{A} an unbounded linear operator is closable, then $\rho(\overline{\mathcal{A}}) = \rho(\mathcal{A})$. If \mathcal{A} is closed then $\rho(\mathcal{A}) = \{\lambda \in \mathbb{C} \mid (\lambda I - \mathcal{A})^{-1} \in \mathcal{L}(X)\}$.

Hints (Exercise): If $\rho(\mathcal{A}) \neq 0$ implies \mathcal{A} is closed. (Show first that if T is a UBLO with $T^{-1} \in \mathcal{L}(X)$ implies T is closed).

Corollary 1.1.1. Let $\mathcal{A} : D(\mathcal{A}) \subset X \longrightarrow X$ be a closed UBLO then $\sigma(\mathcal{A}) = \sigma_p(\mathcal{A}) \cup \sigma_r(\mathcal{A}) \cup \sigma_c(\mathcal{A})$ where

1. $\sigma_p(\mathcal{A}) = \{\lambda \in \mathbb{C} : \ker(\lambda I - \mathcal{A}) \neq \{0\}\}$ (punctual spectrum and λ 's are the eigenvalue).
2. $\sigma_c(\mathcal{A}) = \left\{ \lambda \in \mathbb{C} : \ker(\lambda I - \mathcal{A}) = \{0\}, \overline{\text{Rg}(\lambda I - \mathcal{A})} \subset X \right\}$ (continuous spectrum).
3. $\sigma_r(\mathcal{A}) = \left\{ \lambda \in \mathbb{C} : \lambda I - \mathcal{A} \text{ is injective, } \overline{\text{Rg}(\lambda I - \mathcal{A})} = X, \text{Rg}(\lambda I - \mathcal{A}) \neq X \right\}$ (residual spectrum).

Exercise 3: L

Let

$$X = \ell^2(\mathbb{C}) = \left\{ (x_n)_{n \geq 0} : \sum_n |x_n|^2 < \infty \right\},$$

with $(\mathcal{A} x_n)_{n \geq 1} = \left(\frac{x_n}{1+n} \right)_{n \geq 0}$. Prove that \mathcal{A} is a BLO, injective, $\overline{\text{Rg}(\mathcal{A})} = X$ and $\text{Rg}(\mathcal{A}) \subset X$.

Theorem 1.1.1. If \mathcal{A} is a closed UBLO then $\rho(\mathcal{A})$ is open. If $\mu \in \rho(\mathcal{A})$, then for all $\lambda \in \mathbb{C}$ with $r := |\mu - \lambda|$, $\|(\mu I - \mathcal{A})^{-1}\| < 1$ then $\lambda \in \rho(\mathcal{A})$ and

$$(\lambda I - \mathcal{A})^{-1} = \sum_{n=0}^{\infty} (\mu - \lambda)^n (\mu I - \mathcal{A})^{-(n+1)}$$

To do Question: do you need \mathcal{A} closed?

Theorem 1.1.2 (Resolvent Identity). Let \mathcal{A} be a UBLO. For $\lambda \in \rho(A)$, define the resolvent operator

$$R(\lambda) := (\lambda I - A)^{-1}.$$

Then for all $\lambda, \mu \in \rho(A)$,

$$R(\lambda) - R(\mu) = (\mu - \lambda) R(\mu) R(\lambda).$$

Corollary 1.1.2. The mapping $\lambda \mapsto R(\lambda)$ from $\rho(\mathcal{A})$ into $\mathcal{L}(X)$ is analytic. Moreover,

$$\frac{d^n}{d\lambda^n} (\lambda I - A)^{-1} = (-1)^n n! [(\lambda I - A)^{-1}]^{(n+1)}.$$

1.1.2 Dual Operators

Let $X \cong X^*$ and \mathcal{A} a closed UBLO with $\overline{D(\mathcal{A})} = X$ a dense UBLO.

If X and Y are Banach spaces with duals X^* and Y^* , then for $x \in X$ and $x^* \in X^*$, we define the duality product as $\langle x^*, x \rangle$.

Definition 1.1.4 (Dual Operator of \mathcal{A}). *Let $\mathcal{A} : D(\mathcal{A}) \subset X \rightarrow Y$ (UBLO), be such that $\overline{D(\mathcal{A})} = X$. The dual operator $\mathcal{A}^* : D(\mathcal{A}^*) \subset Y^* \rightarrow X^*$ is a UBLO defined as follows:*

$$D(\mathcal{A}^*) := \{y^* \in Y^* \mid \exists z^* \in X^* \text{ such that } \langle y^*, Ax \rangle = \langle z^*, x^* \rangle \forall x \in D(\mathcal{A})\}.$$

and $y^* \in D(\mathcal{A}^*)$, the element z^* is unique and we define $\mathcal{A}^* y^* := z^*$.

Lemma 1.1.2. *Let X, Y be Banach spaces and let $\mathcal{A} \in \mathcal{L}(X, Y)$. Then $\mathcal{A}^* \in \mathcal{L}(Y^*, X^*)$ and*

$$\|\mathcal{A}^*\|_{\mathcal{L}(Y^*, X^*)} = \|\mathcal{A}\|_{\mathcal{L}(X, Y)}.$$

Lemma 1.1.3. *Let X be a reflexive Banach space with $X = X^*$ and let $\mathcal{A} : D(\mathcal{A}) \subset X \rightarrow X$ be a closedly dense UBLO. Then $\overline{D(\mathcal{A}^*)} = X^*(\cong X)$, and \mathcal{A}^* is closed.*

Theorem 1.1.3. *Let \mathcal{A} be a closedly dense UBLO. Then $\rho(\mathcal{A}) = \rho(\mathcal{A}^*)$ and for all $\lambda \in \rho(A)$,*

$$[(\lambda I - \mathcal{A})^{-1}]^* = (\lambda I - \mathcal{A}^*).$$

Exercise 4: I

1. Let $\mathcal{A} = \frac{d}{dx}$ on $X = L^2(\mathbb{R})$ and $D(\mathcal{A}) = \{f \in X : f' \in L^2(\mathbb{R})\}$. Show the following:

- $\rho(\mathcal{A}) = \mathbb{C} \setminus i\mathbb{R}$ which implies $\sigma(\mathcal{A}) = i\mathbb{R}$.
- \mathcal{A} is a closed unbounded linear operator.
- If $\lambda \in \rho(\mathcal{A})$ then $(\lambda I - \mathcal{A})^{-1} : X \rightarrow D(\mathcal{A})$ is bounded.

For $\Re(\lambda) \neq 0$; show for all $g \in X$, there exists uniquely $f \in D(\mathcal{A})$ such that $(\lambda I - \mathcal{A})f = g$.

For $\Re(\lambda) = 0$; show for all $f_n \in X$ with $\|f_n\|_{\ell^2} = 1$ then $(i\omega I - \mathcal{A})f_n \rightarrow 0$.

2. Do same for $\mathcal{A} = -i \frac{d}{dx}$.