

Farmhannong Agro Weekly - 출처 확인용 PDF

이 파일은 원문 기사 전체를 복제한 보관본이 아니라, 링크 차단/삭제 시에도 출처 확인을 돕기 위한 색인 PDF입니다.

카드 발행일	2026 - 08 - 24
카드 번호	11
출처	UMN Crop News
원문 제목	Preharvest weed control: Scout fields and field edges now to prevent weed seed production
원 URL	https://blog-crop-news.extension.umn.edu/2026/08/preharvest-weed-control-scout-fields.html
제목 검색 URL	https://www.google.com/search?q=%22Preharvest+weed+control%3A+Scout+fields+and+field+edges+now+to+prevent+weed+seed+production%22+UMN+Crop+News
검색 키워드	Minnesota Extension preharvest weed control scouting waterhemp Palmer amaranth kochia lambsquarters mowing 64 to 100% viable seed reduction August 19 2026
PDF 생성일	2026 - 08 - 24 08:54

카드 요약 / 당시 확인 내용

University of Minnesota Extension educators Eric Yu and Dave Nicolai advise scouting fields and field edges before harvest to prevent weed seed production. Key weeds include waterhemp and pigweed species, giant and common ragweed, kochia, common lambsquarters, annual grasses and Palmer amaranth, a noxious weed requiring early detection. Waterhemp is capable of producing hundreds of thousands of seeds per plant and can produce viable seed within 10 days of pollination, while lambsquarters can produce up to 30,000 seeds per plant. Under late summer conditions pigweed species can flower within three weeks with seed maturity three weeks later, and redroot pigweeds continue flowering until frost. Research found mowing before seed maturity reduced viable seed production by 64 to 100%, although results varied among species and timing. Growers should identify weed patches, harvest clean areas first and weedy areas last, and clean equipment between heavily infested and clean fields.