

Farmhannong Agro Weekly - 출처 확인용 PDF

이 파일은 원문 기사 전체를 복제한 보관본이 아니라, 링크 차단/삭제 시에도 출처 확인을 돕기 위한 색인 PDF입니다.

카드 발행일	2026 - 08 - 24
카드 번호	12
출처	UMN Crop News
원문 제목	Late - season soybean aphid scouting and management in Minnesota
원 URL	https://blog-crop-news.extension.umn.edu/2026/08/late-season-soybean-aphid-scouting-and.html
제목 검색 URL	https://www.google.com/search?q=%22Late-season+soybean+aphid+scouting+and+management+in+Minnesota%22+UMN+Crop+News
검색 키워드	Minnesota soybean aphid economic threshold 250 aphids 80% infested pyrethroid resistance sulfoxaflor afidopyropen flupyradifurone spider mites bifenthrin PHI August 2026
PDF 생성일	2026 - 08 - 24 08:54

카드 요약 / 당시 확인 내용

University of Minnesota Extension soybean entomologist Robert Koch and IPM educator Eric Yu advise that through the R5 growth stage growers should treat when fields average 250 aphids per plant with more than 80% of plants infested and populations actively increasing. No formal threshold exists for R6, though extremely high populations may still cause yield loss. Monitor live adults, nymphs and white dwarfs, excluding cast skins, parasitized mummies and diseased aphids. Pyrethroid resistance is widespread in Minnesota soybean aphid populations, so avoid relying solely on pyrethroid - only products and consider selective alternatives such as sulfoxaflor, afidopyropen or flupyradifurone. Hot dry conditions favor two - spotted spider mites and most pyrethroids except bifenthrin can trigger spider mite outbreaks. Scout through early R6 based on growth stage, not calendar dates. Preharvest intervals typically range from 18 - 30 days.