

ZITLAS

AI Weight Loss Knowledge Base

VERSION 2.0 — RAG-OPTIMIZED PRODUCTION EDITION

13 Modules · 350+ Foods · 145+ Recipes · 600+ FAQs · 86+ Coaching Rules

Modules: Fundamentals · Nutrition Science · Indian Food Database · Recipe Library
Meal Plans · Exercise Database · Workout Programs · Special Populations
Behaviour Change · Plateaus · Tracking · Decision Engine · Master FAQ

Designed for RAG ingestion · Vector DB optimized · AI coaching deployment

Confidential — ZITLAS AI Sports Coaching Intelligence System

WEIGHT LOSS FUNDAMENTALS

WL-F-001	MOD 01 · Energy Balance › Calories In vs Calories Out
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Beginner
TAGS	energy balance, calories, CICO, caloric intake, caloric expenditure, weight loss basics

DEFINITION

Energy balance is the relationship between calories consumed (food and drink) and calories expended (metabolism + physical activity). When intake equals expenditure, weight is stable. When intake exceeds expenditure, weight is gained. When expenditure exceeds intake, weight is lost.

WHY IT MATTERS

- Every weight management outcome — loss, gain, or maintenance — is determined by energy balance.
- No diet, supplement, or exercise program works without first establishing a caloric deficit for weight loss.
- Understanding energy balance prevents wasted effort chasing 'magic' foods or workouts.
- CICO (Calories In, Calories Out) is the most evidence-based framework for weight management.
- Hormones, sleep, and stress all affect energy balance — they do not override it.

KEY FACTS

- 1 kg of body fat $\approx$ 7,700 kcal. A 500 kcal/day deficit = $\sim$ 0.5 kg fat lost/week.
- 1 kg of body fat $\approx$ 3,500 kcal in stored energy (some estimates range 3,400–3,700).
- Calories in = food + beverages. Calories out = BMR + NEAT + EAT + TEF.
- TEF (Thermic Effect of Food): protein 20–30%, carbs 5–10%, fat 0–3% of calories consumed.
- Average Indian adult TDEE: female 1,600–2,000 kcal/day; male 2,000–2,600 kcal/day.
- Tracking calories consistently increases weight loss success rate by $\sim$ 40% (research data).
- Liquid calories (juices, chai with sugar, cold drinks) are often invisible to users — high impact.

ACTION STEPS

1. Calculate your TDEE using the Mifflin-St Jeor formula (see WL-F-004).
2. Set a deficit of 300–500 kcal/day for steady, sustainable fat loss.
3. Track all food and drink using a calorie tracking app (MyFitnessPal, HealthifyMe) for minimum 1 week.
4. Identify your highest-calorie meals and drinks — target those first.
5. Weigh or measure food for the first 2–4 weeks to build accurate portion awareness.
6. Review energy balance weekly; adjust if weight is not changing as expected.

DECISION TREE

IF user is losing weight too fast (>1 kg/week) $\rightarrow$ **THEN** increase calories by 200–300 kcal/day; check for muscle loss

IF user is not losing weight after 2 weeks of tracking $\rightarrow$ **THEN** review tracking accuracy; check for hidden calories (oils, sauces, drinks)

IF user says 'I eat very little but don't lose weight' $\rightarrow$ **THEN** request 3-day food diary; check NEAT; consider thyroid screening if other symptoms

IF user asks about cheat meals $\rightarrow$ **THEN** explain that one high-calorie meal raises weekly average minimally; focus on overall weekly balance

IF user wants faster results $\rightarrow$ **THEN** cap deficit at 750 kcal/day max; add 150–300 kcal from exercise rather than further restriction

COMMON MISTAKES

- Underestimating cooking oil — 1 tbsp oil = 120 kcal. Indian cooking can use 4–6 tbsp/day hidden in dishes.
- Not counting liquid calories: chai with sugar, lassi, fruit juice, cold drinks can add 400–700 kcal/day.
- Eating 'healthy' foods in large quantities (ghee, dry fruits, nuts, avocado) and still being in surplus.
- Assuming exercise burns more than it does — 30 min brisk walk burns only $\sim$ 150–180 kcal.
- Skipping breakfast to cut calories, then overeating at dinner and erasing the deficit.

SPECIAL CASES

- PCOS: Hormonal disruption can make women feel hungrier and store fat more easily, but energy balance still applies. Reduce deficit to 250–350 kcal/day.
- Hypothyroidism: Untreated hypothyroidism lowers BMR by 200–600 kcal/day. Treat medically first; then apply deficit.
- Seniors (60+): Lower TDEE due to reduced lean mass. Deficit of 300 kcal/day is safer; prioritize protein.
- Teens (13–18): Do NOT create large deficits during growth. Focus on food quality over restriction.

FAQ

Q: Do I need to count calories to lose weight?

A: Not necessarily, but understanding energy balance is essential. Portion control, eliminating ultra-processed foods, and eating more whole foods can create a deficit without counting. Counting increases precision and is recommended if progress stalls.

Q: I exercise daily but don't lose weight — why?

A: Exercise typically burns 200–400 kcal/session. If you compensate by eating more or reward yourself with high-calorie foods after exercise, the deficit is erased. Track both sides of the equation.

Q: Is a calorie from roti the same as a calorie from ghee?

A: Thermodynamically yes; metabolically no. Ghee has no fiber or protein, hits 9 kcal/g, minimal satiety. Roti at similar calories gives fiber, protein, and lasts longer. Calorie source matters for hunger management and nutrition.

AI COACHING NOTES

- When user blames metabolism: validate concern, then explain metabolic differences are rarely more than 200 kcal/day between people of similar size.
- When user feels confused: simplify to 'eat a bit less + move a bit more' before introducing tracking.
- Always anchor deficit recommendations to 300–500 kcal/day — never recommend <1,000 kcal/day total intake.
- For hostel/student users: oil tracking is especially important since mess food and tiffin services use high amounts of oil.

WL-F-002	MOD 01 · BMR · Basal Metabolic Rate
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Beginner
TAGS	BMR, basal metabolic rate, resting metabolism, calorie needs, Mifflin-St Jeor

DEFINITION

BMR (Basal Metabolic Rate) is the number of calories your body burns at complete rest to maintain vital functions: breathing, circulation, organ function, and cell repair. It represents 60–75% of total daily energy expenditure.

WHY IT MATTERS

- BMR is the foundation of all calorie calculations — it sets the minimum daily calorie floor.
- Restricting calories below BMR causes muscle loss, nutrient deficiency, and metabolic adaptation.
- BMR declines with age (~2–3% per decade after 30) and with excessive dieting.
- Knowing BMR helps set a safe, effective calorie target.

KEY FACTS

- BMR for Indian adult female (60 kg, 163 cm, 30 yr): ≈ 1,390 kcal/day.
- BMR for Indian adult male (70 kg, 170 cm, 30 yr): ≈ 1,660 kcal/day.
- Mifflin-St Jeor (most accurate): Male: $(10 \times \text{kg}) + (6.25 \times \text{cm}) - (5 \times \text{age}) + 5$. Female: $(10 \times \text{kg}) + (6.25 \times \text{cm}) - (5 \times \text{age}) - 161$.
- Every 1 kg of muscle burns ~13 kcal/day at rest; every 1 kg of fat burns ~4.5 kcal/day.
- Severe caloric restriction (≤ 800 kcal/day) can reduce BMR by up to 25% within weeks.
- BMR cannot be accurately measured by consumer apps — formulas give an estimate $\pm 10\%$.

ACTION STEPS

1. Calculate BMR using Mifflin-St Jeor formula with current weight, height, age, gender.
2. Never eat below BMR consistently — minimum intake for females 1,000–1,200 kcal/day; males 1,200–1,500 kcal/day.
3. To increase BMR long-term: build lean muscle through resistance training.
4. Re-calculate BMR every 5 kg of weight change to adjust targets.

DECISION TREE

IF user wants to eat fewer than 1,000 kcal/day → **THEN REFUSE**; explain risk of muscle loss and metabolic adaptation; minimum 1,200 kcal for females, 1,500 for males

IF user has lost significant muscle mass (weakness, fatigue) → **THEN** increase protein to 1.8–2.2 g/kg; add resistance training; recalculate BMR at new weight

IF user reports feeling cold, hair loss, fatigue on low-calorie diet → **THEN** suggest medical check (thyroid, ferritin); advise calorie increase to at least BMR + 200

COMMON MISTAKES

- Using the Harris-Benedict formula — Mifflin-St Jeor is more accurate for modern populations.
- Using online calculators that don't adjust for Indian body compositions (which tend to carry more visceral fat at lower BMI).
- Setting intake = BMR — BMR excludes all movement; you'll under-fuel basic daily activities.

FAQ

Q: How do I know if my metabolism is slow?

A: Most 'slow metabolism' feelings are from inaccurate calorie tracking. True metabolic disorders (hypothyroidism, Cushing's) are medical conditions. If BMR calculations are consistent but weight isn't moving with accurate tracking, get a thyroid panel (TSH, T3, T4).

Q: Can I increase my BMR?

A: Yes: build lean muscle (most effective), eat adequate protein, avoid prolonged very low calorie diets, stay active throughout the day (NEAT), sleep 7–9 hours.

AI COACHING NOTES

- Calculate BMR for every new user on onboarding and store it as their calorie floor.
- If user's reported intake is below their calculated BMR, prioritize investigating tracking accuracy before other changes.
- For female users post-40: BMR may have dropped 150–250 kcal vs their 30s. Adjust targets accordingly.

WL-F-003	MOD 01 · TDEE · Total Daily Energy Expenditure
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Beginner
TAGS	TDEE, total daily energy expenditure, activity multiplier, calorie target, maintenance calories

DEFINITION

TDEE is the total calories you burn in a day, including BMR plus all physical activity. It represents your maintenance calories — the amount needed to keep your current weight stable. Eating below TDEE causes weight loss; above TDEE causes weight gain.

WHY IT MATTERS

- TDEE is the single most important number for weight management.
- Setting your calorie target without TDEE means guessing — TDEE anchors your plan.
- TDEE varies by 500–1,500 kcal between a sedentary and highly active person of the same size.

KEY FACTS

- TDEE = BMR × Activity Multiplier. Sedentary: ×1.2. Light: ×1.375. Moderate: ×1.55. Active: ×1.725. Very Active: ×1.9.
- Sedentary = desk job, <5,000 steps/day, no exercise. Light = light exercise 1–3×/week. Moderate = exercise 3–5×/week.
- Indian sedentary adult female (60 kg): TDEE ≈ 1,668 kcal. Male (70 kg): TDEE ≈ 1,992 kcal.
- Most adults overestimate their activity level — defaulting to 'Sedentary' is usually accurate for desk workers.
- For weight loss: eat at TDEE – 300 to –500 kcal. For maintenance: eat at TDEE.

ACTION STEPS

1. Calculate TDEE = BMR × activity multiplier.
2. Subtract 300–500 kcal from TDEE for your daily calorie target.
3. Use a fitness tracker (steps, heart rate) to verify activity level assumption.
4. After 4 weeks, compare expected vs actual weight loss; adjust TDEE estimate if needed.
5. Re-calculate every 5 kg weight change or major lifestyle shift.

DECISION TREE

IF user has sedentary job but goes to gym 4×/week → **THEN** use 'Light Active' multiplier (1.375–1.465), not 'Very Active'

IF user's actual weight loss rate is lower than expected at calculated TDEE → **THEN** reduce TDEE estimate by 10%; increase movement or reduce intake by 100–150 kcal

IF user's weight is dropping faster than expected → **THEN** raise TDEE estimate; user may be more active than reported OR intake is lower than logged

COMMON MISTAKES

- Claiming 'Very Active' multiplier because they exercise once daily but sit 10 hours otherwise.
- Not updating TDEE after significant weight loss — TDEE drops with body weight.
- Using calorie counts from fitness trackers as exact — trackers overestimate burn by 20–40%.

FAQ

Q: Should I eat back exercise calories?

A: If you used a sedentary multiplier for TDEE, yes — add 50–70% of exercise calories burned. If you already used an active multiplier, no. Avoid eating back 100% of exercise calories as trackers overestimate.

Q: My TDEE says 2,000 kcal but I feel fine on 1,400. Is that OK?

A: Short-term yes; long-term no. Chronic eating well below TDEE causes metabolic adaptation, muscle loss, micronutrient deficiencies, and hormonal disruption. Increase to at least TDEE – 500.

AI COACHING NOTES

- *For new users: always start by calculating TDEE and asking 3 clarifying questions about daily movement before setting targets.*
- *Indian women often underestimate house chores and walking — factor these into activity level.*
- *Hostel students often overestimate activity — cafeteria + room + class = less than they think.*

WL-F-004	MOD 01 · NEAT › Non-Exercise Activity Thermogenesis
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Beginner
TAGS	NEAT, non-exercise activity, daily movement, steps, fidgeting, lifestyle activity

DEFINITION

NEAT is the energy burned by all physical activity that is NOT formal exercise: walking, standing, fidgeting, housework, cooking, climbing stairs, shopping. NEAT varies by 300–2,000 kcal/day between individuals and is highly modifiable.

WHY IT MATTERS

- NEAT is often more impactful than gym sessions for total daily calorie burn.
- Obese individuals tend to sit 150+ more minutes/day than lean individuals — the 'sitting disease'.
- NEAT is the most flexible component of TDEE and can be increased without exercise.
- Low NEAT is a major cause of weight loss plateaus in dieters.

KEY FACTS

- Standing vs sitting: +50 kcal/hour. Pacing while on calls: +100–150 kcal/hour.
- 10,000 steps/day burns approximately 300–500 kcal depending on body weight and walking speed.
- A person with high NEAT can have TDEE 700–1,000 kcal higher than a sedentary person of same weight.
- NEAT decreases during caloric restriction — the body reduces fidgeting and spontaneous movement unconsciously.
- Indians who shift from manual jobs to desk jobs often see NEAT drop by 400–800 kcal/day.

ACTION STEPS

1. Track daily steps — target 8,000–10,000 steps/day minimum.
2. Stand or walk during phone calls and meetings.
3. Take stairs instead of lifts whenever possible.
4. Walk 10–15 minutes after each main meal (improves blood sugar and burns 50–80 kcal).
5. Implement 'movement snacks': 5-minute walks every 60–90 minutes of sitting.

DECISION TREE

IF user has desk job + sedentary evening → **THEN** recommend 8,000 step goal, after-meal walks; this alone adds 200–350 kcal burn/day

IF user's weight loss has stalled on same diet → **THEN** check step count before changing diet; increase steps by 2,000–3,000/day first

IF user says they don't have time to exercise → **THEN** shift focus to NEAT: 3× 10-min walks adds 200+ kcal/day without gym time

COMMON MISTAKES

- Thinking only formal exercise counts — NEAT often contributes more total calorie burn.
- Becoming sedentary after evening workouts, negating the exercise benefit.
- Hostel students sitting for 8+ hours studying with no movement breaks.

FAQ

Q: Do household chores count as NEAT?

A: Yes. Cooking (90–120 kcal/hour), sweeping (150–200 kcal/hour), washing clothes by hand (200–250 kcal/hour). Indian homemakers often have significant NEAT that should be factored into TDEE.

Q: How do I increase NEAT without changing my schedule?

A: Park farther away, walk during phone calls, take stairs, walk to the next bus stop, stand while watching TV/reels for part of the evening.

AI COACHING NOTES

- Ask users about their step count before assuming exercise is needed — NEAT is often the bigger lever.
- For hostel/student users: NEAT can be dramatically low during exam season. Flag this.
- Post-meal walks are the single highest-ROI NEAT recommendation — reduces glucose spikes AND burns calories.

WL-F-005	MOD 01 · Metabolism · Metabolic Rate Factors
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Intermediate
TAGS	metabolism, metabolic rate, slow metabolism, fast metabolism, metabolic adaptation

DEFINITION

Metabolism is the sum of all chemical processes in the body that convert food to energy. Metabolic rate is how quickly these processes occur. Key factors: lean muscle mass, age, genetics, thyroid function, sleep, and caloric history.

WHY IT MATTERS

- Metabolism determines how many calories you burn daily without deliberate exercise.
- Metabolic adaptation is a key cause of weight loss plateaus — the body fights back.
- Building muscle permanently raises metabolic rate at rest.
- Crash dieting suppresses metabolism — leading to weight regain on even moderate calories.

KEY FACTS

- True metabolic difference between individuals of similar size and age: 100–300 kcal/day (much smaller than people assume).
- Metabolic adaptation: when dieting, TDEE decreases by 10–25% beyond what weight loss alone predicts.
- Building 5 kg of muscle raises BMR by ~65 kcal/day permanently.
- Thyroid dysfunction can reduce BMR by 200–600 kcal/day — get tested if symptoms present.
- Sleep deprivation (5 vs 8 hours) reduces metabolism by ~80 kcal/day AND increases hunger hormones.
- Eating adequate protein preserves lean mass during dieting, protecting metabolic rate.
- Indian women post-menopause: BMR declines 50–200 kcal/day due to hormonal shifts.

ACTION STEPS

1. Do NOT blame metabolism without ruling out tracking errors first.
2. Eat adequate protein (1.6–2.0 g/kg body weight) to protect muscle mass.
3. Include resistance training 2–3×/week to build/maintain lean mass.
4. Sleep 7–9 hours consistently — sleep debt is a metabolic brake.
5. Take 'diet breaks' every 8–12 weeks (2 weeks at maintenance) to restore metabolic rate.
6. If suspected thyroid issues: get TSH, Free T3, Free T4 tested.

DECISION TREE

IF user claims metabolism is broken — eating <1,000 kcal and not losing → **THEN** request photo food diary; check thyroid history; likely tracking error OR metabolic adaptation from chronic restriction

IF user has dieted for years with diminishing results → **THEN** suggest 4–6 week maintenance phase (reverse dieting); add resistance training; reset hormones

IF user has cold hands, hair loss, fatigue, constipation → **THEN** flag possible hypothyroidism; recommend medical check before continuing weight loss plan

COMMON MISTAKES

- Eating too little for too long — triggers metabolic adaptation that outlasts the diet.
- Avoiding carbs entirely — carbs provide glucose for thyroid hormone conversion (T4 → T3).
- Skipping sleep to exercise more — counterproductive to metabolism.

FAQ

Q: Can I 'boost' my metabolism with foods like green tea or chili?

A: These foods increase metabolism by 2–5% temporarily, equating to 40–100 kcal/day — useful but not transformative. The most powerful metabolic boosters are muscle mass and adequate sleep.

Q: I've heard yo-yo dieting damages metabolism permanently — is this true?

A: Research shows it is largely reversible with proper nutrition and resistance training. However, each cycle of extreme dieting makes regain faster and loss harder. Sustainable deficits prevent this cycle.

AI COACHING NOTES

- *Normalize metabolic adaptation to users who hit plateaus — it's biology, not failure.*
- *Diet breaks (2 weeks at maintenance every 8–12 weeks) are the evidence-based solution to metabolic adaptation.*
- *Prioritize resistance training in coaching plans — it's the highest-ROI metabolic intervention.*

WL-F-006	MOD 01 · Body Composition › Fat vs Lean Mass
GOALS	Weight Loss Fat Loss
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Intermediate
TAGS	body fat, lean mass, muscle, body composition, fat loss, recomposition

DEFINITION

Body composition refers to the ratio of fat mass to lean mass (muscle, bone, water, organs). Two people at identical body weight can have dramatically different health profiles based on their body composition. Fat loss — reducing body fat percentage — is the actual goal; 'weight loss' is just a proxy.

WHY IT MATTERS

- Losing fat while preserving muscle = better health outcomes, better appearance, higher metabolic rate.
- Losing weight via extreme restriction often results in 40–60% muscle loss — harmful long-term.
- Body fat distribution (visceral vs subcutaneous) matters more than total fat for health risks.
- Indian adults tend to carry more visceral fat at lower BMI — increasing metabolic disease risk.

KEY FACTS

- Healthy body fat % males: 10–20%. Females: 18–28%. Athletic: males 6–13%, females 14–20%.
- Visceral fat (around organs) is metabolically active and strongly associated with T2D, CVD.
- Indians have 3–5% more body fat than Western populations at the same BMI (TOFI phenomenon: Thin Outside, Fat Inside).
- Waist circumference: males >90 cm, females >80 cm = elevated metabolic risk (Indian cutoffs).
- During aggressive dieting, 30–50% of weight lost can be lean mass without adequate protein + resistance training.
- 1 kg of muscle occupies ~1/3 the space of 1 kg of fat — body recomposition changes shape without always changing scale weight.

ACTION STEPS

1. Measure waist circumference weekly — more diagnostic than scale weight alone.
2. Eat 1.6–2.2 g protein/kg body weight to protect lean mass during a deficit.
3. Include resistance training 2–3×/week to minimize muscle loss while dieting.
4. Track progress photos monthly alongside scale weight for full picture.
5. Use DEXA scan (■1,500–3,000) for accurate body fat measurement; or use US Navy formula (see Module 11).

DECISION TREE

IF user loses weight but looks the same or 'flabbier' → **THEN** possible muscle loss from low protein + no resistance training; increase protein, add weights

IF user's scale weight is same but they look/feel better → **THEN** body recomposition happening — fat being replaced by muscle; don't change the plan

IF user has high waist circumference despite normal BMI → **THEN** Indian TOFI pattern; focus on visceral fat reduction: low-GI diet, 150+ min cardio/week, reduce refined carbs

COMMON MISTAKES

- Obsessing over scale weight without tracking measurements, photos, or body fat.
- Extreme caloric restriction without protein/resistance training — losing muscle not fat.
- Assuming BMI alone captures health — especially misleading for Indians with higher visceral fat.

FAQ

Q: Can I lose fat and gain muscle at the same time?

A: Yes — body recomposition is possible, especially for beginners, people returning after a break, and those with significant fat reserves. Requires eating near maintenance (not aggressive deficit) + high protein + resistance training.

Q: I lost 5 kg on the scale but my waist looks the same — why?

A: You may have lost water, glycogen, or muscle rather than fat. Ensure protein ≥ 1.6 g/kg, add resistance training, and measure waist monthly instead of relying on scale alone.

AI COACHING NOTES

- Always track waist + hips + weight together — never just scale weight.
- For female users: emphasize fat loss over weight loss from day 1 to reduce unhealthy restriction behaviors.
- Indian users with normal BMI but high waist: flag visceral fat risk; adjust diet recommendations accordingly.

WL-F-007	MOD 01 · Water Retention › Causes and Management
GOALS	Weight Loss Weight Maintenance
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Beginner
TAGS	water retention, bloating, scale fluctuation, sodium, glycogen, inflammation

DEFINITION

Water retention is the accumulation of excess fluid in body tissues, causing temporary weight increases of 0.5–3 kg on the scale. It is not fat gain and does not indicate diet failure. Common causes: high sodium, hormones, glycogen replenishment, inflammation, medications.

WHY IT MATTERS

- Water weight fluctuations cause confusion, frustration, and abandonment of working diet plans.
- Scale can fluctuate 1–3 kg daily due to water, glycogen, and gut contents — unrelated to fat.
- Understanding water retention prevents panic and unnecessary diet changes.

KEY FACTS

- Each gram of stored glycogen binds ~3–4 g of water. Eating high-carb after low-carb diet = 1–3 kg water weight gain overnight.
- High sodium meal (restaurant, packaged food) can cause 0.5–2 kg of water retention within 24 hours.
- Hormonal water retention: females can retain 1–3 kg in the week before menstruation (luteal phase).
- High inflammation (illness, muscle soreness, injury) increases water retention.
- Drinking more water REDUCES water retention — dehydration signals the body to hold fluid.
- Creatine supplementation causes 0.5–1.5 kg of intramuscular water retention (performance benefit).

ACTION STEPS

1. Weigh daily, first thing in morning, after toilet, before food/water — track 7-day rolling average.
2. Focus on trend over 2–4 weeks, not daily numbers.
3. Reduce sodium to <2,000 mg/day if water retention is problematic (reduce pickles, papad, packaged snacks).
4. Stay hydrated — 2.5–3.5 L water/day reduces paradoxical retention.
5. Track menstrual cycle; expect 1–2 kg retention in week before period — ignore scale that week.

DECISION TREE

IF user gained 1–2 kg after high-carb or salty meal → **THEN** explain glycogen/sodium water retention; not fat gain; will drop within 2–3 days

IF user gained weight during period week → **THEN** normal; track non-period weeks for true trend; typical retention 0.5–3 kg in luteal phase

IF user reports persistent swelling in legs/ankles → **THEN** could be medical: heart, kidney, or venous insufficiency; recommend GP visit

IF user started creatine and gained weight → **THEN** 1–1.5 kg water retention in first 2 weeks is normal and beneficial for performance

COMMON MISTAKES

- Panicking about overnight weight gain after a high-carb meal and cutting calories further.
- Avoiding water to reduce water retention — counterproductive.
- Weighing at different times of day and comparing those readings.

FAQ

Q: Why do I gain 2 kg the day after a festival meal?

A: Mostly glycogen replenishment (from high-carb festive food) plus sodium retention. Not fat. Continue normal eating and it will drop within 3–5 days.

Q: Does drinking water make you bloat?

A: Immediate feeling of fullness yes, but adequate hydration (2.5–3.5 L/day) reduces chronic water retention by signaling kidneys that water supply is sufficient.

AI COACHING NOTES

- *Water retention is the #1 source of user confusion about scale weight — address it proactively in onboarding.*
- *For female users: set up menstrual cycle tracking and exclude period week from trend analysis.*
- *Weekly average weight is more reliable than daily weight — always guide users toward this.*

WL-F-008	MOD 01 · Sustainable Rate of Loss · Safe Weight Loss Speed
GOALS	Weight Loss Fat Loss
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Beginner
TAGS	rate of loss, sustainable weight loss, safe deficit, 0.5 kg per week, 1 kg per week, slow loss

DEFINITION

Sustainable rate of fat loss is 0.25–1 kg per week (1–4 kg/month). Faster loss risks muscle loss, nutrient deficiency, metabolic adaptation, and weight regain. The safest, most maintainable rate is 0.5 kg/week (500 kcal/day deficit).

WHY IT MATTERS

- Studies show slow weight loss (0.5 kg/week) results in 85%+ fat loss vs ~60% for aggressive loss.
- Users who lose rapidly regain weight within 2 years at significantly higher rates.
- Sustainable loss allows habit formation — the behaviors that caused loss are practiced long enough to stick.

KEY FACTS

- Safe rate: 0.5–1% of body weight per week. For 80 kg person: 400–800 g/week.
- Required deficit for 0.5 kg/week fat loss: ~500 kcal/day.
- Required deficit for 1 kg/week fat loss: ~1,000 kcal/day — safe only for those with TDEE >2,500.
- Very Low Calorie Diets (VLCD, <800 kcal/day) should ONLY be used medically under supervision.
- Initial rapid loss in first 1–2 weeks (2–4 kg) is primarily water and glycogen, not fat.

ACTION STEPS

1. Set target: 0.5 kg/week as default. Increase to 0.75 kg only if TDEE >2,200 kcal.
2. Calculate deficit: 500 kcal/day for 0.5 kg/week loss.
3. If losing faster than 1 kg/week without medical supervision: add 200–300 kcal/day.
4. Set a 'minimum floor': never eat below BMR.
5. Expect and communicate to users: loss will slow as weight decreases (TDEE reduces with lower body weight).

DECISION TREE

IF user wants to lose 10 kg in 1 month → **THEN** educate: safely 2–4 kg fat possible; 10 kg requires dangerous restriction; set realistic 12-week goal

IF user is losing 1.5 kg/week consistently → **THEN** check protein intake and resistance training; may be losing muscle; reassess if not doing both

IF user has not lost in 4 weeks with verified tracking → **THEN** true plateau; apply plateau protocols from Module 10

COMMON MISTAKES

- Using week 1 rapid loss (water weight) as the benchmark — then feeling disappointed by slower weeks.
- Setting unrealistic timelines (e.g., 15 kg in 2 months) and abandoning when unachievable.
- Comparing with social media 'fast transformations' — most involve water manipulation, lighting, filters.

FAQ

Q: My friend lost 8 kg in a month on a keto diet — why am I losing slowly?

A: Early rapid keto loss is primarily glycogen + water (each gram glycogen holds 3–4g water). Actual fat loss rate was still ~1–2 kg. Slow, steady loss is more fat and less muscle — better outcome long-term.

AI COACHING NOTES

- *Frame 0.5 kg/week positively: 2 kg/month, 6 kg in 3 months, 24 kg in 1 year is transformative.*
- *When users want to go faster: focus on the process (habits, health) not just speed of the scale.*

WL-F-009	MOD 01 · Weight Loss Myths › Common Misconceptions Debunked
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Beginner
TAGS	weight loss myths, misconceptions, fat burning foods, spot reduction, carbs bad, meal timing myths

DEFINITION

Weight loss myths are widely held beliefs about losing weight that are not supported by evidence. Believing myths leads to unnecessary restriction, ineffective strategies, and eventual abandonment of evidence-based approaches.

WHY IT MATTERS

- Indian weight loss culture is filled with harmful myths (e.g., 'rice is bad', 'eat only salad').
- Acting on myths leads to nutrient deficiencies and damaged relationships with food.
- Debunking myths is foundational coaching work — users cannot succeed if operating on false beliefs.

KEY FACTS

- MYTH: Eating at night makes you fat. FACT: Total daily calories determine weight, not timing of eating.
- MYTH: Carbs make you fat. FACT: Excess calories make you fat. Carbs are the primary fuel for brain and exercise.
- MYTH: Spot reduction works (e.g., crunches reduce belly fat). FACT: Fat is lost systemically, not locally.
- MYTH: Fat-free foods are better for weight loss. FACT: Often loaded with extra sugar; dietary fat is essential.
- MYTH: Sweating more = burning more fat. FACT: Sweat is water loss, not fat. Regained when you drink.
- MYTH: Detox diets cleanse the body. FACT: Liver and kidneys detoxify 24/7; juice cleanses are marketing.
- MYTH: Skipping meals speeds weight loss. FACT: Often leads to overeating later; total deficit is what matters.
- MYTH: Rice must be avoided for weight loss. FACT: Rice can be included in a calorie-controlled plan; it's the quantity and accompaniments that matter.
- MYTH: Protein powder is steroids. FACT: Whey/casein protein is food; no hormonal effect.
- MYTH: More gym hours = more fat loss. FACT: Overtraining without recovery reduces results; 3–5×/week optimal.
- MYTH: Green tea burns significant fat. FACT: Increases metabolism by ~4%; about 80 kcal/day — minor benefit.
- MYTH: Eating less is always better. FACT: Severe restriction causes muscle loss, metabolism slowdown, deficiencies.

ACTION STEPS

1. Identify which myths the user believes during intake — address them before starting the plan.
2. Replace myth with the evidence-based reality in plain language.
3. Reference personal experience: 'Why isn't your current approach working?' to gently challenge myth-based strategies.

DECISION TREE

IF user says 'I don't eat after 7pm' expecting weight loss → **THEN** validate structure but correct causality: calorie deficit causes loss, not the timing

IF user has eliminated all rice/roti from diet → **THEN** explore: are they getting enough carbs from other sources? Is their energy suffering? Reintroduce if needed

IF user believes their fat came from eating fat → **THEN** explain insulin-mediated fat storage from excess calories of any macronutrient; dietary fat is essential

COMMON MISTAKES

- Reinforcing myths (even gently) to avoid conflict with user's existing beliefs.
- Replacing one myth with another restrictive belief (e.g., 'all carbs bad' → 'all sugar bad').

FAQ

Q: Is rice really bad for weight loss?

A: No. 1 cup cooked white rice (200g) has ~260 kcal, 5g protein, 55g carbs, 0.5g fat. The issue is portion size and what accompanies it (heavy gravies, ghee). Rice in a deficit diet with adequate protein is fine.

Q: Are there foods that burn fat?

A: No food 'burns' fat directly. Some foods (protein, high-fiber vegetables) require more energy to digest (TEF) or promote satiety, helping maintain a deficit. No food negates a calorie surplus.

AI COACHING NOTES

- *Make myth-busting a regular part of check-ins — new myths emerge constantly from social media.*
- *For Indian users specifically: address rice, chai, ghee myths — these cause unnecessary food fear.*
- *Use empathy: 'I understand why you'd think that — this message is everywhere. Here's what the research actually shows.'*

WL-F-010	MOD 01 · BMI › Body Mass Index — Uses and Limitations
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Beginner
TAGS	BMI, body mass index, obesity classification, underweight, overweight, Indian BMI

DEFINITION

BMI (Body Mass Index) = $\text{weight (kg)} \div \text{height}^2 \text{ (m}^2\text{)}$. It classifies individuals into weight categories but does not measure body fat directly. It is a screening tool, not a diagnostic measure. For Indians, lower BMI cutoffs apply due to higher metabolic risk at lower weights.

WHY IT MATTERS

- BMI is widely used by doctors and health guidelines — understanding it helps set realistic targets.
- BMI underestimates health risk for Indians — metabolic disease risk is elevated at lower BMIs.
- BMI does not distinguish fat from muscle — a muscular athlete may have 'obese' BMI.

KEY FACTS

- Standard WHO BMI: Underweight <18.5, Normal 18.5–24.9, Overweight 25–29.9, Obese ≥30.
- Indian/Asian BMI cutoffs: Underweight <18.5, Normal 18.5–22.9, Overweight 23–24.9, Obese ≥25.
- Indians face significant metabolic disease risk at BMI 23–25 where Westerners are still in 'normal' range.
- BMI does not measure visceral fat — waist circumference (WC) is more predictive of metabolic risk.
- BMI target for Indian adults: ideally 18.5–22.9 for metabolic health (not just 'normal' by Western standards).
- Losing 5–10% of body weight (regardless of starting BMI) significantly improves blood sugar, BP, cholesterol.

ACTION STEPS

1. Calculate BMI = $\text{weight(kg)} \div (\text{height in metres})^2$.
2. Use Indian cutoffs (not Western): normal 18.5–22.9, overweight 23–24.9.
3. Pair BMI with waist circumference for full risk picture.
4. Set realistic BMI targets: for most Indians, 20–23 is ideal.

DECISION TREE

IF user BMI 23.5 and doctor says 'you're normal' → **THEN** explain Indian cutoff difference; waist circumference more relevant; metabolic labs may be needed

IF user BMI >35 (class II obesity) → **THEN** high-risk category; recommend medical supervision; prioritize mobility-friendly exercise

IF user BMI <18.5 (underweight) → **THEN** do NOT recommend weight loss; refer to nutritionist for healthy weight gain plan

FAQ

Q: My BMI is 26 but I'm muscular — am I actually overweight?

A: BMI doesn't account for muscle. Measure waist circumference and body fat percentage (DEXA or Navy formula) for a more accurate picture. If waist <90 cm (male) or <80 cm (female), risk is likely low.

AI COACHING NOTES

- Do not use BMI as the only metric — it is a starting point, not the whole story.
- For Indian users with BMI 23–25 who feel 'normal': explain elevated metabolic risk at these levels for their population.

WL-F-011	MOD 01 · Hormones and Weight › Insulin, Cortisol, Leptin, Ghrelin
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Intermediate
TAGS	hormones, insulin, cortisol, leptin, ghrelin, thyroid, estrogen, weight loss hormones

DEFINITION

Multiple hormones regulate body weight, appetite, fat storage, and metabolism. Key hormones: insulin (fat storage/release), cortisol (stress + fat storage), leptin (satiety signal), ghrelin (hunger signal), thyroid hormones (metabolic rate), estrogen/testosterone (body composition).

WHY IT MATTERS

- Hormonal imbalances can impair weight loss despite correct diet and exercise.
- Understanding hormones helps explain why stress, poor sleep, and crash dieting sabotage results.
- Some hormonal conditions (PCOS, hypothyroidism) require medical management before weight loss.

KEY FACTS

- **INSULIN:** Released in response to carbohydrate and protein intake. High insulin = fat storage mode. Low insulin = fat burning mode. Insulin resistance = cells don't respond, more insulin secreted, more fat stored.
- **CORTISOL:** Stress hormone. Chronically high cortisol → increased visceral fat, muscle breakdown, sugar cravings. Indians with high-stress jobs/studies are at elevated risk.
- **LEPTIN:** 'Fullness hormone' from fat cells. Overweight individuals can become leptin-resistant — don't feel full despite adequate signals.
- **GHRELIN:** 'Hunger hormone' from stomach. Rises before meals, after poor sleep, and during caloric restriction — explains diet hunger.
- **THYROID:** T3/T4 regulate metabolic rate. Hypothyroidism (underactive) reduces BMR by 200–600 kcal/day. India has high iodine deficiency prevalence → hypothyroidism risk.
- **ESTROGEN:** Low estrogen (post-menopause) promotes visceral fat accumulation. Explains weight gain pattern in women over 50.

ACTION STEPS

1. Manage insulin: reduce refined carbs, eat fiber with meals, don't eat carbs alone, walk after meals.
2. Manage cortisol: 7–9 hours sleep, stress management (yoga, breathing), limit caffeine after 2pm.
3. Restore leptin sensitivity: reduce ultra-processed food, ensure adequate sleep, avoid extreme restriction.
4. Manage ghrelin: eat regular meals (3–4× daily), include protein at every meal, sleep adequately.
5. Check thyroid (TSH, T3, T4) if: weight won't budge despite accurate tracking, fatigue, cold hands, hair loss, constipation.

DECISION TREE

IF user reports strong carb/sugar cravings after meals → **THEN** possible insulin resistance; recommend low-GI diet, post-meal walks, reduce refined carbs; test fasting insulin

IF user reports weight gain despite small food intake + fatigue + cold → **THEN** hypothyroidism flag; recommend GP visit before continuing plan

IF user has high stress job, belly fat predominantly, poor sleep → **THEN** cortisol-driven pattern; prioritize sleep + stress management before calorie restriction

IF user is female 45–55 with new belly fat accumulation → **THEN** perimenopause estrogen decline; adjust to higher protein, resistance training, reduce refined carbs

FAQ

Q: Do I need to 'balance my hormones' to lose weight?

A: For most people, the best way to regulate hormones is through the basics: adequate sleep, stress management, whole foods, regular exercise, and appropriate calorie intake. Supplements marketed as 'hormone balancers' have little evidence.

AI COACHING NOTES

- *Never diagnose hormonal conditions — refer to appropriate medical professional.*
- *Lifestyle interventions (sleep, stress, food quality) improve hormonal balance in most users.*
- *When PCOS or hypothyroidism is confirmed: apply Module 08 Special Populations protocol.*

WL-F-012	MOD 01 · Caloric Deficit › How to Create and Sustain a Deficit
GOALS	Weight Loss Fat Loss
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Beginner
TAGS	caloric deficit, deficit calculation, how to lose weight, calorie target, diet planning

DEFINITION

A caloric deficit occurs when you consume fewer calories than your body burns (TDEE). The deficit forces the body to use stored energy (fat, glycogen, muscle) for fuel. A sustained deficit of 300–500 kcal/day produces 0.35–0.5 kg of fat loss per week.

WHY IT MATTERS

- A caloric deficit is the only way to lose body fat — no exception.
- The source of the deficit (less food, more movement, or both) is flexible — total balance is not.
- Combining moderate food reduction + movement increase is more sustainable than food reduction alone.

KEY FACTS

- 500 kcal/day deficit × 7 days = 3,500 kcal deficit = ~0.5 kg fat loss.
- 300 kcal deficit = 0.3 kg/week. 750 kcal deficit = 0.75 kg/week (safe upper limit for most).
- Best approach: 250–300 kcal from food reduction + 150–200 kcal from increased movement.
- Protein-sparing deficit (high protein + resistance training) preserves muscle during weight loss.
- Sustainable deficit does not produce hunger, fatigue, or irritability beyond the first 1–2 weeks.

ACTION STEPS

1. Step 1: Calculate TDEE.
2. Step 2: Subtract 300–500 kcal from TDEE = daily calorie target.
3. Step 3: Set protein target first: 1.6–2.0 g/kg body weight.
4. Step 4: Fill remaining calories with carbohydrates and fats.
5. Step 5: Track for 2 weeks; adjust if progress differs from expected.

DECISION TREE

IF user is consistently hungry on their deficit → **THEN** increase protein by 20–30g/day; increase fiber; reduce liquid calories; spread meals across 4× daily

IF user deficit is >750 kcal/day → **THEN** reduce to 500 kcal max deficit; add exercise to contribute the balance

IF user feels fine on deficit but scale not moving → **THEN** check tracking accuracy (oils, sauces, drinks); recalculate TDEE; verify no medication causing retention

COMMON MISTAKES

- Creating deficit entirely by cutting food — leads to inadequate nutrition and increased hunger.
- Ignoring cooking oil in calorie counts — 4 tbsp oil in day's cooking = 480 kcal.
- Not tracking accurately for the first week — then assuming diet 'isn't working'.

FAQ

Q: What is the fastest safe way to create a calorie deficit?

A: Combine: reduce daily rice/roti portion by 25%, replace one tea with milk for black/herbal tea, walk 30 min after dinner, reduce cooking oil by 1 tbsp. Together this creates 400–600 kcal deficit without hunger.

AI COACHING NOTES

- *Always calculate the user's specific TDEE and deficit target — not generic 1,200/1,500 kcal.*
- *Create 'deficit menus' for common meals rather than asking users to count every calorie initially.*

NUTRITION SCIENCE

WL-N-001	MOD 02 · Protein › Role in Weight Loss
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Beginner
TAGS	protein, protein intake, muscle preservation, satiety, high protein diet, whey, paneer, dal protein

DEFINITION

Protein is a macronutrient made of amino acids essential for muscle building, repair, enzyme production, and immune function. For weight loss, protein is the most important macro: it preserves lean muscle during a deficit, has the highest satiety, and has the highest thermic effect of food (20–30%).

WHY IT MATTERS

- High protein intake preserves lean muscle mass during caloric restriction — prevents the 'skinny fat' outcome.
- Protein is the most satiating macronutrient — reduces hunger hormones ghrelin and increases PYY.
- Thermic effect: digesting 100 kcal of protein burns 20–30 kcal — almost free calories.
- Protein prevents muscle loss: critical because muscle drives metabolic rate.
- Indian diets are chronically low in protein — average intake is 45–60g/day vs recommended 80–120g/day.

KEY FACTS

- Minimum protein for weight loss: 1.6 g/kg body weight/day.
- Optimal protein for muscle preservation: 2.0–2.4 g/kg/day during deficit.
- For 60 kg female: target 96–120g protein/day. For 75 kg male: 120–150g/day.
- Indian vegetarian protein sources (per 100g): paneer 18g, Greek yogurt 10g, tofu 8g, cooked dal 9g, chana 9g.
- Non-veg sources (per 100g): chicken breast 31g, egg whole 13g, fish (rohu) 19g.
- Protein powder (whey): 24–25g protein per 30g serving. Legitimate supplement, not a drug.
- Spreading protein across 4 meals (20–30g each) is more effective for muscle synthesis than eating it all at once.
- Excess protein (>3.0 g/kg) provides no additional benefit and excess is used for energy.

ACTION STEPS

1. Calculate target: $\text{bodyweight (kg)} \times 1.6\text{--}2.0 = \text{daily protein grams}$.
2. Include at least 20–30g protein in each of 3–4 meals.
3. Prioritize sources: eggs, chicken, fish, paneer, Greek yogurt, legumes + grains combined.
4. If vegetarian/vegan: combine protein sources daily — rice + dal, roti + chana, tofu + seeds.
5. Consider whey protein supplement if consistently struggling to hit targets from food alone.
6. Track protein as the primary macro — once protein is hit, fill remaining calories with carbs and fat.

DECISION TREE

IF user is vegetarian and eating <60g protein/day → **THEN** recommend: 2 tbsp peanut butter + 1 cup Greek yogurt + 100g paneer + 2 cups dal + 1 cup sprouts = ~85-90g protein

IF user is losing weight but feeling weak → **THEN** likely insufficient protein; check intake; target 2.0g/kg with resistance training

IF user asks about protein powder safety → **THEN** clarify: whey is food (derived from milk); no hormonal effects; safe for most; choose reputable brand

IF user is vegan struggling with protein → **THEN** recommend: 150g tofu, 1 cup edamame, 2 cups mixed dal, 30g protein powder, hemp seeds, nutritional yeast

COMMON MISTAKES

- Eating dal at every meal but only consuming 20–25g protein/day total — dal is 9g/100g cooked, a small portion.
- Treating protein powder as dangerous — it is concentrated food protein, not a drug.
- Eating all protein in one meal (e.g., large dinner) — muscle protein synthesis is optimized with spread-out intake.
- Counting legumes as both protein and carbs but not tracking either accurately.

FAQ

Q: Can vegetarians get enough protein to lose fat without supplements?

A: Yes, but it requires deliberate effort. A day with 200g paneer + 2 cups dal + 1 cup Greek yogurt + 1 cup sprouts provides ~80–90g protein. Add protein powder for efficiency if targets remain elusive.

Q: Is eating too much protein bad for kidneys?

A: In healthy individuals, no. High protein diets (up to 2.2 g/kg) are safe. Evidence of harm exists only in individuals with pre-existing kidney disease. Those with CKD should get medical supervision for protein intake.

Q: Do I need protein immediately after a workout?

A: Post-workout anabolic window is wider than previously thought (up to 2 hours). Getting adequate daily protein is more important than exact timing. A protein-rich meal within 2 hours of training is ideal.

AI COACHING NOTES

- Protein is the most commonly under-consumed macro in Indian diets — address this in every plan.
- For vegetarians: build a 'protein menu' of go-to high-protein combinations before discussing calories.
- For hostel students: help them identify protein sources available in mess (eggs are usually available and affordable).
- When users are resistant to protein powder: recommend food-first approach; use powder only as backup.

WL-N-002	MOD 02 · Carbohydrates · Role and Management in Weight Loss
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Beginner
TAGS	carbohydrates, carbs, rice, roti, sugar, glycemic index, low carb, complex carbs, fiber

DEFINITION

Carbohydrates are the body's primary energy source, providing 4 kcal/gram. They are not inherently fattening — excess total calories cause fat gain. Simple carbs (sugar, white bread) digest rapidly; complex carbs (oats, dal, vegetables) digest slowly and provide more sustained energy and satiety.

WHY IT MATTERS

- Carbohydrates are not the enemy — chronic excess is. The brain requires ~130g glucose/day.
- Cutting carbs eliminates many traditional Indian foods unnecessarily, reducing adherence.
- Complex carbohydrates provide fiber, B vitamins, and sustained energy for activity.
- Very low carb diets work through calorie reduction, not carb magic — fat loss occurs because of the deficit created.

KEY FACTS

- Daily carbohydrate needs: 45–65% of total calories for most active individuals.
- For 1,600 kcal diet: 180–260g carbs/day. For 2,000 kcal diet: 225–325g/day.
- Glycemic Index (GI): ranks carbs by blood glucose response. High GI >70, Medium 56–69, Low ≤55.
- Low-GI Indian foods: oats (55), whole wheat roti (62), brown rice (68), basmati rice (58), dal (25–45).
- High-GI Indian foods: white rice (72–80), idli (77), white bread (75), cornflakes (81).
- 1 cup cooked white rice = 45g carbs, 200 kcal. 1 medium roti = 25–30g carbs, 120–150 kcal.
- Fiber (a type of carb) slows glucose absorption, improves satiety, feeds gut bacteria.

ACTION STEPS

1. Don't eliminate carbs — reduce portions and improve quality.
2. Replace refined grains (white rice, maida) with whole grains (brown rice, whole wheat, millets) where possible.
3. Always pair carbs with protein and fiber to slow digestion and stabilize blood sugar.
4. Reduce carb portions at dinner (largest carb meal = most spike; lower activity in evening).
5. Use the plate method: 50% vegetables, 25% protein, 25% carbs for balanced meals.
6. Prioritize: vegetables > legumes > whole grains > white rice/maida.

DECISION TREE

IF user has PCOS or diabetes → **THEN** recommend low-GI carbs exclusively; limit to 45% calories; pair with protein at every meal; post-meal walks

IF user has eliminated all carbs and feels fatigued → **THEN** add 100–150g complex carbs back; fatigue is often from insufficient carbs for brain and activity

IF user asks about keto diet → **THEN** explain: keto works by creating deficit + losing glycogen water; not magic; adherence is hardest part; not needed; consider lower-carb instead

IF user eats 3 bowls of rice at each meal → **THEN** reduce to 1 bowl; add 1 cup vegetables and 100g protein to the meal to replace satiety

COMMON MISTAKES

- Avoiding all rice and roti but eating large quantities of dry fruits, ghee, and nuts (higher calorie per gram).
- Thinking 'multigrain' or 'whole grain' bread is a diet food — still 80–100 kcal per slice.
- Eating fruit juice instead of whole fruit — removes fiber, concentrates sugar.
- Drinking 500ml aerated drink (55g sugar = 220 kcal) and not counting it.

FAQ

Q: Can I eat rice and still lose weight?

A: Yes. 1 bowl (150g cooked) = 200 kcal. Include it in your calorie budget with adequate protein and vegetables. Most Indian success stories include rice in the diet.

Q: Is brown rice better than white rice?

A: Brown rice has more fiber, B vitamins, and minerals. Lower GI (68 vs 72 for white). Good upgrade if available, but white rice in appropriate portions with dal/protein is not a weight loss barrier.

Q: Are fruits too sugary for weight loss?

A: Whole fruits have fiber that slows sugar absorption. 1 medium banana (27g carbs, 105 kcal) has significantly different metabolic effect than 27g table sugar. 2–3 fruits/day is fine for most; those with diabetes should manage portions of high-GI fruits.

AI COACHING NOTES

- *Never tell Indian users to eliminate rice or roti — this creates diet failure and food guilt.*
- *The goal is right quantity + right combination, not elimination.*
- *Coach carb timing: more carbs at breakfast/lunch when activity is higher; less at dinner.*

WL-N-003	MOD 02 · Dietary Fats › Types, Functions, and Weight Loss
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Intermediate
TAGS	dietary fat, fat intake, ghee, oil, omega 3, saturated fat, trans fat, cooking oil, nuts

DEFINITION

Dietary fats are essential macronutrients providing 9 kcal/gram. They are required for hormone production, fat-soluble vitamin absorption (A, D, E, K), brain function, and cell membrane integrity. Fat itself does not cause fat gain — excess total calories do. However, fat is calorie-dense and easy to overconsume.

WHY IT MATTERS

- Fat is 9 kcal/g — more than double protein and carbs (4 kcal/g). A small overestimation creates a large calorie excess.
- Indian cooking uses 3–6 tbsp oil/day — often invisible in calorie counts but adds 360–720 kcal.
- Omega-3 fatty acids reduce inflammation, improve insulin sensitivity, and support fat loss.
- Trans fats (vanaspati, partially hydrogenated oils) increase inflammation and metabolic risk.
- Essential fats cannot be made by the body — must come from diet.

KEY FACTS

- Types: Saturated (solid at room temp: ghee, coconut oil, butter), Monounsaturated (MUFA: olive, mustard, groundnut oil), Polyunsaturated (PUFA: sunflower, omega-3).
- Trans fats: found in vanaspati, commercial fried foods, packaged biscuits — most dangerous; increase LDL, decrease HDL.
- Recommended fat intake: 20–35% of total calories. For 1,600 kcal: 35–62g fat/day.
- 1 tbsp cooking oil = 14g fat = 120–125 kcal. Indians often use 4–6 tbsp/day in cooking = 480–720 kcal.
- Ghee: 112 kcal/tbsp, 12.7g total fat, 7.9g saturated fat. Not a superfood for weight loss despite cultural belief.
- Omega-3 sources (Indian): flaxseed (chia), walnuts, mustard oil, fatty fish (mackerel, sardines, hilsa).
- Omega-6:Omega-3 ratio in Indian diet: 30:1 to 50:1 (ideal 4:1 to 10:1) — chronic inflammation risk.

ACTION STEPS

1. Measure oil with spoons, not by pour — 1–2 tbsp per meal maximum.

2. Switch to cooking methods that use less oil: steaming, air frying, baking, grilling.
3. Use mustard or groundnut oil for everyday cooking (better MUFA/PUFA ratio vs refined oils).
4. Avoid vanaspati and partially hydrogenated fats completely.
5. Include omega-3 sources daily: 1 tsp flaxseed powder, handful walnuts, or fatty fish 2×/week.
6. Reduce ghee to 1 tsp/day maximum for weight loss phase.

DECISION TREE

IF user uses 4+ tbsp oil daily in cooking → **THEN** cutting oil by 2 tbsp = saving 240 kcal/day = 0.25 kg less fat/month without any other change

IF user follows 'keto' and eating unlimited ghee/butter → **THEN** clarify: keto requires calorie deficit too; unlimited fat in keto = calorie surplus = weight gain

IF user says ghee is healthy and eats 3–4 tbsp/day → **THEN** ghee has benefits but is calorically dense; for weight loss limit to 1 tsp/day

COMMON MISTAKES

- Adding ghee to roti + cooking oil in sabzi + butter on bread = 500–700 kcal from fat before accounting for food.
- Eating large quantities of nuts (healthy but 160–180 kcal per 25g handful).
- Switching to 'diet' or 'lite' oil — not significantly different in calories from regular oil.

FAQ

Q: Is ghee good for weight loss?

A: No. Ghee is 112 kcal/tbsp, primarily saturated fat. It has medium-chain fatty acids and fat-soluble vitamins but the calorie density makes it a weight loss obstacle if used in traditional amounts. Limit to 1 tsp/day.

Q: Which cooking oil is best for weight loss?

A: All oils are ~120 kcal/tbsp — none is 'diet-friendly' in large amounts. For health: mustard oil (good omega-3/6 ratio), groundnut oil (MUFA), or coconut oil in small amounts. The quantity matters more than the type.

AI COACHING NOTES

- *Oil reduction is often the single biggest calorie-saving opportunity in Indian diets.*
- *Always ask about cooking methods and oil quantity in dietary assessment.*
- *Teach users to measure oil — most dramatically underestimate how much they use.*

WL-N-004	MOD 02 · Dietary Fiber · Satiety and Gut Health
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Beginner
TAGS	fiber, dietary fiber, satiety, gut health, constipation, weight loss, vegetables, legumes

DEFINITION

Dietary fiber is a type of carbohydrate that is not digested by human enzymes. Soluble fiber (oats, psyllium, legumes) forms a gel, slows absorption, and reduces hunger. Insoluble fiber (vegetables, whole grains) adds bulk to stool and speeds gut transit. Both types are essential for weight management and gut health.

WHY IT MATTERS

- High-fiber foods have low calorie density — you can eat large volumes for fewer calories.
- Fiber slows gastric emptying, keeping you full longer after meals.
- Fermentable fiber feeds beneficial gut bacteria, which influence weight, immunity, and mood.
- High-fiber diet reduces risk of type 2 diabetes, CVD, colon cancer — critical for India (high disease burden).
- Most Indians eat only 10–15g fiber/day vs recommended 25–38g/day.

KEY FACTS

- Recommended fiber: women 25g/day, men 38g/day. Most Indians severely under-consume.
- High-fiber Indian foods (per 100g): rajma (15g), chickpeas (12g), green peas (5g), spinach (2.2g).
- 1 cup cooked oats: 4g fiber. 1 medium apple: 4.4g. 1 cup cooked dal: 15g.
- Psyllium husk (isabgol): 7g fiber per 2 tbsp — best fiber supplement; safe, affordable.
- Fiber adds no net calories — absorbed fiber calories from fermentation are minimal (~2 kcal/g).
- Increase fiber slowly (5g/week) to avoid gas and bloating.

ACTION STEPS

1. Add 1 cup salad (cucumber, tomato, carrot) to every main meal — adds 2–3g fiber, almost zero calories.
2. Eat whole fruit, not juice — restores fiber that juicing removes.
3. Increase dal/legume servings to 2× daily.
4. Add psyllium husk to water or yogurt if struggling to hit 25g target.
5. Replace white bread with whole wheat; white rice with brown rice or oats for some meals.

DECISION TREE

IF user is hungry on their calorie target → **THEN** increase fiber: add vegetables, legumes, psyllium husk; this can reduce hunger by 20–30% without adding calories

IF user reports constipation on weight loss diet → **THEN** increase fiber + water significantly; add psyllium husk 1 tbsp before bed with 300ml water

FAQ

Q: Does fiber have calories?

A: Technically yes, but very few (~0–2 kcal/g from fermentation vs 4 kcal/g for digestible carbs). For practical purposes, high-fiber vegetables and psyllium are essentially calorie-free.

AI COACHING NOTES

- *When user is hungry: add vegetables before reducing calories — often more effective.*
- *Fiber is the most underutilized weight management tool — integrate into every meal plan.*

WL-N-005	MOD 02 · Hydration · Water Intake and Weight Loss
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Beginner
TAGS	hydration, water, water intake, thirst, liquid calories, dehydration, fat loss

DEFINITION

Hydration refers to maintaining adequate fluid balance for physiological function. Water is calorie-free, essential for metabolism, fat oxidation, kidney function, and appetite regulation. Chronic mild dehydration is common and frequently mistaken for hunger.

WHY IT MATTERS

- Thirst is often misidentified as hunger — 500ml water before a meal can reduce meal size by 13%.
- Fat metabolism (lipolysis) requires adequate hydration — dehydration impairs fat burning.
- Adequate water intake is correlated with better weight loss outcomes in studies.
- Replacing caloric beverages (chai with sugar, juices, cold drinks) with water is a high-impact easy win.

KEY FACTS

- Daily water needs: females 2.7 L/day, males 3.7 L/day (all sources including food). In Indian climate: add 0.5–1L for heat and activity.
- 1 glass (250ml) water 30 min before meal reduces intake by 50–75 kcal per meal.
- Dehydration of just 1–2% body weight impairs cognitive function, mood, and exercise performance.
- Liquid calories in Indian diet: 1 sweet chai = 60–80 kcal. 1 glass mango juice = 130 kcal. 300ml cold drink = 130 kcal.
- Cutting 3 sweet chais/day to black/herbal tea saves ~200–240 kcal/day.

ACTION STEPS

1. Drink 500ml water upon waking before any food or drink.
2. Drink 250–350ml water 20–30 min before each main meal.
3. Carry a 1L water bottle; refill 2–3× daily.
4. Replace sugary beverages with plain water, jeera water, or herbal teas.
5. Check urine color: pale yellow = hydrated; dark yellow = drink more water.

DECISION TREE

IF user reports frequent hunger between meals → **THEN** ask about water intake; suggest 1 glass water first; often hunger is thirst

IF user drinks 3–4 sweet chai/day → **THEN** reducing to 1 cup or switching to black tea saves 120–240 kcal/day — high-impact, low-effort change

FAQ

Q: Does drinking more water speed up weight loss?

A: Directly, modestly. Indirectly, significantly — by replacing caloric drinks, reducing false hunger, supporting metabolism and fat oxidation. Aim for 3 L/day.

Q: Can I drink sparkling water on a weight loss diet?

A: Yes. Plain sparkling/fizzy water has zero calories. Flavored sparkling waters — check for added sugars. Avoid sweetened sodas.

AI COACHING NOTES

- *Before adding any supplement or dramatic change: ask about water intake and liquid calories.*
- *For hostel students: sweet chai and cold drinks are often a major hidden calorie source.*

WL-N-006	MOD 02 · Meal Timing › Optimal Eating Schedule
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Intermediate
TAGS	meal timing, intermittent fasting, meal frequency, breakfast, dinner timing, circadian eating

DEFINITION

Meal timing refers to when and how frequently you eat throughout the day. While total daily calories matter most, meal timing influences hunger, energy levels, hormones (insulin, cortisol), and sleep quality. Evidence supports consistency over perfection.

WHY IT MATTERS

- Eating at consistent times regulates hunger hormones, reducing overeating.
- Late-night eating is associated with weight gain — not because of magic, but because high-calorie snacking happens.
- Breakfast skipping often leads to overeating at lunch and dinner, increasing total intake.
- Intermittent fasting (IF) works primarily by creating a time-restricted eating window, reducing total calories.

KEY FACTS

- Optimal meal frequency: 3–4 meals/day for most people. More frequent small meals have no metabolic advantage.
- IF 16:8 (16-hour fast, 8-hour eating window): reduces total intake by 200–500 kcal/day for most people without conscious restriction.
- Eating dinner within 3 hours of sleep disrupts sleep quality and may increase fat storage.
- Eating a large breakfast reduces total daily intake by 30–40 kcal more than eating a large dinner.
- Post-meal walk (10–15 min) reduces post-meal glucose spike by 20–30% and burns 50–80 kcal.
- The 'anabolic window' for protein intake post-workout: within 2 hours (not 30 min as previously believed).

ACTION STEPS

1. Eat within 1–2 hours of waking (unless practicing IF — then first meal within eating window).
2. Space meals 3–5 hours apart for adequate hunger/fullness cycling.
3. Last meal 2–3 hours before sleep — reduces indigestion and sleep disruption.
4. If practicing IF: choose window that fits lifestyle (e.g., 12pm–8pm most common for Indians).
5. Walk 10–15 minutes after each main meal — most impactful timing intervention.

DECISION TREE

IF user wants to try intermittent fasting → **THEN** assess suitability: not recommended for teens, pregnant women, diabetics on medication, history of eating disorders; others: 14:10 or 16:8 both effective

IF user skips breakfast and overeats at night → **THEN** recommend light breakfast (protein + fiber) to reduce evening hunger; track both scenarios and compare

IF user is diabetic on insulin/metformin → **THEN** do NOT recommend extended fasting without doctor approval; meal timing must align with medication schedule

FAQ

Q: Should I eat breakfast if I'm not hungry?

A: For weight loss: eating a protein-rich breakfast helps most people reduce total daily intake. But IF without breakfast is also effective for those who prefer it. Listen to your body AND track outcomes.

Q: Is eating after 8pm really bad?

A: Not inherently. If your diet is appropriate, eating after 8pm doesn't cause unique fat gain. However, most late-night eating is calorie-dense snacking — that's the real issue.

AI COACHING NOTES

- *Don't force a specific meal timing structure — find what fits the user's lifestyle and schedule.*
- *For shift workers: use modified IF that aligns eating window with waking hours (Module 08).*

WL-N-007	MOD 02 · Satiety and Hunger › Managing Hunger on a Deficit
GOALS	Weight Loss Fat Loss
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Intermediate
TAGS	satiety, hunger, fullness, appetite, hunger management, volume eating, satiety index

DEFINITION

Satiety is the feeling of fullness and satisfaction after eating that suppresses appetite until the next meal. Hunger is the physiological drive to eat. On a caloric deficit, hunger increases — managing it is the key to adherence. Protein, fiber, and water have the highest satiety per calorie.

WHY IT MATTERS

- Poor satiety management is the primary reason diets fail — hunger overrides willpower.
- High-satiety foods allow adherence to a deficit without constant deprivation.
- Understanding what drives hunger (hormones, sleep, stress, psychology) prevents reactive eating.

KEY FACTS

- Satiety Index scores (relative to white bread = 100): boiled potatoes 323, fish 225, oatmeal 209, apples 197, eggs 150, lentils 133, white rice 138, brown rice 132, white bread 100, croissant 47.
- Protein increases satiety hormones (GLP-1, PYY) and suppresses ghrelin more than carbs or fat.
- Volume eating: eating large volumes of low-calorie foods (vegetables, water-based soups) reduces hunger without increasing calories significantly.
- Sleep deprivation increases ghrelin (hunger hormone) by 15–28% — explaining post-bad-sleep junk food cravings.
- Eating slowly (20 min per meal) allows satiety hormones to signal fullness — reduces intake by 100–200 kcal/meal.

ACTION STEPS

1. Build meals around high-satiety proteins and high-fiber vegetables first.
2. Drink 500ml water 20–30 min before each meal.
3. Eat slowly: put utensil down between bites; aim for 20 min per meal.
4. Use smaller plates and bowls — reduces perceived portion size and automatic serving amount.
5. Start meals with salad or soup — reduces total meal intake by 100–150 kcal.
6. Sleep 7–9 hours — insufficient sleep is a major hunger driver.

DECISION TREE

IF user is constantly hungry on their calorie target → **THEN** increase protein to 2.0 g/kg; add 1 cup vegetables per meal; add 500ml water before meals; check sleep quality

IF user reports hunger mainly in evenings → **THEN** likely from inadequate protein and fiber at dinner; restructure dinner to protein-first

IF user eats too fast → **THEN** implement 20-min meal timer; teach mindful eating pace; high-fiber foods require more chewing

COMMON MISTAKES

- Restricting calories without prioritizing satiety — always hungry = diet abandonment.
- Eating lunch alone (carbs + oil, no protein or fiber) — ready to eat again in 90 minutes.
- Confusing emotional hunger with physiological hunger — emotional hunger is sudden, craving-specific.

FAQ

Q: Why am I so hungry even on 1,400 kcal?

A: Check: How much protein? (Should be 80–100g+). How much fiber? (25g+). How much water? Are you sleeping 7+ hours? Are you under high stress? Fix these first before adjusting calories.

AI COACHING NOTES

- *Address satiety before addressing calories — a user who isn't hungry can maintain any plan.*
- *The highest-ROI satiety interventions: protein first at every meal, 500ml water pre-meal, vegetables at every meal.*

WL-N-008	MOD 02 · Glycemic Index and Load › Blood Sugar and Weight Loss
GOALS	Weight Loss Fat Loss General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Intermediate
TAGS	glycemic index, GI, glycemic load, GL, blood sugar, insulin, diabetes, PCOS, low GI diet

DEFINITION

Glycemic Index (GI) measures how quickly a carbohydrate-containing food raises blood glucose relative to pure glucose (GI 100). Low GI (≤ 55) foods digest slowly, raise blood sugar gradually, sustain energy, and improve satiety. Glycemic Load (GL) = $GI \times \text{carbohydrate content} \div 100$, and is more practically useful.

WHY IT MATTERS

- Low-GI diets reduce insulin spikes, improve satiety, and are especially beneficial for PCOS and T2D.
- High-GI foods cause rapid blood sugar rise then crash — leading to hunger 60–90 min later.
- Indians eating predominantly high-GI foods (white rice, idli, puri) have elevated T2D and CVD risk.
- Lowering GI of meals is achievable without eliminating traditional foods.

KEY FACTS

- Low GI foods (≤ 55): lentils 25, chickpeas 28, barley 28, basmati rice 58, whole wheat 54, oats 55.
- Medium GI (56–69): brown rice 68, whole wheat roti 62, sweet potato 63, mango 60.
- High GI (≥ 70): white rice 72–80, idli 77, white bread 75, cornflakes 81, watermelon 72.
- GI can be lowered by: adding protein/fat/fiber to meal, vinegar/acidic foods, longer cooking time (paradoxically, slightly overcooked rice is lower GI), cooling and reheating rice.
- PCOS management: low-GI diet reduces insulin resistance, improves testosterone balance, reduces symptoms.
- Combination foods have lower GI than individual components — rice + dal has lower GI than rice alone.

ACTION STEPS

1. Pair all carbohydrates with protein, fat, or fiber to lower effective glycemic load.
2. Eat dal or salad before rice/roti — reduces glucose spike from the meal.
3. Switch to basmati rice (lower GI vs polished white) for everyday use.
4. Use smaller portion of rice/roti + more sabzi and dal to reduce meal GI.
5. Avoid high-GI drinks (fruit juice, sweet chai, cold drinks).
6. Walk 10–15 min after main meals — reduces glucose spike by 20–30%.

DECISION TREE

IF user has PCOS → **THEN** mandate low-GI diet; every carb paired with protein/fat/fiber; refer to Module 08 PCOS protocol

IF user has T2D or prediabetes → **THEN** low-GI diet + post-meal walks + portion control; coordinate with their doctor for medication management

IF user is non-diabetic but has post-meal energy crash/hunger → **THEN** possible high-GI eating pattern; restructure meals with low-GI carbs + protein

FAQ

Q: Is there a way to eat white rice and keep blood sugar stable?

A: Yes: eat it with dal, sabzi, curd (protein/fat/fiber). Eat dal first, then rice. Use smaller portions. Walk after eating. This combination reduces the glycemic impact significantly.

AI COACHING NOTES

- *GI is especially important for PCOS, diabetes, and users with low satiety on current diets.*
- *For the average user without metabolic issues: total calorie balance matters more than GI.*

WL-N-009	MOD 02 · Micronutrients › Vitamins and Minerals for Weight Loss
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Intermediate
TAGS	micronutrients, vitamins, minerals, vitamin D, iron, calcium, B12, magnesium, zinc, iodine

DEFINITION

Micronutrients (vitamins and minerals) are needed in small amounts for metabolism, hormonal function, immune health, and energy production. Deficiencies — extremely common in India — impair weight loss by disrupting energy metabolism, increasing fatigue, and affecting appetite regulation.

WHY IT MATTERS

- India has the world's highest burden of micronutrient deficiencies (vitamin D, B12, iron, iodine).
- Low iron causes anemia → fatigue → reduced NEAT and exercise capacity → slower weight loss.
- Vitamin D deficiency linked to insulin resistance, fatigue, depression (all sabotage weight loss).
- B12 deficiency (very common in vegetarians/vegans) causes fatigue, nerve issues, anemia.
- Severe caloric restriction creates micronutrient deficiencies by reducing dietary variety.

KEY FACTS

- Vitamin D: >70% Indians are deficient. Supplement 2,000–4,000 IU/day. Get blood test (25-OH-D) — aim for >40 ng/mL.
- Vitamin B12: vegetarians/vegans at high risk. Supplement 500–1,000 mcg/day. India's typical dal-rice diet provides near-zero B12.
- Iron: Indian women (30–60% anemic). Signs: fatigue, cold, pale gums. Eat iron-rich foods with vitamin C (e.g., lemon with spinach).
- Calcium: Indian diet often low, especially in adults not drinking milk. 1,000–1,200 mg/day. Sources: dairy, ragi, sesame.
- Magnesium: reduces cortisol, improves sleep, supports glucose metabolism. 300–400 mg/day. Sources: nuts, seeds, dark chocolate.
- Iodine: essential for thyroid function. Use iodized salt. India has endemic iodine deficiency in many regions.
- Zinc: immune function, wound healing, testosterone. Sources: pumpkin seeds, meat, legumes.

ACTION STEPS

1. Test at minimum: vitamin D, vitamin B12, iron/ferritin, TSH once a year.
2. Supplement vitamin D (2,000 IU/day minimum) and B12 (1,000 mcg/day) for all vegetarians.
3. Increase dietary iron: spinach, rajma, horse gram, ragi (pair with vitamin C source).
4. For calcium without dairy: ragi laddoo, sesame chutney, dark leafy vegetables.
5. Ensure iodized salt is being used — not rock salt (which lacks iodine).

DECISION TREE

IF vegetarian/vegan user → **THEN** automatically recommend B12 and vitamin D supplementation; flag zinc monitoring
IF user reports extreme fatigue on diet → **THEN** rule out anemia (iron, B12) and hypothyroidism before attributing to calorie restriction
IF user has poor exercise recovery and low energy → **THEN** check magnesium status; supplement 300mg/day; improve sleep

FAQ

Q: Do I need a multivitamin if I'm eating healthy?

A: For most Indians: vitamin D and B12 supplementation is nearly universal recommendation. A daily multivitamin can fill gaps but isn't a substitute for nutritious food. Focus on food first, supplement what you can't reliably get from diet.

AI COACHING NOTES

- *Micronutrient testing (at minimum vitamin D, B12, iron) at onboarding helps identify barriers to weight loss.*
- *Always supplement B12 and vitamin D for vegetarian/vegan users — these are near-universal deficiencies.*

WL-N-010	MOD 02 · Food Quality › Whole Foods vs Ultra-Processed Foods
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Beginner
TAGS	food quality, whole foods, ultra-processed, UPF, packaged food, junk food, clean eating

DEFINITION

Food quality refers to the nutritional value and processing level of foods. Whole foods (minimal processing) are more nutritious, satiating, and health-promoting. Ultra-processed foods (UPFs) — industrially manufactured with added sugars, refined oils, salt, and additives — promote overconsumption and are strongly linked to obesity.

WHY IT MATTERS

- UPF consumption is the strongest dietary predictor of obesity in large population studies.
- UPFs are engineered for palatability — hyperpalatable combinations of fat, salt, sugar override satiety signals.
- Same calories from whole food vs UPF produce different hunger, hormone, and metabolic responses.
- India's UPF consumption is growing rapidly — biscuits, namkeen, instant noodles, packaged juices.

KEY FACTS

- NOVA food classification: Group 1 (unprocessed: vegetables, eggs), Group 2 (processed: salt, cheese), Group 3 (processed: bread, canned), Group 4 (ultra-processed: biscuits, chips, packaged juices).
- People eating UPF-heavy diet consume spontaneously 500 kcal/day MORE than whole-food eaters at same 'ad libitum' eating (Hall et al., 2019 RCT).
- UPFs provide 56% of calories in the average urban Indian diet (and rising).
- Indian UPF examples: Maggi, Lays, Biscuits (Parle-G, Oreo), packaged fruit juices, Maaza/Frooti, bread, instant oats packets.
- Comparing: 1 bag Lay's (50g) = 270 kcal, 15g fat, no protein. 50g peanuts = 285 kcal, 12g protein, 10g fat. Same calories; radically different satiety and nutrition.

ACTION STEPS

1. 80/20 rule: aim for 80% whole/minimally processed foods, 20% flexible — achievable and sustainable.
2. Read ingredient lists: if first 3 ingredients include sugar, refined flour, or vegetable oil — it's a UPF.
3. Replace packaged snacks with whole food alternatives (nuts, fruit, roasted chana, boiled eggs).
4. Cook majority of meals at home — best control over ingredients and cooking methods.
5. Limit restaurant and packaged food to ≤3 times per week during active weight loss.

DECISION TREE

IF user eats most meals from outside or packaged food → **THEN** first priority: help establish 2–3 home-cooked meals/day; this often creates 400–700 kcal deficit alone

IF user eats 'healthy' but has lots of packaged 'health' products → **THEN** evaluate: protein bars, diet cookies, packaged granola — often calorie-dense UPFs marketed as diet foods

FAQ

Q: Are protein bars healthy?

A: Most commercial protein bars are ultra-processed, with 200–350 kcal, significant sugar and additives. Use only if genuinely unable to get protein from whole foods. Boiled eggs, paneer, roasted chana are better whole-food alternatives.

AI COACHING NOTES

- *Food quality and tracking work together — someone eating 80% whole foods needs less precise tracking.*
- *Don't shame UPF choices — most users have emotional ties to these foods. Gradual displacement strategy works better.*

WL-N-011	MOD 02 · Supplements · Evidence-Based Guide for Weight Loss
GOALS	Weight Loss Fat Loss General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Intermediate
TAGS	supplements, protein powder, creatine, fat burner, whey, multivitamin, omega 3, supplement guide

DEFINITION

Dietary supplements are products taken to complement the diet, including vitamins, minerals, protein powders, herbs, and marketed 'fat burners'. For weight loss, very few supplements have robust evidence. Most marketed 'fat burners' and detox products are ineffective and potentially harmful.

WHY IT MATTERS

- Supplement industry is a multi-billion dollar market with extensive misleading marketing.
- Understanding which supplements have evidence prevents wasted money and potential harm.
- A few supplements (protein, creatine, vitamin D, B12, omega-3) have genuine evidence and safety data.

KEY FACTS

- EVIDENCE LEVEL — STRONG: Whey/plant protein powder (convenience for hitting protein targets), Creatine monohydrate (performance and muscle preservation), Vitamin D (deficiency is near-universal), B12 (essential for vegetarians/vegans), Omega-3 (1–3g EPA+DHA/day).

- EVIDENCE LEVEL — MODERATE: Caffeine (100–200mg pre-workout: +5–11% performance, modest fat oxidation), Psyllium husk (fiber, satiety, blood sugar), Magnesium glycinate (sleep, insulin sensitivity).
- EVIDENCE LEVEL — WEAK/NONE: Green tea extract (minor ~4%), Raspberry ketones (human evidence lacking), CLA (marginal at best), Garcinia cambogia (no meaningful effect in trials), Detox teas (marketing; laxative effect).
- HARMFUL/AVOID: Ephedra-containing products, synephrine, high-dose herbal diuretics, unverified 'fat burner' blends.
- Whey protein: 24–25g protein/scoop, 110–130 kcal. Derived from milk. Not a steroid, not hormonal. Safe for all ages.
- Creatine: 3–5g/day. Not a steroid. Increases strength output by 5–15%, allowing more training volume → more muscle → higher metabolic rate. Causes ~1–1.5 kg water retention (intramuscular, beneficial).

ACTION STEPS

1. Priority 1: Vitamin D 2,000 IU/day (everyone in India).
2. Priority 2: Vitamin B12 1,000 mcg/day (all vegetarians/vegans).
3. Priority 3: Omega-3 1–2g EPA+DHA/day (if not eating fatty fish 2×/week).
4. Priority 4: Whey or plant protein supplement if consistently under protein target from food.
5. Optional: Creatine monohydrate 3–5g/day if doing resistance training.
6. Avoid: all 'fat burner', 'detox', 'slimming' products. Report to user: these don't work and may be harmful.

DECISION TREE

IF user wants to buy a 'fat burner' → **THEN** explain: no fat burner meaningfully works; save money; achieve deficit through diet + exercise; most contain caffeine (get from coffee) or stimulants

IF user asks about protein powder safety → **THEN** explain it's concentrated food, not a drug; choose reputable brand with FSSAI certification; whey for milk-tolerant, plant protein for vegans/lactose-intolerant

IF user asks about detox tea/weight loss tea → **THEN** explain: primarily laxative and diuretic effect; causes water loss, not fat loss; can cause electrolyte imbalance

FAQ

Q: Does creatine make you fat?

A: No. Creatine causes 1–1.5 kg of intramuscular water retention in the first 2 weeks. This is not fat. Long-term, creatine improves strength, allows more training, builds more muscle — net positive for body composition.

Q: Which protein powder is best for Indians?

A: Whey concentrate (affordable, effective): MuscleBlaze, MyProtein, Optimum Nutrition are reputable. For vegetarians who avoid whey: pea protein, soy protein, or rice+pea blend. Avoid 'mass gainers' if goal is fat loss.

AI COACHING NOTES

- Always address supplement use early in intake — many users spend ■3,000–8,000/month on ineffective products.
- The supplement priority stack for weight loss: (1) vitamin D, (2) B12 (if veg), (3) protein powder if needed, (4) creatine if training, (5) omega-3 if no fish.
- Never recommend proprietary blends or products making dramatic claims.

WL-N-012	MOD 02 · Calorie Density › Volume Eating Strategy
GOALS	Weight Loss Fat Loss
USERS	Adult · Teen · Senior · Male · Female · Student · Office Worker · Vegetarian · Non-Vegetarian · Vegan
DIFF	Beginner
TAGS	calorie density, volume eating, low calorie density, energy density, satiety per calorie

DEFINITION

Calorie density is the number of calories per gram of food. Low-calorie-density foods (vegetables, broth soups) allow large volumes for few calories. High-calorie-density foods (oils, dry fruits, cheese) deliver many calories in small amounts. Eating low-calorie-density foods increases meal satisfaction without increasing calories.

WHY IT MATTERS

- Volume of food in the stomach is a key satiety signal — low-calorie-dense foods maximize volume.
- Building meals around low-density foods allows more food, less hunger, and lower calorie intake.
- Most Indian traditional diet fails because it's calorie-dense (oil-heavy cooking, sweets).

KEY FACTS

- Calorie density by category (kcal/100g): oils 900, ghee 900, dry fruits 500–600, paneer 260, chicken breast 165, cooked rice 130, cooked dal 120, potatoes 80, most vegetables 20–50, cucumber 16, watermelon 30.

- Replace 100g oil (900 kcal) in cooking with water, stock, or air fryer — saves enormous calories.
- Replacing 1 cup rice (200 kcal) with 3 cups salad (50–75 kcal) as meal filler — significantly reduces overall meal density.

ACTION STEPS

1. Start every meal with low-calorie-density foods: salad, soup, raita.
2. Fill half the plate with vegetables before adding rice/roti.
3. Use water-based cooking methods: boiling, steaming, air frying instead of deep frying.
4. Swap calorie-dense snacks for low-density options: chips → cucumber + hummus, dry fruits → whole fruit.

FAQ

Q: Can I eat as much vegetables as I want?

A: Yes. Non-starchy vegetables (spinach, cucumber, tomato, capsicum, cabbage, carrot) average 20–40 kcal/100g. You'd struggle to eat 500g of salad for 100–200 kcal. Go for it.

AI COACHING NOTES

- *Calorie density education is one of the highest-leverage teaching moments — shifts a user's entire food relationship.*
 - *Frame it positively: 'You can eat MORE of these foods' rather than restricting.*
-

COMPLETE INDIAN FOOD DATABASE

WL-FD-000	MOD 03 · Food Database Guide · How to Use This Database
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student
DIFF	Beginner
TAGS	food database, calorie reference, Indian foods, macro tracking, calorie counting

DEFINITION

This database contains 350+ Indian foods with complete nutritional profiles. Each entry includes serving size, calories, macronutrients (protein, carbs, fat, fiber), Glycemic Index (GI), Satiety Rating (H/M/L), Weight Loss Rating, and practical tips for use in meal planning.

WHY IT MATTERS

- Accurate calorie tracking requires reliable Indian food data — most apps undercount Indian foods.
- Satiety rating helps choose foods that keep you full within a calorie budget.
- GI rating guides food choices for PCOS, diabetes, and those with blood sugar regulation issues.
- Weight Loss (WL) rating provides a quick decision tool for coaches and users.

KEY FACTS

- WL RATING KEY: Excellent = high volume per calorie, high protein/fiber, highly recommended. Good = nutritious, include regularly. Moderate = include in controlled portions. Limit = occasional treat, minimize during active fat loss.
- SATIETY KEY: H=High (stays full 3-4h), M=Medium (stays full 2-3h), L=Low (hungry again in 1-2h).
- GI KEY: ≤55 Low, 56–69 Medium, ≥70 High.
- Calorie values are per the listed serving size — not per 100g unless specified.
- Cooking method significantly changes calorie content: 1 cup boiled dal = 116 kcal; 1 cup dal tadka (1 tbsp oil) = 165 kcal.

AI COACHING NOTES

- Always verify home cooking oil amounts — Indian cooking uses more oil than standard references assume.
- Restaurant portions are typically 1.5–2× the servings listed here.
- Satiety rating determines food sequence: eat H-satiety foods first (dal, curd, salad) before M/L foods (rice, roti).

WL-FD-001	MOD 03 · Food Database · Rice and Grain Products
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student
DIFF	Beginner
TAGS	rice, grains, roti, oats, bajra, ragi, calories in rice, brown rice vs white rice

Rice and Grain Products

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
White Rice (cooked)	1 cup/200g	260	5	57	0.4	0.6	72	M	Moderate	Eat with dal+sabzi; control portion
Brown Rice (cooked)	1 cup/200g	220	5	46	1.8	3.5	68	M-H	Good	More fiber, lower GI than white rice
Basmati Rice (cooked)	1 cup/200g	240	4.4	53	0.5	0.6	58	M	Good	Lower GI than regular white rice
Red Rice (cooked)	1 cup/200g	215	4.9	45	1.2	2.0	55	M-H	Good	High antioxidants; good for PCOS
Matta Rice (Kerala, cooked)	1 cup/200g	220	5.2	46	0.8	1.8	60	M	Good	Parboiled; retains more nutrients
Poha (cooked)	1 cup/200g	150	3.5	33	1.0	1.5	70	M	Good	Light, easy to digest; good breakfast

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Idli (steamed)	2 medium/100g	130	3.8	27	0.5	0.8	77	M	Good	Fermented = better bioavailability; watch coconut chutney
Dosa (plain)	1 medium/75g	120	3.5	22	1.8	0.5	75	M	Good	Oil added for cooking; get oil-light version
Pongal	1 cup/200g	230	7	36	8	2	65	M	Moderate	Ghee+rice combo; moderate portion
Upma (sooji)	1 cup/200g	200	5	30	6	2	65	M	Moderate	High in refined semolina; ok if portion-controlled
Oats (rolled, cooked)	1 cup/200g	150	5	27	2.5	4	55	H	Excellent	Best breakfast grain; high beta-glucan fiber
Ragi/Finger Millet (cooked)	1 cup/200g	180	4	38	1.5	2	68	M-H	Good	High calcium; good for women
Jowar (sorghum, cooked)	1 cup/200g	220	7.5	44	2	4.5	55	H	Good	High fiber; gluten-free
Bajra (pearl millet, cooked)	1 cup/200g	230	8	42	3.5	4.5	54	H	Good	High protein for a grain; great for winters
Quinoa (cooked)	1 cup/200g	222	8	39	3.5	5	53	H	Excellent	Complete protein; all amino acids; costly but excellent
Broken Wheat/Dalia (cooked)	1 cup/200g	150	5.5	31	0.8	4.5	46	H	Excellent	High fiber, low GI; excellent weight loss grain

WL-FD-002 MOD 03 · Food Database · Breads and Rotis

GOALS Weight Loss | Fat Loss | Weight Maintenance | General Health

USERS Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student

DIFF Beginner

TAGS roti calories, chapati, paratha, naan, bread, whole wheat roti

Breads and Rotis

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Whole Wheat Roti (no oil)	1 medium/35g	85	3	17	0.5	2.5	62	M	Good	Better than maida; whole grain preferred
Whole Wheat Roti (with ghee, 1 tsp)	1 medium/35g	125	3	17	5	2.5	62	M	Moderate	Ghee adds 40 kcal; limit to 1 tsp
Multigrain Roti	1 medium/35g	90	3.5	16	1.5	3	60	M-H	Good	Best roti option for weight loss
Maida Roti/Chapati	1 medium/35g	100	2.8	21	1	0.5	70	L	Moderate	Lower nutrition than whole wheat; occasional ok
Paratha (plain, 1 tsp oil)	1 medium/60g	180	4	28	6	2	65	M	Moderate	Double the calories of roti; limit portion
Stuffed Aloo Paratha (1 tsp oil)	1 medium/100g	280	5	45	9	3	65	M-H	Moderate	Filling but calorie-dense; 1 only
Puri (deep fried)	2 medium/60g	240	4	30	12	1.5	70	L	Limit	High fat from frying; use only occasionally
Bhatura (deep fried)	1 medium/100g	350	7	55	12	2	72	L	Limit	Very calorie-dense; restaurant portion often 2× listed
Naan (plain, tandoor)	1 naan/80g	260	8	48	5	2	70	M	Moderate	Restaurant naan often buttered; request plain
Butter Naan	1 naan/80g	320	8	48	12	2	70	L	Limit	Adds 60+ kcal of butter; avoid for weight loss
Thepla (methi)	2 thepla/80g	200	6	28	7	3	60	M	Moderate	Portable; good travel food; made with whole wheat

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Makki di Roti	1 medium/60g	150	3	30	3	2.5	65	M	Good	Gluten-free; pairs with mustard greens (sarson)
Bread (white)	2 slices/60g	160	5	30	2	1	75	L	Moderate	High GI; better than biscuits but not ideal
Brown Bread (whole wheat)	2 slices/60g	150	6	28	2.5	3	65	M	Good	Verify '100% whole wheat' — most are mixed

WL-FD-003 MOD 03 · Food Database › Dals and Legumes

GOALS Weight Loss | Fat Loss | Weight Maintenance | General Health

USERS Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student

DIFF Beginner

TAGS dal, legumes, dal protein, rajma, chana, moong dal, masoor dal, protein sources

Dals and Legumes

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Masoor Dal (cooked, no oil)	1 cup/200g	116	9	20	0.4	8	26	H	Excellent	Best weight loss dal; high protein, high fiber
Moong Dal (cooked, no oil)	1 cup/200g	105	7.5	19	0.4	7.6	25	H	Excellent	Easy to digest; high protein; excellent for all
Moong Dal (sprouted)	1 cup/150g	60	4.3	11	0.4	1.4	25	H	Excellent	Raw form; higher enzyme activity; zero cooking
Toor/Arhar Dal (cooked)	1 cup/200g	126	8.7	23	0.5	6.7	22	H	Excellent	Most commonly consumed dal in India
Chana Dal (cooked)	1 cup/200g	180	10.5	30	2.5	8.5	11	H	Excellent	Lowest GI of all dals; highest satiety
Urad Dal (cooked)	1 cup/200g	133	8.9	24	0.5	6	43	H	Excellent	Rich in B vitamins; high protein
Rajma (cooked)	1 cup/200g	225	15.4	40	0.9	13.1	28	H	Excellent	Best protein + fiber combo; excellent WL food
Chickpeas/Chana (cooked)	1 cup/200g	269	14.5	45	4.2	12.5	28	H	Excellent	Top food for satiety and nutrition
Black Chana (cooked)	1 cup/200g	230	15.3	38	2.8	12.5	32	H	Excellent	High iron, protein; excellent for anemic women
Moth Dal (cooked)	1 cup/200g	210	13	36	1.5	10	38	H	Excellent	Less common but nutritionally excellent
Peas (green, cooked)	1 cup/160g	125	8.6	22.5	0.4	8.6	48	H	Excellent	Excellent protein and fiber source
Soybean (cooked)	1 cup/200g	298	28.6	17	15.4	10.3	15	H	Excellent	Highest plant protein; complete amino acid profile
Edamame (cooked)	1 cup/155g	189	18.5	15.8	8	8	18	H	Excellent	Young soybean; excellent protein snack
Masoor Dal (cooked, tadka)	1 cup/200g	165	9	21	4	8	26	H	Good	Add 1 tsp oil tadka: adds 40-50 kcal
Dal Makhani (restaurant)	1 cup/200g	300	10	28	16	7	30	M-H	Moderate	High fat from cream/butter; limit portion

WL-FD-004 MOD 03 · Food Database › Vegetables

GOALS Weight Loss | Fat Loss | Weight Maintenance | General Health

USERS Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student

DIFF Beginner

TAGS vegetables, sabzi, spinach, palak, bhindi, vegetable calories, low calorie foods

Vegetables

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Spinach (raw)	100g	23	2.9	3.6	0.4	2.2	15	H	Excellent	Iron + vitamin C; extremely low calorie
Spinach (cooked)	100g	41	5.4	6.4	0.7	4.3	15	H	Excellent	Cook with minimum oil
Broccoli (cooked)	100g	35	2.4	7	0.4	2.6	15	H	Excellent	High in sulforaphane; anti-cancer properties
Cauliflower (cooked)	100g	25	2	5	0.3	2	15	H	Excellent	Versatile; low calorie rice substitute possible
Cabbage (cooked)	100g	29	1.3	6.1	0.1	2.5	10	H	Excellent	Liver health; extremely filling per calorie
Cucumber (raw)	100g	16	0.7	3.6	0.1	0.5	15	H	Excellent	95% water; best low-calorie snack
Tomato (raw)	100g	18	0.9	3.9	0.2	1.2	30	M	Excellent	Lycopene; vitamin C; use freely
Onion (raw)	100g	40	1.1	9.3	0.1	1.7	20	M	Excellent	Use freely; adds flavor without calories
Bottle Gourd/Lauki	100g	14	0.6	3.4	0.02	0.5	15	M	Excellent	One of the lowest calorie vegetables
Ridge Gourd/Turai	100g	18	0.7	4.1	0.1	0.7	20	M	Excellent	High water content; excellent for weight loss
Bitter Gourd/Karela	100g	17	1.0	3.7	0.17	2.8	15	M-H	Excellent	Lowers blood sugar; excellent for diabetics
Drumstick/Moringa	100g pods	37	2.1	8.5	0.1	3.2	20	M	Excellent	Very high in vitamins; traditional superfood
Capsicum/Bell Pepper (red)	100g	31	1	6	0.3	2.1	15	M	Excellent	High vitamin C; free to use
Eggplant/Brinjal	100g	25	1	6	0.2	3	15	M	Excellent	Low calorie; often cooked in oil — watch cooking fat
Bhindi/Okra	100g	33	1.9	7.5	0.2	3.2	20	H	Excellent	High fiber (mucilage); regulates blood sugar
French Beans	100g	31	1.8	7.1	0.2	3.4	15	H	Excellent	High fiber; excellent for gut health
Carrot (raw)	100g	41	0.9	9.6	0.2	2.8	47	M	Excellent	Beta-carotene; good snack option
Beetroot (cooked)	100g	44	1.7	9.6	0.2	2	64	M	Good	Medium GI; excellent iron and folate
Sweet Potato (cooked)	100g	86	1.6	20	0.1	3	63	M-H	Good	Good carb source; more nutritious than white potato
Potato (boiled)	100g	77	2	17	0.1	1.8	80	H	Moderate	High GI when hot; cooling increases resistant starch, lowers GI
Potato (fried)	100g	312	3.5	36	17	2.7	85	L	Limit	Oil absorption triples calories; avoid for weight loss
Corn/Maize (boiled)	100g	96	3.4	21	1.5	2	55	M	Good	Good fiber; avoid adding butter or cheese
Green Peas (cooked)	100g	84	5.8	15	0.4	5.7	48	H	Excellent	High protein and fiber for a vegetable
Fenugreek/Methi (leaves)	100g	49	4.4	9	0.9	2.7	25	H	Excellent	Lowers blood sugar; excellent for PCOS, diabetes
Coriander leaves	100g	23	2.1	3.7	0.5	2.8	15	H	Excellent	Use generously; adds flavor for very few calories
Palak (cooked, no oil)	100g	41	5.4	6.4	0.7	4.3	15	H	Excellent	Iron-rich; best with lemon juice to enhance iron absorption

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Drumstick leaves (cooked)	100g	64	6.8	10.4	1.7	4	20	H	Excellent	Moringa: extremely high in calcium, iron, vitamins
Cluster beans/Guar	100g	16	1.2	2.9	0.1	1.7	15	H	Excellent	Galactomannan fiber; excellent for blood sugar

WL-FD-005	MOD 03 · Food Database · Fruits
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student
DIFF	Beginner
TAGS	fruits, fruit calories, banana calories, mango, apple, papaya, diabetic fruit

Fruits

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Apple (medium)	1 medium/150g	78	0.4	21	0.2	3.7	39	M-H	Good	High fiber; great snack; low GI
Banana (medium)	1 medium/120g	105	1.3	27	0.4	3.1	51	M-H	Good	Quick energy; post-workout suitable; avoid 3+ daily
Guava	1 medium/150g	99	2.2	23	1.4	8.9	30	H	Excellent	Highest fiber of common fruits; excellent for WL
Papaya	1 cup/150g	60	0.7	15	0.4	2.5	60	M	Excellent	Digestive enzymes; very low calorie; excellent choice
Mango (Alphonso)	1 cup/150g	99	1.4	25	0.6	2.6	60	M	Moderate	High natural sugars; limit to 1 cup/day when in season
Orange	1 medium/130g	62	1.2	15.4	0.2	3.1	43	M	Excellent	Vitamin C; eat whole; not juice
Pear	1 medium/150g	101	0.6	27	0.2	5.5	38	H	Excellent	Very high fiber; excellent satiety
Pomegranate (seeds)	1 cup/175g	144	2.9	32.5	2.1	7	53	M	Good	High antioxidants; moderate calorie; portion control
Grapes (green)	1 cup/150g	104	1.1	27	0.2	1.4	59	L-M	Moderate	Easy to overeat; limit to 1 cup; low fiber
Watermelon	1 cup/152g	46	0.9	11.5	0.2	0.6	72	L-M	Good	Very low calorie but high GI; fine in weight loss context
Strawberry	1 cup/152g	49	1	11.7	0.5	3	41	M	Excellent	Low calorie, high antioxidants; excellent for WL
Pineapple	1 cup/165g	82	0.9	21.6	0.2	2.3	66	M	Good	Bromelain enzyme; moderate GI; ok in portions
Chikoo/Sapota	1 medium/100g	83	0.4	20	1.1	5.3	55	M	Moderate	High sugar; limited to 1 per day
Coconut (fresh, scraped)	50g	177	1.7	7.6	17	4.7	45	H	Moderate	Very high calorie from fat; limited use in WL phase
Dates (dried)	3 dates/30g	80	0.7	21.4	0.1	1.9	42	M	Moderate	Natural sweetener; high sugar; limit to 2-3 per day
Amla (Indian Gooseberry)	2 medium/60g	22	0.5	5.1	0.1	1.7	30	H	Excellent	Vitamin C powerhouse; anti-inflammatory
Jamun (Java Plum)	1 cup/150g	75	0.7	19	0.2	0.6	25	M	Excellent	Lowers blood sugar; excellent for diabetes and PCOS

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Kiwi	2 medium/150g	93	1.8	22.5	0.8	4.5	52	M-H	Excellent	High vitamin C; better sleep quality (evening consumption)
Litchi	10 litchis/100g	66	0.8	16.5	0.4	1.3	57	L	Moderate	Seasonal; limit as high in natural sugars
Custard Apple/Sitaphal	1 medium/100g	94	2.1	23.6	0.4	2.4	60	M	Moderate	High natural sugar; small portion only

WL-FD-006 MOD 03 · Food Database › Dairy Products

GOALS Weight Loss | Fat Loss | Weight Maintenance | General Health

USERS Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student

DIFF Beginner

TAGS dairy, paneer, curd, milk, greek yogurt, ghee, paneer calories

Dairy Products

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Milk (full fat, 3.5%)	1 cup/240ml	149	8	12	8	0	31	M	Moderate	High protein but high fat; switch to low-fat for WL
Milk (low fat, 1.5%)	1 cup/240ml	107	8.2	12.5	2.5	0	31	M-H	Good	Good protein source; better fat profile for WL
Milk (toned, 3%)	1 cup/240ml	130	7.5	11.7	5	0	31	M	Good	Most commonly available in India
Curd/Dahi (full fat)	1 cup/240g	149	8.5	11.4	8	0	35	M-H	Good	Probiotics; excellent WL food; watch added sugar
Curd/Dahi (low fat)	1 cup/240g	110	9	12	3	0	35	H	Excellent	Best everyday dairy for WL; high protein
Greek Yogurt	1 cup/200g	133	22.9	8.3	0.7	0	11	H	Excellent	Highest protein dairy; best choice for protein intake
Paneer (regular)	100g	265	18.3	3.4	20.8	0	25	H	Moderate	High protein but calorie dense; max 100g per serving
Paneer (low fat)	100g	165	20	4.5	7.5	0	25	H	Good	Better option for WL while maintaining protein
Tofu (firm)	100g	76	8.1	1.9	4.7	0.3	15	M-H	Excellent	Excellent vegan protein; very low calorie
Tofu (silken)	100g	55	4.8	2.9	2.7	0	15	M	Excellent	Lower protein but smooth texture; good in smoothies
Buttermilk/Chaas (plain)	1 cup/240ml	55	3.5	5.5	1.8	0	35	M	Excellent	Perfect weight loss beverage; probiotic; filling
Butter	1 tbsp/14g	102	0.1	0	11.5	0	0	L	Limit	Pure fat; adds minimal nutrition vs calories
Ghee	1 tsp/5g	45	0	0	5	0	0	L	Limit	Limit to 1 tsp/day max during WL phase
Khoya/Mawa	50g	213	9	25	8.5	0	55	M	Limit	Used in sweets; very calorie dense
Lassi (plain, 1 cup)	240ml	130	6	18	4	0	40	M	Moderate	Control sugar; salted lassi better for WL than sweet
Lassi (sweet, 1 cup)	240ml	210	6	32	5.5	0	55	L	Limit	High sugar; better to have plain curd
Cheese (cheddar)	30g	120	7.4	0.4	9.9	0	0	M	Moderate	Calorie dense; limit to 30g serving
Cottage Cheese/Paneer substitute	100g	98	11.1	3.4	4.3	0	15	H	Good	Lower fat than regular paneer; good WL option

WL-FD-007	MOD 03 · Food Database · Eggs and Non-Vegetarian
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student
DIFF	Beginner
TAGS	egg, chicken, fish, non-veg protein, chicken calories, egg protein

Eggs and Non-Vegetarian

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Egg (whole, boiled)	1 large/50g	78	6.3	0.6	5.3	0	0	H	Excellent	Best WL protein; nutrient-dense; affordable
Egg whites only	3 whites/99g	52	10.9	0.7	0.2	0	0	M	Excellent	Pure protein; very low calorie; good post-workout
Egg (scrambled, no oil)	2 eggs/100g	148	10	1.6	11	0	0	H	Good	Cook with non-stick; no added oil needed
Egg (fried, 1 tsp oil)	1 egg/55g	115	6.3	0.6	9.3	0	0	M	Moderate	Added cooking oil adds 40 kcal
Chicken Breast (grilled, no skin)	100g	165	31	0	3.6	0	0	H	Excellent	Best non-veg WL food; highest protein density
Chicken Breast (roasted, no skin)	100g	175	33	0	4	0	0	H	Excellent	Lean; no carbs; excellent satiety
Chicken Thigh (cooked, no skin)	100g	209	26	0	11	0	0	H	Good	More fat than breast but still good protein source
Chicken (curry, with gravy)	100g	180	20	5	9	0	0	H	Good	Depends heavily on oil and cream used; verify preparation
Chicken Tandoori	100g	150	24	3	5	0	0	H	Excellent	Best restaurant chicken option for WL
Egg Bhurji (1 tsp oil)	2 eggs + veg	220	13	4	16	0.8	0	H	Moderate	Oil adds significant calories; use minimal
Fish - Rohu (cooked)	100g	97	18.5	0	2.7	0	0	H	Excellent	Lean freshwater fish; very popular in India
Fish - Pomfret (cooked)	100g	95	19	0	2.5	0	0	H	Excellent	Low fat; high protein; excellent choice
Fish - Mackerel/Bangda	100g	158	20	0	8.6	0	0	H	Excellent	Fatty fish; excellent omega-3 source
Fish - Sardine/Pedvey	100g	142	17	0	8	0	0	H	Excellent	Omega-3 rich; affordable; excellent nutrition
Fish - Hilsa	100g	271	19.4	0	22	0	0	H	Moderate	Highest omega-3 in Indian fish; calorie-dense
Fish - Tuna (canned in water)	100g	116	25.5	0	1	0	0	H	Excellent	Ready to use; excellent protein; watch sodium
Prawn/Shrimp (cooked)	100g	85	18	0.9	1.2	0	0	H	Excellent	Very lean; high protein; excellent WL food
Mutton (lean cut, cooked)	100g	250	25.6	0	16	0	0	H	Moderate	High protein but calorie-dense from fat; 100g limit
Mutton Keema (lean)	100g	185	20	2	11	0	0	H	Moderate	Good protein; watch added oil in preparation
Liver (chicken, cooked)	100g	172	24	3	7	0	0	H	Excellent	Highest iron content; B12 rich; excellent for anemia

WL-FD-008	MOD 03 · Food Database · Snacks
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student

DIFF	Beginner
TAGS	snacks, healthy snacks, makhana, roasted chana, nuts, weight loss snacks

Snacks

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Roasted Chana (whole)	30g (small handful)	116	7.2	17	2.1	5	28	H	Excellent	Best Indian snack for WL; high protein + fiber
Makhana/Fox Nuts (plain)	30g (15–20 pieces)	106	4	20	0.5	0.4	50	M-H	Good	Low fat; roast without oil; watch flavored versions
Peanuts (roasted)	30g (small handful)	167	7.5	5.3	14	2.3	14	H	Moderate	Healthy fat but calorie-dense; limit to 30g serving
Almonds	15 nuts/18g	103	3.8	3.4	8.9	1.9	15	H	Moderate	Excellent nutrition; easy to overeat; measure portion
Walnuts	4 halves/14g	92	2.2	1.9	9.2	0.9	15	H	Moderate	Best omega-3 nut; measure; calorie-dense
Cashews	6 nuts/15g	87	2.2	4.6	7	0.5	22	M	Moderate	Less protein than other nuts; easy to overeat
Peanut Butter	1 tbsp/16g	94	4	3	8	1	14	H	Moderate	High satiety; measure carefully; no-sugar versions
Sprouts (mixed, raw)	1 cup/100g	35	3.8	6	0.4	2	22	H	Excellent	No-cook protein + fiber; excellent hostel option
Boiled Egg	1 large/50g	78	6.3	0.6	5.3	0	0	H	Excellent	Most convenient protein snack
Dhokla (steamed)	100g/3 pieces	155	6.2	27	3.5	2	60	M-H	Good	Fermented; good protein from besan; watch chutney calories
Khakra (whole wheat)	2 pieces/30g	110	3.5	19	2.5	1.7	60	M	Good	Light; portable; better than biscuits
Thepla (homemade)	1 thepla/50g	120	3.5	18	3.8	2	60	M	Good	Travel-friendly; good fiber
Murmura/Puffed Rice (plain)	1 cup/15g	60	1.2	13.5	0.2	0.1	70	L	Moderate	Low calorie but low satiety and high GI; add sprouts or chana
Corn (bhutta, grilled)	1 cob/100g	96	3.4	21	1.5	2	55	M	Good	Good snack without butter; adds fiber
Cucumber + Hummus	100g + 30g	83	3	11	3	3	25	H	Excellent	Low calorie; high satiety; excellent WL snack combo
Popcorn (air popped)	1 cup/8g	31	1	6	0.4	1.2	55	M-H	Good	Large volume for very few calories; avoid butter/oil
Biscuits (Parle-G)	2 biscuits/14g	68	1	10	2.8	0.3	70	L	Limit	Low nutrition; easy to overeat; not satiating
Namkeen (mixed)	30g	145	3	16	8	1.5	65	L	Limit	High fat + salt; minimal satiety; avoid for WL
Chips (Lay's)	30g (small pack)	162	2	16	10	1.3	72	L	Limit	High fat, salt; engineered for overconsumption
Roasted Peanuts + Jaggery Chikki	1 bar/30g	140	4	17	7	1.5	55	M	Moderate	Better than chocolate; reasonable snack; limit 1 bar

WL-FD-009	MOD 03 · Food Database · Street Foods
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student
DIFF	Beginner
TAGS	street food, vada pav, pani puri, samosa, chaat, street food calories

Street Foods

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Vada Pav	1 piece/130g	280	7	40	12	3	65	M	Limit	Calorie-dense; good protein from vada but deep fried
Pani Puri (6 pieces)	6 puris + filling	250	5	38	10	2	68	L	Limit	Calorie and sodium dense; occasional treat
Bhel Puri (plate)	1 plate/150g	200	5	35	7	3	65	M	Moderate	Better option if made without sev; add protein (chana)
Sev Puri (plate)	1 plate/150g	280	6	38	13	2.5	65	L	Limit	High fat from sev; calorie-dense for portion
Pav Bhaji	1 serving/300g	450	10	65	18	6	65	M-H	Limit	High butter; portion control critical
Samosa (1 piece)	1 samosa/80g	220	4	28	11	2.5	70	L	Limit	Deep fried; high refined carb and fat
Kachori	1 piece/80g	270	5	32	14	2	65	L	Limit	Very calorie-dense; occasional treat only
Aloo Tikki	2 tikkis/120g	230	4.5	32	10	2.5	70	M	Limit	Pan fried; better than deep fried; homemade with less oil
Chole Bhature	1 serving/350g	680	18	95	26	10	68	H	Limit	Occasional treat; extremely calorie dense as a meal
Dahi Puri (6 pieces)	6 puris	290	8	42	10	2.5	62	M	Moderate	Curd adds protein; better than pani puri for nutrition
Misal Pav	1 serving/300g	380	13	58	12	8	60	H	Moderate	High fiber from usal; better street food option
Poha (street stall)	1 plate/200g	200	4	40	5	2	68	M	Moderate	Watch oil; homemade with 1 tsp oil = better choice
Manchurian (dry, 1 plate)	150g	280	6	36	13	2	65	L	Limit	Deep fried; high sodium; occasional treat
Spring Roll (fried, 2 pieces)	2 rolls/120g	280	5	34	14	2	68	L	Limit	Deep fried wrapper; high calorie for size
Rava Dosa (crispy)	1 large/100g	180	4	28	6	1	72	L-M	Moderate	Thinner = less filling but fewer calories than regular dosa

WL-FD-010 MOD 03 · Food Database · Restaurant Foods

GOALS Weight Loss | Fat Loss | Weight Maintenance | General Health

USERS Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student

DIFF Beginner

TAGS restaurant food, butter chicken, biryani, paneer butter masala, restaurant calories

Restaurant Foods

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Butter Chicken (200g + rice)	1 serving/400g	680	35	55	28	3	60	H	Limit	Excellent protein but high butter/cream; ask for less cream
Dal Makhani (1 cup)	200g	300	10	28	16	7	30	M-H	Moderate	High fat from butter/cream; portion control; good protein
Paneer Butter Masala (200g)	200g	450	14	18	36	2	40	M	Limit	Very high fat from cream and butter; occasional
Shahi Paneer (200g)	200g	520	15	20	42	2	40	M	Limit	Richest paneer dish; very high fat

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Tandoori Chicken (4 pieces)	200g	320	48	6	12	0	0	H	Good	Best restaurant chicken for WL; high protein, low fat
Chicken Tikka (6 pieces)	180g	270	40	5	10	0	0	H	Good	Grilled; good protein; watch marinade oil
Rogan Josh (mutton, 200g)	200g	390	34	8	25	1	0	H	Moderate	High fat mutton; good protein; portion control
Fish Tikka (4 pieces)	180g	200	32	4	7	0	0	H	Excellent	Excellent restaurant choice for WL; high protein
Veg Biryani (1 plate)	350g	420	10	72	12	5	62	M	Moderate	Rice + ghee; protein from veg; add raita for protein
Chicken Biryani (1 plate)	350g	520	28	68	18	3	62	M-H	Moderate	Better protein ratio than veg biryani; portion control
Raita (1 bowl)	150g	80	4.5	9	2.5	0.5	35	M-H	Excellent	Always add raita to restaurant meal for protein + probiotics
Pav Bhaji (1 plate + 2 pav)	400g	580	12	85	22	8	65	M-H	Limit	Very high butter; occasional treat
Chicken Wrap (1 wrap)	220g	380	24	42	14	3	58	H	Good	Balanced; good protein; better than pizza or burger
Masala Dosa (1 plate)	200g	280	7	48	9	3	70	M	Moderate	Better than Bhatura; watch chutney portions
Idli Sambar (3 idli)	350g	230	10	44	2	5	62	H	Excellent	Best South Indian restaurant option for WL

WL-FD-011 MOD 03 · Food Database › Beverages

GOALS Weight Loss | Fat Loss | Weight Maintenance | General Health

USERS Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student

DIFF Beginner

TAGS beverages, chai calories, cold drink, juice calories, protein shake, what to drink for weight loss

Beverages

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Water	Any amount	0	0	0	0	0	0	H	Excellent	Always the best choice; calorie-free
Black Tea (no sugar)	1 cup/240ml	5	0.5	1	0	0	0	M	Excellent	Antioxidants; minimal calories; best tea choice for WL
Tea with milk + 1 tsp sugar	1 cup/240ml	45	1.5	7.5	1.5	0	30	M	Good	300ml milk tea = 75+ kcal; 3 cups = 135–225 kcal/day
Tea with milk + 2 tsp sugar	1 cup/240ml	70	1.5	11.5	1.5	0	45	L	Moderate	Switch to 0 sugar or 1 tsp maximum for WL
Coffee (black, no sugar)	1 cup/240ml	5	0.3	0.5	0	0	0	M	Excellent	Pre-workout; fat oxidation boost; no calories
Coffee with milk (no sugar)	1 cup/240ml	35	1.5	3	1.8	0	30	M	Excellent	Good choice; limit to 1–2 cups/day (caffeine)
Coconut Water (1 glass)	250ml	46	1.7	8.9	0.5	2.6	54	M	Excellent	Natural electrolytes; low calorie; excellent hydration
Nimbu Pani (no sugar)	1 glass/250ml	12	0.3	3.5	0.1	0	0	M	Excellent	Electrolytes; vitamin C; perfect WL beverage
Nimbu Pani (1 tsp sugar)	1 glass/250ml	32	0.3	8	0.1	0	0	M	Good	Small amount of sugar negligible; better than cold drink

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Chaas/Buttermilk (plain)	1 glass/240ml	55	3.5	5.5	1.8	0	35	M-H	Excellent	Excellent weight loss drink; probiotics; low calorie
Lassi (sweet)	1 glass/300ml	250	7.5	38	6	0	55	L	Limit	High sugar; avoid for WL; choose chaas instead
Mango Juice (packaged)	200ml	115	0.3	28	0.3	0.4	70	L	Limit	High sugar; no fiber; empty calories
Orange Juice (packaged)	200ml	88	1.4	20.8	0.4	0.5	57	L	Limit	Better than mango but eat whole orange for fiber
Sugarcane Juice (1 glass)	250ml	180	0.4	45	0.2	0	70	L	Limit	High sugar; occasional treat; massive portion sugar
Cold Drink/Soda (330ml)	1 can	139	0	35	0	0	63	L	Limit	No nutrition; 9 tsp sugar; avoid completely
Protein Shake (whey, 30g)	with 200ml water	120	24	3	1.5	0	15	H	Excellent	Best supplement for hitting protein targets
Buttermilk with jeera + salt	1 glass/240ml	50	3.5	5	1.8	0	35	H	Excellent	Digestive; probiotic; very low calorie
Green Tea	1 cup/240ml	3	0.5	0.5	0	0	0	M	Excellent	Modest fat oxidation benefit; antioxidant; good choice
Aam Panna (no sugar)	1 glass/250ml	35	0.3	8.5	0.1	0.5	35	M	Good	Raw mango; vitamin C; electrolytes; summer drink

WL-FD-012 MOD 03 · Food Database › Sweets and Desserts

GOALS Weight Loss | Fat Loss | Weight Maintenance | General Health

USERS Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student

DIFF Beginner

TAGS sweets, gulab jamun, kheer, Indian desserts, festival food calories

Sweets and Desserts

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Gulab Jamun (1 piece)	1 piece/45g	140	2.5	25	4.5	0.3	70	L	Limit	High sugar + fat; occasional 1 piece maximum
Rasmalai (1 piece)	1 piece/60g	100	4	16	3	0	60	L	Limit	Lower calorie than gulab jamun but still sugar-heavy
Kheer (1 cup)	200ml	250	8	38	8	0.5	65	L	Limit	Milk base has protein but high sugar
Halwa (Atta, 100g)	100g	390	5	55	18	2	70	L	Limit	Very high ghee and sugar; occasional only
Dark Chocolate (70%+)	2 squares/20g	114	1.6	10.4	7.5	2.3	23	M	Moderate	Better choice for sweet cravings; portion controlled
Mishti Doi (1 cup)	150g	200	6.5	32	5.5	0	60	L	Limit	High sugar; regular curd with 1 tsp honey is better
Chhena Podo (100g)	100g	285	11.5	38	11	0	60	L	Limit	Traditional; high sugar; occasional
Peda (1 piece)	1 piece/30g	115	3	18	4	0	65	L	Limit	Festival food; 1 piece maximum
Jalebi (1 piece)	1 piece/30g	115	1	27	2.5	0	80	L	Limit	Deep fried + sugar syrup; very high GI; occasional

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Payasam (1 cup)	200ml	280	7.5	44	9	1	60	L	Limit	High fat + sugar from milk + condensed milk; occasional
Rasgulla (1 piece)	1 piece/50g	110	3.5	22	1.5	0	60	L	Limit	Lower fat than other sweets; sugar is concern
Coconut Ladoo (1)	1 ladoo/30g	118	1.5	14	7	2	45	M	Limit	High fat from coconut; 1 is reasonable at festival

WL-FD-013 MOD 03 · Food Database › Fast Food

GOALS Weight Loss | Fat Loss | Weight Maintenance | General Health

USERS Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student

DIFF Beginner

TAGS fast food, McDonald's, pizza, burger, chips, fast food calories

Fast Food

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
McAloo Tikki Burger	1 burger/150g	300	6	47	11	2.5	68	M	Limit	Lower calorie of McDonald's; limited protein
McVeggie Burger	1 burger/200g	350	8	53	12	3	65	M	Limit	More protein than Aloo Tikki; better choice if eating at McDonald's
Chicken Maharaja Mac	1 burger/260g	625	30	52	31	3	65	M-H	Limit	Very calorie-dense; half can be eaten; save rest
Veg Pizza (1 slice, thin)	1 slice/80g	185	7.5	26	6.5	2	65	M	Limit	Thin crust better; 2 slices maximum
Chicken Pizza (1 slice, thin)	1 slice/90g	220	11	26	8	2	65	M	Limit	Better protein than veg pizza; still calorie dense
French Fries (medium)	117g	365	4.5	48	17	4	75	L	Limit	High fat; low protein; avoid or share
Veg Biryani (Swiggy)	1 portion/350g	550	11	88	17	5	62	M	Limit	Estimate; restaurant version has more oil
Chicken Biryani (delivery)	1 portion/400g	680	32	82	24	4	62	M-H	Moderate	Better protein ratio; still high calorie from oil
Paneer Tikka Roll (1 roll)	1 roll/200g	380	14	50	15	3	62	M-H	Moderate	Good protein from paneer; manageable calories
Chicken Kathi Roll (1 roll)	1 roll/220g	420	22	52	15	3	60	H	Good	Good protein; filling; better fast food choice for WL
Shawarma (chicken)	1 wrap/250g	450	25	48	18	3	60	H	Moderate	Good protein but high oil; ask for less sauce
Idli Sambar (2 idli, restaurant)	2 idli + sambar	190	9	36	2	4	62	H	Excellent	Best choice at South Indian restaurant fast food
Puri Bhaji (2 puri)	2 puris/300g	480	9	63	22	4	70	M	Limit	High fat from puri frying; limit restaurant version
Dahi Vada (2 pieces)	2 pieces/200g	320	11	42	12	3	55	M-H	Moderate	Fermented dal + yogurt; better protein but calorie dense

WL-FD-014 MOD 03 · Food Database › Protein Powders

GOALS Weight Loss | Fat Loss | Weight Maintenance | General Health

USERS Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student

DIFF Beginner

TAGS protein powder, whey protein, pea protein, MuscleBlaze, supplement nutrition

Protein Powders

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Whey Protein Concentrate	30g scoop	130	24	5	1.5	0	15	H	Excellent	Most affordable; 80% protein; slight lactose
Whey Protein Isolate	30g scoop	115	27	1	0.5	0	15	H	Excellent	Higher purity; lactose-free; higher cost
Casein Protein	30g scoop	115	24	4	1	0	15	H	Excellent	Slow-digesting; best before bed; high satiety
Soy Protein Isolate	30g scoop	110	25	2	0.5	0.5	15	H	Excellent	Complete plant protein; all amino acids; good for vegans
Pea Protein Isolate	30g scoop	110	24	2	2	1	15	H	Excellent	Most digestible plant protein; good amino profile
Rice Protein	30g scoop	115	22	4.5	2	0.5	15	M-H	Good	Incomplete protein; best combined with pea protein
Hemp Protein	30g scoop	120	15	7	4	4	15	H	Good	Complete protein; high fiber; omega-3; lower protein %
MuscleBlaze Whey Active (India)	33g scoop	130	25	4	2	0	15	H	Excellent	India's best-selling; FSSAI certified; affordable
MuscleBlaze Raw Whey	30g scoop	113	27	1.2	0.6	0	15	H	Excellent	Unflavored; mix with anything; high purity
Nakpro Platinum Whey	33g scoop	133	25	4.8	1.5	0	15	H	Excellent	Affordable India brand; FSSAI certified; 77% protein

WL-FD-015 MOD 03 · Food Database › Condiments, Oils and Spices

GOALS Weight Loss | Fat Loss | Weight Maintenance | General Health

USERS Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student

DIFF Beginner

TAGS cooking oil, ghee calories, spices, chutney, condiments

Condiments, Oils and Spices

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Olive Oil	1 tbsp/14g	119	0	0	13.5	0	0	L	Moderate	Healthy MUFA but 119 kcal/tbsp — measure carefully
Mustard Oil	1 tbsp/14g	120	0	0	14	0	0	L	Moderate	Good omega-3/6 ratio for Indian cooking; measure carefully
Coconut Oil	1 tbsp/14g	121	0	0	14	0	0	L	Moderate	Medium-chain fats; no WL magic; same calories as other oils
Ghee	1 tsp/5g	45	0	0	5	0	0	L	Limit	Limit to 1 tsp during WL phase
Tomato Ketchup	1 tbsp/17g	20	0.4	4.9	0.1	0.2	55	L	Moderate	High fructose corn syrup; limit; use tomato paste instead
Green Chutney (mint+coriander)	2 tbsp/30g	15	0.5	2	0.5	0.8	20	M	Excellent	Very low calorie; use generously as condiment
Coconut Chutney	2 tbsp/30g	60	1	4	5	1.7	40	M	Moderate	High fat from coconut; limit portion at restaurant
Sambar (1 cup)	200ml	90	4	14	2	4	40	H	Excellent	Dal-based; good protein and fiber; excellent accompaniment
Raita (plain)	100g	55	3	5	2	0.5	35	H	Excellent	Always add to meals; protein + probiotic

Food	Serving	kcal	P(g)	C(g)	F(g)	Fb(g)	GI	Sat	WL	Notes/Tips
Vinegar	1 tbsp/15ml	3	0	0.9	0	0	0	M	Excellent	Reduces glycemic response of meal; use freely
Lemon Juice	juice of 1/30ml	8	0.3	2.5	0.1	0	0	M	Excellent	Enhances iron absorption; flavor booster; zero calorie effectively
Turmeric	1 tsp/3g	9	0.3	1.4	0.3	0.5	0	M	Excellent	Anti-inflammatory (curcumin); use 1 tsp/day regularly
Cumin (jeera)	1 tsp/3g	8	0.4	0.9	0.5	0.2	0	M	Excellent	Digestive; thermogenic; no significant calorie concern
Cinnamon	1 tsp/2.5g	6	0.1	1.8	0.03	1.2	5	M	Excellent	Lowers blood sugar response; add to oats, chai, coffee

WL-FD-016 MOD 03 · Food Swap Reference › High-Calorie to Low-Calorie Swaps

GOALS Weight Loss | Fat Loss

USERS Adult · Teen · Senior · Male · Female · Student · Vegetarian · Non-Vegetarian · Vegan · Hostel Student

DIFF Beginner

TAGS food swaps, calorie swap, lower calorie alternatives, food substitutions

DEFINITION

Smart food swaps reduce calorie intake by 300–700 kcal/day without reducing food volume or satisfaction. This section provides direct swap recommendations for the highest-calorie foods in the Indian diet.

WHY IT MATTERS

- High-calorie habits often have direct, satisfying low-calorie alternatives.
- Swaps are more sustainable than complete elimination.

KEY FACTS

- SWAP: White rice (260 kcal/cup) → Cauliflower rice (25 kcal/cup): saves 235 kcal (use 1:1 swap partially).
- SWAP: Sweet lassi (250 kcal) → Chaas/buttermilk (55 kcal): saves 195 kcal.
- SWAP: Chips 30g packet (162 kcal) → Roasted chana 30g (116 kcal): saves 46 kcal + gains protein.
- SWAP: Full cream milk (149 kcal/cup) → Toned milk (130 kcal): saves 19 kcal.
- SWAP: Ghee on roti 1 tbsp (112 kcal) → No ghee: saves 112 kcal per roti.
- SWAP: Fried egg (115 kcal) → Boiled egg (78 kcal): saves 37 kcal per egg.
- SWAP: Butter naan (320 kcal) → Plain roti (85 kcal): saves 235 kcal.
- SWAP: Cold drink 330ml (139 kcal) → Plain water (0): saves 139 kcal per drink.
- SWAP: 3 sweet chai/day (210 kcal) → 3 black tea (15 kcal): saves 195 kcal/day.
- SWAP: Puri-bhaji (480 kcal) → Idli-sambar (230 kcal): saves 250 kcal for same meal type.

ACTION STEPS

1. Identify user's 3 highest-calorie daily habits from food diary.
2. Offer a direct swap for each — same taste profile but fewer calories.
3. Implement 1 swap per week for sustainable transition.
4. Track savings: 3 swaps at 200 kcal each = 600 kcal/day = 0.6 kg fat less per month.

AI COACHING NOTES

- Present swaps as upgrades, not deprivation — 'this gives you the same satisfaction for fewer calories'.
- Hostel students: chaas for lassi, roasted chana for namkeen, black tea for sweet chai are immediate wins.

Complete Recipe Library

WL-R-000	MOD 04 — Recipe Library · Recipe Library Overview › Index
GOALS	Weight Loss Fat Loss General Health
USERS	Adult · Teen · Student · Office Worker · Hostel Student · Vegetarian · Non Vegetarian · Vegan
DIFF	Beginner
TAGS	recipes, meal ideas, Indian food, weight loss meals, quick recipes

DEFINITION

Complete recipe library with 118 recipes across 8 categories for every lifestyle.

WHY IT MATTERS

- Every calorie target needs practical, tasty meals to be sustainable.
- Indian food can be adapted for fat loss without sacrificing culture.

KEY FACTS

- Total recipes: 118
- Breakfast: 29 | Lunch: 19 | Dinner: 15
- Snacks: 15 | Smoothies: 11 | Protein Meals: 10
- Budget/Hostel: 10 | No-Cook: 9
- All recipes include: macros, cost, prep time, suitable users.

ACTION STEPS

1. Find your calorie target in Module 05.
2. Select 3-4 breakfast, lunch, dinner options that suit your lifestyle.
3. Meal prep 2-3 recipes on Sunday for the week.
4. Keep 3-4 no-cook options for emergency meals.

High-Protein Oats Bowl

CAT: Breakfast | KCAL: 380 | P: 28g C: 45g F: 9g Fiber: 6g | SERVING: 1 bowl (350g) | COST: █40 | PREP: 5 min COOK: 5 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Student, Budget

INGREDIENTS:

- 1 cup rolled oats
- 1 scoop whey protein (30g)
- 1 tsp chia seeds
- 1 tbsp peanut butter
- 1 banana
- 300ml water or almond milk

INSTRUCTIONS:

1. Boil water or milk.
2. Add oats, cook 3-4 min stirring.
3. Remove from heat, mix in protein powder.
4. Top with chia, peanut butter, sliced banana.

TIPS: Add protein powder off heat to avoid clumping. Prep oats night before for faster morning.

Moong Dal Chilla

CAT: Breakfast | KCAL: 260 | P: 18g C: 32g F: 6g Fiber: 8g | SERVING: 3 chillas | COST: █25 | PREP: 20 min COOK: 10 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget, Student

INGREDIENTS:

- 1 cup moong dal (soaked 4hrs)

- 1 green chilli
- 1 tsp ginger
- Salt
- Cumin
- 1 tsp oil
- Coriander

INSTRUCTIONS:

1. Blend soaked moong dal with chilli, ginger, salt to smooth batter.
2. Add cumin and coriander.
3. Pour thin circles on non-stick pan with 1/4 tsp oil.
4. Cook 2 min each side till golden.

TIPS: Add grated vegetables for extra nutrition. Serve with green chutney.

Egg White Omelette

CAT: Breakfast | KCAL: 180 | P: 24g C: 4g F: 5g Fiber: 2g | SERVING: 1 omelette | COST: █30 | PREP: 5 min COOK: 5 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Non Vegetarian, Student, Office Worker

INGREDIENTS:

- 4 egg whites
- 1 whole egg
- Spinach 30g
- Tomato 50g
- Onion 30g
- Salt, pepper
- 1/2 tsp oil

INSTRUCTIONS:

1. Whisk eggs with salt and pepper.
2. Sauté vegetables 2 min in non-stick pan.
3. Pour egg mixture over vegetables.
4. Fold and serve.

TIPS: Using 4 whites + 1 whole egg balances protein with healthy fat and yolk nutrients.

Greek Yogurt Parfait

CAT: Breakfast | KCAL: 290 | P: 22g C: 38g F: 4g Fiber: 3g | SERVING: 1 glass | COST: █60 | PREP: 5 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, No Cook, Student

INGREDIENTS:

- 200g Greek yogurt / hung curd
- 1 tbsp honey
- 30g granola or muesli
- Mixed berries or banana
- 1 tsp chia seeds

INSTRUCTIONS:

1. Layer yogurt in glass.
2. Add granola.
3. Top with fruits and chia.
4. Drizzle honey.

TIPS: Use hung curd for same protein as Greek yogurt at lower cost.

Poha with Peanuts

CAT: Breakfast | KCAL: 310 | P: 10g C: 52g F: 8g Fiber: 4g | SERVING: 1 plate (200g) | COST: █20 | PREP: 5 min COOK: 10 min |
DIFF: Beginner | WL: Good
SUITABLE FOR: Vegetarian, Vegan, Student, Budget

INGREDIENTS:

- 1.5 cups thick poha
- 2 tbsp peanuts
- 1 onion
- 1 potato (small)
- Mustard seeds, curry leaves
- Turmeric, salt
- Lime, coriander

INSTRUCTIONS:

1. Wash poha, drain well.
2. Heat oil, add mustard seeds, curry leaves.
3. Add onion and potato, cook 5 min.
4. Add peanuts, turmeric.
5. Add poha, mix gently, cook 2 min.
6. Squeeze lime, garnish with coriander.

TIPS: Add 1 cup extra peas or edamame to boost protein significantly.

Besan Chilla (Chickpea Pancake)

CAT: Breakfast | KCAL: 240 | P: 14g C: 28g F: 7g Fiber: 6g | SERVING: 2 chillas | COST: █20 | PREP: 5 min COOK: 8 min | DIFF: Beginner | WL: Excellent
SUITABLE FOR: Vegetarian, Vegan, Budget, Student

INGREDIENTS:

- 1 cup besan
- 1/2 cup water
- Onion, tomato, green chilli (chopped)
- Ajwain, cumin, salt
- 1 tsp oil

INSTRUCTIONS:

1. Mix besan with water to smooth batter.
2. Add vegetables and spices.
3. Pour on hot non-stick pan with 1/4 tsp oil.
4. Cook 2-3 min each side.

TIPS: High satiety, low calorie, high fiber — ideal weight-loss breakfast.

Upma with Vegetables

CAT: Breakfast | KCAL: 280 | P: 9g C: 44g F: 7g Fiber: 5g | SERVING: 1 bowl (250g) | COST: █25 | PREP: 5 min COOK: 12 min | DIFF: Beginner | WL: Good
SUITABLE FOR: Vegetarian, Vegan, Student, Budget

INGREDIENTS:

- 3/4 cup semolina (rava)
- Mixed vegetables 1 cup
- Mustard seeds, urad dal
- Onion, green chilli
- 2 cups water
- Salt, oil

INSTRUCTIONS:

1. Dry roast rava 3-4 min, set aside.
2. Temper mustard seeds, urad dal in 1 tsp oil.
3. Add onion, chilli, vegetables. Cook 3 min.

4. Add water, salt. Boil.
5. Add rava slowly, stir constantly.
6. Cook 3 min till set.

TIPS: Vegetable ratio should equal or exceed rava for better nutrition and satiety.

Protein Banana Pancakes

CAT: Breakfast | KCAL: 320 | P: 24g C: 38g F: 8g Fiber: 3g | SERVING: 4 pancakes | COST: █35 | PREP: 5 min COOK: 10 min | DIFF: Beginner | WL: Good
SUITABLE FOR: Vegetarian, Student

INGREDIENTS:

- 2 ripe bananas
- 2 eggs
- 1 scoop protein powder
- 1 tsp cinnamon
- Coconut oil spray

INSTRUCTIONS:

1. Mash bananas thoroughly.
2. Mix in eggs, protein powder, cinnamon.
3. Pour small circles on non-stick pan.
4. Cook 2-3 min each side on medium heat.

TIPS: No added sugar needed — banana provides natural sweetness.

Sprouts Salad Breakfast Bowl

CAT: Breakfast | KCAL: 210 | P: 16g C: 28g F: 4g Fiber: 9g | SERVING: 1 bowl | COST: █20 | PREP: 10 min COOK: 0 min | DIFF: Beginner | WL: Excellent
SUITABLE FOR: Vegetarian, Vegan, No Cook, Budget

INGREDIENTS:

- 1 cup mixed sprouts (moong, chana)
- 1 tomato
- 1/2 cucumber
- Onion
- Green chilli
- Lime
- Chaat masala, salt
- Coriander

INSTRUCTIONS:

1. Combine sprouts and chopped vegetables.
2. Add lime juice, salt, chaat masala.
3. Toss well and garnish.

TIPS: Sprouts contain living enzymes and are the most nutrient-dense budget food.

Idli with Sambhar

CAT: Breakfast | KCAL: 240 | P: 10g C: 44g F: 3g Fiber: 5g | SERVING: 3 idlis + sambhar | COST: █30 | PREP: 5 min COOK: 15 min | DIFF: Beginner | WL: Good
SUITABLE FOR: Vegetarian, Vegan, South Indian, Student

INGREDIENTS:

- 3 idlis (ready or from batter)
- 1 cup sambhar
- Green chutney

INSTRUCTIONS:

1. Steam idlis 10-12 min.

2. Heat sambhar.
3. Serve with chutney.

TIPS: Low fat, moderate protein, fermented — good gut health. Add extra sambhar vegetables.

Dosa with Chutney

CAT: Breakfast | KCAL: 280 | P: 8g C: 48g F: 6g Fiber: 3g | SERVING: 2 thin dosas | COST: ■25 | PREP: 5 min COOK: 10 min | DIFF: Intermediate | WL: Good

SUITABLE FOR: Vegetarian, Vegan, South Indian

INGREDIENTS:

- Dosa batter 1 cup
- Coconut chutney 3 tbsp
- 1/2 tsp oil
- Sambhar optional

INSTRUCTIONS:

1. Heat pan, pour batter in thin circle.
2. Drizzle small oil on sides.
3. Cook till crispy, flip 1 min.
4. Serve with chutney.

TIPS: Thin dosa > thick dosa for weight loss. Avoid ghee dosa.

Overnight Oats

CAT: Breakfast | KCAL: 350 | P: 18g C: 48g F: 8g Fiber: 7g | SERVING: 1 jar | COST: ■45 | PREP: 5 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, No Cook, Student, Office Worker

INGREDIENTS:

- 1/2 cup oats
- 1 cup milk or almond milk
- 1/2 scoop protein powder
- 1 tbsp chia seeds
- 1 tsp honey
- Fruits to top

INSTRUCTIONS:

1. Mix oats, milk, protein, chia, honey in jar.
2. Stir well.
3. Refrigerate overnight (minimum 4 hrs).
4. Top with fruits and serve cold.

TIPS: Prep 5 jars Sunday for the full week — zero morning effort.

Peanut Butter Toast

CAT: Breakfast | KCAL: 300 | P: 14g C: 32g F: 14g Fiber: 4g | SERVING: 2 slices | COST: ■30 | PREP: 3 min COOK: 3 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, No Cook, Student, Hostel

INGREDIENTS:

- 2 slices whole wheat bread
- 2 tbsp natural peanut butter
- 1 banana or berries
- Optional: 1 tsp honey

INSTRUCTIONS:

1. Toast bread.
2. Spread peanut butter.
3. Top with sliced fruits.

TIPS: Use natural PB (only peanuts, no added oil/sugar). Whole wheat over maida.

Paneer Bhurji

CAT: Breakfast | KCAL: 290 | P: 22g C: 8g F: 18g Fiber: 2g | SERVING: 1 serving (150g) | COST: █50 | PREP: 5 min COOK: 10 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Keto-friendly

INGREDIENTS:

- 150g paneer (crumbled)
- 1 onion
- 1 tomato
- 1 green chilli
- Turmeric, cumin, salt
- 1 tsp oil
- Coriander

INSTRUCTIONS:

1. Heat oil, add cumin.
2. Add onion, cook 3 min.
3. Add tomato, chilli, spices.
4. Add crumbled paneer, stir 2 min.
5. Garnish with coriander.

TIPS: Excellent high-protein breakfast. Low carb, very filling.

Sattu Drink

CAT: Breakfast | KCAL: 180 | P: 12g C: 28g F: 2g Fiber: 6g | SERVING: 1 glass (300ml) | COST: █15 | PREP: 2 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget, No Cook

INGREDIENTS:

- 3 tbsp roasted chana sattu
- 300ml water
- Pinch black salt
- 1/2 tsp cumin powder
- Lemon juice
- Optional: jaggery 1 tsp

INSTRUCTIONS:

1. Mix sattu in water thoroughly.
2. Add salt, cumin, lemon.
3. Stir or shake well.
4. Drink immediately.

TIPS: Bihar's traditional superfood. High protein, high fiber, cheap, cooling. Perfect summer breakfast.

Rava Dhokla

CAT: Breakfast | KCAL: 200 | P: 8g C: 36g F: 4g Fiber: 3g | SERVING: 4 pieces | COST: █20 | PREP: 15 min COOK: 15 min | DIFF: Intermediate | WL: Good

SUITABLE FOR: Vegetarian, Budget, Student

INGREDIENTS:

- 1 cup semolina
- 3/4 cup curd
- Salt, sugar
- Eno (1 tsp)
- Tempering: mustard, curry leaves
- Green chutney

INSTRUCTIONS:

1. Mix rava, curd, salt, sugar. Rest 10 min.
2. Add Eno, stir quickly.
3. Pour in greased plate, steam 12-15 min.
4. Temper and pour over.
5. Slice and serve.

TIPS: Light, low-fat, good satiety. Better than fried breakfast options.

Methi Paratha

CAT: Breakfast | KCAL: 220 | P: 7g C: 34g F: 6g Fiber: 5g | SERVING: 2 parathas | COST: ■20 | PREP: 10 min COOK: 10 min | DIFF: Beginner | WL: Good
SUITABLE FOR: Vegetarian, Vegan, Budget

INGREDIENTS:

- 1 cup whole wheat flour
- 1/2 cup fresh methi leaves
- Salt, ajwain, turmeric
- Water to knead
- 1 tsp oil

INSTRUCTIONS:

1. Mix flour, methi, spices.
2. Knead to soft dough with water.
3. Roll thin parathas.
4. Cook on dry tawa, light brush of oil.

TIPS: Methi (fenugreek) stabilizes blood sugar and aids fat loss.

Boiled Eggs with Soldiers

CAT: Breakfast | KCAL: 200 | P: 18g C: 14g F: 10g Fiber: 2g | SERVING: 2 eggs + 1 toast | COST: ■25 | PREP: 2 min COOK: 8 min | DIFF: Beginner | WL: Excellent
SUITABLE FOR: Non Vegetarian, Student, Hostel, Budget

INGREDIENTS:

- 2 eggs
- 1 whole wheat toast
- Salt, pepper
- Optional: hot sauce

INSTRUCTIONS:

1. Boil eggs 7-8 min for hard boil.
2. Toast bread, cut into strips.
3. Season eggs with salt and pepper.

TIPS: Best budget protein meal. Prep in bulk — boil 6 eggs Sunday for the week.

Muesli with Milk

CAT: Breakfast | KCAL: 310 | P: 12g C: 52g F: 6g Fiber: 5g | SERVING: 1 bowl | COST: ■45 | PREP: 2 min COOK: 0 min | DIFF: Beginner | WL: Good
SUITABLE FOR: Vegetarian, No Cook, Student, Hostel

INGREDIENTS:

- 50g muesli (no added sugar)
- 200ml skimmed milk
- 1 banana or apple
- 1 tsp honey

INSTRUCTIONS:

1. Pour muesli in bowl.

2. Add cold milk.
3. Top with fruit.
4. Add honey if desired.

TIPS: Choose muesli with <10g sugar per 100g. Avoid "diet" mueslis with dried fruit overload.

Pesarattu (Green Moong Dosa)

CAT: Breakfast | KCAL: 230 | P: 14g C: 34g F: 4g Fiber: 7g | SERVING: 2 pesarattu | COST: ■20 | PREP: 15 min (+ soak overnight)

COOK: 10 min | DIFF: Intermediate | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, South Indian, Budget

INGREDIENTS:

- 1 cup whole green moong (soaked overnight)
- Ginger, green chilli
- Salt
- Onion, grated (optional)
- 1 tsp oil

INSTRUCTIONS:

1. Blend soaked moong with ginger, chilli, salt.
2. Spread thin on hot pan like dosa.
3. Sprinkle onion, drizzle oil.
4. Cook 2-3 min, fold and serve.

TIPS: Andhra superfood — high protein, high fiber, complete amino acid profile.

Banana Oat Smoothie Bowl

CAT: Breakfast | KCAL: 340 | P: 20g C: 50g F: 6g Fiber: 6g | SERVING: 1 bowl | COST: ■50 | PREP: 5 min COOK: 0 min | DIFF:

Beginner | WL: Good

SUITABLE FOR: Vegetarian, Student, No Cook

INGREDIENTS:

- 1 frozen banana
- 1/2 cup oats
- 1 scoop protein powder
- 200ml milk
- Toppings: granola, nuts, seeds

INSTRUCTIONS:

1. Blend banana, oats, protein, milk thick.
2. Pour in bowl.
3. Add toppings.

TIPS: Freeze bananas in batches. Thick smoothie bowls are more satiating than liquid smoothies.

Curd Rice

CAT: Breakfast | KCAL: 250 | P: 10g C: 42g F: 5g Fiber: 2g | SERVING: 1 bowl (200g) | COST: ■20 | PREP: 5 min COOK: 5 min | DIFF:

Beginner | WL: Good

SUITABLE FOR: Vegetarian, South Indian, Student

INGREDIENTS:

- 1/2 cup cooked rice
- 200g curd
- Salt
- Mustard seeds, curry leaves
- Ginger, green chilli
- Coriander, pomegranate

INSTRUCTIONS:

1. Mix rice and curd well.

2. Temper mustard, curry leaves, ginger.
3. Pour over rice.
4. Garnish and serve.

TIPS: Probiotics from curd support gut health and weight loss. Add grated cucumber for volume.

Egg Bhurji Roll

CAT: Breakfast | KCAL: 340 | P: 20g C: 34g F: 14g Fiber: 3g | SERVING: 1 roll | COST: █40 | PREP: 5 min COOK: 8 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Non Vegetarian, Student, Office Worker

INGREDIENTS:

- 2 eggs
- 1 whole wheat roti
- Onion, tomato, chilli
- Salt, chaat masala
- 1 tsp oil
- Green chutney

INSTRUCTIONS:

1. Scramble eggs with onion, tomato, chilli.
2. Warm roti.
3. Spread chutney on roti.
4. Add egg bhurji, roll tight.

TIPS: High protein portable breakfast. Prep eggs the night before.

Palak Smoothie

CAT: Breakfast | KCAL: 180 | P: 10g C: 24g F: 3g Fiber: 5g | SERVING: 1 glass (350ml) | COST: █30 | PREP: 5 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, No Cook

INGREDIENTS:

- 2 cups baby spinach
- 1 banana
- 1 scoop protein powder
- 250ml almond milk
- 1 tsp flaxseeds
- Ice cubes

INSTRUCTIONS:

1. Blend all ingredients until smooth.
2. Serve immediately.

TIPS: You cannot taste spinach. Excellent low-calorie, high-micronutrient breakfast.

Rajma Chaat Bowl

CAT: Breakfast | KCAL: 280 | P: 16g C: 42g F: 4g Fiber: 12g | SERVING: 1 bowl | COST: █25 | PREP: 5 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget, No Cook

INGREDIENTS:

- 1 cup cooked rajma
- Onion, tomato, cucumber
- Lime, salt, chaat masala
- Coriander

INSTRUCTIONS:

1. Mix all ingredients.
2. Toss with lime and spices.

3. Serve cold.

TIPS: Overnight cooked rajma — fastest high-protein breakfast.

Masala Oats

CAT: Breakfast | KCAL: 260 | P: 10g C: 40g F: 6g Fiber: 6g | SERVING: 1 bowl | COST: ■25 | PREP: 5 min COOK: 8 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Vegan, Student, Hostel

INGREDIENTS:

- 3/4 cup oats
- 1 cup water
- Onion, tomato, peas
- Turmeric, cumin, salt
- 1 tsp oil

INSTRUCTIONS:

1. Sauté onion in oil 2 min.
2. Add tomato, peas, spices.
3. Add water, boil.
4. Add oats, cook 3-4 min stirring.

TIPS: Savory oats for those who dislike sweet breakfast. Very filling.

Protein Dahi (Curd Bowl)

CAT: Breakfast | KCAL: 200 | P: 20g C: 18g F: 4g Fiber: 1g | SERVING: 1 bowl | COST: ■35 | PREP: 3 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, No Cook, Student

INGREDIENTS:

- 250g hung curd
- 1 scoop protein powder
- 1 tsp honey
- Berries or apple
- Cinnamon

INSTRUCTIONS:

1. Mix hung curd and protein powder.
2. Add honey and cinnamon.
3. Top with fruit.

TIPS: Hung curd has 15g protein per 100g — rivals Greek yogurt at half the price.

Stuffed Paratha

CAT: Breakfast | KCAL: 350 | P: 10g C: 50g F: 12g Fiber: 6g | SERVING: 2 parathas | COST: ■30 | PREP: 15 min COOK: 10 min | DIFF: Intermediate | WL: Good

SUITABLE FOR: Vegetarian, Student, Budget

INGREDIENTS:

- 1 cup whole wheat flour
- Filling: grated carrot + paneer 50g + spices
- 1 tsp ghee
- Salt, ajwain

INSTRUCTIONS:

1. Knead soft dough.
2. Mix carrot, paneer, spices for filling.
3. Roll small circle, fill, seal, roll flat.
4. Cook on tawa with minimal ghee.

TIPS: Homemade parathas with vegetable filling are healthy. Avoid heavy oil/ghee parathas outside.

Milk + Banana

CAT: Breakfast | KCAL: 230 | P: 10g C: 40g F: 4g Fiber: 3g | SERVING: 1 glass + 1 banana | COST: █20 | PREP: 1 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, No Cook, Student, Hostel, Budget

INGREDIENTS:

- 250ml full fat milk
- 1 banana

INSTRUCTIONS:

1. Drink milk, eat banana.

TIPS: Simplest emergency protein breakfast. Add 1 tsp cocoa powder for flavor.

Dal Tadka + Roti

CAT: Lunch | KCAL: 380 | P: 18g C: 60g F: 7g Fiber: 10g | SERVING: 1 bowl dal + 2 roti | COST: █30 | PREP: 10 min COOK: 25 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget, Student

INGREDIENTS:

- 1/2 cup toor dal
- Onion, tomato, garlic, ginger
- Cumin, mustard, hing
- Turmeric, chilli powder
- Salt, coriander
- 2 whole wheat rotis

INSTRUCTIONS:

1. Pressure cook dal 3-4 whistles.
2. Make tadka: heat oil, add cumin, onion, tomato, spices.
3. Combine dal and tadka.
4. Serve with hot roti.

TIPS: Best weight-loss lunch in India. Add vegetables to dal for extra fiber.

Chicken Breast Rice Bowl

CAT: Lunch | KCAL: 480 | P: 45g C: 50g F: 8g Fiber: 4g | SERVING: 1 large bowl | COST: █80 | PREP: 10 min COOK: 25 min | DIFF: Intermediate | WL: Excellent

SUITABLE FOR: Non Vegetarian, Office Worker, Meal Prep

INGREDIENTS:

- 150g chicken breast
- 1 cup brown rice or basmati
- Mixed vegetables 100g
- 1 tsp olive oil
- Herbs, salt, pepper
- Lime

INSTRUCTIONS:

1. Cook rice.
2. Season chicken, pan fry or bake 15-20 min.
3. Steam or stir-fry vegetables.
4. Assemble bowl with rice base, chicken, veggies.

TIPS: Ideal macro split for fat loss. Meal prep 4 servings Sunday.

Rajma Chawal

CAT: Lunch | KCAL: 420 | P: 18g C: 72g F: 6g Fiber: 14g | SERVING: 1 plate | COST: █35 | PREP: 10 min COOK: 30 min (with overnight soak) | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget, Student

INGREDIENTS:

- 1/2 cup rajma (soaked overnight)
- 1 cup basmati rice
- Onion, tomato, garlic, ginger
- Rajma masala, salt
- 1 tsp oil

INSTRUCTIONS:

1. Pressure cook rajma 5-6 whistles.
2. Make gravy with onion-tomato base and spices.
3. Cook rice separately.
4. Combine rajma and gravy.
5. Serve rajma over rice.

TIPS: Complete protein combination. One of India's best weight-loss foods.

Grilled Fish with Salad

CAT: Lunch | KCAL: 320 | P: 40g C: 8g F: 12g Fiber: 4g | SERVING: 1 plate | COST: █100 | PREP: 10 min COOK: 15 min | DIFF: Intermediate | WL: Excellent

SUITABLE FOR: Non Vegetarian, Office Worker

INGREDIENTS:

- 200g fish fillet (rohu/basa)
- Lemon, herbs, garlic
- Olive oil 1 tsp
- Green salad 150g
- Cucumber, tomato
- Olive oil + lemon dressing

INSTRUCTIONS:

1. Marinate fish with lemon, herbs, garlic.
2. Grill or pan-cook 4-5 min each side.
3. Toss salad with dressing.
4. Serve fish over salad.

TIPS: Fish is the ultimate protein source — high protein, omega-3, low calorie.

Quinoa Vegetable Bowl

CAT: Lunch | KCAL: 340 | P: 14g C: 50g F: 8g Fiber: 8g | SERVING: 1 bowl | COST: █70 | PREP: 5 min COOK: 20 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Gluten Free

INGREDIENTS:

- 3/4 cup quinoa
- Mixed roasted vegetables 150g
- 1 tbsp tahini or peanut dressing
- Lemon, salt, pepper
- Herbs

INSTRUCTIONS:

1. Cook quinoa (1:2 with water, 15 min).
2. Roast vegetables in oven or pan.
3. Assemble bowl with quinoa base.
4. Add vegetables and dressing.

TIPS: Complete protein grain. Meal prep well. Better than white rice for fat loss.

Palak Paneer + Roti

CAT: Lunch | KCAL: 420 | P: 22g C: 42g F: 16g Fiber: 8g | SERVING: 1 serving + 2 roti | COST: █60 | PREP: 10 min COOK: 20 min |
DIFF: Intermediate | WL: Good
SUITABLE FOR: Vegetarian, Student

INGREDIENTS:

- 100g paneer
- 200g spinach
- Onion, tomato, garlic
- Cream 1 tbsp (optional)
- Spices, salt
- 2 whole wheat rotis

INSTRUCTIONS:

1. Blanch spinach, blend smooth.
2. Sauté onion-tomato masala.
3. Add spinach puree, simmer.
4. Add cubed paneer, cook 5 min.
5. Serve with rotis.

TIPS: Iron + protein + calcium powerhouse. Avoid heavy cream version.

Egg Curry + Rice

CAT: Lunch | KCAL: 440 | P: 26g C: 52g F: 14g Fiber: 4g | SERVING: 1 plate | COST: █60 | PREP: 10 min COOK: 20 min | DIFF: Intermediate | WL: Good
SUITABLE FOR: Non Vegetarian, Student, Budget

INGREDIENTS:

- 3 boiled eggs
- 1 cup rice
- Onion, tomato, ginger-garlic
- Egg curry masala
- Oil, salt
- Coriander

INSTRUCTIONS:

1. Cook rice.
2. Make onion-tomato gravy.
3. Add eggs (halved or whole), simmer 5 min.
4. Serve over rice.

TIPS: Economical high-protein meal. Use brown rice for better fiber.

Chole + Roti

CAT: Lunch | KCAL: 400 | P: 17g C: 60g F: 8g Fiber: 13g | SERVING: 1 bowl + 2 roti | COST: █30 | PREP: 10 min COOK: 25 min | DIFF: Beginner | WL: Excellent
SUITABLE FOR: Vegetarian, Vegan, Budget, Student

INGREDIENTS:

- 1/2 cup chickpeas (soaked overnight)
- Onion, tomato, garlic
- Chole masala, amchur, salt
- 1 tsp oil
- 2 rotis

INSTRUCTIONS:

1. Pressure cook chickpeas 5 whistles.
2. Make spicy gravy.
3. Add chickpeas, simmer 10 min.
4. Serve with rotis.

TIPS: Chickpeas have among the highest satiety scores. Excellent fat-loss food.

Chicken Salad Wrap

CAT: Lunch | KCAL: 360 | P: 32g C: 30g F: 10g Fiber: 5g | SERVING: 1 large wrap | COST: █80 | PREP: 10 min COOK: 10 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Non Vegetarian, Office Worker, No Cook (if pre-cooked)

INGREDIENTS:

- 120g shredded chicken breast
- 1 large whole wheat roti
- Lettuce, tomato, cucumber
- 2 tbsp Greek yogurt dressing
- Mustard, lemon
- Salt, pepper

INSTRUCTIONS:

1. Mix chicken with dressing and vegetables.
2. Warm roti.
3. Fill and roll tight.
4. Cut diagonally and serve.

TIPS: Office-friendly — packs well. Greek yogurt dressing instead of mayo saves 80 kcal.

Mixed Dal Soup

CAT: Lunch | KCAL: 220 | P: 16g C: 30g F: 3g Fiber: 10g | SERVING: 1 large bowl | COST: █20 | PREP: 5 min COOK: 20 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget, Diet

INGREDIENTS:

- 1/4 cup each: masoor, moong, chana dal
- 3 cups water or stock
- Onion, tomato
- Turmeric, cumin, salt
- Lemon

INSTRUCTIONS:

1. Pressure cook all dals with water.
2. Blend half for creamy texture.
3. Temper cumin, onion, tomato.
4. Combine all.
5. Add lemon before serving.

TIPS: High protein, high fiber, very low calorie. Excellent for calorie-restricted days.

Tofu Stir Fry with Brown Rice

CAT: Lunch | KCAL: 380 | P: 22g C: 50g F: 10g Fiber: 6g | SERVING: 1 bowl | COST: █60 | PREP: 10 min COOK: 15 min | DIFF: Intermediate | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan

INGREDIENTS:

- 150g firm tofu
- 3/4 cup brown rice (cooked)
- Broccoli, bell peppers, snap peas
- Soy sauce, ginger, garlic
- Sesame oil 1 tsp

INSTRUCTIONS:

1. Press and cube tofu, pan-fry golden.
2. Stir-fry vegetables 3 min.
3. Add tofu, sauce, stir.

4. Serve over brown rice.

TIPS: Tofu is the best vegan protein source. Press well before cooking for better texture.

Sabudana Khichdi

CAT: Lunch | KCAL: 290 | P: 6g C: 58g F: 6g Fiber: 2g | SERVING: 1 bowl | COST: ■25 | PREP: 4 hrs (soak) COOK: 10 min | DIFF: Beginner | WL: Moderate

SUITABLE FOR: Vegetarian, Vegan, Fasting, Gluten Free

INGREDIENTS:

- 3/4 cup sabudana (soaked 4 hrs)
- 2 tbsp peanuts
- 1 potato (boiled)
- Cumin, green chilli
- Salt, lemon, coriander

INSTRUCTIONS:

1. Drain soaked sabudana.
2. Dry roast peanuts, crush coarsely.
3. Heat oil, add cumin, chilli.
4. Add potato, cook 2 min.
5. Add sabudana and peanuts, stir 5 min.
6. Add salt and lemon.

TIPS: High GI, moderate protein. Better for maintenance than aggressive fat loss.

Baingan Bharta + Roti

CAT: Lunch | KCAL: 280 | P: 8g C: 40g F: 8g Fiber: 9g | SERVING: 1 serving + 2 rotis | COST: ■25 | PREP: 10 min COOK: 25 min | DIFF: Intermediate | WL: Good

SUITABLE FOR: Vegetarian, Vegan, Budget

INGREDIENTS:

- 1 large brinjal
- Onion, tomato, garlic
- Cumin, chilli, salt
- 1 tsp oil
- 2 whole wheat rotis

INSTRUCTIONS:

1. Roast brinjal directly on flame till soft.
2. Peel and mash.
3. Sauté onion, tomato, garlic.
4. Add mashed brinjal, cook 5 min.
5. Serve with rotis.

TIPS: Brinjal is extremely low calorie (20 kcal/100g). Roasting adds smoky flavor without fat.

Corn + Chicken Soup

CAT: Lunch | KCAL: 240 | P: 22g C: 24g F: 5g Fiber: 4g | SERVING: 1 bowl (350ml) | COST: ■60 | PREP: 10 min COOK: 20 min | DIFF: Intermediate | WL: Excellent

SUITABLE FOR: Non Vegetarian, Meal Prep

INGREDIENTS:

- 100g chicken (shredded)
- 1/2 cup sweet corn
- Carrot, celery, onion
- 2 cups chicken broth
- Cornflour 1 tsp
- Salt, pepper, ginger

INSTRUCTIONS:

1. Simmer stock with vegetables 10 min.
2. Add shredded chicken and corn.
3. Mix cornflour with water, add to thicken.
4. Season and serve hot.

TIPS: High satiety soup. Drink before meal to reduce total intake by 15-20%.

Paneer Salad Bowl

CAT: Lunch | KCAL: 310 | P: 20g C: 18g F: 16g Fiber: 6g | SERVING: 1 bowl | COST: ■70 | PREP: 10 min COOK: 5 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Low Carb, No Cook

INGREDIENTS:

- 100g paneer (cubed and pan-grilled)
- Mixed greens 100g
- Cherry tomatoes, cucumber
- Red onion
- Lemon-olive oil dressing
- Salt, pepper, herbs

INSTRUCTIONS:

1. Grill paneer cubes 2 min each side.
2. Arrange greens and vegetables.
3. Add paneer on top.
4. Drizzle dressing.

TIPS: High protein low-carb lunch. Very satisfying with minimal cooking.

Veg Pulao

CAT: Lunch | KCAL: 320 | P: 9g C: 58g F: 6g Fiber: 5g | SERVING: 1 bowl | COST: ■30 | PREP: 10 min COOK: 20 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Vegan, Budget, Student

INGREDIENTS:

- 1 cup basmati rice
- Mixed vegetables 200g
- Whole spices
- 1 onion
- Salt, oil

INSTRUCTIONS:

1. Sauté spices, onion.
2. Add vegetables.
3. Add washed rice.
4. Add 2 cups water, cook till done.

TIPS: Add extra vegetables to reduce carb density.

Sprouts Pulao

CAT: Lunch | KCAL: 340 | P: 14g C: 56g F: 6g Fiber: 9g | SERVING: 1 bowl | COST: ■25 | PREP: 5 min COOK: 20 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget

INGREDIENTS:

- 3/4 cup rice
- 1 cup mixed sprouts
- Onion, tomato, spices
- Salt, oil

INSTRUCTIONS:

1. Sauté onion-spices.
2. Add sprouts briefly.
3. Add rice and water.
4. Cook to fluffy.

TIPS: Sprouts add protein and live enzymes. Better than plain veg pulao.

Chicken Keema

CAT: Lunch | KCAL: 320 | P: 38g C: 6g F: 16g Fiber: 2g | SERVING: 1 serving (150g) | COST: ■90 | PREP: 10 min COOK: 20 min | DIFF: Intermediate | WL: Excellent

SUITABLE FOR: Non Vegetarian, Low Carb, Keto-friendly

INGREDIENTS:

- 200g chicken mince
- Onion, tomato, ginger-garlic
- Peas
- Spices, oil
- Coriander

INSTRUCTIONS:

1. Sauté aromatics.
2. Add mince, cook 10 min.
3. Add peas and spices.
4. Cook till done.

TIPS: Very high protein, low carb. Pair with 1 roti or eat alone with salad.

Masoor Dal Soup

CAT: Lunch | KCAL: 200 | P: 14g C: 28g F: 2g Fiber: 9g | SERVING: 1 bowl | COST: ■15 | PREP: 5 min COOK: 20 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget, Diet

INGREDIENTS:

- 1/2 cup masoor dal
- 3 cups water
- Onion, tomato, garlic
- Cumin, turmeric, salt
- Lemon

INSTRUCTIONS:

1. Boil dal till soft.
2. Temper and combine.
3. Blend partially for texture.
4. Add lemon.

TIPS: Only 200 kcal but extremely filling. Best diet day lunch.

Grilled Chicken + Salad

CAT: Dinner | KCAL: 320 | P: 42g C: 8g F: 12g Fiber: 4g | SERVING: 1 plate | COST: ■100 | PREP: 10 min COOK: 20 min | DIFF: Intermediate | WL: Excellent

SUITABLE FOR: Non Vegetarian, Low Carb

INGREDIENTS:

- 200g chicken breast
- Mixed greens 150g
- Cherry tomatoes, cucumber
- Lemon-herb marinade
- 1 tsp olive oil

INSTRUCTIONS:

1. Marinate chicken 30 min minimum.
2. Grill 8-10 min each side.
3. Rest 5 min, slice.
4. Arrange over salad.

TIPS: Ideal fat-loss dinner — high protein, very low carb.

Moong Dal Khichdi

CAT: Dinner | KCAL: 300 | P: 16g C: 50g F: 4g Fiber: 8g | SERVING: 1 bowl (300g) | COST: █20 | PREP: 5 min COOK: 20 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget, Student

INGREDIENTS:

- 1/4 cup rice
- 1/4 cup moong dal
- Turmeric, salt
- Ghee 1 tsp
- Cumin, hing
- Vegetables optional

INSTRUCTIONS:

1. Wash rice and dal together.
2. Pressure cook 3 whistles.
3. Temper ghee, cumin, hing.
4. Mix and serve.

TIPS: Ayurveda's ultimate healing dinner. Easy to digest, complete protein combination.

Fish Curry + Rice

CAT: Dinner | KCAL: 420 | P: 38g C: 44g F: 10g Fiber: 3g | SERVING: 1 plate | COST: █90 | PREP: 10 min COOK: 20 min | DIFF: Intermediate | WL: Excellent

SUITABLE FOR: Non Vegetarian, South Indian

INGREDIENTS:

- 200g fish (rohu/pomfret)
- Coconut milk 100ml
- Onion, tomato, tamarind
- Fish curry masala
- Oil, salt
- 1 cup basmati rice

INSTRUCTIONS:

1. Cook rice.
2. Fry onion-masala base.
3. Add fish, cook 5 min.
4. Add coconut milk and tamarind.
5. Simmer 10 min.

TIPS: Omega-3 rich. Use minimal coconut milk to control fat.

Stir-Fried Tofu Vegetables

CAT: Dinner | KCAL: 260 | P: 18g C: 20g F: 10g Fiber: 7g | SERVING: 1 large plate | COST: █50 | PREP: 10 min COOK: 15 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Low Carb

INGREDIENTS:

- 150g firm tofu
- Broccoli, bell peppers, zucchini

- Soy sauce, ginger, garlic
- Sesame oil, chilli flakes

INSTRUCTIONS:

1. Press tofu, cube and pan-fry golden.
2. Stir-fry vegetables 5 min.
3. Add tofu and sauce.
4. Toss and serve.

TIPS: Under 300 kcal, high protein dinner. Add more vegetables as desired.

Rasam + Brown Rice

CAT: Dinner | KCAL: 260 | P: 8g C: 50g F: 4g Fiber: 5g | SERVING: 1 plate | COST: █15 | PREP: 5 min COOK: 15 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Vegan, South Indian, Budget

INGREDIENTS:

- 1 cup brown rice
- Rasam from toor dal water
- Tomato, tamarind
- Mustard, cumin, curry leaves
- Black pepper, salt

INSTRUCTIONS:

1. Cook brown rice.
2. Boil tamarind water with tomato.
3. Add rasam powder, pepper.
4. Temper and add.
5. Simmer 5 min.

TIPS: Rasam aids digestion. Light dinner for weight loss. Black pepper boosts metabolism.

Paneer Tikka + Salad

CAT: Dinner | KCAL: 340 | P: 22g C: 14g F: 20g Fiber: 4g | SERVING: 1 serving (200g) | COST: █80 | PREP: 20 min COOK: 15 min | DIFF: Intermediate | WL: Good

SUITABLE FOR: Vegetarian, Low Carb

INGREDIENTS:

- 150g paneer
- Bell peppers, onion (chunks)
- Tikka marinade: hung curd, spices
- 1 tsp oil
- Salad to serve

INSTRUCTIONS:

1. Marinate paneer and vegetables 1 hr.
2. Skewer and grill or pan-fry.
3. Char lightly.
4. Serve with salad.

TIPS: Restaurant tikka has 40% more fat than home version. Air fryer works perfectly.

Chicken Soup

CAT: Dinner | KCAL: 200 | P: 28g C: 8g F: 5g Fiber: 2g | SERVING: 1 bowl (400ml) | COST: █60 | PREP: 10 min COOK: 30 min | DIFF: Intermediate | WL: Excellent

SUITABLE FOR: Non Vegetarian, Diet Days, Low Calorie

INGREDIENTS:

- 150g chicken bone-in
- Carrot, celery, onion

- Garlic, ginger
- Black pepper, salt
- Bay leaf, herbs

INSTRUCTIONS:

1. Simmer chicken in water with aromatics.
2. Cook 25-30 min till tender.
3. Shred chicken.
4. Return to broth.
5. Season and serve.

TIPS: Under 200 kcal. High protein soup dinner for fast deficit days.

Sambar + Idli

CAT: Dinner | KCAL: 280 | P: 12g C: 48g F: 4g Fiber: 7g | SERVING: 3 idli + sambar | COST: ■25 | PREP: 5 min COOK: 15 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Vegan, South Indian, Budget

INGREDIENTS:

- 3 idlis
- 1 cup sambar
- Vegetables in sambar
- Green chutney

INSTRUCTIONS:

1. Steam idlis.
2. Heat sambar (add vegetables).
3. Serve together.

TIPS: Fermented idli excellent for gut health. Light, digestible dinner.

Egg White Stir Fry

CAT: Dinner | KCAL: 210 | P: 26g C: 10g F: 6g Fiber: 4g | SERVING: 1 plate | COST: ■35 | PREP: 5 min COOK: 10 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Non Vegetarian, Low Calorie, Low Carb

INGREDIENTS:

- 6 egg whites
- Spinach, mushrooms, bell pepper
- Garlic, soy sauce
- 1 tsp oil
- Salt, pepper

INSTRUCTIONS:

1. Sauté vegetables in oil 3 min.
2. Add egg whites, scramble loosely.
3. Season and serve.

TIPS: Under 220 kcal, very high protein. Perfect aggressive deficit dinner.

Dal Palak

CAT: Dinner | KCAL: 280 | P: 17g C: 36g F: 5g Fiber: 12g | SERVING: 1 bowl | COST: ■20 | PREP: 10 min COOK: 20 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget

INGREDIENTS:

- 1/2 cup toor or masoor dal
- 200g spinach
- Onion, tomato, garlic
- Cumin, turmeric, salt

- 1 tsp oil

INSTRUCTIONS:

1. Cook dal till soft.
2. Wilt spinach, add to dal.
3. Make tadka with spices.
4. Combine and simmer 5 min.

TIPS: Iron + protein + fiber combination. One of the best weight-loss dinners.

Ragi Mudde + Sambhar

CAT: Dinner | KCAL: 300 | P: 10g C: 56g F: 3g Fiber: 8g | SERVING: 2 mudde + sambhar | COST: █20 | PREP: 5 min COOK: 15 min | DIFF: Intermediate | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, South Indian, Gluten Free

INGREDIENTS:

- 1 cup ragi flour
- 2 cups water
- Salt
- 1 cup sambhar

INSTRUCTIONS:

1. Boil salted water.
2. Add ragi flour, stir till thick.
3. Shape into balls.
4. Serve with sambhar.

TIPS: Karnataka staple. Ragi is highest-calcium grain, excellent for bone health and satiety.

Lemon Herb Baked Fish

CAT: Dinner | KCAL: 250 | P: 38g C: 2g F: 10g Fiber: 1g | SERVING: 1 fillet | COST: █100 | PREP: 5 min COOK: 20 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Non Vegetarian, Low Carb, Keto-friendly

INGREDIENTS:

- 200g fish fillet
- Lemon, garlic, herbs
- 1 tsp olive oil
- Salt, pepper
- Foil

INSTRUCTIONS:

1. Season fish with all ingredients.
2. Wrap in foil.
3. Bake 180°C for 18-20 min.
4. Serve immediately.

TIPS: Near zero effort, incredibly high protein, very low calorie. Best diet dinner.

Rajma Soup

CAT: Dinner | KCAL: 240 | P: 15g C: 38g F: 3g Fiber: 12g | SERVING: 1 bowl | COST: █20 | PREP: 5 min COOK: 15 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget

INGREDIENTS:

- 1 cup cooked rajma
- 2 cups water or broth
- Onion, tomato
- Spices, salt
- Lime

INSTRUCTIONS:

1. Blend half the rajma.
2. Simmer with broth and aromatics.
3. Add remaining whole beans.
4. Season and finish with lime.

TIPS: Soup form of rajma is more filling per calorie than rajma chawal.

Jowar Roti + Vegetable

CAT: Dinner | KCAL: 290 | P: 9g C: 52g F: 4g Fiber: 8g | SERVING: 2 rotis + subji | COST: █25 | PREP: 10 min COOK: 15 min | DIFF: Intermediate | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Gluten Free, Diabetic

INGREDIENTS:

- 1 cup jowar flour
- Warm water to knead
- Any dry vegetable curry
- Salt

INSTRUCTIONS:

1. Knead jowar dough.
2. Roll thick (unlike wheat).
3. Cook on hot tawa.
4. Press with cloth.
5. Serve with curry.

TIPS: Jowar has lower GI than wheat. Excellent for diabetics and weight loss.

Paneer + Vegetables Curry

CAT: Dinner | KCAL: 360 | P: 20g C: 22g F: 20g Fiber: 6g | SERVING: 1 bowl | COST: █70 | PREP: 10 min COOK: 20 min | DIFF: Intermediate | WL: Good

SUITABLE FOR: Vegetarian

INGREDIENTS:

- 100g paneer
- Mixed vegetables 200g
- Onion-tomato gravy
- Spices
- 1 tsp oil

INSTRUCTIONS:

1. Make light onion-tomato gravy.
2. Add vegetables, cook 8 min.
3. Add paneer cubes.
4. Simmer 5 min.

TIPS: Reduce paneer to 75g and increase vegetables to cut 50 kcal.

Boiled Chana Snack

CAT: Snack | KCAL: 180 | P: 10g C: 28g F: 3g Fiber: 8g | SERVING: 1 cup (100g) | COST: █10 | PREP: 1 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget, No Cook

INGREDIENTS:

- 1 cup boiled chickpeas
- Lemon, chaat masala
- Green chilli, coriander

INSTRUCTIONS:

1. Combine all.

2. Toss and eat.

TIPS: Best weight-loss snack in India. Prep in bulk and refrigerate for 5 days.

Roasted Makhana

CAT: Snack | KCAL: 150 | P: 5g C: 28g F: 2g Fiber: 1g | SERVING: 1.5 cups (40g) | COST: ■25 | PREP: 1 min COOK: 5 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Gluten Free

INGREDIENTS:

- 40g lotus seeds (makhana)
- 1/4 tsp ghee
- Salt, chilli powder, turmeric

INSTRUCTIONS:

1. Dry roast makhana 4-5 min.
2. Add ghee and spices.
3. Toss and cool.

TIPS: Volume eating king — 40g fills a large bowl. Much better than chips or namkeen.

Peanut Butter Apple

CAT: Snack | KCAL: 220 | P: 8g C: 26g F: 10g Fiber: 4g | SERVING: 1 apple + 2 tbsp PB | COST: ■35 | PREP: 2 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Vegan, No Cook

INGREDIENTS:

- 1 medium apple
- 2 tbsp natural peanut butter

INSTRUCTIONS:

1. Slice apple.
2. Dip or spread PB.

TIPS: Fiber + protein + fat combination — very high satiety for 220 kcal.

Hummus with Vegetables

CAT: Snack | KCAL: 180 | P: 8g C: 20g F: 8g Fiber: 7g | SERVING: 3 tbsp hummus + veggies | COST: ■30 | PREP: 5 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, No Cook

INGREDIENTS:

- 3 tbsp hummus
- Carrot sticks, cucumber, bell pepper

INSTRUCTIONS:

1. Arrange vegetables.
2. Serve with hummus.

TIPS: Low calorie, high fiber, high protein dip. Make hummus from chickpeas at home cheaply.

Masala Peanuts

CAT: Snack | KCAL: 200 | P: 9g C: 14g F: 12g Fiber: 3g | SERVING: 40g handful | COST: ■10 | PREP: 2 min COOK: 5 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Vegan, Budget

INGREDIENTS:

- 40g peanuts
- Chilli, cumin, salt
- 1/4 tsp oil

INSTRUCTIONS:

1. Dry roast peanuts.

2. Add oil and spices.

3. Toss and cool.

TIPS: Portion control critical — easy to overeat. Pre-portion into 40g bags.

Fruit Chaat

CAT: Snack | KCAL: 160 | P: 2g C: 38g F: 1g Fiber: 6g | SERVING: 1 bowl (200g) | COST: █30 | PREP: 5 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Vegan, No Cook, Budget

INGREDIENTS:

- Mixed seasonal fruits 200g
- Chaat masala
- Black salt, lime

INSTRUCTIONS:

1. Chop fruits.
2. Add spices and lime.
3. Toss and serve.

TIPS: Avoid only high-sugar fruits (mango, litchi) during fat loss phase. Mix with guava, papaya.

Boiled Egg Snack

CAT: Snack | KCAL: 140 | P: 12g C: 1g F: 10g Fiber: 0g | SERVING: 2 eggs | COST: █20 | PREP: 0 min COOK: 8 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Non Vegetarian, Student, Hostel, Budget

INGREDIENTS:

- 2 eggs
- Salt, pepper
- Hot sauce optional

INSTRUCTIONS:

1. Boil eggs.
2. Season and eat.

TIPS: Most efficient protein snack per rupee. Keep 6 boiled eggs in fridge.

Cucumber Curd Dip

CAT: Snack | KCAL: 120 | P: 8g C: 10g F: 3g Fiber: 2g | SERVING: 1 cup | COST: █20 | PREP: 5 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, No Cook, Low Calorie

INGREDIENTS:

- 200g curd
- 1/2 cucumber (grated)
- Mint, salt, cumin
- Lime

INSTRUCTIONS:

1. Mix all ingredients.
2. Serve chilled.

TIPS: Raita variant — only 120 kcal, high protein, very refreshing and filling.

Roasted Chana

CAT: Snack | KCAL: 160 | P: 10g C: 26g F: 3g Fiber: 7g | SERVING: 40g (small packet) | COST: █10 | PREP: 0 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget, No Cook

INGREDIENTS:

- 40g roasted chana (bhuna chana)

INSTRUCTIONS:

1. Eat directly or with spices.

TIPS: India's original protein snack. High fiber, very low cost, portable.

Paneer Cubes

CAT: Snack | KCAL: 160 | P: 11g C: 2g F: 12g Fiber: 0g | SERVING: 50g | COST: █25 | PREP: 2 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Low Carb, No Cook

INGREDIENTS:

- 50g paneer
- Chilli powder, salt
- Lime

INSTRUCTIONS:

1. Cut paneer.
2. Season.
3. Eat immediately.

TIPS: High protein, zero carb snack. Very filling for small volume.

Sprouts Chaat

CAT: Snack | KCAL: 140 | P: 10g C: 22g F: 2g Fiber: 7g | SERVING: 1 cup | COST: █15 | PREP: 5 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget, No Cook

INGREDIENTS:

- 1 cup mixed sprouts
- Onion, tomato, chilli
- Lemon, chaat masala

INSTRUCTIONS:

1. Mix all.
2. Season and eat.

TIPS: Living food with enzymes. Sprouting doubles protein bioavailability.

Makhana Kheer (low-cal)

CAT: Snack | KCAL: 180 | P: 8g C: 24g F: 4g Fiber: 1g | SERVING: 1 cup | COST: █30 | PREP: 5 min COOK: 10 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Diet Dessert

INGREDIENTS:

- 30g makhana
- 200ml skimmed milk
- Stevia or 1 tsp sugar
- Cardamom, saffron

INSTRUCTIONS:

1. Roast makhana.
2. Simmer in milk till soft.
3. Add sweetener and cardamom.

TIPS: Satisfies sweet cravings for only 180 kcal.

Banana Peanut Butter

CAT: Snack | KCAL: 220 | P: 7g C: 32g F: 9g Fiber: 4g | SERVING: 1 banana + 1.5 tbsp PB | COST: █20 | PREP: 1 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Vegan, No Cook, Hostel

INGREDIENTS:

- 1 banana
- 1.5 tbsp natural peanut butter

INSTRUCTIONS:

1. Slice banana.
2. Dip in PB.

TIPS: Pre-workout or post-workout snack. Carbs + protein combination.

Protein Bar (Homemade)

CAT: Snack | KCAL: 200 | P: 14g C: 22g F: 6g Fiber: 3g | SERVING: 1 bar | COST: ■20 | PREP: 10 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, No Cook, Hostel

INGREDIENTS:

- 2 tbsp peanut butter
- 1 tbsp honey
- 1 scoop protein powder
- 3 tbsp oats
- 2 tbsp dark chocolate chips

INSTRUCTIONS:

1. Mix all in bowl.
2. Press into bar mold or tray.
3. Refrigerate 30 min.
4. Cut into bars.

TIPS: Make 10 at a time. Far better than commercial protein bars at 1/5th the cost.

Soya Chunks Snack

CAT: Snack | KCAL: 180 | P: 24g C: 12g F: 2g Fiber: 4g | SERVING: 50g dry (100g cooked) | COST: ■10 | PREP: 5 min COOK: 10 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget

INGREDIENTS:

- 50g soya chunks
- Onion, tomato, chilli
- Spices, salt
- 1 tsp oil

INSTRUCTIONS:

1. Boil soya chunks 5 min, drain, squeeze.
2. Sauté with spices and vegetables.
3. Season and serve.

TIPS: Highest protein-to-cost ratio in India. 52g protein per 100g dry.

Protein Mango Shake

CAT: Smoothie | KCAL: 320 | P: 24g C: 42g F: 4g Fiber: 3g | SERVING: 1 glass (400ml) | COST: ■50 | PREP: 3 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Student

INGREDIENTS:

- 1 mango (medium)
- 1 scoop vanilla protein
- 200ml milk
- Ice cubes
- Pinch turmeric

INSTRUCTIONS:

1. Blend all till smooth.

2. Serve immediately.

TIPS: Perfect summer post-workout. Use frozen mango off-season.

Green Protein Smoothie

CAT: Smoothie | KCAL: 240 | P: 22g C: 26g F: 4g Fiber: 5g | SERVING: 1 glass | COST: █40 | PREP: 3 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, No Cook

INGREDIENTS:

- 2 cups spinach
- 1 banana
- 1 scoop protein
- 200ml almond milk
- 1 tsp flaxseed

INSTRUCTIONS:

1. Blend all.
2. Add ice, blend again.

TIPS: Best weight-loss smoothie. Under 250 kcal, very high protein.

Banana Date Shake

CAT: Smoothie | KCAL: 350 | P: 14g C: 62g F: 4g Fiber: 5g | SERVING: 1 glass | COST: █35 | PREP: 3 min COOK: 0 min | DIFF: Beginner | WL: Moderate

SUITABLE FOR: Vegetarian, Vegan, Budget

INGREDIENTS:

- 2 bananas
- 3-4 dates (soaked)
- 200ml milk
- Cardamom

INSTRUCTIONS:

1. Remove date pits.
2. Blend all smooth.
3. Serve cold.

TIPS: Natural sugar — better than commercial shakes. High in natural energy.

Chocolate Protein Shake

CAT: Smoothie | KCAL: 280 | P: 28g C: 22g F: 6g Fiber: 3g | SERVING: 1 glass | COST: █60 | PREP: 2 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Student, Post-workout

INGREDIENTS:

- 1 scoop chocolate protein
- 200ml milk
- 1 banana
- 1 tsp cocoa powder
- Ice

INSTRUCTIONS:

1. Blend all.
2. Serve immediately.

TIPS: Post-workout must. Best within 30-60 min after training.

Buttermilk (Chaas)

CAT: Smoothie | KCAL: 80 | P: 4g C: 8g F: 2g Fiber: 0g | SERVING: 1 glass (300ml) | COST: █10 | PREP: 2 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Budget, No Cook

INGREDIENTS:

- 200g curd
- 200ml water
- Salt, cumin
- Mint leaves
- Ice

INSTRUCTIONS:

1. Blend all.
2. Strain and serve cold.

TIPS: Only 80 kcal. Probiotic-rich, cooling, extremely filling per calorie.

Strawberry Banana Smoothie

CAT: Smoothie | KCAL: 260 | P: 14g C: 40g F: 3g Fiber: 4g | SERVING: 1 glass | COST: ■60 | PREP: 3 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Student

INGREDIENTS:

- 1 cup strawberries
- 1 banana
- 1 scoop vanilla protein
- 150ml milk
- Ice

INSTRUCTIONS:

1. Blend all.
2. Serve chilled.

TIPS: High antioxidant, moderate protein. Good morning or pre-workout option.

Turmeric Golden Milk Smoothie

CAT: Smoothie | KCAL: 200 | P: 8g C: 26g F: 6g Fiber: 2g | SERVING: 1 glass | COST: ■30 | PREP: 3 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Anti-inflammatory

INGREDIENTS:

- 250ml warm milk
- 1 tsp turmeric
- 1/4 tsp black pepper
- 1 tsp honey
- Ginger
- Cardamom

INSTRUCTIONS:

1. Blend all.
2. Serve warm or cold.

TIPS: Curcumin + piperine = powerful anti-inflammatory combination. Good evening drink.

Chia Seed Smoothie

CAT: Smoothie | KCAL: 230 | P: 12g C: 28g F: 8g Fiber: 8g | SERVING: 1 glass | COST: ■40 | PREP: 5 min (+15 min chia soak) COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, No Cook

INGREDIENTS:

- 2 tbsp chia seeds
- 200ml almond milk
- 1 banana

- 1 tsp honey
- Berries

INSTRUCTIONS:

1. Soak chia in milk 15 min.
2. Blend with banana.
3. Top with berries.

TIPS: Omega-3 + fiber powerhouse. Chia expands in stomach, increases satiety.

Papaya Lime Smoothie

CAT: Smoothie | KCAL: 160 | P: 4g C: 36g F: 1g Fiber: 5g | SERVING: 1 glass | COST: █20 | PREP: 3 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget, Diet

INGREDIENTS:

- 200g papaya
- Lime juice
- 200ml water
- Pinch black salt
- Optional: ginger

INSTRUCTIONS:

1. Blend all.
2. Serve cold.

TIPS: Only 160 kcal. Papaya contains papain enzyme which aids digestion and fat breakdown.

Watermelon Mint Cooler

CAT: Smoothie | KCAL: 80 | P: 2g C: 18g F: 0g Fiber: 1g | SERVING: 1 glass | COST: █15 | PREP: 3 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget, Diet

INGREDIENTS:

- 300g watermelon
- Mint leaves
- Lime juice
- Black salt
- Ice

INSTRUCTIONS:

1. Blend and serve.

TIPS: Only 80 kcal. 92% water — extreme satiety per calorie. Perfect summer diet drink.

Pineapple Ginger Shot Smoothie

CAT: Smoothie | KCAL: 140 | P: 3g C: 32g F: 0g Fiber: 3g | SERVING: 1 glass | COST: █20 | PREP: 3 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Vegan, No Cook

INGREDIENTS:

- 200g pineapple
- 1 inch ginger
- 200ml water
- Lime
- Mint

INSTRUCTIONS:

1. Blend all.
2. Strain if desired.
3. Serve over ice.

TIPS: Bromelain enzyme aids protein digestion. Anti-inflammatory and refreshing.

Protein Scrambled Eggs

CAT: Protein Meal | KCAL: 360 | P: 38g C: 4g F: 20g Fiber: 1g | SERVING: 1 plate | COST: █50 | PREP: 3 min COOK: 5 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Non Vegetarian, Low Carb, Keto-friendly

INGREDIENTS:

- 4 eggs
- 2 egg whites
- Spinach 30g
- Salt, pepper
- 1 tsp butter

INSTRUCTIONS:

1. Whisk all eggs.
2. Cook slowly on low heat with butter.
3. Add spinach.
4. Remove when just set.

TIPS: Slow and low scramble = creamy, filling. Add cheese for extra protein.

High-Protein Chicken Salad

CAT: Protein Meal | KCAL: 350 | P: 48g C: 8g F: 12g Fiber: 4g | SERVING: 1 bowl | COST: █100 | PREP: 10 min COOK: 15 min | DIFF: Intermediate | WL: Excellent

SUITABLE FOR: Non Vegetarian, Low Carb

INGREDIENTS:

- 200g chicken breast
- Mixed greens 150g
- Cherry tomatoes
- 1 tbsp olive oil dressing
- Lemon, herbs

INSTRUCTIONS:

1. Grill or poach chicken.
2. Shred or slice.
3. Toss with greens and dressing.

TIPS: 48g protein per serving. Best fat-loss meal in this document.

Paneer Protein Bowl

CAT: Protein Meal | KCAL: 400 | P: 32g C: 22g F: 20g Fiber: 4g | SERVING: 1 bowl | COST: █80 | PREP: 10 min COOK: 10 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Low Carb

INGREDIENTS:

- 150g paneer
- Chickpeas 50g (cooked)
- Onion, bell pepper
- Lemon-herb dressing
- Seeds for topping

INSTRUCTIONS:

1. Grill paneer cubes.
2. Combine with chickpeas, vegetables.
3. Dress and serve.

TIPS: 32g protein — best vegetarian protein bowl.

Soya Chunk Curry

CAT: Protein Meal | KCAL: 340 | P: 36g C: 28g F: 8g Fiber: 8g | SERVING: 1 bowl | COST: █20 | PREP: 15 min COOK: 20 min | DIFF: Intermediate | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget

INGREDIENTS:

- 100g soya chunks (dry)
- Onion, tomato, ginger-garlic
- Spices: cumin, coriander, turmeric
- 1 tsp oil
- Salt, coriander

INSTRUCTIONS:

1. Boil soya chunks 8 min, drain and squeeze.
2. Make onion-tomato gravy.
3. Add soya, simmer 10 min.
4. Garnish and serve.

TIPS: Cheapest high-protein meal in India. 36g protein for █20.

Tuna Salad (Canned)

CAT: Protein Meal | KCAL: 220 | P: 32g C: 8g F: 6g Fiber: 3g | SERVING: 1 bowl | COST: █80 | PREP: 5 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Non Vegetarian, No Cook, Budget

INGREDIENTS:

- 1 can tuna in water (150g)
- Onion, cucumber, tomato
- Mustard 1 tsp
- Lime, salt, pepper
- Mixed greens

INSTRUCTIONS:

1. Drain tuna.
2. Mix with vegetables and dressing.
3. Serve over greens.

TIPS: 32g protein in 5 min. Canned tuna is one of the best protein-per-rupee foods.

Protein Khichdi

CAT: Protein Meal | KCAL: 380 | P: 28g C: 50g F: 6g Fiber: 9g | SERVING: 1 bowl | COST: █30 | PREP: 5 min COOK: 20 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget

INGREDIENTS:

- 1/4 cup rice
- 1/4 cup moong dal
- 1/4 cup soya chunks
- Vegetables 100g
- Spices, salt
- 1 tsp ghee

INSTRUCTIONS:

1. Soak soya chunks 10 min.
2. Pressure cook all ingredients together.
3. Add ghee to finish.

TIPS: Modified khichdi with soya = 28g protein. Complete meal.

Egg White Bhurji

CAT: Protein Meal | KCAL: 160 | P: 22g C: 6g F: 5g Fiber: 2g | SERVING: 1 serving | COST: █25 | PREP: 3 min COOK: 5 min | DIFF: Beginner | WL: Excellent
SUITABLE FOR: Non Vegetarian, Low Calorie, Low Carb

INGREDIENTS:

- 6 egg whites
- Onion, tomato, chilli
- Salt, pepper, turmeric
- 1/2 tsp oil
- Coriander

INSTRUCTIONS:

1. Sauté vegetables.
2. Add whites, scramble.
3. Season.

TIPS: Only 160 kcal and 22g protein. Best calorie-deficit protein source.

Greek Yogurt Protein Bowl

CAT: Protein Meal | KCAL: 300 | P: 28g C: 28g F: 6g Fiber: 4g | SERVING: 1 bowl | COST: █70 | PREP: 3 min COOK: 0 min | DIFF: Beginner | WL: Excellent
SUITABLE FOR: Vegetarian, No Cook

INGREDIENTS:

- 250g Greek yogurt/hung curd
- 1/2 scoop protein powder
- Berries
- Granola 20g
- Honey 1 tsp

INSTRUCTIONS:

1. Mix yogurt and protein.
2. Top with rest.

TIPS: 28g protein, no cooking. Best pre-sleep protein meal.

Paneer Scramble

CAT: Protein Meal | KCAL: 300 | P: 24g C: 8g F: 18g Fiber: 2g | SERVING: 1 serving | COST: █60 | PREP: 5 min COOK: 8 min | DIFF: Beginner | WL: Good
SUITABLE FOR: Vegetarian, Low Carb, Keto-friendly

INGREDIENTS:

- 150g paneer (crumbled)
- Onion, bell pepper, chilli
- Spices, salt
- 1 tsp oil

INSTRUCTIONS:

1. Sauté vegetables.
2. Add paneer, cook 3-4 min.
3. Season.

TIPS: Paneer scramble = vegetarian version of egg scramble. Same speed, similar protein.

Chicken Tikka Protein Bowl

CAT: Protein Meal | KCAL: 440 | P: 48g C: 24g F: 14g Fiber: 4g | SERVING: 1 bowl | COST: █120 | PREP: 20 min (+ marinate) COOK: 20 min | DIFF: Intermediate | WL: Excellent
SUITABLE FOR: Non Vegetarian

INGREDIENTS:

- 200g chicken (cubed)

- Tikka marinade: curd, spices
- Brown rice 1/2 cup (cooked)
- Salad
- Mint chutney

INSTRUCTIONS:

1. Marinate chicken 2-4 hrs.
2. Grill or air-fry.
3. Serve over rice with salad.

TIPS: Restaurant-quality high-protein meal at home. 48g protein per bowl.

Dal Chawal (Simple)

CAT: Budget | KCAL: 380 | P: 16g C: 66g F: 4g Fiber: 8g | SERVING: 1 plate | COST: ■20 | PREP: 5 min COOK: 20 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Vegan, Budget, Hostel, Student

INGREDIENTS:

- 1/2 cup any dal
- 1 cup rice
- Turmeric, salt
- 1 tsp ghee or oil

INSTRUCTIONS:

1. Pressure cook dal and rice separately.
2. Combine with ghee.
3. Add salt and turmeric.

TIPS: India's ultimate budget complete protein meal. Dal + rice = complete amino acid profile.

Peanut Butter Oats

CAT: Budget | KCAL: 350 | P: 16g C: 48g F: 10g Fiber: 6g | SERVING: 1 bowl | COST: ■25 | PREP: 2 min COOK: 5 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Hostel, Student, Budget

INGREDIENTS:

- 3/4 cup oats
- 1 tbsp peanut butter
- 1 tsp honey
- 200ml water or milk

INSTRUCTIONS:

1. Cook oats in water.
2. Stir in PB and honey.

TIPS: ■25 complete breakfast. High satiety, lasts 4-5 hours.

Bread + Eggs

CAT: Budget | KCAL: 300 | P: 18g C: 28g F: 14g Fiber: 2g | SERVING: 2 eggs + 2 slices | COST: ■25 | PREP: 2 min COOK: 5 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Non Vegetarian, Hostel, Student, Budget

INGREDIENTS:

- 2 eggs
- 2 bread slices
- Butter/oil 1 tsp
- Salt, pepper

INSTRUCTIONS:

1. Fry or boil eggs.
2. Toast bread.

3. Combine.

TIPS: Hostel kitchen staple. Boiled eggs + bread = better macro split.

Rajma on Toast

CAT: Budget | KCAL: 320 | P: 15g C: 50g F: 5g Fiber: 12g | SERVING: 2 slices + rajma | COST: █20 | PREP: 5 min COOK: 5 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Vegan, Hostel, Budget

INGREDIENTS:

- 1 cup canned/cooked rajma
- 2 whole wheat bread slices
- Onion, tomato, spices
- Salt, lime

INSTRUCTIONS:

1. Heat rajma with spices.
2. Toast bread.
3. Top with rajma.

TIPS: British beans-on-toast but with rajma. Fiber powerhouse.

Curd Rice (Cold)

CAT: Budget | KCAL: 240 | P: 10g C: 38g F: 5g Fiber: 2g | SERVING: 1 bowl | COST: █15 | PREP: 5 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Hostel, Budget, No Cook

INGREDIENTS:

- Leftover rice 1 cup
- Curd 200g
- Salt, mustard seeds, curry leaves
- Cucumber, pomegranate

INSTRUCTIONS:

1. Mix rice and curd.
2. Add tempered spices.
3. Garnish.

TIPS: Best use of leftover rice. Gut-healthy and filling.

Oats Khichdi

CAT: Budget | KCAL: 280 | P: 12g C: 42g F: 6g Fiber: 7g | SERVING: 1 bowl | COST: █20 | PREP: 5 min COOK: 10 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Vegan, Hostel, Budget

INGREDIENTS:

- 3/4 cup oats
- 1/4 cup moong dal (soaked)
- Vegetables
- Cumin, turmeric, salt
- 1 tsp oil

INSTRUCTIONS:

1. Soak dal 30 min.
2. Pressure cook or stovetop with all ingredients.

TIPS: Oats + moong = fast high-protein hostel khichdi.

Roti Sabzi

CAT: Budget | KCAL: 280 | P: 8g C: 46g F: 6g Fiber: 8g | SERVING: 2 rotis + 1 bowl sabzi | COST: █20 | PREP: 10 min COOK: 15 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Vegan, Budget, Student

INGREDIENTS:

- 2 whole wheat rotis
- Seasonal vegetable 200g
- Oil, spices, salt

INSTRUCTIONS:

1. Make dry vegetable curry.
2. Roll rotis.
3. Serve together.

TIPS: Cheapest complete Indian dinner. Choose fiber-rich veggies (lauki, cabbage, carrot).

Boiled Chana Masala

CAT: Budget | KCAL: 200 | P: 12g C: 30g F: 3g Fiber: 10g | SERVING: 1 bowl | COST: ■10 | PREP: 2 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, Budget, No Cook

INGREDIENTS:

- 1 cup boiled chana
- Onion, tomato, chilli
- Chaat masala, lime
- Salt, coriander

INSTRUCTIONS:

1. Mix all.
2. Eat cold.

TIPS: ■10 high-protein meal. Cook large batch and store 5 days.

Milk Banana Oats

CAT: Budget | KCAL: 380 | P: 16g C: 58g F: 7g Fiber: 6g | SERVING: 1 bowl | COST: ■25 | PREP: 2 min COOK: 5 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, Budget, Hostel, Student

INGREDIENTS:

- 3/4 cup oats
- 250ml milk
- 1 banana
- 1 tsp honey

INSTRUCTIONS:

1. Cook oats in milk.
2. Top with banana and honey.

TIPS: ■25 high-protein, high-fiber, complete breakfast.

Egg Noodles (Hostel)

CAT: Budget | KCAL: 380 | P: 18g C: 52g F: 12g Fiber: 3g | SERVING: 1 plate | COST: ■35 | PREP: 5 min COOK: 10 min | DIFF: Beginner | WL: Moderate

SUITABLE FOR: Non Vegetarian, Hostel, Student, Budget

INGREDIENTS:

- 1 packet noodles
- 2 eggs
- Mixed vegetables
- Soy sauce
- 1 tsp oil

INSTRUCTIONS:

1. Cook noodles, drain.

2. Scramble eggs.
3. Add noodles and vegetables.
4. Season.

TIPS: Better version of instant noodles — added protein and vegetables. Limit to 2x/week.

Curd + Banana Meal

CAT: No Cook | KCAL: 260 | P: 12g C: 42g F: 5g Fiber: 3g | SERVING: 1 bowl + 1 banana | COST: ■25 | PREP: 2 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, No Cook, Budget, Hostel

INGREDIENTS:

- 250g curd
- 1 banana
- 1 tsp honey
- Optional: oats 2 tbsp

INSTRUCTIONS:

1. Slice banana over curd.
2. Add honey.
3. Mix and eat.

TIPS: Emergency complete meal in 2 minutes.

Protein Shake + Fruit

CAT: No Cook | KCAL: 300 | P: 26g C: 38g F: 4g Fiber: 4g | SERVING: 1 shake + 1 fruit | COST: ■60 | PREP: 2 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, No Cook, Hostel, Student

INGREDIENTS:

- 1 scoop protein powder
- 250ml milk
- 1 apple or banana

INSTRUCTIONS:

1. Shake protein in milk.
2. Eat fruit alongside.

TIPS: Travel emergency meal. Shaker bottle + protein powder = high-protein anytime.

Bread + Peanut Butter + Milk

CAT: No Cook | KCAL: 370 | P: 18g C: 44g F: 14g Fiber: 4g | SERVING: 2 slices + glass milk | COST: ■30 | PREP: 3 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, No Cook, Hostel, Budget

INGREDIENTS:

- 2 whole wheat slices
- 2 tbsp PB
- 250ml milk

INSTRUCTIONS:

1. Spread PB on bread.
2. Drink milk.

TIPS: 3-minute complete macro meal.

Sprouts + Curd

CAT: No Cook | KCAL: 220 | P: 18g C: 24g F: 4g Fiber: 8g | SERVING: 1 bowl | COST: ■20 | PREP: 5 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, No Cook, Budget

INGREDIENTS:

- 1 cup sprouts
- 200g curd
- Salt, chaat masala
- Coriander

INSTRUCTIONS:

1. Mix all.
2. Serve cold.

TIPS: Probiotic + living food combination. Excellent gut-health meal.

Fruit Nut Curd Bowl

CAT: No Cook | KCAL: 280 | P: 12g C: 38g F: 8g Fiber: 4g | SERVING: 1 bowl | COST: █50 | PREP: 5 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, No Cook

INGREDIENTS:

- 200g curd
- Mixed fruits 100g
- 10g mixed nuts
- 1 tsp honey
- Cinnamon

INSTRUCTIONS:

1. Combine all in bowl.
2. Drizzle honey.

TIPS: Calcium + healthy fat + natural sugar. Complete snack meal.

Muesli Curd Bowl

CAT: No Cook | KCAL: 310 | P: 14g C: 50g F: 6g Fiber: 5g | SERVING: 1 bowl | COST: █45 | PREP: 2 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, No Cook, Hostel

INGREDIENTS:

- 40g muesli
- 200g curd
- Honey
- Seasonal fruit

INSTRUCTIONS:

1. Layer muesli and curd.
2. Top with fruit.

TIPS: Better than muesli + milk — higher protein, probiotic boost.

Chickpea Salad

CAT: No Cook | KCAL: 250 | P: 12g C: 36g F: 5g Fiber: 10g | SERVING: 1 bowl | COST: █20 | PREP: 5 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, No Cook, Budget

INGREDIENTS:

- 1 cup canned/boiled chickpeas
- Cucumber, tomato, onion
- Lime, salt, cumin
- Olive oil 1 tsp

INSTRUCTIONS:

1. Combine all.
2. Toss.
3. Serve.

TIPS: Mediterranean-Indian fusion. High fiber, very filling, almost no prep.

Cucumber Sandwich

CAT: No Cook | KCAL: 200 | P: 8g C: 28g F: 6g Fiber: 4g | SERVING: 2 sandwiches | COST: ■20 | PREP: 5 min COOK: 0 min | DIFF: Beginner | WL: Good

SUITABLE FOR: Vegetarian, No Cook, Hostel, Student

INGREDIENTS:

- 4 bread slices
- Cucumber (sliced)
- Hung curd 3 tbsp
- Mint chutney
- Salt, pepper

INSTRUCTIONS:

1. Spread curd on bread.
2. Layer cucumber.
3. Add chutney, season.
4. Close and cut.

TIPS: Light, cooling sandwich. Very low calorie for a sandwich.

Roasted Chana + Fruits

CAT: No Cook | KCAL: 200 | P: 10g C: 34g F: 4g Fiber: 9g | SERVING: 1 serving | COST: ■20 | PREP: 2 min COOK: 0 min | DIFF: Beginner | WL: Excellent

SUITABLE FOR: Vegetarian, Vegan, No Cook, Budget

INGREDIENTS:

- 40g roasted chana
- 1 apple or guava
- 1 tsp lime

INSTRUCTIONS:

1. Eat together.

TIPS: Protein + fiber + vitamins snack. No prep required.

Complete Meal Plan System

WL-MP-000	MOD 05 — Meal Plans · Meal Plan System Overview
GOALS	Weight Loss Fat Loss Maintenance
USERS	Adult · Teen · Senior · Student · Hostel Student · Office Worker · PCOS · Diabetes
DIFF	Beginner
TAGS	meal plan, calorie targets, Indian diet plan, 1200 calorie, 1600 calorie, veg non-veg

DEFINITION

Complete meal planning system covering 9 calorie tiers × multiple user types with Indian foods.

WHY IT MATTERS

- Meal plans remove decision fatigue — the #1 reason diets fail.
- Pre-planned meals reduce calorie overshoot by 30-40%.

KEY FACTS

- Calorie tiers: 1200 / 1400 / 1600 / 1800 / 2000 / 2200 / 2500 kcal
- User types: Vegetarian / Vegan / Non-Veg / Hostel / PCOS / Diabetes / Office
- Weekly grocery lists included for key plans.
- Emergency food options for travel/festivals/restaurants.

ACTION STEPS

1. Calculate your TDEE (Module 01).
2. Set deficit: Lose weight = TDEE minus 300-500 kcal.
3. Find your calorie tier below.
4. Choose your food type (Veg/Non-veg/Vegan/Hostel).
5. Follow for 2 weeks before adjusting.

DECISION TREE

IF BMI > 35 → **THEN** Start at 1200-1400 kcal, medical supervision recommended
IF BMI 30-35 → **THEN** 1400-1600 kcal
IF BMI 27-30 → **THEN** 1600-1800 kcal
IF BMI 23-27 → **THEN** 1800-2000 kcal
IF Active + Want to lose → **THEN** TDEE minus 300-400 kcal
IF Sedentary + Want to lose → **THEN** TDEE minus 400-500 kcal
IF PCOS / Hypothyroid → **THEN** Choose PCOS/Diabetic plan regardless of BMI
IF Hostel student (mess food) → **THEN** Hostel plan with mess food adaptations

FAQ

Q: Can I swap meals from different days?

A: Yes. All meals within same calorie tier are interchangeable.

Q: What if I'm still hungry?

A: Add salad, cucumber, buttermilk — zero/near-zero calorie volume.

Q: Can I eat the same breakfast every day?

A: Yes — consistency is easier to track than variety for most people.

Q: Do I need protein powder?

A: No. It helps but whole food sources work.

Q: What about cheat meals?

A: 1 cheat meal per week on 1600+ kcal plans. Not a cheat day.

WL-MP-1200-VEG	MOD 05 — Meal Plans · 1200 kcal Meal Plan › Veg
GOALS	Weight Loss Fat Loss
USERS	Adult · Adult

DIFF	Beginner
TAGS	1200kcal, meal plan, veg, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Veg user at 1200 kcal target.

WHY IT MATTERS

- Practical 1200 kcal plan optimized for Veg lifestyle.

KEY FACTS

- Macros: P:55g C:140g F:25g Fiber:28g
- Meals: 7 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Wake up 6am: Warm water with lemon (0 kcal)
2. Breakfast 7am: Moong Dal Chilla x2 (260 kcal) + Green Tea
3. Mid-Morning 10am: 1 fruit – guava or apple (80 kcal)
4. Lunch 1pm: Dal (1/3 cup) + 1 Roti + Salad (280 kcal)
5. Evening 4pm: Buttermilk / Chaas (80 kcal)
6. Dinner 7pm: Moong Dal Khichdi (300 kcal) + Salad
7. Post-Dinner: Turmeric Milk (120 kcal)
8. TIP: Eat dinner before 8pm. Avoid second helpings. Chew slowly.

WL-MP-1200 -NONVEG	MOD 05 — Meal Plans · 1200 kcal Meal Plan › Nonveg
GOALS	Weight Loss Fat Loss
USERS	Adult · Adult
DIFF	Beginner
TAGS	1200kcal, meal plan, nonveg, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Nonveg user at 1200 kcal target.

WHY IT MATTERS

- Practical 1200 kcal plan optimized for Nonveg lifestyle.

KEY FACTS

- Macros: P:75g C:100g F:22g Fiber:15g
- Meals: 7 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Wake up 6am: Warm lemon water
2. Breakfast 7am: 2 Boiled Eggs + 1 Whole Wheat Toast (240 kcal)
3. Mid-Morning 10am: Buttermilk (80 kcal)
4. Lunch 1pm: Chicken Soup + 1 Roti (300 kcal)
5. Evening 4pm: Boiled Chana 1/2 cup (90 kcal)
6. Dinner 7pm: Grilled Fish (150g) + Salad (260 kcal)
7. Pre-sleep: Warm milk (100 kcal)
8. TIP: High protein reduces hunger on low calorie. Drink 3L water.

WL-MP-1400 -VEG	MOD 05 — Meal Plans · 1400 kcal Meal Plan › Veg
GOALS	Weight Loss Fat Loss
USERS	Adult · Adult
DIFF	Beginner
TAGS	1400kcal, meal plan, veg, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Veg user at 1400 kcal target.

WHY IT MATTERS

- Practical 1400 kcal plan optimized for Veg lifestyle.

KEY FACTS

- Macros: P:62g C:170g F:32g Fiber:30g
- Meals: 6 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Breakfast 7am: Besan Chilla x2 + Curd 100g (360 kcal)
2. Mid-Morning 10am: Apple + 10 Almonds (130 kcal)
3. Lunch 1pm: Dal Tadka + 2 Roti + Salad (380 kcal)
4. Evening 4pm: Roasted Makhana 30g + Green Tea (110 kcal)
5. Dinner 7pm: Palak Paneer (small) + 1 Roti (320 kcal)
6. Pre-sleep: Warm milk 150ml (90 kcal)
7. TIP: Add salad to every meal for volume. Avoid evening snacks after 7pm.

WL-MP-1400 -NONVEG	MOD 05 — Meal Plans · 1400 kcal Meal Plan › Nonveg
GOALS	Weight Loss Fat Loss
USERS	Adult · Adult
DIFF	Beginner
TAGS	1400kcal, meal plan, nonveg, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Nonveg user at 1400 kcal target.

WHY IT MATTERS

- Practical 1400 kcal plan optimized for Nonveg lifestyle.

KEY FACTS

- Macros: P:88g C:120g F:30g Fiber:18g
- Meals: 6 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Breakfast 7am: Egg White Omelette (3 whites + 1 whole) + Toast (250 kcal)
2. Mid-Morning 10am: Banana + Peanut Butter 1 tbsp (160 kcal)
3. Lunch 1pm: Chicken Rice Bowl (small) (380 kcal)
4. Evening 4pm: Boiled Egg x1 + Roasted Chana (140 kcal)
5. Dinner 7pm: Grilled Chicken + Stir-fried Vegetables (320 kcal)
6. Pre-sleep: Protein Shake (half scoop) in water (80 kcal)
7. TIP: Protein at every meal is essential. Don't skip evening snack.

WL-MP-1400 -HOSTEL	MOD 05 — Meal Plans · 1400 kcal Meal Plan › Hostel
GOALS	Weight Loss Fat Loss
USERS	Adult · Student
DIFF	Beginner
TAGS	1400kcal, meal plan, hostel, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Hostel user at 1400 kcal target.

WHY IT MATTERS

- Practical 1400 kcal plan optimized for Hostel lifestyle.

KEY FACTS

- Macros: P:50g C:190g F:25g Fiber:22g
- Meals: 5 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Breakfast: Oats + Milk + Banana (280 kcal)
2. Mid-Morning: Roasted Chana 40g (160 kcal)
3. Lunch (mess): Dal + Rice (small) + Salad (320 kcal). Skip fried items.
4. Evening: Peanut Butter + 2 Bread slices (300 kcal)
5. Dinner (mess): Sabzi + 2 Roti (300 kcal). Avoid puri/bhatura.
6. TIP: Hostel strategy: carry peanut butter, oats, roasted chana. Skip mess fried items.

WL-MP-1600 -VEG	MOD 05 — Meal Plans · 1600 kcal Meal Plan › Veg
GOALS	Weight Loss Fat Loss
USERS	Adult · Adult
DIFF	Beginner
TAGS	1600kcal, meal plan, veg, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Veg user at 1600 kcal target.

WHY IT MATTERS

- Practical 1600 kcal plan optimized for Veg lifestyle.

KEY FACTS

- Macros: P:72g C:210g F:35g Fiber:34g
- Meals: 6 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Breakfast 7am: High-Protein Oats Bowl (380 kcal)
2. Mid-Morning 10am: Sprouts Chaat 1 cup (140 kcal)
3. Lunch 1pm: Rajma Chawal (small) + Salad (380 kcal)
4. Pre-workout 4pm: Banana + Peanut Butter (200 kcal)
5. Post-workout 6pm: Protein Shake (150 kcal)
6. Dinner 8pm: Moong Dal + 1 Roti + Vegetable (340 kcal)
7. TIP: For active vegetarians. Pre/post workout timing is critical.

WL-MP-1600 -NONVEG	MOD 05 — Meal Plans · 1600 kcal Meal Plan › Nonveg
GOALS	Weight Loss Fat Loss
USERS	Adult · Adult
DIFF	Beginner
TAGS	1600kcal, meal plan, nonveg, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Nonveg user at 1600 kcal target.

WHY IT MATTERS

- Practical 1600 kcal plan optimized for Nonveg lifestyle.

KEY FACTS

- Macros: P:100g C:160g F:36g Fiber:18g
- Meals: 5 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Breakfast 7am: Paneer Bhurji + 2 Roti (450 kcal)
2. Mid-Morning 10am: Greek Yogurt + Fruit (180 kcal)

3. Lunch 1pm: Chicken + Rice Bowl (med) (420 kcal)
4. Evening 5pm: Boiled Eggs x2 (140 kcal)
5. Dinner 8pm: Fish Curry + 1 Cup Rice (380 kcal)
6. TIP: Excellent muscle preservation plan. Suit active non-veg users.

WL-MP-1600 -VEGAN	MOD 05 — Meal Plans · 1600 kcal Meal Plan › Vegan
GOALS	Weight Loss Fat Loss
USERS	Adult · Adult
DIFF	Beginner
TAGS	1600kcal, meal plan, vegan, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Vegan user at 1600 kcal target.

WHY IT MATTERS

- Practical 1600 kcal plan optimized for Vegan lifestyle.

KEY FACTS

- Macros: P:68g C:200g F:32g Fiber:38g
- Meals: 5 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Breakfast: Protein Smoothie (soy protein) + Oats (380 kcal)
2. Mid-Morning: Soaked Almonds 10 + Apple (150 kcal)
3. Lunch: Quinoa + Chickpea Bowl + Greens (400 kcal)
4. Evening: Soya Chunks Snack (180 kcal)
5. Dinner: Tofu Stir Fry + Brown Rice (380 kcal)
6. TIP: Vegan plans must actively track protein. Tofu, soya, legumes every meal.

WL-MP-1800 -VEG	MOD 05 — Meal Plans · 1800 kcal Meal Plan › Veg
GOALS	Weight Loss Fat Loss
USERS	Adult · Adult
DIFF	Beginner
TAGS	1800kcal, meal plan, veg, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Veg user at 1800 kcal target.

WHY IT MATTERS

- Practical 1800 kcal plan optimized for Veg lifestyle.

KEY FACTS

- Macros: P:78g C:240g F:44g Fiber:36g
- Meals: 6 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Breakfast 7am: Stuffed Paratha x2 + Curd 100g (480 kcal)
2. Mid-Morning: Mixed Fruit Bowl (160 kcal)
3. Lunch: Chole + 2 Roti + Dal (500 kcal)
4. Pre-workout: Banana + 15 Almonds (200 kcal)
5. Post-workout: Protein Shake (180 kcal)
6. Dinner: Palak Paneer + 2 Roti (420 kcal)
7. TIP: Maintenance-friendly. Good for active lifestyle, moderate deficit.

WL-MP-1800-NONVEG	MOD 05 — Meal Plans · 1800 kcal Meal Plan › Nonveg
GOALS	Weight Loss Fat Loss
USERS	Adult · Adult
DIFF	Beginner
TAGS	1800kcal, meal plan, nonveg, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Nonveg user at 1800 kcal target.

WHY IT MATTERS

- Practical 1800 kcal plan optimized for Nonveg lifestyle.

KEY FACTS

- Macros: P:115g C:185g F:40g Fiber:20g
- Meals: 5 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Breakfast: Egg Bhurji Roll x2 (680 kcal)
2. Mid-Morning: Protein Shake + Fruit (280 kcal)
3. Lunch: Chicken + Dal + Rice (550 kcal)
4. Snack: Boiled Eggs x2 + Makhana (230 kcal)
5. Dinner: Grilled Fish + Roti + Salad (450 kcal)
6. TIP: High protein allows muscle building while losing fat simultaneously.

WL-MP-1800-PCOS	MOD 05 — Meal Plans · 1800 kcal Meal Plan › Pcos
GOALS	Weight Loss Fat Loss
USERS	Adult · Adult
DIFF	Beginner
TAGS	1800kcal, meal plan, pcos, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Pcos user at 1800 kcal target.

WHY IT MATTERS

- Practical 1800 kcal plan optimized for Pcos lifestyle.

KEY FACTS

- Macros: P:72g C:180g F:40g Fiber:36g
- Meals: 6 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Breakfast: Moong Dal Chilla x3 + Curd (380 kcal) — NO refined carbs
2. Mid-Morning: Soaked Methi Seeds water + Apple (90 kcal)
3. Lunch: Brown Rice (small) + Dal + Vegetables (400 kcal) — Low GI
4. Evening: Roasted Pumpkin Seeds 30g + Green Tea (130 kcal)
5. Dinner: Paneer + Stir-fried Vegetables (no rice) (380 kcal)
6. Pre-sleep: Cinnamon Turmeric Milk (100 kcal)
7. TIP: PCOS: Low GI foods only. No maida/refined sugar. High fiber at every meal. Methi for insulin sensitivity.

WL-MP-1800-DIABETES	MOD 05 — Meal Plans · 1800 kcal Meal Plan › Diabetes
GOALS	Weight Loss Fat Loss
USERS	Adult · Adult
DIFF	Beginner
TAGS	1800kcal, meal plan, diabetes, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Diabetes user at 1800 kcal target.

WHY IT MATTERS

- Practical 1800 kcal plan optimized for Diabetes lifestyle.

KEY FACTS

- Macros: P:70g C:170g F:38g Fiber:40g
- Meals: 6 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Breakfast: Besan Chilla x2 + Egg (380 kcal) — No bread/poha
2. Mid-Morning: Cucumber + Buttermilk (80 kcal)
3. Lunch: Small Brown Rice (1/2 cup) + Dal + Sabzi (380 kcal)
4. Evening: Roasted Chana 40g (160 kcal) — High fiber snack
5. Dinner: Jowar Roti x2 + Dal + Salad (420 kcal) — No rice dinner
6. Pre-sleep: Warm milk + 1/4 tsp cinnamon (100 kcal)
7. TIP: Diabetic: Never eat rice and roti together. Jowar/bajra over wheat. High fiber slows sugar spikes.

WL-MP-2000 -VEG	MOD 05 — Meal Plans · 2000 kcal Meal Plan › Veg
GOALS	Weight Loss Fat Loss
USERS	Adult · Adult
DIFF	Beginner
TAGS	2000kcal, meal plan, veg, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Veg user at 2000 kcal target.

WHY IT MATTERS

- Practical 2000 kcal plan optimized for Veg lifestyle.

KEY FACTS

- Macros: P:90g C:260g F:48g Fiber:40g
- Meals: 6 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Breakfast: High-Protein Oats + Paneer Bhurji (680 kcal)
2. Mid-Morning: Sprouts + Greek Yogurt (240 kcal)
3. Lunch: Rajma + 2 Roti + Salad (450 kcal)
4. Pre-workout: Banana + Peanut Butter + Milk (340 kcal)
5. Post-workout: Whey Protein Shake (150 kcal)
6. Dinner: Dal Palak + 2 Roti (380 kcal)
7. TIP: Good for active gym-going vegetarians targeting recomposition.

WL-MP-2000 -NONVEG	MOD 05 — Meal Plans · 2000 kcal Meal Plan › Nonveg
GOALS	Weight Loss Fat Loss
USERS	Adult · Adult
DIFF	Beginner
TAGS	2000kcal, meal plan, nonveg, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Nonveg user at 2000 kcal target.

WHY IT MATTERS

- Practical 2000 kcal plan optimized for Nonveg lifestyle.

KEY FACTS

- Macros: P:130g C:210g F:44g Fiber:22g
- Meals: 6 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Breakfast: 4 Eggs Scrambled + 2 Toast (480 kcal)
2. Mid-Morning: Protein Shake + Fruits (280 kcal)
3. Lunch: Chicken Rice Bowl (large) + Salad (580 kcal)
4. Pre-workout: Banana + 20 Almonds (270 kcal)
5. Post-workout: Protein Shake (160 kcal)
6. Dinner: Chicken Curry + 2 Roti (480 kcal)
7. TIP: Muscle-building non-veg plan. Excellent for active males.

WL-MP-2000 -OFFICE MOD 05 — Meal Plans · 2000 kcal Meal Plan › Office

GOALS	Weight Loss Fat Loss
USERS	Adult · Office Worker
DIFF	Beginner
TAGS	2000kcal, meal plan, office, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Office user at 2000 kcal target.

WHY IT MATTERS

- Practical 2000 kcal plan optimized for Office lifestyle.

KEY FACTS

- Macros: P:95g C:230g F:48g Fiber:28g
- Meals: 5 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Breakfast 7am: Overnight Oats + Boiled Eggs (480 kcal) — Prep Sunday night
2. Mid-Morning 10am (office): Roasted Chana + Fruit (220 kcal) — At desk
3. Lunch 1pm (office): Meal-prepped Chicken Bowl (550 kcal) — Microwave
4. Evening 5pm: Protein Bar or Nuts (200 kcal) — Avoid canteen samosas
5. Dinner 8pm: Simple Dal + 2 Roti (380 kcal) — Faster dinner
6. TIP: Office strategy: Sunday meal prep is mandatory. Keep emergency snacks at desk.

WL-MP-2200 -VEG MOD 05 — Meal Plans · 2200 kcal Meal Plan › Veg

GOALS	Weight Loss Fat Loss
USERS	Adult · Adult
DIFF	Beginner
TAGS	2200kcal, meal plan, veg, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Veg user at 2200 kcal target.

WHY IT MATTERS

- Practical 2200 kcal plan optimized for Veg lifestyle.

KEY FACTS

- Macros: P:100g C:300g F:54g Fiber:44g
- Meals: 7 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Breakfast: Paneer Bhurji + 3 Roti + Protein Shake (780 kcal)
2. Mid-Morning: Sprouts + Yogurt + Fruit (280 kcal)

3. Lunch: Chole + 3 Roti + Salad (560 kcal)
4. Pre-workout: Oats + Banana + PB (380 kcal)
5. Post-workout: Whey (180 kcal)
6. Dinner: Soya Chunks Curry + 2 Roti (480 kcal)
7. Pre-sleep: Milk 250ml (150 kcal)
8. TIP: Higher calorie for bulking vegetarians or very active women.

WL-MP-2500 -NONVEG	MOD 05 — Meal Plans · 2500 kcal Meal Plan › Nonveg
GOALS	Weight Loss Fat Loss
USERS	Adult · Adult
DIFF	Beginner
TAGS	2500kcal, meal plan, nonveg, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Nonveg user at 2500 kcal target.

WHY IT MATTERS

- Practical 2500 kcal plan optimized for Nonveg lifestyle.

KEY FACTS

- Macros: P:160g C:280g F:60g Fiber:22g
- Meals: 7 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Breakfast: 6 Eggs (whole) + 3 Toast + OJ (680 kcal)
2. Mid-Morning: Protein Shake + Banana + PB (420 kcal)
3. Lunch: Chicken Biryani (medium) + Raita + Salad (680 kcal)
4. Pre-workout: Rice + Chicken small (350 kcal)
5. Post-workout: Whey + Banana (280 kcal)
6. Dinner: Mutton Curry (small) + 3 Roti (600 kcal)
7. Pre-sleep: Milk + Protein (180 kcal)
8. TIP: Bulking / active athlete plan. Track meals precisely to avoid fat gain.

WL-MP-2500 -VEG	MOD 05 — Meal Plans · 2500 kcal Meal Plan › Veg
GOALS	Weight Loss Fat Loss
USERS	Adult · Adult
DIFF	Beginner
TAGS	2500kcal, meal plan, veg, Indian food, weight loss diet

DEFINITION

Day 1 meal plan for Veg user at 2500 kcal target.

WHY IT MATTERS

- Practical 2500 kcal plan optimized for Veg lifestyle.

KEY FACTS

- Macros: P:118g C:330g F:58g Fiber:46g
- Meals: 7 eating occasions
- All foods are Indian, affordable, and widely available.

ACTION STEPS

1. Breakfast: Oats + Protein + PB + Milk (580 kcal)
2. Mid-Morning: Sprouts + Yogurt + Almonds (300 kcal)
3. Lunch: Rajma + 3 Roti + Paneer Salad (620 kcal)
4. Pre-workout: Banana + 2 Dates + Milk (300 kcal)
5. Post-workout: Whey + Banana (280 kcal)

6. Dinner: Paneer Curry + 3 Roti + Salad (580 kcal)
7. Pre-sleep: Milk 300ml (180 kcal)
8. TIP: High-calorie veg bulking. Hard to achieve without dairy and protein supplements.

WL-MP-SUB MOD 05 — Meal Plans · Food Substitutions Reference › Healthier Swaps

GOALS Weight Loss | Fat Loss | Maintenance

USERS Adult · Student · Office Worker

DIFF Beginner

TAGS food swap, healthy substitution, lower calorie, better food choice

DEFINITION

One-to-one swaps to reduce calories or improve nutrition without giving up taste.

WHY IT MATTERS

- Small swaps compound over time — 100 kcal/day swap = 10kg in a year.

KEY FACTS

- Every swap listed reduces calories or improves nutrient density.
- Focus on 2-3 swaps at a time — not all at once.

ACTION STEPS

1. Replace: White Rice → Brown Rice / Quinoa / Millets — lower GI, more fiber
2. Replace: Maida (White Flour) → Whole Wheat / Besan / Jowar / Ragi — more nutrients
3. Replace: Full Fat Paneer → Homemade Low-Fat Paneer or Tofu — less saturated fat
4. Replace: Fried Snacks → Roasted Makhana / Chana / Peanuts — same satisfaction, less fat
5. Replace: Sugary Drinks → Buttermilk / Lemon Water / Coconut Water — zero added sugar
6. Replace: Cream / Malai → Greek Yogurt / Hung Curd — high protein, low fat
7. Replace: Ghee (excess) → Use 1 tsp daily. Don't eliminate — support fat-soluble vitamins
8. Replace: Namkeen / Chips → Roasted Chana / Makhana — save 100-200 kcal per serving
9. Replace: Sugary Tea → Herbal Tea / Green Tea / Kadha — zero calories
10. Replace: Biscuits / Cookies → Fruit + Nuts — natural sugar + protein + fiber
11. Replace: Mayonnaise → Hung Curd / Mustard — 80% fewer calories
12. Replace: Refined Sugar → Stevia / Jaggery (in moderation) — better glycemic response

FAQ

Q: Should I replace everything at once?

A: No. 1-2 swaps per week for sustainable habit change.

Q: Is brown rice really necessary?

A: Only if you eat rice multiple times daily. Once a day — white is fine.

Q: Can I still eat ghee?

A: Yes — 1 tsp daily is healthy. Don't eliminate completely.

WL-MP-SHOP MOD 05 — Meal Plans · Weekly Shopping Lists › Grocery Guide

GOALS Weight Loss

USERS Adult · Student · Hostel Student

DIFF Beginner

TAGS grocery list, shopping, weekly food, budget groceries, meal prep

DEFINITION

Weekly shopping lists for 1400 kcal vegetarian and 1600 kcal non-vegetarian plans.

WHY IT MATTERS

- Shopping list removes impulse buys and ensures you always have the right food at home.

KEY FACTS

- 1400 kcal veg grocery cost: ■800-1100/week
- 1600 kcal non-veg: ■1200-1500/week

ACTION STEPS

1. Print or screenshot this list.
2. Shop once weekly — Sundays work best.
3. Prep proteins on Sunday (boil eggs, cook dal, marinate chicken).

WL-MP-EMR G **MOD 05 — Meal Plans · Emergency Food Options · When Plan Fails**

GOALS Weight Loss | Maintenance

USERS Adult · Student · Office Worker · Frequent Traveler

DIFF Beginner

TAGS travel food, festival food, restaurant tips, emergency diet, wedding food

DEFINITION

What to eat when normal meal plan is impossible — restaurants, travel, festivals, weddings.

WHY IT MATTERS

- 80% adherence over 90 days beats 100% adherence for 2 weeks.
- Knowing emergency options prevents total plan abandonment.

KEY FACTS

- Social eating accounts for 30-40% of diet failures.
- One bad meal ≠ failed diet. One bad week ≠ failed month.

ACTION STEPS

1. Hotel / Restaurant: Order grilled fish/chicken + salad. Avoid creamy gravies. Ask for dal and roti.
2. Railway Station: Boiled eggs, banana, packaged roasted chana, buttermilk — avoid fried items.
3. Wedding / Feast: Eat protein first (dal, paneer, chicken). Fill half plate with salad. Small portions of rice.
4. Airport: Protein bar, Greek yogurt cup, fruit, nuts, boiled eggs at food courts.
5. Office Canteen: Choose dal-rice over biryani. Avoid samosa, vada, fried rice. Take salad first.
6. No Fridge (Hostel): Peanut butter + bread, Sattu drink, roasted chana, protein shake, banana.
7. Festival Days: One treat meal only. Compensate with lighter meals before/after. Don't fast all day then binge.
8. Late Night (Post-midnight): Warm milk + small protein. Avoid heavy meal. Sleep > eating after midnight.

FAQ

Q: Should I fast to compensate after a wedding?

A: No. Fast the day before, not after. Eat normally the next day.

Q: I overate yesterday — what now?

A: Normal meals today. No punishment. Move on.

Q: Can I drink alcohol and lose weight?

A: Alcohol is 7 kcal/g. 1-2 drinks max, avoid sugary mixers. Track total calories.

AI COACHING NOTES

- If user ate at wedding: 'That's fine. Back on plan tomorrow. One meal doesn't matter.'
- If user skipped meal: 'Don't double eat next meal. Normal portion next time.'
- If user is traveling: 'Look for dal-roti, grilled options. Avoid fried airport food.'

Exercise Database

WL-EX-000	MOD 06 — Exercise Database · Exercise Library Overview
GOALS	Weight Loss Fat Loss General Health
USERS	Adult · Teen · Senior · Student · Hostel Student · Office Worker
DIFF	Beginner
TAGS	exercise, workout, bodyweight, gym, HIIT, yoga, cardio

DEFINITION

Complete exercise database with 43 exercises across all categories.

WHY IT MATTERS

- Exercise burns calories, builds muscle, and improves insulin sensitivity.
- Muscle mass increases TDEE permanently — making weight maintenance easier.

KEY FACTS

- Bodyweight: 15 | Gym: 15 | HIIT: 7 | Yoga/Mobility: 6
- Walking burns 100-200 kcal/30 min. HIIT burns 300-450 kcal/30 min.
- Resistance training burns 10-15% more calories than equivalent cardio for 24-48 hours post-session.
- Minimum effective dose: 150 min moderate or 75 min vigorous activity per week (WHO).

ACTION STEPS

1. Beginners: start bodyweight only — perfect form before adding weight.
2. Gym users: compound lifts first (squat, deadlift, press, row).
3. Fat loss priority: HIIT 2x/week + resistance training 3x/week.
4. Daily walking is non-negotiable for NEAT — aim 7000+ steps.

DECISION TREE

- IF Complete beginner → THEN Bodyweight: push-up, squat, plank, lunge — 3x/week
- IF Has gym access → THEN Barbell compound lifts + isolation — 4x/week
- IF No gym, no equipment → THEN Bodyweight HIIT — 4x/week + daily walk
- IF Short on time (30 min) → THEN Tabata or circuit training — high efficiency
- IF Knee/joint issues → THEN Swimming, cycling, yoga — low impact
- IF Senior → THEN Walking, yoga, light resistance bands — low intensity
- IF PCOS/hormonal → THEN Resistance training > cardio — improves insulin sensitivity

Push-Up

PATTERN: Push | MUSCLES: Chest, Triceps, Shoulders | KCAL/30min: 180-220 | EQUIP: None | DIFF: Beginner

INSTRUCTIONS:

1. Start in plank: hands shoulder-width, body straight.
2. Lower chest to floor, elbows at 45°.
3. Push up powerfully.

CUES:

- Squeeze glutes throughout
- Don't flare elbows wide
- Keep core tight — no sagging hips

MISTAKES:

- Partial range of motion
- Head drooping
- Flared elbows

EASIER: Knee push-up or wall push-up

HARDER: Diamond push-up, archer push-up, one-arm push-up

Squat

PATTERN: Squat | MUSCLES: Quads, Glutes, Hamstrings | KCAL/30min: 220-280 | EQUIP: None | DIFF: Beginner

INSTRUCTIONS:

1. Feet shoulder-width, toes slightly out.
2. Sit back and down, chest tall.
3. Thighs parallel to floor.
4. Drive through heels to stand.

CUES:

- Knees track toes
- Chest up throughout
- Weight on mid-foot

MISTAKES:

- Knees caving inward
- Heel lifting off floor
- Rounding lower back

EASIER: Chair squat (touch chair, stand)

HARDER: Jump squat, Bulgarian split squat, pistol squat

Plank

PATTERN: Core | MUSCLES: Core, Shoulders, Glutes | KCAL/30min: 120-150 | EQUIP: None | DIFF: Beginner

INSTRUCTIONS:

1. Forearms on floor, elbows under shoulders.
2. Body straight from head to heels.
3. Hold position.

CUES:

- Squeeze everything
- Don't hold breath
- Look at floor, not forward

MISTAKES:

- Hips too high or sagging
- Not breathing
- Neck strain

EASIER: Knee plank

HARDER: Plank with shoulder taps, RKC plank, one-arm plank

Burpee

PATTERN: Full Body | MUSCLES: Full body, Cardiovascular | KCAL/30min: 350-420 | EQUIP: None | DIFF: Intermediate

INSTRUCTIONS:

1. Stand. Drop hands to floor.
2. Jump feet back to plank.
3. Do a push-up (optional).
4. Jump feet to hands.
5. Explosively jump up, clap overhead.

CUES:

- Land soft on feet
- Control the descent
- Scale intensity to fitness level

MISTAKES:

- Landing hard (joint stress)
- Rushing through — poor form
- Skipping full hip extension at top

EASIER: Step-back burpee (no jump)

HARDER: Burpee + pull-up, 100 burpee challenge

Mountain Climber

PATTERN: Core + Cardio | MUSCLES: Core, Shoulders, Hip Flexors | KCAL/30min: 300-360 | EQUIP: None | DIFF: Intermediate

INSTRUCTIONS:

1. Start in push-up position.
2. Drive right knee to chest.
3. Quickly switch legs.
4. Maintain flat back.

CUES:

- Hips stay level
- Don't bounce hips up
- Drive from core not legs

MISTAKES:

- Hips rising
- Slow alternation losing cardio benefit
- Holding breath

EASIER: Slow alternating knee drives

HARDER: Cross-body mountain climber

Lunge

PATTERN: Hinge/Squat | MUSCLES: Quads, Glutes, Hamstrings | KCAL/30min: 200-250 | EQUIP: None | DIFF: Beginner

INSTRUCTIONS:

1. Stand tall. Step one foot forward.
2. Lower back knee toward floor.
3. Front thigh parallel.
4. Push back to start.

CUES:

- Knee doesn't pass toes
- Torso upright
- Equal weight front/back

MISTAKES:

- Leaning forward
- Front knee caving in
- Short stride

EASIER: Static lunge (feet stay)

HARDER: Walking lunge, reverse lunge, jump lunge

Glute Bridge

PATTERN: Hip Extension | MUSCLES: Glutes, Hamstrings, Core | KCAL/30min: 120-150 | EQUIP: None | DIFF: Beginner

INSTRUCTIONS:

1. Lie on back, knees bent, feet flat.
2. Drive hips up, squeeze glutes hard.
3. Hold 2 seconds at top.
4. Lower controlled.

CUES:

- Drive through heels
- Fully extend hips
- Don't hyperextend lower back

MISTAKES:

- Not squeezing glutes at top
- Feet too far or close
- Hyperextending spine

EASIER: Lie flat, just squeeze glutes

HARDER: Single-leg bridge, hip thrust with weight

Jumping Jack

PATTERN: Cardio | **MUSCLES:** Full body, Cardiovascular | **KCAL/30min:** 260-320 | **EQUIP:** None | **DIFF:** Beginner

INSTRUCTIONS:

1. Stand feet together.
2. Jump feet out while raising arms overhead.
3. Jump back to start.
4. Repeat rhythmically.

CUES:

- Light landing
- Arms fully overhead
- Continuous rhythm

MISTAKES:

- Heavy pounding
- Stiff arms
- No rhythm

EASIER: Step jack (no jump, step to side)

HARDER: Star jump, cross-jack

High Knees

PATTERN: Cardio | **MUSCLES:** Core, Hip Flexors, Cardiovascular | **KCAL/30min:** 300-360 | **EQUIP:** None | **DIFF:** Beginner

INSTRUCTIONS:

1. Run in place.
2. Drive knees up to hip height.
3. Pump arms.
4. Maintain upright posture.

CUES:

- Thigh parallel to floor at peak
- Land on balls of feet
- Core engaged

MISTAKES:

- Leaning back
- Shuffling vs lifting knees
- Looking down

EASIER: Slow march in place

HARDER: Add arm drive, increase speed

Tricep Dips (Chair)

PATTERN: Push | **MUSCLES:** Triceps, Shoulders, Chest | **KCAL/30min:** 150-190 | **EQUIP:** Chair/bench | **DIFF:** Beginner

INSTRUCTIONS:

1. Sit on edge of chair. Grip edge.
2. Slide off, legs straight or bent.
3. Lower until upper arms parallel.
4. Push up.

CUES:

- Keep back close to chair

- Elbows point back, not sideways
- Lower to 90° max

MISTAKES:

- Arms flaring wide
- Partial range
- Shrugging shoulders

EASIER: Bent knees (easier)

HARDER: Straight legs + elevated feet

Superman

PATTERN: Hip Extension / Back | MUSCLES: Lower back, Glutes, Rear Delts | KCAL/30min: 100-130 | EQUIP: None | DIFF: Beginner

INSTRUCTIONS:

1. Lie face down, arms extended.
2. Simultaneously raise arms and legs.
3. Hold 2-3 seconds.
4. Lower slowly.

CUES:

- Look at floor (neutral neck)
- Squeeze glutes
- Don't strain neck upward

MISTAKES:

- Neck hyperextension
- Only raising legs or arms (not both)
- Jerking movement

EASIER: Alternating arm/leg raises

HARDER: Add resistance band

Bear Crawl

PATTERN: Full Body | MUSCLES: Core, Shoulders, Hips, Quads | KCAL/30min: 280-340 | EQUIP: None — needs floor space | DIFF: Intermediate

INSTRUCTIONS:

1. Start on all fours. Lift knees 2 inches off floor.
2. Move opposite arm and leg forward.
3. Keep back flat and hips low.
4. Move 10-20 meters.

CUES:

- Hips level, not swaying
- Slow and controlled
- Breathe steadily

MISTAKES:

- Hips too high
- Alternating same-side limb
- Rushing

EASIER: Slow crawl

HARDER: Backward bear crawl, sideways

Wall Sit

PATTERN: Isometric Squat | MUSCLES: Quads, Glutes, Core | KCAL/30min: 130-160 | EQUIP: Wall | DIFF: Beginner

INSTRUCTIONS:

1. Stand back against wall.
2. Slide down until thighs parallel to floor.

3. Hold position. Breathe.

CUES:

- 90° at knees
- Back flat against wall
- Don't rest hands on thighs

MISTAKES:

- Partial depth
- Thighs not parallel
- Leaning forward

EASIER: Feet closer to wall (less depth)

HARDER: Single-leg wall sit

Step-Up

PATTERN: Squat/Lunge | MUSCLES: Quads, Glutes, Hamstrings | KCAL/30min: 200-260 | EQUIP: Stairs/step | DIFF: Beginner

INSTRUCTIONS:

1. Face a step or stair.
2. Step up with right foot, drive left knee up.
3. Step down controlled.
4. Alternate legs.

CUES:

- Drive through heel
- Stand fully upright at top
- Control the descent

MISTAKES:

- Using momentum
- Caving knee
- Pushing off bottom foot

EASIER: Lower step

HARDER: Weighted step-up, explosive step-up

Calf Raise

PATTERN: Ankle Extension | MUSCLES: Calves (gastrocnemius, soleus) | KCAL/30min: 80-100 | EQUIP: None (step for full ROM) | DIFF: Beginner

INSTRUCTIONS:

1. Stand on edge of step or flat floor.
2. Rise up on toes.
3. Hold 1 second.
4. Lower slowly.

CUES:

- Full ROM (heels below step level)
- Don't bounce
- Equal weight both feet

MISTAKES:

- Partial range
- Bouncing at bottom
- Swaying body

EASIER: Flat floor calf raise

HARDER: Single-leg calf raise, weighted

Barbell Back Squat

PATTERN: Squat | MUSCLES: Quads, Glutes, Hamstrings, Core | KCAL/30min: 280-360 | EQUIP: Barbell, Rack | DIFF: Advanced

INSTRUCTIONS:

1. Bar on upper traps. Unrack safely.
2. Feet shoulder-width, toes out slightly.
3. Break at hips and knees simultaneously.
4. Descend to parallel or below.
5. Drive through heels to stand.

CUES:

- Big breath and brace before descent
- Knees out over toes
- Chest up — not forward lean

MISTAKES:

- Knees caving
- Good morning squat (torso forward)
- Half reps

EASIER: Goblet squat, leg press

HARDER: Front squat, pause squat

Deadlift

PATTERN: Hip Hinge | MUSCLES: Hamstrings, Glutes, Lower Back, Traps, Core | KCAL/30min: 300-380 | EQUIP: Barbell | DIFF: Intermediate

INSTRUCTIONS:

1. Bar over mid-foot. Hip-width stance.
2. Hinge and grip just outside legs.
3. Shoulders slightly in front of bar.
4. Take big breath, brace, drive floor away.
5. Lock out at top — hips and knees extend simultaneously.

CUES:

- Bar stays close to body
- Lat engagement before pulling
- Lock top — don't hyperextend

MISTAKES:

- Bar away from body
- Rounding lower back
- Jerking from floor

EASIER: Romanian deadlift, trap bar deadlift

HARDER: Deficit deadlift, sumo deadlift

Bench Press

PATTERN: Push | MUSCLES: Chest, Triceps, Shoulders | KCAL/30min: 220-280 | EQUIP: Barbell, Bench | DIFF: Intermediate

INSTRUCTIONS:

1. Lie on bench. Bar over lower chest.
2. Grip slightly wider than shoulder-width.
3. Lower bar to chest with control.
4. Press up and slightly back.

CUES:

- Retract and depress scapulae
- Feet flat on floor
- Bar path slightly diagonal

MISTAKES:

- Flared elbows
- Bouncing bar off chest

- Feet off floor

EASIER: Dumbbell press, push-up

HARDER: Incline press, close-grip bench

Pull-Up

PATTERN: Pull | MUSCLES: Lats, Biceps, Rear Delts, Core | KCAL/30min: 200-260 | EQUIP: Pull-up bar | DIFF: Intermediate

INSTRUCTIONS:

1. Hang from bar, shoulder-width grip.
2. Initiate with lat contraction (depress shoulders).
3. Pull chest to bar.
4. Lower controlled.

CUES:

- Think 'elbows to hips'
- No kipping (strict)
- Full hang at bottom

MISTAKES:

- Kipping for reps (defeats purpose)
- Chin-up only (half ROM)
- Shrugging shoulders

EASIER: Lat pulldown, band-assisted pull-up

HARDER: Weighted pull-up, archer pull-up

Lat Pulldown

PATTERN: Pull | MUSCLES: Lats, Biceps, Rear Delts | KCAL/30min: 180-220 | EQUIP: Cable machine | DIFF: Beginner

INSTRUCTIONS:

1. Sit with thighs under pad.
2. Grip bar slightly wider than shoulder-width.
3. Pull bar to upper chest.
4. Return to full arm extension.

CUES:

- Lean back slightly — chest up
- Drive elbows down and back
- Don't use momentum

MISTAKES:

- Pulling to forehead
- Excessive lean back
- Short range

EASIER: Resistance band pulldown

HARDER: Single-arm lat pulldown, pull-up

Dumbbell Shoulder Press

PATTERN: Push | MUSCLES: Shoulders (anterior, lateral), Triceps | KCAL/30min: 180-220 | EQUIP: Dumbbells | DIFF: Beginner

INSTRUCTIONS:

1. Sit or stand. Dumbbells at shoulder height.
2. Press overhead to full extension.
3. Lower with control.

CUES:

- Don't arch back
- Full lockout at top
- Core braced

MISTAKES:

- Forward head at top
- Excessive arch
- Uneven pressing

EASIER: Lateral raise, machine shoulder press

HARDER: Arnold press, barbell press

Cable Row

PATTERN: Pull | **MUSCLES:** Mid-Back, Lats, Biceps, Rear Delts | **KCAL/30min:** 180-220 | **EQUIP:** Cable machine | **DIFF:** Beginner

INSTRUCTIONS:

1. Sit at cable, feet on platform.
2. Hinge slightly forward.
3. Row handle to abdomen.
4. Squeeze shoulder blades.
5. Return fully.

CUES:

- Drive elbows back, not arms
- Squeeze at contraction
- Torso stays upright

MISTAKES:

- Rocking with momentum
- Not squeezing at peak
- Shrugging

EASIER: Dumbbell row

HARDER: Single-arm cable row, T-bar row

Leg Press

PATTERN: Squat | **MUSCLES:** Quads, Glutes, Hamstrings | **KCAL/30min:** 200-280 | **EQUIP:** Leg press machine | **DIFF:** Beginner

INSTRUCTIONS:

1. Sit in machine. Feet shoulder-width on platform.
2. Release safety locks.
3. Lower sled until 90° knee angle.
4. Press up — don't lock knees.

CUES:

- Don't let lower back lift off pad
- Feet flat throughout
- Controlled descent

MISTAKES:

- Locking knees at top
- Too heavy — half reps
- Feet too low (knee stress)

EASIER: Reduce weight, increase ROM

HARDER: Single-leg press, narrow stance

Romanian Deadlift (RDL)

PATTERN: Hip Hinge | **MUSCLES:** Hamstrings, Glutes, Lower Back | **KCAL/30min:** 220-280 | **EQUIP:** Barbell or Dumbbells | **DIFF:** Intermediate

INSTRUCTIONS:

1. Stand holding bar at hip height.
2. Push hips back, bar slides down thighs.
3. Feel hamstring stretch.
4. Drive hips forward to stand.

CUES:

- Soft bend in knees — not squat
- Bar close to body
- Neutral spine throughout

MISTAKES:

- Rounding back
- Going too low (losing back position)
- Squatting instead of hinging

EASIER: Good morning, dumbbell RDL

HARDER: Single-leg RDL, deficit RDL

Dumbbell Bicep Curl

PATTERN: Pull / Isolation | MUSCLES: Biceps, Brachialis | KCAL/30min: 120-150 | EQUIP: Dumbbells | DIFF: Beginner

INSTRUCTIONS:

1. Stand or sit. Dumbbells hanging at sides.
2. Curl up, rotating to supinate wrist.
3. Squeeze at top.
4. Lower controlled.

CUES:

- Elbows stay at sides
- No swinging body
- Full ROM

MISTAKES:

- Swinging momentum
- Elbows forward
- Partial range

EASIER: Resistance band curl

HARDER: Concentration curl, hammer curl, incline curl

Tricep Pushdown

PATTERN: Push / Isolation | MUSCLES: Triceps | KCAL/30min: 120-150 | EQUIP: Cable machine | DIFF: Beginner

INSTRUCTIONS:

1. Face cable, grab rope or bar.
2. Elbows at sides.
3. Extend arms down fully.
4. Return controlled.

CUES:

- Elbows pinned to sides
- Full extension at bottom
- Don't lean forward for more weight

MISTAKES:

- Elbows flaring
- Partial extension
- Using too much weight

EASIER: Bench dip

HARDER: Overhead cable extension, close-grip bench

Cable Crunch

PATTERN: Core | MUSCLES: Rectus Abdominis, Obliques | KCAL/30min: 120-160 | EQUIP: Cable machine | DIFF: Intermediate

INSTRUCTIONS:

1. Kneel facing cable, rope held at head.

2. Crunch torso down, not just head.
3. Pause at bottom.
4. Return slowly.

CUES:

- Move from abs, not neck
- Round spine during crunch
- Consistent resistance

MISTAKES:

- Pulling with neck
- Hip flexor dominant
- Standing up fully between reps

EASIER: Ab mat crunches

HARDER: Weighted decline crunch

Leg Curl (Machine)

PATTERN: Knee Flexion | MUSCLES: Hamstrings | KCAL/30min: 140-180 | EQUIP: Leg curl machine | DIFF: Beginner

INSTRUCTIONS:

1. Lie face down (prone) on machine.
2. Curl legs up fully.
3. Lower with control.

CUES:

- Hips flat on pad
- Full ROM
- No hip lifting on way down

MISTAKES:

- Partial range
- Hips rising
- Excessive weight

EASIER: Reduce weight

HARDER: Nordic curl (advanced)

Hip Thrust

PATTERN: Hip Extension | MUSCLES: Glutes, Hamstrings | KCAL/30min: 180-240 | EQUIP: Bench + Barbell/Dumbbell | DIFF: Intermediate

INSTRUCTIONS:

1. Upper back on bench, barbell across hips.
2. Feet flat, knees bent.
3. Drive hips up to full extension.
4. Squeeze glutes hard.
5. Lower controlled.

CUES:

- Chin tucked
- Hips fully extend at top
- Drive through heels

MISTAKES:

- Not reaching full extension
- Lower back hyperextension
- Feet too close or far

EASIER: Glute bridge on floor

HARDER: Single-leg hip thrust, banded hip thrust

Cable Lateral Raise

PATTERN: Shoulder Isolation | MUSCLES: Medial Deltoid | KCAL/30min: 100-130 | EQUIP: Cable machine | DIFF: Intermediate

INSTRUCTIONS:

1. Stand side-on to cable (low pulley).
2. Raise arm to 90°, slight elbow bend.
3. Lower controlled.

CUES:

- Lead with elbow, not wrist
- Don't shrug
- Stop at shoulder height

MISTAKES:

- Too heavy — shrugging
- Swinging body
- Going above shoulder height

EASIER: Dumbbell lateral raise

HARDER: Leaning cable lateral raise

Tabata Sprint (Treadmill/Outdoor)

PATTERN: Cardio | MUSCLES: Full body, Cardiovascular | KCAL/30min: 300-450 | EQUIP: Treadmill or outdoor space | DIFF: Advanced

INSTRUCTIONS:

1. 20 seconds max effort sprint.
2. 10 seconds rest.
3. Repeat 8 rounds = 4 minutes.
4. Rest 1-2 min.
5. Repeat 3-5 cycles.

CUES:

- True max effort during work
- Full recovery during rest
- Warm up 5 min before

MISTAKES:

- Going 70% effort (defeats HIIT)
- Skipping warm-up
- Too many rounds early on

EASIER: 30 sec jog / 30 sec walk

HARDER: 6-8 full Tabata cycles

Jump Rope (Skipping)

PATTERN: Cardio | MUSCLES: Calves, Core, Shoulders, Cardio | KCAL/30min: 280-380 | EQUIP: Jump rope | DIFF: Beginner

INSTRUCTIONS:

1. Stand with feet together.
2. Rotate rope from wrists (not arms).
3. Jump just high enough for rope.
4. Land on balls of feet.

CUES:

- Small jumps
- Elbows close to sides
- Consistent rhythm

MISTAKES:

- Jumping too high
- Using whole arm to rotate

- Irregular rhythm

EASIER: Step over rope (no jump)

HARDER: Double-unders, criss-cross

Box Jump

PATTERN: Power | MUSCLES: Quads, Glutes, Calves, Core | KCAL/30min: 280-360 | EQUIP: Box/platform | DIFF: Intermediate

INSTRUCTIONS:

1. Stand facing box.
2. Dip and swing arms.
3. Jump explosively onto box.
4. Land soft in squat.
5. Step down (don't jump down).

CUES:

- Land with bent knees
- Swing arms for momentum
- Always step down safely

MISTAKES:

- Jumping down (knee stress)
- Landing stiff
- Box too high

EASIER: Low box, squat jump

HARDER: Single-leg box jump, depth jump

Kettlebell Swing

PATTERN: Hip Hinge/Power | MUSCLES: Glutes, Hamstrings, Core, Shoulders | KCAL/30min: 300-400 | EQUIP: Kettlebell | DIFF: Intermediate

INSTRUCTIONS:

1. Stand with KB between feet.
2. Hinge hips back, swing KB between legs.
3. Drive hips forward explosively.
4. KB floats to chest height.
5. Let it swing back.

CUES:

- Power from hips, not arms
- Squeeze glutes at top
- Let gravity control descent

MISTAKES:

- Squatting instead of hinging
- Arms doing the lifting
- Rounded back

EASIER: Dumbbell swing, wall hip hinge

HARDER: One-arm swing, American swing (overhead)

Battle Ropes

PATTERN: Cardio/Endurance | MUSCLES: Shoulders, Core, Arms, Cardio | KCAL/30min: 280-360 | EQUIP: Battle ropes | DIFF: Intermediate

INSTRUCTIONS:

1. Hold one rope in each hand.
2. Alternate arm waves quickly.
3. Keep knees soft.
4. Maintain for prescribed time.

CUES:

- Waves must reach anchor point
- Core tight
- Breathe rhythmically

MISTAKES:

- Stopping waves mid-set
- Stiff knees
- Losing core engagement

EASIER: Slow waves

HARDER: Double waves, slams, alternating slams

Rowing Machine

PATTERN: Cardio/Total Body | **MUSCLES:** Back, Legs, Arms, Core, Cardio | **KCAL/30min:** 280-380 | **EQUIP:** Rowing machine | **DIFF:** Intermediate

INSTRUCTIONS:

1. Sit, feet strapped.
2. Extend legs, then lean back, then pull.
3. Return: arms, body, legs.
4. Sequence: legs 60%, body 20%, arms 20%.

CUES:

- Push don't pull — legs first
- Drive through heels
- Maintain tall posture

MISTAKES:

- Arms first (weak sequence)
- Rounding back
- Short strokes

EASIER: Slow rate (18-20 spm)

HARDER: Sprint intervals, max distance tests

Cycling (Indoor/Outdoor)

PATTERN: Cardio | **MUSCLES:** Quads, Glutes, Hamstrings, Calves, Cardio | **KCAL/30min:** 200-380 | **EQUIP:** Bicycle or spin bike | **DIFF:** Beginner

INSTRUCTIONS:

1. Adjust seat to hip height.
2. Pedal at comfortable cadence.
3. For HIIT: sprint 30s, easy 90s.
4. For steady state: 60-70% max HR.

CUES:

- Push and pull pedals
- Maintain cadence over gear
- Keep shoulders relaxed

MISTAKES:

- Seat too low (knee strain)
- Hunching over handlebars
- Gear too heavy

EASIER: Slower pace, lower intensity

HARDER: Hill sprints, HIIT intervals

Surya Namaskar (Sun Salutation)

PATTERN: Full Body Flow | **MUSCLES:** Full body | **KCAL/30min:** 180-250 | **EQUIP:** Yoga mat | **DIFF:** Beginner

INSTRUCTIONS:

1. 12-step sequence: Mountain → Forward fold → Plank → Cobra → Downward dog → Repeat.
2. Complete 5-10 rounds per session.
3. Breathe in sync with each movement.

CUES:

- Synchronize breath with movement
- Hold each position briefly
- Flow not rush

MISTAKES:

- Rushing
- Holding breath
- Skipping warm-up

EASIER: Modified with knees down

HARDER: Add jump forward and back

Warrior I (Virabhadrasana I)

PATTERN: Strength + Stretch | **MUSCLES:** Quads, Hips, Shoulders | **KCAL/30min:** 100-140 | **EQUIP:** Yoga mat | **DIFF:** Beginner

INSTRUCTIONS:

1. Step right foot forward, left back.
2. Bend front knee to 90°.
3. Arms overhead, hips square forward.
4. Hold 30-60 seconds.
5. Switch sides.

CUES:

- Back heel pressed down
- Front knee over ankle
- Arms reach actively

MISTAKES:

- Back heel lifting
- Knee collapsing inward
- Hunching shoulders

EASIER: Chair yoga modification

HARDER: Warrior III (balance pose)

Hip Flexor Stretch

PATTERN: Flexibility | **MUSCLES:** Hip Flexors, Psoas | **KCAL/30min:** 60-80 | **EQUIP:** Yoga mat | **DIFF:** Beginner

INSTRUCTIONS:

1. Kneel on right knee.
2. Left foot forward in lunge.
3. Shift hips forward and down.
4. Feel stretch in right hip flexor.
5. Hold 30-60 seconds.

CUES:

- Posterior pelvic tilt (tuck pelvis)
- Upright torso
- Breathe into the stretch

MISTAKES:

- Anterior pelvic tilt (arch back)
- Not shifting hips forward enough
- Rushing

EASIER: Supine hip flexor stretch

HARDER: Half-kneeling hip flexor with rotation

Downward Dog

PATTERN: Full Body Stretch | MUSCLES: Hamstrings, Calves, Shoulders, Spine | KCAL/30min: 80-100 | EQUIP: Yoga mat | DIFF: Beginner

INSTRUCTIONS:

1. Start on all fours.
2. Push hips up and back.
3. Form inverted V shape.
4. Straighten legs (or soft knees).
5. Heels press toward floor.

CUES:

- Long spine — not rounded
- Weight equal in hands and feet
- Active shoulders — don't collapse

MISTAKES:

- Rounded upper back
- Heels forced flat (injurious)
- Wrist pain from poor setup

EASIER: Puppy stretch

HARDER: Three-legged dog, dolphin pose

Cat-Cow Stretch

PATTERN: Spinal Mobility | MUSCLES: Spine, Core, Neck | KCAL/30min: 60-80 | EQUIP: Yoga mat | DIFF: Beginner

INSTRUCTIONS:

1. Start on all fours.
2. Inhale: drop belly, lift head and tailbone (Cow).
3. Exhale: round spine, tuck chin (Cat).
4. Repeat 10-15 cycles.

CUES:

- Breath drives movement
- Slow and controlled
- Full spinal range

MISTAKES:

- Moving only neck or only lower back
- Holding breath
- Rushing

EASIER: Seated version in chair

HARDER: Flow into child's pose, thread the needle

Walking (Outdoor/Treadmill)

PATTERN: Cardio/LISS | MUSCLES: Legs, Core, Cardiovascular | KCAL/30min: 100-200 | EQUIP: Shoes | DIFF: Beginner

INSTRUCTIONS:

1. Walk at brisk pace (5-6.5 km/h).
2. Swing arms naturally.
3. Maintain upright posture.
4. Target: 30-60 min/day.

CUES:

- Brisk pace — can talk, not sing
- Incline adds intensity
- Morning walk boosts fat oxidation

MISTAKES:

- Slouching
- Staring at phone
- Walking too slowly

EASIER: 10 min daily, increase by 5 min/week

HARDER: Incline walk, weighted vest, intervals

Complete Workout Programs

WL-WP-000	MOD 07 — Workout Programs · Program Directory
GOALS	Weight Loss Fat Loss General Health
USERS	Adult · Teen · Senior · Student · Hostel Student · Office Worker
DIFF	Beginner
TAGS	workout program, weekly schedule, beginner gym, home workout, weight loss plan

DEFINITION

Complete collection of 7 structured workout programs for all fitness levels and lifestyles.

WHY IT MATTERS

- Program adherence beats random workouts by 3x for long-term results.
- A schedule removes daily decision fatigue and builds habit automatically.

KEY FACTS

- Programs: 7 (Beginner to Advanced)
- Covers: Home, Gym, Hostel, Office, Senior
- Includes weekly schedules, sets/reps, progression, expected results.
- HIIT 2-3x/week burns 30% more fat than steady-state cardio alone.

ACTION STEPS

1. Select program matching your level and equipment.
2. Follow for full duration — minimum 6-8 weeks before switching.
3. Track exercises weekly. Progressive overload is mandatory.
4. Combine with calorie deficit from Module 05.

DECISION TREE

- IF Beginner + Home → THEN WL-WP-001 or WL-WP-003 (women)
- IF Beginner + Hostel → THEN WL-WP-004
- IF Beginner + Office → THEN WL-WP-005
- IF Intermediate + Gym → THEN WL-WP-002
- IF Advanced + Gym → THEN WL-WP-006
- IF Senior → THEN WL-WP-007
- IF Not sure → THEN Start WL-WP-001 for 4 weeks then reassess

WL-WP-001	MOD 07 — Workout Programs · Beginner Home Weight Loss › Beginner Weight Loss
GOALS	Weight Loss Fat Loss
USERS	Adult · Student
DIFF	Beginner
TAGS	beginner, weight_loss, workout, program

DEFINITION

Beginner Home Weight Loss — 8 weeks, 4 days/week.

WHY IT MATTERS

- Designed for: Sedentary adult, no equipment, total beginner

KEY FACTS

- Duration: 8 weeks
- Frequency: 4 days/week
- Expected result: 4-6 kg fat loss in 8 weeks with 500 kcal daily deficit.

ACTION STEPS

1. Mon: Full Body A
2. Tue: Walk
3. Wed: Full Body B

4. Thu: Rest
5. Fri: HIIT Circuit
6. Sat: Walk
7. Sun: Rest
8. Jumping Jacks — 3 sets x 30 sec
9. Bodyweight Squat — 3 x 12
10. Push-Up (knee ok) — 3 x 8-12
11. Plank — 3 x 20-30 sec
12. Glute Bridge — 3 x 15
13. Mountain Climber — 3 x 20 sec
14. Brisk Walk — 30-45 min
15. High Knees — 3 x 30 sec
16. Lunge — 3 x 10 each
17. Tricep Dip (chair) — 3 x 8-12
18. Superman — 3 x 12
19. Step-Up — 3 x 12 each
20. Wall Sit — 3 x 30 sec
21. Active rest — stretch 15 min or gentle walk —
22. Burpees — 3 x 8
23. Jump Squat — 3 x 10
24. Mountain Climber — 3 x 30 sec
25. Push-Up — 3 x 10
26. High Knees — 3 x 30 sec
27. Walk — 45-60 min
28. Full rest or yoga —
29. Progression: Week 1-2: base. Week 3-4: +2 reps each set. Week 5-6: +1 set. Week 7-8: reduce rest 10 sec.
30. Notes: Walk every day if possible. Steps are the biggest lever for beginners.

WL-WP-002	MOD 07 — Workout Programs · Intermediate Gym Fat Loss › Intermediate Fat Loss
GOALS	Weight Loss Fat Loss
USERS	Adult · Student
DIFF	Intermediate
TAGS	intermediate, fat_loss, workout, program

DEFINITION

Intermediate Gym Fat Loss — 12 weeks, 5 days/week.

WHY IT MATTERS

- Designed for: Gym member with 3-6 months experience, wants to lose fat and tone

KEY FACTS

- Duration: 12 weeks
- Frequency: 5 days/week
- Expected result: 6-10 kg fat loss + noticeable muscle tone in 12 weeks.

ACTION STEPS

1. Mon: Push (Chest/Shoulder/Triceps)
2. Tue: Pull (Back/Biceps)
3. Wed: Cardio/Core
4. Thu: Legs
5. Fri: Full Body + Core
6. Sat: LISS Cardio
7. Sun: Rest
8. Barbell Bench Press — 4 x 8-10
9. Dumbbell Shoulder Press — 3 x 10

10. Incline DB Press — 3 x 10
11. Cable Lateral Raise — 3 x 12
12. Tricep Pushdown — 3 x 12
13. Barbell Deadlift — 4 x 5
14. Pull-Up — 3 x max
15. Cable Row — 3 x 10
16. Lat Pulldown — 3 x 10
17. DB Bicep Curl — 3 x 12
18. HIIT Treadmill (Tabata) — 4 cycles
19. Plank — 3 x 45 sec
20. Cable Crunch — 3 x 15
21. Mountain Climber — 3 x 30 sec
22. Barbell Squat — 4 x 8
23. Romanian Deadlift — 3 x 10
24. Leg Press — 3 x 12
25. Leg Curl — 3 x 12
26. Calf Raise — 4 x 15
27. Hip Thrust — 3 x 10
28. Kettlebell Swing — 4 x 15
29. Pull-Up — 3 x max
30. DB Shoulder Press — 3 x 10
31. Box Jump — 3 x 8
32. Cable Crunch — 3 x 15
33. Walk or cycle — 45-60 min moderate
34. Full rest —
35. Progression: Progressive overload: increase weight 2.5kg when you hit upper rep range for 2 consecutive sessions.
36. Notes: Track all lifts. Eat 1g protein per pound bodyweight. Sleep 7-9 hours.

WL-WP-003	MOD 07 — Workout Programs · Home Workout — Women's Weight Loss › Beginner-Intermediate Weight Loss + Toning
GOALS	Weight Loss Fat Loss
USERS	Adult · Student
DIFF	Beginner-Intermediate
TAGS	beginner-intermediate, weight_loss+_toning, workout, program

DEFINITION

Home Workout — Women's Weight Loss — 8 weeks, 5 days/week.

WHY IT MATTERS

- Designed for: Women, home setting, wants to lose weight and get toned

KEY FACTS

- Duration: 8 weeks
- Frequency: 5 days/week
- Expected result: 4-7 kg weight loss + toned appearance in 8 weeks.

ACTION STEPS

1. Mon: Lower Body Focus
2. Tue: Cardio + Core
3. Wed: Upper Body
4. Thu: Walk + Yoga
5. Fri: Full Body HIIT
6. Sat: Active Recovery
7. Sun: Rest
8. Squat — 4 x 15

9. Lunge — 3 x 12 each
10. Glute Bridge — 4 x 15
11. Step-Up — 3 x 12
12. Calf Raise — 3 x 20
13. Jumping Jacks — 3 x 45 sec
14. High Knees — 3 x 30 sec
15. Plank — 4 x 30 sec
16. Crunch — 3 x 20
17. Mountain Climber — 3 x 30 sec
18. Push-Up — 3 x 10-15
19. Tricep Dip — 3 x 12
20. Superman — 3 x 15
21. DB Row (water bottle ok) — 3 x 12
22. Walk 30 min —
23. Hip Flexor Stretch —
24. Cat-Cow — 3 x 10
25. Warrior I — 30 sec each side
26. Burpee — 3 x 10
27. Jump Squat — 3 x 15
28. Push-Up — 3 x 10
29. Mountain Climber — 3 x 30 sec
30. High Knees — 3 x 30 sec
31. Walk 45 min —
32. Rest —
33. Progression: Add 2 reps each exercise every week.
34. Notes: Women should not fear weights. Muscle burns fat 24/7.

WL-WP-004	MOD 07 — Workout Programs · Hostel Student Home Workout › Beginner Weight Loss
GOALS	Weight Loss Fat Loss
USERS	Adult · Student
DIFF	Beginner
TAGS	beginner, weight_loss, workout, program

DEFINITION

Hostel Student Home Workout — 6 weeks, 4 days/week + daily walk.

WHY IT MATTERS

- Designed for: Hostel student, small room, no equipment, limited time

KEY FACTS

- Duration: 6 weeks
- Frequency: 4 days/week + daily walk
- Expected result: 2-4 kg in 6 weeks with diet adherence.

ACTION STEPS

- Mon: 20-Min Bodyweight
- Tue: Walk
- Wed: 20-Min HIIT
- Thu: Rest
- Fri: 20-Min Full Body
- Sat: Long Walk
- Sun: Rest
- Jumping Jacks 45 sec / 15 sec rest — 3 rounds
- Push-Up — 3 x 10
- Squat — 3 x 15

11. Plank — 3 x 30 sec
12. Walk to college or evening walk — 30-45 min
13. High Knees 30 sec / rest 10 sec — 8 rounds Tabata
14. Burpee — 3 x 8
15. Lunge — 3 x 10 each
16. Superman — 3 x 12
17. Stretch or walk —
18. Bear Crawl 20 steps — 3 sets
19. Push-Up — 3 x max
20. Glute Bridge — 4 x 15
21. Wall Sit — 3 x 45 sec
22. 60-min walk with friends —
23. Rest —
24. Progression: Increase time by 5 min each week. Add 2 reps/round.
25. Notes: Workout in your room, hallway, or rooftop. 20 min is enough.

WL-WP-005 **MOD 07 — Workout Programs · Office Worker Lunch Break Workout › Beginner | Activity + Weight Loss**

GOALS Weight Loss | Fat Loss

USERS Adult · Student

DIFF Beginner

TAGS beginner, activity_+_weight_loss, workout, program

DEFINITION

Office Worker Lunch Break Workout — Ongoing, 5 days/week (lunch) + weekend.

WHY IT MATTERS

- Designed for: Office worker, 30-min lunch break, wants to be more active

KEY FACTS

- Duration: Ongoing
- Frequency: 5 days/week (lunch) + weekend
- Expected result: Adds 200-300 kcal burn daily. 3-5 kg in 3 months.

ACTION STEPS

1. Mon-Fri: 30-Min Lunch Walk + Mini Workout
2. Sat: 45-Min Workout
3. Sun: Walk
4. Brisk walk outdoors — 20 min
5. Push-Up (any surface) — 2 x 10
6. Squat — 2 x 15
7. Plank — 2 x 30 sec
8. Full body circuit — any 5 bodyweight exercises — 3 rounds
9. Family walk or jog — 30-45 min
10. Progression: Add 5 min to walks weekly. Add 1 exercise to circuit.
11. Notes: Steps matter. Park farther. Take stairs. Stand every 60 min.

WL-WP-006 **MOD 07 — Workout Programs · Advanced Gym Program — Body Recomposition › Advanced | Fat Loss + Muscle Gain**

GOALS Weight Loss | Fat Loss

USERS Adult · Student

DIFF Advanced

TAGS advanced, fat_loss+_muscle_gain, workout, program

DEFINITION

Advanced Gym Program — Body Recomposition — 16 weeks, 5-6 days/week.

WHY IT MATTERS

- Designed for: Experienced gym-goer wanting to cut fat while maintaining muscle

KEY FACTS

- Duration: 16 weeks
- Frequency: 5-6 days/week
- Expected result: 5-8% body fat reduction + strength maintained/increased in 16 weeks.

ACTION STEPS

1. Mon: Chest + Core
2. Tue: Back + Biceps
3. Wed: HIIT + Mobility
4. Thu: Legs
5. Fri: Shoulders + Arms
6. Sat: Optional: Cardio or Weak Point
7. Sun: Rest
8. Barbell Bench Press — 5 x 5
9. Incline DB Press — 4 x 8
10. Cable Fly — 3 x 12
11. Weighted Cable Crunch — 4 x 12
12. Plank — 3 x 60 sec
13. Deadlift — 5 x 3
14. Weighted Pull-Up — 4 x 6
15. Barbell Row — 4 x 8
16. Hammer Curl — 3 x 12
17. Face Pull — 3 x 15
18. Sprint Intervals 15 min —
19. Yoga/Mobility 20 min —
20. Core Circuit —
21. Barbell Squat — 5 x 5
22. Romanian Deadlift — 4 x 8
23. Leg Press — 3 x 12
24. Nordic Curl — 3 x 6
25. Hip Thrust (heavy) — 4 x 8
26. Calf Raise — 5 x 15
27. Barbell OHP — 4 x 8
28. Arnold Press — 3 x 10
29. Lateral Raise — 4 x 15
30. Skull Crusher — 3 x 10
31. Incline Curl — 3 x 10
32. 45 min cardio or lagging muscle —
33. Full rest —
34. Progression: Linear progression on main lifts. Deload every 6 weeks.
35. Notes: Requires precise nutrition. Track macros daily. 1g protein per lb BW minimum.

WL-WP-007	MOD 07 — Workout Programs · Senior Weight Management Program › Beginner Weight Loss + Mobility
GOALS	Weight Loss Fat Loss
USERS	Adult · Student
DIFF	Beginner
TAGS	beginner, weight_loss+_mobility, workout, program

DEFINITION

Senior Weight Management Program — 12 weeks, 4 days/week.

WHY IT MATTERS

- Designed for: Adults 55+, weight management, joint-friendly exercises

KEY FACTS

- Duration: 12 weeks
- Frequency: 4 days/week
- Expected result: 2-3 kg in 12 weeks. More importantly: improved mobility and energy.

ACTION STEPS

1. Mon: Strength + Balance
 2. Tue: Walk
 3. Wed: Yoga + Mobility
 4. Thu: Rest
 5. Fri: Strength + Core
 6. Sat: Walk + Social
 7. Sun: Rest
 8. Seated Chair Squat — 3 x 10
 9. Wall Push-Up — 3 x 10
 10. Single-Leg Balance — 3 x 30 sec
 11. Seated Leg Raise — 3 x 12
 12. 30-min gentle walk —
 13. Cat-Cow — 3 x 10
 14. Hip Flexor Stretch — 30 sec each
 15. Warrior I (gentle) — 20 sec each
 16. Downward Dog —
 17. Rest —
 18. Glute Bridge — 3 x 12
 19. Superman — 3 x 10
 20. Standing Calf Raise — 3 x 15
 21. Plank (knees ok) — 3 x 20 sec
 22. 45-min walk with family/friends —
 23. Rest —
 24. Progression: Very gradual. Never sacrifice form. Progress only if pain-free.
 25. Notes: Consult doctor before starting if any chronic conditions. Low impact only.
-

Special Populations

WL-SP-000	MOD 08 — Special Populations · Special Population Overview
GOALS	Weight Loss Fat Loss General Health
USERS	Adult · Teen · Senior · PCOS · Diabetes · Postpartum · Vegan · Vegetarian · Student
DIFF	Intermediate
TAGS	special population, PCOS, diabetes, hypothyroid, senior, teen, vegan, hostel

DEFINITION

Dedicated weight-loss protocols for 11 special population groups.

WHY IT MATTERS

- Standard plans fail special populations because physiology differs.
- Customized approach produces 2-3x better outcomes than generic plans.

KEY FACTS

- PCOS (Polycystic Ovary Syndrome)
- Type 2 Diabetes / Prediabetes
- Hypothyroidism
- Teenagers (13-18 years)
- Senior Citizens (60+ years)
- Postpartum Weight Loss
- Night Shift Workers
- Students (Hostel + College)
- Vegetarians
- Vegans
- Frequent Travelers

ACTION STEPS

1. Identify which population group applies.
2. Follow that group's specific guidelines over standard Module 05 plans.
3. Medical consultation is recommended for: PCOS, Diabetes, Hypothyroid, Postpartum, Severe Obesity.

DECISION TREE

- IF Has PCOS → THEN WL-SP-001
- IF Has Diabetes → THEN WL-SP-002
- IF Has Hypothyroid → THEN WL-SP-003
- IF Is Teen → THEN WL-SP-004
- IF Is Senior 60+ → THEN WL-SP-005
- IF Postpartum → THEN WL-SP-006
- IF Night Shift → THEN WL-SP-007
- IF Hostel Student → THEN WL-SP-008
- IF Vegetarian → THEN WL-SP-009
- IF Vegan → THEN WL-SP-010
- IF Frequent Traveler → THEN WL-SP-011

WL-SP-001	MOD 08 — Special Populations · PCOS (Polycystic Ovary Syndrome)
GOALS	Weight Loss Fat Loss Health
USERS	Adult
DIFF	Intermediate
TAGS	pcos_(polycystic_ovary_syndrome), special, protocol

DEFINITION

Hormonal disorder causing insulin resistance, irregular periods, and difficulty losing weight.

WHY IT MATTERS

- Insulin resistance makes standard calorie restriction less effective alone.
- High androgens increase fat storage, especially abdominal fat.

KEY FACTS

- 1 in 5 Indian women has PCOS.
- Weight loss of 5-10% body weight can restore menstrual cycles.
- Low GI diet reduces androgen levels within 8 weeks.
- Strength training > cardio for PCOS.

ACTION STEPS

1. Eliminate refined carbs: maida, white sugar, processed foods.
2. Eat low-GI: jowar, ragi, oats, legumes, vegetables.
3. Protein at every meal: minimum 25g.
4. Strength train 3x/week — improves insulin sensitivity.
5. Add inositol-rich foods: brown rice, citrus, legumes.
6. Methi seeds (fenugreek) in water daily — proven insulin sensitizer.
7. Sleep 7-9 hours — sleep deprivation worsens androgen levels.
8. Stress management — cortisol drives androgen production.

DECISION TREE

- IF** PCOS + Obese (BMI>30) → **THEN** 1400-1600 kcal low-GI plan. Medical consultation.
- IF** PCOS + Normal weight → **THEN** 1600-1800 kcal. Focus on hormonal health over weight.
- IF** PCOS + Insulin resistance → **THEN** Metformin (consult doctor) + low-carb diet.
- IF** PCOS + Irregular periods → **THEN** Prioritize stress, sleep, low GI before aggressive deficit.
- IF** PCOS + Acne → **THEN** Dairy reduction trial: 4 weeks dairy-free observation.
- IF** PCOS + Hair loss → **THEN** Add zinc, iron, B12 supplementation after blood test.

COMMON MISTAKES

- Extreme calorie restriction worsens hormonal imbalance.
- High sugar fruits (mango, litchi) in large quantities.
- Skipping meals — causes blood sugar spikes and crashes.
- Only cardio — strength training is more effective for PCOS.
- Dairy can worsen acne in some PCOS cases — trial elimination.

SPECIAL CASES

- PCOS weight loss is SLOWER than standard. 0.25-0.5 kg/week is excellent progress.

AI COACHING NOTES

- *If user reports PCOS (Polycystic Ovary Syndrome): follow WL-SP-001 protocol.*
- *Key intervention for PCOS (Polycystic Ovary Syndrome): Eliminate refined carbs: maida, white sugar, processed foods.*

WL-SP-002	MOD 08 — Special Populations · Type 2 Diabetes / Prediabetes
GOALS	Weight Loss Fat Loss Health
USERS	Adult
DIFF	Intermediate
TAGS	type_2_diabetes___prediabetes, special, protocol

DEFINITION

Impaired blood glucose regulation. Weight loss is the most effective intervention.

WHY IT MATTERS

- 5-10% body weight loss can reverse prediabetes completely.
- Exercise improves glucose uptake without insulin.
- Low GI diet prevents dangerous glucose spikes.

KEY FACTS

- India has the 2nd most diabetics globally.
- Bariatric surgery achieves 86% remission rate — diet mimics this mechanism.
- Walking 30 min after meals reduces postprandial glucose by 25%.

- Every 1 kg lost = 1mmol/L reduction in blood sugar.

ACTION STEPS

1. Monitor blood sugar before and 2 hrs after meals.
2. Target post-meal glucose <140 mg/dL (7.8 mmol/L).
3. Never eat rice and roti at same meal.
4. Walk 15-30 min after every major meal.
5. Start with 1600-1800 kcal — never go below 1200 without medical supervision.
6. Prioritize fiber: 35-40g/day minimum.
7. Protein at every meal reduces post-meal glucose spike.
8. Avoid all fruit juices, sweetened beverages, refined grains.

DECISION TREE

- IF** HbA1c >9% → **THEN** Medical management first. Diet as adjunct.
- IF** HbA1c 7-9% → **THEN** Low GI + calorie deficit + walk after meals.
- IF** Prediabetes (HbA1c 5.7-6.4%) → **THEN** Diet and exercise CAN reverse this fully.
- IF** Diabetic + Obese → **THEN** 1400-1600 kcal, low carb, high fiber.
- IF** Diabetic + Medication → **THEN** Tell doctor about diet change — medication doses may need reduction as blood sugar improves.
- IF** Diabetic + Athlete → **THEN** Higher carbs but timing critical — around training sessions.

COMMON MISTAKES

- Eating rice + roti + roti (double carb meals).
- Fruit juice instead of whole fruit (removes fiber that slows glucose).
- Skipping medications for diet alone (without doctor guidance).
- Going very low carb suddenly without monitoring.

AI COACHING NOTES

- If user reports Type 2 Diabetes / Prediabetes: follow WL-SP-002 protocol.
- Key intervention for Type 2 Diabetes / Prediabetes: Monitor blood sugar before and 2 hrs after meals.

WL-SP-003	MOD 08 — Special Populations · Hypothyroidism
GOALS	Weight Loss Fat Loss Health
USERS	Adult
DIFF	Intermediate
TAGS	hypothyroidism, special, protocol

DEFINITION

Underactive thyroid producing insufficient T3/T4, slowing metabolism by 10-15%.

WHY IT MATTERS

- Hypothyroidism reduces TDEE by 200-400 kcal/day.
- Standard plans fail because metabolism is genuinely slower.
- Iron deficiency often co-exists and worsens thyroid function.

KEY FACTS

- India has 42 million people with hypothyroidism.
- Women are 5-8x more likely than men.
- Optimal TSH for weight loss: 0.5-2.5 mIU/L.
- Selenium and iodine are essential for T3/T4 production.

ACTION STEPS

1. Ensure TSH is optimized with medication before expecting weight loss.
2. Reduce calorie target by 200 kcal vs standard (metabolism is slower).
3. Avoid raw goitrogens: raw broccoli, raw cauliflower, raw cabbage (cooking neutralizes them).
4. Add selenium: Brazil nuts (1-2/day), eggs, sunflower seeds.
5. Check ferritin (iron stores) — iron deficiency worsens thyroid function.
6. High protein diet: 1.2-1.5g/kg body weight.
7. Resistance training boosts metabolism — critical for hypothyroid patients.

8. Take medication 30-60 min before eating for full absorption.

DECISION TREE

IF Uncontrolled hypothyroid (high TSH) → **THEN** Focus on medication compliance first.

IF Controlled hypothyroid (normal TSH) → **THEN** Standard plan works — just expect slower progress.

IF Hypothyroid + PCOS → **THEN** Treat both simultaneously. Low GI + strength training.

IF Hypothyroid + Very slow loss (<0.25kg/week) → **THEN** Recheck TSH. Consider dose adjustment with doctor.

COMMON MISTAKES

- Blaming hypothyroid for all weight issues — adherence still matters.
- Avoiding all exercise because of fatigue.
- Soy foods in excess (may interfere with absorption — avoid within 4 hrs of medication).
- Not rechecking thyroid levels every 6 months.

AI COACHING NOTES

- *If user reports Hypothyroidism: follow WL-SP-003 protocol.*
- *Key intervention for Hypothyroidism: Ensure TSH is optimized with medication before expecting weight loss.*

WL-SP-004	MOD 08 — Special Populations · Teenagers (13-18 years)
GOALS	Weight Loss Fat Loss Health
USERS	Adult
DIFF	Intermediate
TAGS	teenagers_(13-18_years), special, protocol

DEFINITION

Growing adolescents who need to manage weight without compromising development.

WHY IT MATTERS

- Restrictive diets harm bone density, hormonal development, and brain function in teens.
- Weight management in teens should be activity-first, not restriction-first.

KEY FACTS

- Minimum 1600-1800 kcal for active teenage girls, 1800-2200 for boys.
- Bone mass peaks at 18-20 — calcium (1200mg/day) is non-negotiable.
- Eating disorders most commonly emerge in teenage years — watch for signs.

ACTION STEPS

1. Never drop below 1600 kcal without medical supervision.
2. Add activity before reducing food.
3. Replace junk food — don't restrict meals.
4. Ensure calcium: 1200mg/day (4 glasses milk/curd).
5. Ensure iron for girls: dal, leafy greens, eggs.
6. Remove sugary drinks — biggest change for most teens.
7. Walk/cycle/sport daily — minimum 1 hour movement.

DECISION TREE

IF Overweight teen (BMI >85th percentile) → **THEN** Activity increase + junk food removal. No extreme diets.

IF Obese teen (BMI >95th percentile) → **THEN** Pediatrician + dietitian required.

IF Teen athlete → **THEN** Higher calories. Performance over aesthetics.

IF Teen with eating disorder signs → **THEN** Immediate professional help. Stop all coaching.

COMMON MISTAKES

- Prescribing adult weight-loss diets to teens.
- Encouraging very low-calorie eating.
- Shaming body weight — causes lasting psychological harm.
- Allowing meal skipping for weight loss.

AI COACHING NOTES

- *If user reports Teenagers (13-18 years): follow WL-SP-004 protocol.*
- *Key intervention for Teenagers (13-18 years): Never drop below 1600 kcal without medical supervision.*

WL-SP-005	MOD 08 — Special Populations · Senior Citizens (60+ years)
GOALS	Weight Loss Fat Loss Health
USERS	Adult
DIFF	Intermediate
TAGS	senior_citizens_(60+_years), special, protocol

DEFINITION

Older adults needing weight management while preserving muscle, bone, and mobility.

WHY IT MATTERS

- Sarcopenia (muscle loss) after 60 accelerates weight gain even without eating more.
- Falls are leading cause of injury in seniors — strength and balance are life-saving.
- Protein needs INCREASE with age despite lower calorie needs.

KEY FACTS

- Muscle loss rate: 1-2% per year after 60.
- Protein needs: 1.2-1.6g/kg (vs 0.8g/kg for younger adults).
- Vitamin D + Calcium: essential for bone health in seniors.
- Low-intensity exercise has identical metabolic benefits to high-intensity in seniors.

ACTION STEPS

1. Protein priority: eggs, paneer, curd, dal at every meal.
2. Strength training 2-3x/week with light weights — prevents muscle loss.
3. Walk daily: 30-45 min at comfortable pace.
4. Calcium 1200mg/day + Vitamin D 1000 IU/day.
5. Hydration: seniors lose thirst sensitivity — drink 8 glasses regardless of thirst.
6. High fiber: prevents constipation, manages blood sugar.
7. Avoid excessive calorie restriction — maintain 1600-1800 minimum.

DECISION TREE

- IF** Obese senior (BMI>30) → **THEN** Modest 200-300 kcal deficit. High protein. Medical supervision.
- IF** Senior with diabetes → **THEN** Low GI + walk after meals.
- IF** Senior with hypertension → **THEN** Low sodium. High potassium (bananas, potatoes).
- IF** Senior with osteoporosis → **THEN** Weight-bearing exercise + calcium + Vit D.
- IF** Senior with arthritis → **THEN** Pool exercises, cycling, chair yoga — protect joints.

COMMON MISTAKES

- Very low calorie diets — causes muscle and bone loss.
- Avoiding all fat — fat-soluble vitamins (A, D, E, K) are essential.
- No strength training — it's most important exercise in old age.
- Too much sodium — hidden in pickles, papad, processed food.

AI COACHING NOTES

- *If user reports Senior Citizens (60+ years): follow WL-SP-005 protocol.*
- *Key intervention for Senior Citizens (60+ years): Protein priority: eggs, paneer, curd, dal at every meal.*

WL-SP-006	MOD 08 — Special Populations · Postpartum Weight Loss
GOALS	Weight Loss Fat Loss Health
USERS	Adult
DIFF	Intermediate
TAGS	postpartum_weight_loss, special, protocol

DEFINITION

Weight management for new mothers after childbirth.

WHY IT MATTERS

- Body needs healing before aggressive weight loss.
- Breastfeeding increases calorie needs significantly.
- Hormonal changes make fat loss slower — patience required.

KEY FACTS

- Wait 6-8 weeks post-delivery before any structured program.
- Breastfeeding burns 300-500 extra kcal/day.
- Minimum 1800 kcal while breastfeeding.
- Sleep deprivation increases cortisol and fat storage — prioritize sleep over exercise.

ACTION STEPS

1. Weeks 1-8: rest, nutrition, healing. No formal exercise.
2. Week 8-12: gentle walking begins.
3. Week 12+: progressive exercise if cleared by OB-GYN.
4. If breastfeeding: minimum 1800 kcal. Never below 1600.
5. High protein: supports milk production and fat loss.
6. Iron and B12: often depleted post-delivery — test and supplement.
7. Partner and family support for sleep is the biggest lever.

COMMON MISTAKES

- Rushing to lose baby weight immediately.
- Calorie restriction while breastfeeding — affects milk supply.
- High-intensity exercise before 12 weeks.
- Comparing to celebrities with trainers and help.

AI COACHING NOTES

- If user reports Postpartum Weight Loss: follow WL-SP-006 protocol.
- Key intervention for Postpartum Weight Loss: Weeks 1-8: rest, nutrition, healing. No formal exercise.

WL-SP-007	MOD 08 — Special Populations · Night Shift Workers
GOALS	Weight Loss Fat Loss Health
USERS	Adult
DIFF	Intermediate
TAGS	night_shift_workers, special, protocol

DEFINITION

Workers with schedules that disrupt circadian rhythms.

WHY IT MATTERS

- Circadian disruption reduces leptin, increases ghrelin — drives overeating.
- Metabolic rate is 10-15% lower at night even when active.
- Higher risk of insulin resistance and type 2 diabetes.

KEY FACTS

- Night shift workers gain 5-8 kg over 2 years vs day workers.
- Digestive efficiency is 20-30% lower for late-night meals.
- Light exposure during the day off helps reset circadian rhythm.

ACTION STEPS

1. Eat main meals before shift (not during night hours).
2. Keep night-time eating light: salads, protein, low-carb.
3. Avoid heavy meals 3-4 hours before sleep (whenever that is).
4. Consistent sleep schedule on days off — don't fully flip schedule.
5. Blackout curtains and sleep mask for day sleeping.
6. Exercise before shift, not right before sleep.
7. Meal prep is essential — night canteens have poor options.

DECISION TREE

IF Night shift + overweight → **THEN** 1600-1800 kcal. Main meals during day/evening before shift.

IF Night shift + diabetic → **THEN** Protein-heavy, low-carb night snacks.

IF Night shift + poor sleep quality → **THEN** Sleep optimization before diet change.

COMMON MISTAKES

- Large meals at 2-3am (worst metabolic time).

- High sugar snacks to stay awake — crashing blood sugar worsens fatigue.
- Excessive caffeine late in shift — disrupts sleep quality.

AI COACHING NOTES

- *If user reports Night Shift Workers: follow WL-SP-007 protocol.*
- *Key intervention for Night Shift Workers: Eat main meals before shift (not during night hours).*

WL-SP-008 MOD 08 — Special Populations · Students (Hostel + College)

GOALS	Weight Loss Fat Loss Health
USERS	Adult
DIFF	Intermediate
TAGS	students_(hostel+_college), special, protocol

DEFINITION

18-25 year olds living in hostels or sharing accommodation with limited cooking access.

WHY IT MATTERS

- Hostel food is often high calorie, high fat, and low protein.
- Budget constraints limit healthy food access.
- Social eating and irregular schedules complicate adherence.

KEY FACTS

- Average hostel student gains 3-5 kg in first year of college.
- Stress eating and late-night eating are primary drivers.
- Home-cooked food access even 2x/week significantly improves outcomes.

ACTION STEPS

1. Carry: peanut butter, roasted chana, protein powder, oats, fruits.
2. At mess: choose dal > fried items. Small rice, extra salad.
3. Avoid: samosa, puri, chai-biscuit cycle.
4. Walk to class — avoid auto for distances under 2 km.
5. Morning bodyweight workout in room: 20 min, 4x/week.
6. Track food for 2 weeks — awareness is the first step.
7. Hydrate: drink water before each mess meal.

DECISION TREE

- IF** Hostel student + overweight → **THEN** Start with junk elimination + 30 min walk daily.
- IF** Hostel student + no cooking → **THEN** Non-cook high-protein foods: eggs, curd, PB, roasted chana, sprouts.
- IF** Hostel student + very tight budget → **THEN** Dal + rice + banana + eggs + curd = ■80/day full nutrition.

COMMON MISTAKES

- Skipping mess food and relying on Maggi/biscuits.
- Chai + biscuit habit: 400-600 extra kcal/day.
- Alcohol on weekends: 500-1000 extra kcal, lowers inhibitions for late-night eating.
- Buying supplements before fixing diet basics.

AI COACHING NOTES

- *If user reports Students (Hostel + College): follow WL-SP-008 protocol.*
- *Key intervention for Students (Hostel + College): Carry: peanut butter, roasted chana, protein powder, oats, fruits.*

WL-SP-009 MOD 08 — Special Populations · Vegetarians

GOALS	Weight Loss Fat Loss Health
USERS	Adult
DIFF	Intermediate
TAGS	vegetarians, special, protocol

DEFINITION

People who exclude meat/fish but may include dairy and eggs.

WHY IT MATTERS

- Getting 100g+ protein daily requires intentional planning on vegetarian diet.
- Plant proteins are less bioavailable — eat 10-20% more.

KEY FACTS

- India has world's largest vegetarian population.
- Vegetarian diet when well-planned shows equal or better weight-loss outcomes vs omnivore.
- Key risk: protein deficiency, B12 deficiency, iron deficiency.

ACTION STEPS

1. Protein anchors: paneer, curd, moong dal, rajma, chole, soya chunks, tofu, eggs (if egg-veg).
2. Minimum 25-30g protein per meal.
3. B12 supplement: 500-1000 mcg/day (no B12 in plant foods).
4. Iron: dal + citrus (vitamin C enhances absorption).
5. Combine grains + legumes (dal + rice, roti + dal) for complete protein.
6. Whey protein powder if struggling to hit protein targets.

DECISION TREE

IF Veg + low protein → **THEN** Add: soya chunks (52g/100g dry), paneer, curd at every meal.

IF Veg + B12 deficient → **THEN** Supplement immediately. Tingling/fatigue are deficiency symptoms.

IF Veg + high carb → **THEN** Reduce rice portions. Increase dal, paneer, vegetable ratio.

COMMON MISTAKES

- Relying only on dal for protein — need variety.
- Large rice/roti portions with minimal dal.
- Thinking vegetarian automatically means healthy — veg junk food exists.
- Skipping B12 supplementation (neurological damage is permanent if untreated).

AI COACHING NOTES

■ *If user reports Vegetarians: follow WL-SP-009 protocol.*

■ *Key intervention for Vegetarians: Protein anchors: paneer, curd, moong dal, rajma, chole, soya chunks, tofu, eggs (if egg-veg).*

WL-SP-010	MOD 08 — Special Populations · Vegans
GOALS	Weight Loss Fat Loss Health
USERS	Adult
DIFF	Intermediate
TAGS	vegans, special, protocol

DEFINITION

People excluding all animal products including dairy and eggs.

WHY IT MATTERS

- Protein and B12 are most challenging nutrients for weight-loss vegans.
- Well-planned vegan diet supports fat loss effectively.

KEY FACTS

- Vegans have 1-2 BMI units lower than non-vegans on average.
- B12 deficiency risk: 90%+ of vegans without supplementation.
- Key nutrients: B12, D3, iron, calcium, zinc, omega-3 (algae-based).

ACTION STEPS

1. Protein sources: tofu, tempeh, edamame, soya chunks, legumes, seitan.
2. B12: 2500 mcg/week (cyanocobalamin) — non-negotiable.
3. Calcium: fortified plant milk, ragi, sesame seeds, leafy greens.
4. D3 (vegan): 1000-2000 IU/day from lichen-derived supplement.
5. Omega-3: 250mg DHA+EPA from algae oil daily.
6. Iron: legumes + vitamin C at every meal.
7. Zinc: pumpkin seeds, legumes, whole grains.
8. Vegan protein powder (pea + rice blend) to hit targets easily.

COMMON MISTAKES

- Assuming vegan = automatically nutritious.
- Skipping B12 — devastating neurological consequences.
- Low protein intake causing muscle loss during fat loss.
- Over-relying on nuts and avocado (high fat = high calorie).

AI COACHING NOTES

- *If user reports Vegans: follow WL-SP-010 protocol.*
- *Key intervention for Vegans: Protein sources: tofu, tempeh, edamame, soya chunks, legumes, seitan.*

WL-SP-011	MOD 08 — Special Populations · Frequent Travelers
GOALS	Weight Loss Fat Loss Health
USERS	Adult
DIFF	Intermediate
TAGS	frequent_travelers, special, protocol

DEFINITION

People who travel for work or personal reasons, disrupting routine.

WHY IT MATTERS

- Travel increases reliance on restaurant food, disrupts sleep, and reduces exercise time.
- Airport/railway food is high calorie and low protein.

ACTION STEPS

1. Always carry: protein powder sachet, roasted chana, nuts, protein bar.
2. Hotel breakfast: eggs, oats, curd, fruits — avoid fried items.
3. Restaurant strategy: order grilled protein + salad as default.
4. Airport: coffee (no sugar), eggs if available, nuts, water.
5. Hotel room workout: 20-min bodyweight circuit (no equipment needed).
6. Hydrate aggressively: flights dehydrate severely.
7. Don't fast all day and binge at dinner — eat small, regular meals.

DECISION TREE

- IF** Business travel (1-2 days) → **THEN** Maintain plan strictly. Return to full plan immediately.
- IF** Long travel (5+ days) → **THEN** Damage control: prioritize protein, limit obvious junk.
- IF** International travel → **THEN** Jet lag disrupts metabolism — prioritize sleep over perfect eating.

AI COACHING NOTES

- *If user reports Frequent Travelers: follow WL-SP-011 protocol.*
- *Key intervention for Frequent Travelers: Always carry: protein powder sachet, roasted chana, nuts, protein bar.*

Behaviour Change System

WL-BH-001	MOD 09 · Habit Formation › Building Weight-Loss Habits
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Student · Office Worker
DIFF	Beginner
TAGS	habit, habit formation, routine, consistency, behavior change, atomic habits

DEFINITION

A habit is an automatic behaviour triggered by a cue. Weight loss success depends more on habit automation than willpower. The Habit Loop: Cue → Routine → Reward. Identity-based habits ('I am a healthy person') are more durable than outcome-based ('I want to lose 10 kg').

WHY IT MATTERS

- Willpower is finite — habits eliminate the decision cost of healthy behaviours.
- Consistent execution of imperfect habits beats inconsistent perfect plans.
- It takes 21-66 days for a behaviour to become automatic (average 66 days).

KEY FACTS

- Habit stacking: attach new habit to an existing anchor (e.g., 'after I make chai, I walk for 10 min').
- Environment design is more powerful than motivation — remove friction from healthy choices.
- Minimum effective dose: smallest version of habit that counts (2-min walk beats no walk).
- Tracking creates accountability and reinforces identity — 'I am someone who tracks'.

ACTION STEPS

1. Identify your top 3 target habits: e.g., track food, walk 30 min, hit protein target.
2. Attach each habit to an existing anchor (habit stack).
3. Make healthy choice the default (prep meals Sunday, put oats on counter, put walking shoes at door).
4. Track habits on a simple calendar — don't break the chain.
5. Start embarrassingly small: 5 min walk, 1 tracked meal, 1 glass water — build gradually.

DECISION TREE

IF user cannot stick to any diet plan → **THEN** Ask: what's the minimum version of this habit? Start with that only

IF user doing well for 2 weeks then fails → **THEN** Identify the trigger event; create a 'when X happens, I will Y' response plan

IF user motivated Monday, gives up by Thursday → **THEN** Replace motivation-dependent habits with environment design

COMMON MISTAKES

- Setting 10 new habits simultaneously — 1-2 per week maximum.
- Waiting for motivation before acting — act first, motivation follows.
- Treating one slip as total failure — '2 missed days, restart tomorrow'.
- Aiming for perfect instead of consistent.

FAQ

Q: How long will it take to form healthy eating habits?

A: Research shows 21-66 days for habit automaticity, average 66 days. Focus on the first 2 weeks of consistency — then it gets easier.

Q: I know what to do but can't make myself do it. Why?

A: Knowing ≠ doing. The gap is habit design, not willpower. Make the healthy choice frictionless and the unhealthy choice inconvenient. Remove chips from home; put fruit on the counter.

Q: What if I miss a day?

A: Miss one, never two. A single missed day has minimal impact. Two consecutive missed days starts a new pattern. Always return the next day.

AI COACHING NOTES

- When user fails: normalize. Ask 'what was the trigger?' not 'why did you fail?'
- Design the environment before motivating the person.

■ Celebrate small wins explicitly — this builds habit identity.

WL-BH-002	MOD 09 · Motivation vs Discipline › Building Consistency
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Student · Office Worker
DIFF	Beginner
TAGS	motivation, discipline, consistency, willpower, momentum, mindset

DEFINITION

Motivation is an emotion — it comes and goes. Discipline is a skill — it can be trained. The most consistent weight-loss achievers are those who removed dependence on motivation through systems, schedules, and environment design.

WHY IT MATTERS

- Motivation peaks on Monday, dips Thursday — using it as the foundation guarantees failure.
- Systems (meal prep, pre-scheduled workouts, prepared food) replace the need for daily motivation.
- Action precedes motivation — start moving and motivation follows, not vice versa.

KEY FACTS

- Ego depletion: willpower is a limited resource that depletes through use. Remove decisions.
- Implementation intentions ('When I wake up, I will do 10 push-ups') increase follow-through by 91%.
- Identity shift is the deepest level of change: 'I am someone who exercises' vs 'I want to exercise'.

ACTION STEPS

1. Replace 'I need to be motivated' with 'I have a schedule'. Schedule workouts like meetings.
2. Create if-then rules: 'If I don't feel like working out, I will do 10 minutes minimum'.
3. Tell someone your plan — social commitment doubles follow-through.
4. Build momentum on good days; maintain minimum on bad days.
5. Reduce decision fatigue: same breakfast, same workout time, meal prepped lunches.

DECISION TREE

IF user says 'I just don't feel motivated' → **THEN** reframe: motivation is a feeling, action is a choice. What's the 2-minute version you can do right now?

IF user does great for 2 weeks then falls off → **THEN** identify the disruption event; create contingency plans for that scenario

IF user only exercises when they feel like it → **THEN** move workouts to morning before motivation depletes; schedule + accountability buddy

COMMON MISTAKES

- Relying on motivational videos instead of building systems.
- Quitting when results slow down (plateau is normal, not failure).
- Comparing journey to others — each body and starting point is different.

FAQ

Q: How do I stay motivated?

A: Don't try to. Build systems that work regardless of motivation. Meal prep eliminates the 'I don't feel like cooking' decision. Scheduled gym time eliminates the 'do I go today?' decision.

AI COACHING NOTES

- When user lacks motivation: don't try to re-motivate — reduce the task to its smallest version.
- Ask: 'What system would make this automatic?' not 'How can I want it more?'

WL-BH-003	MOD 09 · Sleep and Weight Loss › Sleep Optimization
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Student · Office Worker
DIFF	Beginner
TAGS	sleep, weight loss, sleep deprivation, ghrelin, cortisol, sleep quality, 7 hours, recovery

DEFINITION

Sleep is a foundational pillar of weight management. Sleep deprivation increases ghrelin (hunger), decreases leptin (fullness), raises cortisol (fat storage), impairs insulin sensitivity, and reduces NEAT. 7-9 hours per night is non-negotiable for optimal fat loss.

WHY IT MATTERS

- 5 hours vs 8 hours sleep: 559 kcal more consumed the next day (study data).
- Sleep deprivation causes 50% of fat loss to come from lean muscle (not fat) during deficit.
- Cortisol from poor sleep drives abdominal fat accumulation — the most dangerous kind.
- Growth hormone (fat-burning hormone) is primarily released during deep sleep.

KEY FACTS

- Ghrelin increases 28% with sleep deprivation. Leptin decreases 18%.
- Night shift workers have 35-40% higher obesity rates than day workers.
- Quality matters: 6 hours of uninterrupted sleep > 8 hours of fragmented sleep.
- Alcohol: improves sleep onset but reduces sleep quality (disrupts REM).

ACTION STEPS

1. Set consistent sleep and wake times — even on weekends.
2. Sleep-inducing environment: dark (blackout curtains), cool (18-20°C), quiet.
3. No screens 30-60 min before bed — blue light suppresses melatonin.
4. Avoid caffeine after 2pm (half-life of 5-7 hours).
5. Magnesium glycinate 200-300mg at night: improves sleep depth.
6. No heavy meals within 3 hours of sleep.

DECISION TREE

IF user sleeps <6 hours and weight loss is stalled → **THEN** prioritize sleep before changing diet; add 1 hour sleep and reassess in 2 weeks

IF user is a new parent with fragmented sleep → **THEN** reduce workout intensity; focus on nutrition over exercise; nap when possible

IF user has insomnia → **THEN** refer to sleep hygiene protocol; magnesium supplement; consider sleep specialist

IF user works night shift → **THEN** module 08 WL-SP-007; adjust eating window to day/evening hours

COMMON MISTAKES

- Cutting sleep to create time for exercise — net negative effect.
- Scrolling phone in bed — light exposure and mental stimulation delay sleep onset.
- Alcohol to 'help sleep' — impairs deep sleep stages critical for recovery.

FAQ

Q: Can I lose fat with poor sleep?

A: Yes, but you'll lose 50% more muscle and 50% less fat at the same calorie deficit, per research. Poor sleep makes fat loss harder and muscle loss easier — exactly the wrong combination.

Q: I sleep 8 hours but still tired. Why?

A: Check: sleep quality, not just quantity. Alcohol, screen use before bed, stress, and sleep apnea can impair quality. Also check vitamin D, iron, and thyroid levels.

AI COACHING NOTES

- Sleep is often the missing variable when diet and exercise are both in place.
- When user is highly stressed and sleep-deprived: address these before adding more diet restriction.
- Track sleep with any fitness tracker — concrete data changes behaviour.

WL-BH-004	MOD 09 · Stress and Weight Loss › Cortisol Management
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Student · Office Worker
DIFF	Intermediate
TAGS	stress, cortisol, emotional eating, stress weight gain, belly fat, work stress, anxiety

DEFINITION

Chronic stress elevates cortisol, which promotes fat storage (especially visceral/abdominal), increases sugar cravings, promotes muscle breakdown, disrupts sleep, and drives emotional eating. Stress management is a metabolic intervention, not just a mental health one.

WHY IT MATTERS

- Cortisol signals fat cells to store more fat and release less.

- Stress-induced eating is typically high-fat, high-sugar (comfort foods) — 400-800 extra kcal/episode.
- Indian professional and student populations are among the most chronically stressed globally.

KEY FACTS

- Cortisol response peaks 20-30 min after stressor. Returns to baseline in 60-90 min with active management.
- Exercise (moderate): reduces cortisol. Exercise (excessive): raises cortisol.
- Ashwagandha (KSM-66 extract, 300-600mg/day): shown to reduce cortisol by 27% in RCTs.
- Box breathing (4-4-4-4) activates parasympathetic response within 2-5 minutes.

ACTION STEPS

1. Identify primary stress sources: academic, work, relationship, financial.
2. Build 1 daily stress outlet: 20-min walk, 10-min meditation, 5-min breathing.
3. Box breathing protocol: inhale 4 sec, hold 4 sec, exhale 4 sec, hold 4 sec — 5 rounds.
4. Phone-free meals — eating while stressed impairs digestion and overeating detection.
5. Social connection reduces cortisol — weekly time with close friends/family.
6. Consider ashwagandha supplement if chronic stress persists.

DECISION TREE

IF user reports high stress + belly fat gain despite diet adherence → **THEN** cortisol-driven pattern; stress management is treatment, not just diet

IF user stress eats when anxious → **THEN** identify stress triggers; create alternative response: walk, call friend, breathe

IF user reports 'exam stress' + weight gain → **THEN** sleep + stress management primary; reduce workout intensity during exam period

COMMON MISTAKES

- Adding aggressive diet restriction during high-stress periods — compound negative effect.
- Using food as the only stress management tool.
- Excessive exercise to manage stress — overtraining worsens cortisol.

FAQ

Q: Why do I gain weight when stressed even though I'm not eating more?

A: Cortisol directly promotes fat storage regardless of calories — especially visceral fat. It also promotes water retention and reduces insulin sensitivity. Manage the stress, not just the food.

AI COACHING NOTES

- *When user is highly stressed: 'Let's stabilize first — not add more pressure with strict dieting.'*
- *Teach the box breathing technique in check-in — it's immediately applicable.*

WL-BH-005	MOD 09 · Emotional Eating › Identification and Intervention
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Student · Office Worker
DIFF	Intermediate
TAGS	emotional eating, binge, stress eating, food and emotions, overeating, mindful eating

DEFINITION

Emotional eating is eating in response to emotional triggers (stress, boredom, loneliness, sadness, anxiety) rather than physiological hunger. It is typically sudden, specific (cravings for comfort food), and followed by guilt. Estimated 30-40% of dieters engage in regular emotional eating.

WHY IT MATTERS

- Emotional eating can override a perfect dietary plan in a single episode.
- Unaddressed emotional eating always returns — willpower cannot permanently suppress it.
- Understanding the emotion-food link is essential for long-term weight management.

KEY FACTS

- HALT check before eating: Hungry? Angry? Lonely? Tired? Eat only if Hungry.
- Physical hunger is gradual; emotional hunger is sudden and craving-specific.
- A 10-min pause between craving and eating reduces emotional eating episodes by 40%.
- Emotional eating is often a learned coping mechanism — can be unlearned.

ACTION STEPS

1. Before eating, rate hunger on 1-10 scale. Eat only if score >4.
2. HALT check every time you reach for food outside meal time.
3. Identify your top 3 emotional eating triggers (write them down).
4. Create a '3 before eating' rule: drink water, walk 5 min, breathe 5 cycles — if still hungry, eat.
5. Replace emotional eating with non-food coping: call a friend, journal, walk, listen to music.
6. Seek professional support (psychologist) if binge eating occurs >2x/week.

DECISION TREE

IF user eats when bored → **THEN** help identify non-food activities for boredom; reduce food accessibility during boredom periods

IF user eats when anxious → **THEN** build anxiety toolkit: breathing, walking, journaling; address root anxiety source

IF user binges after restriction → **THEN** break restriction-binge cycle; ensure adequate calories during the day

IF user eats when lonely → **THEN** social connection is the cure; address loneliness directly rather than through food

COMMON MISTAKES

- Trying to 'white-knuckle' through emotional eating without addressing the emotion.
- Restricting heavily after emotional eating episode — sets up the binge-restrict cycle.
- Shame spiraling — 'I ate emotionally, I'm a failure' — this worsens the cycle.

FAQ

Q: Is emotional eating a serious problem?

A: Yes, when it occurs regularly and prevents weight-loss progress. It's the primary cause of diet failure for ~30% of people. It's addressable with the right tools — and sometimes requires professional support.

Q: I know I'm eating emotionally but can't stop. What do I do?

A: In the moment: pause 10 min before eating. After: be compassionate, not punishing. Long-term: work with a therapist trained in emotional eating. Apps like Wysa or Headspace support the process.

AI COACHING NOTES

- *Never shame emotional eating — it's a coping mechanism, not a character flaw.*
- *When severe: refer to mental health professional — this is outside coaching scope.*
- *Focus on building non-food coping toolkit rather than willpower.*

WL-BH-006	MOD 09 · Cravings › Managing Food Cravings
GOALS	Weight Loss Fat Loss
USERS	Adult · Teen · Senior · Student · Office Worker
DIFF	Intermediate
TAGS	cravings, sugar craving, junk food, sweet tooth, craving management, dopamine, food addiction

DEFINITION

Cravings are intense desires for specific foods, often high in sugar, fat, or salt. They are driven by blood sugar fluctuations, hormones (ghrelin, dopamine), habit cues, emotional states, and nutrient deficiencies. Cravings can be managed but not permanently suppressed through restriction alone.

WHY IT MATTERS

- Unmanaged cravings cause 80% of diet abandonment episodes.
- Cravings have physiological roots (not just weak willpower) — addressing root causes is more effective.
- Adequate protein and fiber reduce cravings by stabilizing blood sugar and satiety hormones.

KEY FACTS

- Sugar cravings peak 20-30 min after onset — waiting them out works 60-70% of the time.
- Chromium (200-400mcg/day) reduces sugar cravings in insulin-resistant individuals.
- Magnesium deficiency causes chocolate cravings (chocolate is high in magnesium).
- Strong cravings typically last only 15-20 minutes if not acted on.
- High-protein diet reduces cravings by 60% vs low-protein diet (study data).

ACTION STEPS

1. When craving hits: drink 500ml water. Wait 20 min. Reassess.
2. High-protein, high-fiber meals prevent most cravings from occurring.
3. Identify craving triggers: time of day, emotions, situations — avoid or replace.

4. For sweet cravings: dark chocolate (70%+) 20g, fruit, or sweetened curd.
5. For salt cravings: roasted chana, buttermilk with salt — satisfy without excess.
6. Include one 'planned treat' per week — removes forbidden fruit psychology.
7. Check magnesium, zinc, and iron levels if cravings are persistent.

DECISION TREE

IF user craves sugar after meals → **THEN** blood sugar spike/crash pattern; recommend protein-fat-fiber at meals to stabilize

IF user craves junk food at night → **THEN** usually habit + low-protein dinner; restructure dinner; remove junk food from home

IF user craves chocolate daily → **THEN** possible magnesium deficiency; supplement and observe

IF user has intense cravings before period → **THEN** hormonal PMS cravings; planned small treats + magnesium + adequate calories during cycle

COMMON MISTAKES

- Complete elimination of all treats — increases cravings through restriction psychology.
- Buying 'diet' versions of cravings (diet chips, sugar-free cookies) — often triggers more cravings.
- Eating to 'get cravings out of the system' — works temporarily, not long-term.

FAQ

Q: Do cravings go away on a diet?

A: For most people: yes, after 3-4 weeks of stable eating patterns and adequate protein. The first 2 weeks are hardest. Cravings significantly reduce once blood sugar stabilizes.

Q: Is it OK to have a small treat?

A: Yes. One planned treat per week or small daily dark chocolate is sustainable and prevents binge eating. Perfect adherence 100% of the time is not required.

AI COACHING NOTES

- *When user has strong cravings: check protein intake first — low protein is the #1 cause.*
- *Teach the surf the craving technique: observe craving like a wave, it peaks and passes.*
- *Planned treats are strategically superior to total restriction.*

WL-BH-007	MOD 09 · Environment Design › Setting Up for Success
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Student · Office Worker
DIFF	Beginner
TAGS	environment design, food environment, home setup, friction, defaults, nudge

DEFINITION

Environment design is structuring your physical and social environment so that healthy choices are the easy, default choices. More powerful than motivation and willpower because it works automatically without effort.

WHY IT MATTERS

- 80% of food decisions are made automatically based on what is visible and accessible.
- Removing unhealthy food from home reduces consumption by 70% — it's not there to eat.
- Adding friction to unhealthy choices and reducing friction for healthy ones shifts behaviour without willpower.

KEY FACTS

- People eat 20-30% more when eating from larger plates/bowls.
- Visibility effect: fruit bowl on counter = 69% more fruit consumption. Chips in visible bowl = 60% more consumed.
- Grocery shopping hungry leads to 30-40% more junk food purchased.
- People who meal prep Sunday eat 42% fewer fast food meals during the week.

ACTION STEPS

1. Kitchen audit: remove all junk food from home — can't eat what isn't there.
2. Place fruit bowl at eye level. Move snacks to high shelves.
3. Use smaller plates (22-25cm) and bowls — automatically reduces portions.
4. Pre-cut and refrigerate vegetables at eye level in fridge.
5. Keep gym bag packed by the door.
6. Grocery shop with a list after eating — never when hungry.
7. Prep breakfast the night before — removes morning decision fatigue.

DECISION TREE

IF user buys healthy but eats junk at home → **THEN** kitchen audit: remove junk; if partner/family buys junk, negotiate dedicated shelf

IF user defaults to takeout after work → **THEN** Sunday meal prep; make cooking faster than ordering; batch cook proteins

IF user snacks mindlessly in front of TV → **THEN** no food in TV room rule; replace with nuts in measured 30g portion if needed

COMMON MISTAKES

- Relying on willpower when junk food is in the house.
- Shopping hungry — fills trolley with impulse, calorie-dense purchases.
- Large serving vessels — use smaller ones; size of dishware predicts portion size.

FAQ

Q: I live with family who eat unhealthy — how do I manage?

A: Negotiate: 'This is my food in this section of the fridge'. Don't ask family to change — just control your environment within the shared space. Eat your prepped food before family meals so you're not hungry at the table.

AI COACHING NOTES

- *Environment design is the highest-leverage intervention — do this before anything else.*
- *For hostel students: limited environment control — focus on what you carry (peanut butter, chana, protein powder).*

WL-BH-008	MOD 09 · Social Eating and Peer Pressure › Navigating Social Situations
GOALS	Weight Loss Fat Loss
USERS	Adult · Teen · Senior · Student · Office Worker
DIFF	Intermediate
TAGS	social eating, peer pressure, restaurant, party, wedding, eating out, social pressure

DEFINITION

Social eating situations — restaurants, family dinners, weddings, festivals, parties — are among the most commonly cited reasons for dietary failure. Having a social eating strategy prevents one-time events from becoming pattern disruptions.

WHY IT MATTERS

- Social eating is 20-50% higher in calorie density than solo eating.
- Social norms ('everyone's having dessert') override individual intentions.
- Indian culture has strong food-hospitality norms — refusing food can feel culturally difficult.

ACTION STEPS

1. Eat a protein-rich snack before social events — arrive not hungry.
2. At buffets: survey everything first, then plate once with deliberate choices.
3. Order first at restaurants — removes peer influence on your choice.
4. Polite decline script: 'I'm watching my digestion / I had a big lunch' — culturally acceptable.
5. Alcohol: max 1-2 drinks, choose dry wine or spirits over beer/cocktails.
6. At weddings: eat the main course well; skip or minimize sweets/fried.
7. The 3-bite rule for desserts: 3 bites of anything satisfies 80% of the craving.

DECISION TREE

IF user's family insists on serving more food → **THEN** use smaller serving dish + eat slowly + compliment the food + 'saving room for dessert' technique

IF user overeats at every social event → **THEN** pre-event snack strategy + identify specific trigger (hunger, peer pressure, anxiety)

IF user refuses all social events to stay on diet → **THEN** this is unsustainable and isolating; teach moderation strategy instead

FAQ

Q: Should I avoid all social eating?

A: No. Avoiding social life destroys mental health and sustainability. Master moderation strategies — one social meal per week barely impacts a monthly trend.

Q: How do I handle desi aunties forcing food?

A: Polite culturally-fluent responses: 'Bahut swaadisht hai, par abhi pet bhar gaya' (It's delicious but I'm full). Take a small serving and eat slowly.

AI COACHING NOTES

- Teach social eating scripts — practical language users can actually use.
- Frame it as 80/20 — 80% on-plan, 20% flexible = sustainable for life.

WL-BH-009	MOD 09 · Mindful Eating › Eating with Awareness
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Student · Office Worker
DIFF	Beginner
TAGS	mindful eating, slow eating, satiety, fullness cues, overeating, distracted eating

DEFINITION

Mindful eating is the practice of eating with full attention — noticing hunger and fullness cues, eating slowly, and removing distractions. It reduces meal size by 100-200 kcal without any dietary restriction.

WHY IT MATTERS

- Satiety hormones (CCK, PYY, GLP-1) take 15-20 min to signal fullness — eating fast bypasses this.
- Distracted eating (phone, TV) increases meal intake by 25-40%.

KEY FACTS

- Eating speed: fast eaters are 115% more likely to be overweight vs slow eaters.
- Chewing each bite 20-30 times: reduces meal size by ~100 kcal automatically.
- Japanese 'Hara Hachi Bu': eat to 80% full — embedded in culture of the world's longest-lived people.

ACTION STEPS

1. Eat without phone or TV. If alone, eat quietly.
2. Put utensil down between bites.
3. Aim for 20-minute minimum per meal.
4. Rate hunger before and every 5 min during meal (1-10 scale). Stop at 7-8.
5. Start every meal with a glass of water and 2-minute pause.

DECISION TREE

IF user always feels hungry immediately after eating → **THEN** speed eating likely; reduce meal pace; check serving adequacy
IF user eats while working at desk → **THEN** this is distracted eating; remove screen during eating; even 10 min is better

FAQ

Q: Is it really true that eating slowly helps weight loss?

A: Yes. Studies show slow eaters consume 100-200 kcal less per meal. Over a year, this equals 10-20 kg difference without any other change.

AI COACHING NOTES

- Mindful eating is often more powerful than calorie counting for habitual overeaters.
- Phone at meals is the most common barrier — address it directly.

WL-BH-010	MOD 09 · Consistency Over Perfection › The 80/20 Rule
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Adult · Teen · Senior · Student · Office Worker
DIFF	Beginner
TAGS	consistency, 80/20, perfectionism, all or nothing, flexibility, sustainable diet

DEFINITION

80/20 principle in weight management: 80% adherence to plan + 20% flexibility = long-term success. Perfectionistic all-or-nothing thinking is the primary cause of diet abandonment.

WHY IT MATTERS

- Pursuing 100% compliance leads to guilt on any deviation → abandonment.
- 80% adherence over 1 year produces better results than 100% for 3 weeks.
- Flexibility prevents binge-restrict cycles that undermine net progress.

KEY FACTS

- 4-5 on-plan days per week is sufficient for meaningful fat loss.

- One off-plan meal does not derail a week. One off-plan week barely derails a month.
- People who allow flexibility lose 10-15% more weight than strict dieters over 12+ months.

ACTION STEPS

1. Reframe from 'did I fail today' to 'how is my weekly average?'
2. Pre-plan one flexible meal per week — not a cheat day, a planned enjoyment.
3. After an off-plan event: don't compensate by restricting next day — just resume normally.
4. Track adherence as a percentage weekly, not as pass/fail daily.

DECISION TREE

IF user ate 'badly' and wants to fast to compensate → **THEN** NEVER recommend compensation fasting; 'just resume normal meals tomorrow'

IF user has all-or-nothing thinking → **THEN** CBT-style reframe: 'Does one tile make a mosaic? No. Does one meal make a diet? No.'

FAQ

Q: I ate terribly this weekend — should I do a detox?

A: No. Detoxes don't work. Your liver does that automatically. Just return to your normal plan Monday. The weekend damage is usually 2-4 kg water weight and 3,000-5,000 extra kcal — significant but recoverable with 1-2 weeks of normal eating.

AI COACHING NOTES

- 80/20 framing reduces abandonment by 50% in coaching practice.
- Normalize deviation before it happens — 'You will have off days. This is what to do when that happens...'

WL-BH-011	MOD 09 · Time Management for Weight Loss · Busy Schedule Solutions
GOALS	Weight Loss Fat Loss Weight Maintenance General Health
USERS	Office Worker · Student · Hostel Student · Adult
DIFF	Intermediate
TAGS	time management, busy lifestyle, meal prep, quick meals, no time to exercise, time-efficient

DEFINITION

Lack of time is the most commonly cited barrier to healthy eating and exercise. Time management strategies can create 30-60 min of health-focused time in even the busiest schedules.

WHY IT MATTERS

- 'No time' is usually 'it's not a priority yet' — once health impacts work capacity, priorities shift.
- Most healthy habits take <30 min/day — the return on investment is enormous.

KEY FACTS

- Average Indian adult watches 2-4 hours of TV/reels daily — health behaviours can replace some of this.
- Meal prep Sunday: 2 hours → saves 45 min daily → net time gain.
- 20 min workout is sufficient for health benefits — longer is better but not mandatory.

ACTION STEPS

1. Sunday meal prep: cook grains, proteins, chop vegetables — assemble meals in <5 min daily.
2. Phone workout timer: 20 min bodyweight circuit during lunch break.
3. Walk during calls — convert 30 min phone calls to walking time.
4. Batch cook: make 4x dal, freeze portions. Make large salads for 3 days.
5. 5-min meals: protein shake + fruit, curd + oats, PB + roti, boiled eggs.

DECISION TREE

IF user says 'I have no time to cook' → **THEN** ask: do you have 2 hours Sunday? That's all you need (meal prep). During week: 5-min no-cook options.

IF user says 'I have no time to exercise' → **THEN** 20 min bodyweight workout at home. Walk 15 min at lunch. This requires no commute time.

FAQ

Q: How can I eat healthy if I'm too busy to cook?

A: Meal prep strategy: Sunday 2 hours → ready meals all week. Or: 5-minute no-cook staples: overnight oats, Greek yogurt + fruit, boiled eggs (batch cook 6 at once), bread + peanut butter + banana, sprouts + curd.

AI COACHING NOTES

- *Never accept 'I have no time' at face value — help them find it.*
 - *Audit their schedule: where are screen-time hours? Can 30 min be redirected?*
-

Plateaus and Troubleshooting

WL-PL-000	MOD 10 · Plateau Overview › Why Plateaus Happen
GOALS	Weight Loss Fat Loss
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian
DIFF	Intermediate
TAGS	plateau, stalled weight loss, no weight loss, why not losing, adaptive thermogenesis

DEFINITION

A weight-loss plateau is a period of ≥ 2 -3 weeks with no meaningful scale change (< 0.5 kg) despite consistent adherence. It is a normal physiological response to fat loss — not a failure. The body adapts by reducing TDEE through metabolic adaptation, reduced NEAT, and hormonal changes.

WHY IT MATTERS

- Plateaus are not caused by 'eating too little' — they are caused by the body's adaptation to the deficit.
- As weight decreases, TDEE decreases — the same deficit no longer exists.
- NEAT unconsciously reduces during dieting — fidgeting, posture, and movement decrease.

KEY FACTS

- Metabolic adaptation: after 12+ weeks of dieting, TDEE can be 300-500 kcal below what formula predicts.
- Plateau is expected at: 4-6 weeks, 10-12 weeks, and after every 5 kg of loss.
- Common plateau causes: tracking drift, TDEE decrease, hormonal adaptation, water retention masking fat loss.
- Most 'plateaus' are caused by calorie tracking errors, not metabolic adaptation.

ACTION STEPS

1. Step 1: Verify tracking accuracy for 7 days (weigh and measure everything).
2. Step 2: Confirm scale trend (7-day average, not single day reading).
3. Step 3: Check protein (≥ 1.6 g/kg), step count (≥ 7000 /day).
4. Step 4: If genuine plateau after verification: apply break protocol.

DECISION TREE

IF no weight loss for 2 weeks → **THEN** verify tracking accuracy first; do not immediately change plan

IF no weight loss for 3-4 weeks with accurate tracking → **THEN** apply plateau protocol: recalculate TDEE, increase steps 2000/day, refeed or diet break

IF losing cm but not kg on scale → **THEN** body recomposition; not a plateau; maintain plan

IF weight going up then down with no net progress → **THEN** water retention masking fat loss; check weekly average over 3-4 weeks

IF weight stable but body looks different → **THEN** recomposition; celebrate this; don't panic

COMMON MISTAKES

- Drastic calorie cuts at plateau — worsens metabolic adaptation.
- Adding excessive cardio — increases cortisol and muscle breakdown.
- Giving up after 2-week plateau — normal and temporary.

FAQ

Q: I haven't lost weight in 3 weeks. Is my metabolism broken?

A: Unlikely. In order: 1) Check tracking accuracy (oil, sauces, drinks). 2) Check step count. 3) Recalculate TDEE at new weight. 4) If all correct — try a 1-week refeed at maintenance, then resume.

AI COACHING NOTES

- Reassure before changing anything — most 'plateaus' resolve with tracking accuracy fix.
- Never cut calories aggressively at plateau — this is the most common error.

WL-PL-001	MOD 10 · Diet Break Protocol › Refeed and Full Diet Break
GOALS	Weight Loss Fat Loss
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian
DIFF	Intermediate

DEFINITION

A diet break is a planned 1-2 week period at maintenance calories, used to restore metabolic rate, hormonal function, and psychological adherence before resuming a caloric deficit.

WHY IT MATTERS

- Leptin drops 50% within 1 week of aggressive dieting — reducing satiety and metabolic rate.
- A 2-week maintenance break restores leptin and other hormones within 7-14 days.
- Research shows: dieters who took 2-week breaks every 2 weeks lost more fat than continuous dieters.

KEY FACTS

- Diet break frequency: every 8-12 weeks of continuous deficit.
- Maintenance = TDEE (recalculated at current weight).
- Scale will increase 0.5-1.5 kg during break (glycogen + water) — this is NOT fat gain.
- After diet break: resume deficit and fat loss often accelerates.

ACTION STEPS

1. Calculate current TDEE at current body weight.
2. Eat at TDEE for 1-2 weeks (not above).
3. Maintain all exercise and activity.
4. Accept the scale increase — it's glycogen and water, not fat.
5. Resume deficit after break; fat loss typically resumes faster.

DECISION TREE

IF user has been in deficit for 10+ weeks with slowing results → **THEN** recommend 2-week diet break at maintenance

IF user is mentally exhausted from dieting → **THEN** diet break is both physical and psychological reset

IF user is scared of eating maintenance → **THEN** explain glycogen water gain; show expected 0.5-1.5 kg which reverses when deficit resumes

WL-PL-002	MOD 10 · Calorie Tracking Drift · Fixing Hidden Calories
GOALS	Weight Loss Fat Loss
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian
DIFF	Beginner
TAGS	calorie tracking, food diary, hidden calories, tracking errors, oil tracking, portion control

DEFINITION

Calorie tracking drift is the gradual underestimation of calories over time. Research shows most self-reported food diaries undercount by 20-40%. Drift is the most common cause of apparent plateaus.

WHY IT MATTERS

- Most plateaus are caused by tracking drift, not metabolic adaptation.
- Oil, sauces, tasting while cooking, and restaurant food are the most underestimated sources.
- Re-calibrating tracking is the highest-ROI plateau intervention.

KEY FACTS

- Average underreporting: 20% of actual calories.
- Worst offenders: cooking oil (pour instead of measure), restaurant food (2x actual vs home cooking), sauces, ghee, nuts.
- 1 tbsp extra oil per day = 120 extra kcal = 0.5 kg fat per month.

ACTION STEPS

1. 7-day accurate tracking: weigh all food on scale; measure oil with spoon; count nuts.
2. Log restaurant meals at 1.5-2x the stated calories.
3. Photograph meals and review — often reveals portion errors.
4. Track liquid calories first: chai, lassi, juice, coconut water.
5. Use kitchen scale for 2 weeks — recalibrates portion awareness.

DECISION TREE

IF user hit plateau, good tracking discipline → **THEN** check oil quantity, restaurant frequency, and post-workout extra eating

IF user tracks meals but plateau >4 weeks → **THEN** 7-day weighed food diary; usually reveals 300-500 kcal undercount

COMMON MISTAKES

- Estimating portions by eye after months of tracking — errors accumulate.
- Not logging cooking oil because 'it's just a teaspoon' — measure it.
- Forgetting snacks, tastes, and drinks in the diary.

WL-PL-003	MOD 10 · Weight Regain · Prevention and Recovery
GOALS	Weight Maintenance Weight Loss
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian
DIFF	Intermediate
TAGS	weight regain, yo-yo, rebound, after diet, maintaining weight loss, long term maintenance

DEFINITION

Weight regain is the return of lost weight after a diet ends. 80% of dieters regain within 2 years. Primary causes: reverting to old habits, insufficient muscle mass, no maintenance plan, and biological drive to restore pre-diet weight.

WHY IT MATTERS

- Without active maintenance habits, the default environment and habits recreate original weight.
- Muscle loss during dieting reduces metabolic rate — making regain easier on same calories.
- The body has a 'weight set point' that actively resists change — long-term maintenance requires permanent habit change.

KEY FACTS

- Only 20% of dieters maintain >10% weight loss for 1+ year.
- Successful long-term maintainers share: daily weigh-in, high protein, regular exercise, consistent meal patterns.
- Weight cycling (yo-yo): each cycle makes fat loss harder and regain easier.
- Muscle mass is the most powerful defense against regain — prioritize resistance training.

ACTION STEPS

1. Set a maintenance range: your goal weight ± 2 kg.
2. Weigh daily — intervene when trending toward upper boundary.
3. Maintain resistance training — non-negotiable for metabolic rate defense.
4. Keep tracking habits even in maintenance (weekly if not daily).
5. Identify regain triggers: specific social situations, seasonal patterns, stress.

DECISION TREE

IF user regained 3-5 kg → **THEN** don't panic; return to tracking + 300 kcal deficit; most is water weight

IF user regained most of lost weight → **THEN** restart from Module 01; identify what changed; address root cause

IF user is afraid of maintenance (prefers deficit) → **THEN** validate concern; teach maintenance as a skill; 100-200 kcal below calculated maintenance is sustainable long-term

COMMON MISTAKES

- Stopping all weight tracking after reaching goal.
- Abandoning exercise after reaching goal weight.
- Returning to pre-diet eating patterns exactly.

FAQ

Q: How do I keep weight off after losing it?

A: The 5 maintenance principles: 1) Daily weigh-in (catch regain early). 2) Maintain protein intake. 3) Keep resistance training. 4) Stay somewhat active (8000+ steps). 5) Allow ± 2 kg range — intervene if trending toward top.

WL-PL-004	MOD 10 · Low Motivation Plateau · When Motivation Disappears
GOALS	Weight Loss Fat Loss
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian
DIFF	Beginner
TAGS	motivation, plateau motivation, giving up, burnout, diet fatigue, re-motivation

DEFINITION

Motivational plateau is a period where adherence drops, not due to physical barriers, but due to psychological fatigue, slow progress, or changed priorities. Most common at weeks 4-8 and after major life events.

WHY IT MATTERS

- Motivation peaks at start when novelty is high — sustaining requires deeper systems.
- Results slow naturally after initial rapid loss — this creates a motivation dip.

ACTION STEPS

1. Reconnect with WHY: why did you start? Write it down; re-read.
2. Progress audit: measure waist, take progress photo, list non-scale victories.
3. Change something: new recipe, new workout, new walking route — novelty re-engages.
4. Set a near-term micro-goal (2 weeks max) — smaller targets are more motivating.
5. Accountability: share goals with someone; weekly check-in.
6. Rest if needed: 1-week maintenance break is better than giving up.

DECISION TREE

- IF user says 'I've stopped caring' → THEN identify: life stressor? Slow results? Burnout? Address root cause before plan change
- IF results have genuinely slowed → THEN celebrate total progress; zoom out; 2 kg/month over a year = 24 kg transformation
- IF user comparing to faster results of others → THEN normalize individual variation; refocus on personal progress only

WL-PL-005	MOD 10 · Travel and Diet › Maintaining Progress During Travel
GOALS	Weight Loss Fat Loss
USERS	Adult · Office Worker · Frequent Traveler
DIFF	Intermediate
TAGS	travel, diet travel, business travel, airport food, hotel food, maintain diet traveling

DEFINITION

Travel disrupts routines, food access, and exercise habits. Without a strategy, travel derails weight-loss progress. With a plan, travel can be navigated with minimal impact.

ACTION STEPS

1. Pack: protein powder sachets, roasted chana, nuts, protein bars.
2. Hotel breakfast: eggs, oats, fruits, curd — avoid pastries, fried items, buffet overload.
3. Room workout: 20-min bodyweight circuit requires no equipment.
4. Restaurant strategy: grilled protein + salad as default order.
5. Stay hydrated — flights dehydrate severely, mimicking hunger.
6. Accept 1-2 kg scale increase from water and sodium — not fat.

DECISION TREE

- IF 1-2 day business trip → THEN maintain plan strictly; carry emergency snacks
- IF 5+ day extended travel → THEN damage control mode: prioritize protein, limit obvious junk, move daily
- IF international travel → THEN jet lag disrupts hunger signals; prioritize sleep and hydration over perfect eating

COMMON MISTAKES

- Abandoning all healthy habits 'because I'm traveling'.
- Airport fast food habit — arrive early to find healthier options.
- Overcompensating on return with excessive restriction.

WL-PL-006	MOD 10 · Festival and Wedding Navigation › High-Calorie Social Events
GOALS	Weight Loss Fat Loss
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian
DIFF	Beginner
TAGS	wedding diet, festival eating, Diwali diet, Eid diet, party food, social eating strategy

DEFINITION

Indian festivals (Diwali, Eid, Holi, Navratri, Onam) and weddings involve high-calorie, high-fat, high-sugar foods in social settings with strong cultural pressure to eat. Strategy is required to navigate without derailing progress or missing cultural joy.

KEY FACTS

- Average wedding meal: 1,000-1,500 kcal (1-2x daily budget in one meal).

- Diwali period: average 2-5 kg gain from sweets + fried snacks.
- 1 week of festival eating does not ruin months of progress — but uncontrolled festival season (3-4 weeks) can.

ACTION STEPS

1. Pre-event: eat protein-rich meal before arriving — don't arrive starving.
2. Survey all food before taking anything — make conscious choices.
3. Protein first: dal, chicken, paneer — eat these before rice/bread.
4. Sweets: 1-2 pieces only. Use the 3-bite rule.
5. Skip or minimize fried appetizers.
6. Alcohol: max 1-2 drinks, dry wine or spirits over cocktails/beer.
7. Day after: normal eating — no restriction, no punishment.

DECISION TREE

IF wedding season (1-2 months) → **THEN** set maintenance target, not fat-loss; track loosely; avoid full abandonment

IF major festival (Diwali) → **THEN** make 1-2 days of celebration, not 2-week celebration

IF user overate at event → **THEN** reassure; 'back to normal tomorrow'; one event doesn't matter

COMMON MISTAKES

- Skipping meals all day before a wedding to 'save calories' — arrives famished, overeats.
- All-or-nothing: 'I'll restart after festival season' — often leads to months of abandonment.
- Guilt-driven over-restriction post-event.

FAQ

Q: How do I avoid gaining during Diwali?

A: Pre-event strategy: 1) Don't receive sweets at home (give away immediately). 2) Make or gift healthier sweets (dry fruits, dark chocolate). 3) One plate rule at parties. 4) Stay active — extra 20 min walk daily during festival week.

WL-PL-007	MOD 10 · Eating Out Strategy › Restaurant and Takeout Guide
GOALS	Weight Loss Fat Loss
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian
DIFF	Beginner
TAGS	eating out, restaurant diet, takeout, ordering healthy, menu strategy

DEFINITION

Restaurant and takeout food averages 60-100% more calories than home-cooked equivalent due to more oil, butter, cream, sugar, and larger portions. An eating-out strategy allows enjoyment without derailment.

KEY FACTS

- Restaurant food uses 2-4x more oil than home cooking for same dish.
- Restaurant portions are typically 1.5-2x appropriate serving size.
- Menu label calories are typically accurate $\pm 30\%$ — always estimate 25% higher.

ACTION STEPS

1. Order first at the table — before hearing others' orders.
2. Request: 'less oil', 'sauce on the side', 'no butter on naan'.
3. Best choices: tandoori dishes, dal, grilled fish/chicken, idli-sambar, raita.
4. Worst choices: creamy gravies, fried appetizers, biryani with heavy oil, desserts.
5. Portion control: take half in takeaway box immediately for most restaurant main courses.
6. Eat salad/raita first — reduces main course intake by 15%.

DECISION TREE

IF user eats restaurant food 3+ times/week → **THEN** restaurant eating is likely the primary calorie source; strategy is critical

IF user doesn't know what to order → **THEN** default: any grilled protein + dal/salad + 1 roti or small rice

IF user feels social pressure to order more → **THEN** 'sharing starters' as cover; order main only; skip dessert — most groups don't notice

AI COACHING NOTES

- Create a personal 'restaurant go-to list' for their 5 most visited restaurants.
- Teach the 'good enough' order — doesn't need to be perfect, just better.

WL-PL-008	MOD 10 · Poor Adherence Troubleshooting › When Nothing Is Working
GOALS	Weight Loss Fat Loss
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian
DIFF	Intermediate
TAGS	adherence, consistency, not following plan, giving up, why diet fails, re-starting

DEFINITION

Poor adherence — inability to consistently follow the plan — is the most common reason weight-loss efforts fail. Before changing the plan, identify why adherence is failing.

ACTION STEPS

1. Adherence audit: for 1 week, note exactly when and why you deviate from plan.
2. Categorize deviation: hunger? social? emotional? habit? convenience?
3. Fix the root cause: if hunger → protein up. If social → social strategy. If habit → environment design.
4. Simplify the plan: a simpler plan you follow beats a perfect plan you don't.
5. Reduce the target: is the calorie target too aggressive? Increase by 200 kcal and reassess.

DECISION TREE

IF user deviates due to hunger → **THEN** plan is too calorie-restricted; increase protein and fiber before cutting calories further
IF user deviates at social events → **THEN** social eating strategy module (WL-BH-008)
IF user deviates when stressed → **THEN** stress management module (WL-BH-004); emotional eating module (WL-BH-005)
IF user deviates out of boredom → **THEN** environment design module (WL-BH-007); HALT check protocol
IF user deviates due to plan complexity → **THEN** simplify: 3-5 meal templates only; no detailed counting

AI COACHING NOTES

- *Don't add more rules when adherence is poor — remove rules.*
- *Meet the user where they are, not where they should be.*
- *The best plan is the one they'll actually follow.*

WL-PL-009	MOD 10 · Rapid Weight Loss Then Regain › Breaking the Yo-Yo Pattern
GOALS	Weight Loss Fat Loss
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian
DIFF	Intermediate
TAGS	yo-yo dieting, rapid loss, quick fix, crash diet, rebound weight, sustainable loss

DEFINITION

Yo-yo dieting is repeated cycles of rapid weight loss followed by regain. Each cycle worsens metabolism (lower TDEE), reduces muscle mass, and psychologically reinforces failure identity.

WHY IT MATTERS

- Rapid loss uses extreme restriction → muscle loss → lower TDEE → faster regain.
- The rebound is biological — not a willpower failure.

ACTION STEPS

1. Stop the current 'quick fix' approach.
2. Calculate true TDEE and set a modest 300-500 kcal deficit.
3. Commit to minimum 6-month timeline for meaningful change.
4. Add resistance training to protect/rebuild muscle.
5. Target 0.5 kg/week — this feels slow but is life-changing over 1 year.
6. Address the psychology of needing fast results.

DECISION TREE

IF user has yo-yo dieted 3+ times → **THEN** reverse diet protocol: start at current maintenance, gradually decrease over 4-6 weeks
IF user wants fast results and resists slow approach → **THEN** show: 0.5 kg/week = 26 kg/year vs 5 kg quick fix repeated 4x = 20 kg with muscle loss and worsening metabolism

AI COACHING NOTES

- *Reframe slow loss as the powerful choice — not the weak one.*
- *Acknowledge previous yo-yo without blame — the plans were wrong, not the person.*

Progress Tracking Systems

WL-TR-000	MOD 11 · Tracking Overview › What to Track and Why
GOALS	Weight Loss Fat Loss Weight Maintenance
USERS	Adult · Teen · Student · Office Worker
DIFF	Beginner
TAGS	tracking, progress, weight tracking, measurements, body fat, habit tracker

DEFINITION

Weight management tracking is the systematic measurement of progress indicators to identify trends, catch issues early, and maintain accountability. Scale weight is ONE of many indicators — measurements, photos, strength, and habits provide the full picture.

WHY IT MATTERS

- Regular tracking correlates with 2-3x better weight-loss outcomes vs non-trackers.
- Scale weight fluctuates 1-3 kg daily from water and glycogen — trends matter, not daily readings.
- Body measurements and photos reveal fat loss that the scale misses (recomposition).

KEY FACTS

- Track: weight, waist, hips, strength, steps, habits, photos.
- Frequency: daily weight (use weekly average), weekly measurements, monthly photos.
- Best time: morning after toilet, before food/water, consistent days.

ACTION STEPS

1. Set up tracking tools: scale, measuring tape, camera, habit tracker (app or paper).
2. Day 1: record starting weight, waist, hip, bicep measurements + front/side photos.
3. Weekly: weigh daily and take 7-day average; measure waist/hips.
4. Monthly: progress photos + strength benchmarks.

WL-TR-001	MOD 11 · Scale Weight Tracking › Correct Use of the Weighing Scale
GOALS	Weight Loss Fat Loss Weight Maintenance
USERS	Adult · Teen · Student · Office Worker
DIFF	Beginner
TAGS	weight tracking, scale, weigh yourself, morning weight, 7 day average

DEFINITION

The weighing scale measures total body mass including fat, muscle, water, food, and glycogen. Daily fluctuations of 1-3 kg are normal and expected. The 7-day rolling average removes noise and reveals actual fat loss trend.

WHY IT MATTERS

- Daily weight fluctuates: water retention, sodium, hormones, gut content — not fat.
- Most diet frustration comes from reacting to daily fluctuations instead of weekly trends.
- 7-day average removes noise; 4-week trend reveals true rate of loss.

KEY FACTS

- Weigh at same time, same conditions: morning, after toilet, before food/water.
- Expect +0.5-2 kg after salty meal, high-carb day, or before period — all water.
- True fat loss rate: 0.25-1 kg/week. Slower is normal and often better.
- If 4-week trend is flat with accurate tracking: genuine plateau — apply WL-PL protocols.

ACTION STEPS

1. Weigh every morning: after toilet, before eating, same clothing.
2. Log daily weight in app (Happy Scale, MacroFactor) or spreadsheet.
3. Calculate 7-day average weekly — compare averages, not daily readings.
4. Review 4-week trend monthly. Expected loss: 1-4 kg/month depending on target.

DECISION TREE

IF daily weight up 2 kg from yesterday → **THEN** water retention from sodium/carbs/hormones; check weekly average, not today

IF scale not moved for 2 weeks (verified tracking) → **THEN** apply WL-PL-001 plateau protocol

IF scale moving but waist not reducing → **THEN** could be losing from elsewhere or early phase; check waist + photos monthly

COMMON MISTAKES

- Weighing multiple times a day — creates anxiety and meaningless data.
- Comparing Monday weight to Friday weight after weekend eating.
- Stopping weighing when progress slows — this disconnects accountability.

FAQ

Q: Should I weigh daily or weekly?

A: Daily weight + weekly average is most accurate. Weekly only can miss important patterns. Never use single-day readings as decisions.

WL-TR-002	MOD 11 · Body Measurements › Waist, Hip, and Body Tracking
GOALS	Weight Loss Fat Loss Weight Maintenance
USERS	Adult · Teen · Student · Office Worker
DIFF	Beginner
TAGS	body measurements, waist measurement, hip measurement, measuring tape, inches lost, cm lost

DEFINITION

Body measurements track changes in size that scale weight misses — especially during body recomposition (fat loss with muscle gain). Waist circumference is the single best predictor of metabolic health improvement.

WHY IT MATTERS

- Measurements change even when scale is static — most common during recomposition.
- Waist reduction = visceral fat reduction = most important health improvement.
- Indian cutoffs: waist >90 cm male, >80 cm female = elevated metabolic risk.

KEY FACTS

- Key measurements: waist (narrowest), hips (widest), mid-thigh, upper arm.
- Measure weekly at same time (morning, fasted).
- Expected rate of change: 0.5-1.5 cm waist reduction per month with 500 kcal deficit.
- Waist-to-height ratio: waist should be less than half your height (e.g., 170 cm tall → <85 cm waist).

ACTION STEPS

1. Using measuring tape: measure at same body landmarks, same tension (snug but not tight).
2. Record: waist, hips, mid-thigh, upper arm. Date all entries.
3. Take measurements weekly on the same day.
4. Compare month-to-month rather than week-to-week for clean trend.

AI COACHING NOTES

- *When scale stalls: check measurements — often shows progress the scale misses.*
- *Waist reduction is the most meaningful health metric — celebrate it explicitly.*

WL-TR-003	MOD 11 · Progress Photos › Visual Progress Documentation
GOALS	Weight Loss Fat Loss Weight Maintenance
USERS	Adult · Teen · Student · Office Worker
DIFF	Beginner
TAGS	progress photos, before after, body transformation, visual progress, photo tracking

DEFINITION

Progress photos are monthly photographs taken under standardized conditions to document visual body composition changes that measurements and scale weight may not capture.

WHY IT MATTERS

- Body appearance changes before scale changes are dramatic — photos capture this.
- Most people underestimate their own progress; photos provide objective evidence.

- Motivation tool: looking back at 3-month journey is profoundly encouraging.

ACTION STEPS

1. Photo protocol: same time (morning, fasted), same lighting, same clothing (swimwear/sports), same camera distance, front + side + back.
2. Take monthly — not more frequently.
3. Compare month 1 vs month 3 side-by-side — subtle monthly changes become dramatic quarterly.
4. Store privately for personal use.

FAQ

Q: I don't see progress in the mirror even though I've lost 5 kg. Why?

A: Body dysmorphia and familiarity with your own appearance make progress invisible in real-time. Photos from 3 months ago are the most powerful comparison. Also: 5 kg of fat is the size of 5 bags of sugar — substantial, even if hard to see on yourself.

WL-TR-004	MOD 11 · Body Fat Estimation › Methods and Accuracy
GOALS	Weight Loss Fat Loss Weight Maintenance
USERS	Adult · Teen · Student · Office Worker
DIFF	Intermediate
TAGS	body fat percentage, body fat measurement, DEXA, skinfold, BIA, Navy formula

DEFINITION

Body fat percentage is the proportion of total weight that is adipose tissue. It is more meaningful than scale weight for assessing health and body composition. Methods vary in accuracy and accessibility.

KEY FACTS

- Methods (most to least accurate): DEXA scan ($\pm 1\text{-}2\%$), Hydrostatic weighing ($\pm 2\%$), Bod Pod ($\pm 2\%$), US Navy formula ($\pm 3\text{-}4\%$), Skinfold calipers ($\pm 3\text{-}5\%$), BIA (bioelectrical impedance) scale ($\pm 5\text{-}8\%$).
- DEXA scan: ■ 1,500-3,000 in major Indian cities. Most accurate accessible method.
- US Navy Formula (Males): $86.010 \times \log(\text{waist} - \text{neck}) - 70.041 \times \log(\text{height}) + 36.76$
- US Navy Formula (Females): $163.205 \times \log(\text{waist} + \text{hip} - \text{neck}) - 97.684 \times \log(\text{height}) - 78.387$
- BIA scales (Omron, Tanita): affected by hydration — consistent conditions essential.

ACTION STEPS

1. Baseline: DEXA scan or Navy formula at start.
2. Repeat every 3 months.
3. Use same method each time — cross-comparison between methods is invalid.
4. Target healthy ranges: male 10-20%, female 18-28%.

AI COACHING NOTES

- Navy formula is free, reasonably accurate, and repeatable — recommend over BIA scales.
- Direction of change matters more than absolute number.

WL-TR-005	MOD 11 · Strength and Fitness Tracking › Performance Benchmarks
GOALS	Weight Loss Fat Loss Weight Maintenance
USERS	Adult · Teen · Student · Office Worker
DIFF	Beginner
TAGS	strength tracking, fitness benchmarks, push-up test, exercise progress, performance

DEFINITION

Strength and fitness benchmarks track exercise capacity improvements, which indicate muscle preservation during fat loss and general health improvement.

WHY IT MATTERS

- Strength preservation during fat loss = muscle being maintained.
- Strength gains on a deficit indicate body recomposition — optimal outcome.
- Fitness benchmarks provide tangible, non-scale progress measures.

KEY FACTS

- Basic benchmarks for adults: Push-ups in 1 min (males ≥ 30 , females ≥ 15 = good). Wall sit (≥ 60 sec = good). 1.5 km run time (males < 7 min, females < 9 min = good).
- Strength improvements often precede visible body changes by 4-6 weeks.

ACTION STEPS

1. Monthly benchmark: push-up max, plank hold, 1-km walk/run time.
2. Track gym lifts in a notebook: weight $\times$ reps for each main exercise.
3. Progressive overload marker: if you can't add weight or reps, recovery or nutrition is limiting.

WL-TR-006	MOD 11 · Step and NEAT Tracking › Daily Activity Monitoring
GOALS	Weight Loss Fat Loss Weight Maintenance
USERS	Adult · Teen · Student · Office Worker
DIFF	Beginner
TAGS	step count, NEAT, activity tracking, 10000 steps, fitness tracker, daily movement

DEFINITION

Step tracking measures daily NEAT — the most variable and underappreciated component of total calorie burn. Most adults need to dramatically increase daily steps for meaningful NEAT impact.

KEY FACTS

- Minimum: 7,000 steps/day for health. 10,000+ for weight loss.
- Every 1,000 additional steps/day burns ~40-50 additional kcal.
- Difference between 5,000 and 12,000 steps/day: ~280-350 kcal/day difference.
- Affordable step trackers: Mi Band (■2,000), Realme Band (■1,500), or free phone pedometer.

ACTION STEPS

1. Set a daily step goal: 7,000 (beginner), 8,500 (intermediate), 10,000 (active).
2. Track daily and review weekly average.
3. If consistently below goal: add 1 x 15-20 min walk to routine.
4. Increase goal by 1,000 steps/day every 2 weeks.

WL-TR-007	MOD 11 · Habit Tracking › Daily Behaviour Monitoring
GOALS	Weight Loss Fat Loss Weight Maintenance
USERS	Adult · Teen · Student · Office Worker
DIFF	Beginner
TAGS	habit tracker, daily habits, consistency tracker, don't break the chain, adherence

DEFINITION

Habit tracking is the daily recording of key behaviours (tracking food, hitting protein, exercising, sleeping 7+ hours) to build consistency through visual accountability.

WHY IT MATTERS

- Visual streak (don't break the chain) is a powerful motivation tool.
- Habit tracking reveals patterns: 'I consistently fail on weekends' → targeted intervention.
- Tracking habits creates intrinsic accountability without external enforcement.

ACTION STEPS

1. Choose 3-5 key habits: protein target, step goal, food tracked, exercise, sleep 7+ hrs.
2. Use a paper grid or app (Streaks, HabitMinder, Notion).
3. Review weekly: which habit is consistently missed? Focus there next week.
4. Celebrate 7-day streaks — small rewards build momentum.

AI COACHING NOTES

- *Start with 2-3 habits maximum — not 10.*
- *Missing 1 day: acceptable. Missing 2 consecutive days: trigger intervention.*

WL-TR-008	MOD 11 · Weekly Review › End-of-Week Self-Assessment
GOALS	Weight Loss Fat Loss Weight Maintenance

USERS	Adult · Teen · Student · Office Worker
DIFF	Beginner
TAGS	weekly review, check-in, self assessment, weekly progress, accountability, reflection

DEFINITION

A weekly review is a 10-minute structured self-assessment at week's end to evaluate progress, identify friction points, and plan the coming week.

KEY FACTS

- Weekly reviewers are 40% more likely to maintain healthy behaviours vs non-reviewers.
- The review creates course corrections before problems compound.

ACTION STEPS

1. Every Sunday (10 min): Record: 7-day average weight, waist (monthly), steps average, adherence %.
2. Review: What went well? What was hard? What do I change next week?
3. Set: 1 food goal, 1 movement goal, 1 behaviour goal for next week.
4. Celebrate: 1 non-food reward for any progress.

AI COACHING NOTES

- Guide users through weekly review in each check-in session.
- Focus on what went well first — builds positive momentum.

WL-TR-009	MOD 11 · Progress Interpretation · Reading Your Data Correctly
GOALS	Weight Loss Fat Loss Weight Maintenance
USERS	Adult · Teen · Student · Office Worker
DIFF	Intermediate
TAGS	interpreting progress, weight trend, no progress, good progress, reading data, non scale victory

DEFINITION

Correct interpretation of tracking data prevents unnecessary plan changes and false alarms. Most perceived 'problems' are normal fluctuations or expected adaptation — proper interpretation prevents emotional decision-making.

KEY FACTS

- Normal daily weight fluctuation: 0.5-3 kg.
- Expected 4-week loss: 1-4 kg (depending on deficit size).
- Good progress indicators: waist down, strength maintained, energy good, clothes fitting better.
- Non-scale victories: better sleep, more energy, improved mood, better mobility, lower resting heart rate.

ACTION STEPS

1. Monthly interpretation matrix: scale + measurements + photos + strength + energy = full picture.
2. Scale stable but measurements down: recomposition — celebrate this.
3. Scale stable, measurements stable, 4-week verified: true plateau — apply WL-PL protocols.
4. Scale up, measurements down: muscle gain with fat loss — optimal outcome.

DECISION TREE

- IF** scale up + waist down → **THEN** recomposition; excellent outcome; maintain plan
- IF** scale stable + waist down → **THEN** fat loss exactly offset by muscle gain; very good; maintain
- IF** scale down + waist same → **THEN** losing weight but may be muscle; check protein, add resistance training
- IF** scale stable + waist stable + accurate tracking for 4 weeks → **THEN** genuine plateau; WL-PL-001 diet break protocol
- IF** scale up + waist up + poor tracking → **THEN** tracking error plus food excess; fix tracking first

FAQ

Q: I've lost 3 kg but look the same — what's happening?

A: 3 kg is actually significant. Check: same time of day for mirror comparison? Progress photos from week 1 vs now are more objective than mirror. Also: fat distribution change takes time to become visually obvious.

Q: My waist went down 3 cm but scale is same. Should I be worried?

A: Absolutely not — this is the best possible outcome. You're losing fat and gaining muscle simultaneously (recomposition). Don't change anything.

AI COACHING NOTES

- *Teach users to look at all data points before reacting.*
 - *Non-scale victories are as important as scale victories — document and celebrate them.*
-

AI Coach Decision Engine

WL-DE-000	MOD 12 · Decision Engine Overview › How to Use This Module
GOALS	Weight Loss Fat Loss General Health
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes
DIFF	Intermediate
TAGS	AI coach, decision engine, coaching rules, if then, coaching logic, AI weight loss

DEFINITION

The AI Coach Decision Engine contains 100+ IF→THEN coaching rules organized by category. Each rule maps a user condition to the correct coaching response. This module enables consistent, evidence-based coaching across all user types and scenarios.

WHY IT MATTERS

- Coaching rules prevent inconsistent advice across different users.
- Rules encode clinical expertise for AI retrieval.
- Structured decision logic produces better outcomes than free-form advice.

KEY FACTS

- Rule categories: Intake (15), BMI (8), Exercise (12), Plateau (9), Hormonal/Medical (14), Emotional/Psychological (10), Lifestyle (9), Supplements (9).
- Total rules in this module: 86+
- Rules are designed to be retrieved independently by RAG system.

ACTION STEPS

1. Match user's presenting situation to closest rule.
2. Apply the THEN action consistently.
3. Layer multiple rules when user has multiple conditions.
4. Escalate to medical professional for any medical condition.

WL-DE-001	MOD 12 — Decision Engine · Intake and Nutrition Rules
GOALS	Weight Loss Fat Loss General Health
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes
DIFF	Intermediate
TAGS	intake, nutrition, calories, protein, tracking

DEFINITION

Coaching rules for: Intake and Nutrition Rules. 15 IF→THEN rules.

WHY IT MATTERS

- Consistent rule application ensures evidence-based coaching.
- Rules are designed for RAG retrieval on user-specific conditions.

KEY FACTS

- IF: User calories below 1200 (female) or 1500 (male) → THEN: STOP. Increase to minimum floor. Risk: muscle loss, deficiency, rebound.
- IF: User protein below 1.0g/kg → THEN: Increase protein before any other macro adjustment. Add 20-30g/day.
- IF: User eating 1 meal/day → THEN: Redistribute to 3-4 meals. Single-meal pattern causes muscle loss and overeating.
- IF: User tracking calories with app but plateau >3 weeks → THEN: 7-day verified food diary. Most 'plateaus' are hidden calories.
- IF: User consuming 3+ sweet chais/day → THEN: Replace 2 with black/herbal tea. Saves 200-300 kcal/day instantly.
- IF: User eating lots of cooking oil (4+ tbsp/day) → THEN: Reduce to 2 tbsp/day. Saves 240-480 kcal/day.
- IF: User relying on restaurant/takeout >4x/week → THEN: Introduce Sunday meal prep. First change — highest impact.

- IF: User skips breakfast → THEN: Either introduce protein breakfast OR validate IF approach — not both.
- IF: User reports constant hunger on plan → THEN: Increase protein +20g, increase fiber +10g, increase water intake. Do not cut calories further.
- IF: User eating ultra-processed food daily → THEN: Identify 3 UPFs to replace with whole food equivalents. One per week.
- IF: User has adequate calories but low protein (<80g) → THEN: Protein redistribution: hit protein target first, fill rest with carbs/fat.
- IF: User drinks sugary beverages (cold drinks, juice, lassi) → THEN: Liquid calories first. Cut these before food changes.
- IF: User has high fat intake from nuts/seeds (>60g/day) → THEN: Measure portions. 30g nuts = 170 kcal. Portion control, not elimination.
- IF: User eating very low fat (<30g/day) → THEN: Increase to 40-50g. Required for fat-soluble vitamins and hormonal health.
- IF: User on very low carb diet and fatigued → THEN: Increase carbs to 100-150g/day minimum. Brain and thyroid need carbs.

WL-DE-002	MOD 12 — Decision Engine · BMI-Based Protocols
GOALS	Weight Loss Fat Loss General Health
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes
DIFF	Intermediate
TAGS	BMI, obesity, weight classification, BMI protocol

DEFINITION

Coaching rules for: BMI-Based Protocols. 8 IF→THEN rules.

WHY IT MATTERS

- Consistent rule application ensures evidence-based coaching.
- Rules are designed for RAG retrieval on user-specific conditions.

KEY FACTS

- IF: BMI < 18.5 (Underweight) → THEN: DO NOT prescribe deficit. Refer to nutritionist for healthy weight gain.
- IF: BMI 18.5-22.9 (Normal, Indian standard) → THEN: Maintenance or mild deficit (200-300 kcal). Focus on body composition.
- IF: BMI 23-24.9 (Overweight, Indian standard) → THEN: 300-400 kcal deficit. Diet + exercise. No medical intervention needed.
- IF: BMI 25-29.9 (Obese Class I) → THEN: 400-500 kcal deficit. High protein, resistance training. Monthly doctor check-in.
- IF: BMI 30-34.9 (Obese Class II) → THEN: 500-600 kcal deficit. Medical evaluation for comorbidities. Professional supervision.
- IF: BMI 35-39.9 (Obese Class III) → THEN: Medical supervision required. Structured program. Consider bariatric evaluation.
- IF: BMI ≥40 (Severe Obesity) → THEN: Medical and psychological team required. Bariatric surgery evaluation. Not self-managed.
- IF: Normal BMI but high waist circumference → THEN: TOFI pattern. Focus on visceral fat: low GI, cardio, stress management.

WL-DE-003	MOD 12 — Decision Engine · Exercise Rules
GOALS	Weight Loss Fat Loss General Health
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes
DIFF	Intermediate
TAGS	exercise, gym, workout, cardio, resistance training

DEFINITION

Coaching rules for: Exercise Rules. 12 IF→THEN rules.

WHY IT MATTERS

- Consistent rule application ensures evidence-based coaching.
- Rules are designed for RAG retrieval on user-specific conditions.

KEY FACTS

- IF: User does zero exercise → THEN: Start with walking only — 20-30 min/day. Do not add gym yet.
- IF: User exercises but no resistance training → THEN: Add 2x/week resistance training. Most important for long-term metabolism.
- IF: User exercises 6-7x/week and plateau → THEN: Overtraining possible. Add 1-2 rest days. Recovery is when results happen.
- IF: User hates exercise → THEN: Focus on activity (NEAT) vs exercise. Walking, household chores, dancing count.
- IF: User exercises but eats back all calories → THEN: Recalculate TDEE with active multiplier; remove post-workout eating.
- IF: User has joint pain → THEN: Move to low-impact: swimming, cycling, yoga, resistance bands. Protect joints.
- IF: User is gym beginner → THEN: Start with 3 days/week. Full body compound movements only for 8 weeks.
- IF: User is intermediate gym-goer → THEN: Move to push/pull/legs split or upper/lower. Add progressive overload.
- IF: User only does cardio → THEN: Add resistance training. Cardio burns during workout; resistance training burns after.
- IF: User does only HIIT → THEN: Add 1-2 steady state cardio + resistance training. HIIT-only increases cortisol excessively.
- IF: Student with no gym access → THEN: 20 min bodyweight circuit program (WL-WP-004). No equipment needed.
- IF: Office worker with no time → THEN: Lunch walk 20 min + 20 min evening workout. Total: 40 min fragmented — valid.

WL-DE-004	MOD 12 — Decision Engine · Plateau Rules
GOALS	Weight Loss Fat Loss General Health
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes
DIFF	Intermediate
TAGS	plateau, no weight loss, stall, metabolic adaptation

DEFINITION

Coaching rules for: Plateau Rules. 9 IF→THEN rules.

WHY IT MATTERS

- Consistent rule application ensures evidence-based coaching.
- Rules are designed for RAG retrieval on user-specific conditions.

KEY FACTS

- IF: No weight loss for 2 weeks with verified tracking → THEN: Recalculate TDEE at current weight. Increase steps by 2000/day.
- IF: No weight loss for 4 weeks with verified tracking + increased steps → THEN: 2-week diet break at maintenance. Then resume deficit.
- IF: Weight fluctuating 1-2 kg but no net loss (4 weeks) → THEN: Water retention masking loss. Check waist measurements. Use weekly average.
- IF: Waist reducing but scale stable → THEN: Recomposition. Maintain plan. This is an excellent outcome.
- IF: All metrics stalled (6+ weeks) → THEN: Comprehensive review: tracking accuracy, stress, sleep, hormones, medications.
- IF: Lost 5+ kg then plateau → THEN: Expected. Body adapted. Increase NEAT by 2000 steps, minor diet tweak — not big cuts.
- IF: Weight increasing despite deficit → THEN: Re-verify tracking (7-day weighed diary). Check for medication changes, thyroid, water retention.
- IF: User reports plateau but hasn't verified tracking → THEN: Always verify tracking FIRST before any plan changes.
- IF: User on aggressive deficit (>750 kcal) plateauing → THEN: Metabolic adaptation. Reduce deficit to 500 kcal. Add diet break.

WL-DE-005	MOD 12 — Decision Engine · Hormonal and Medical Rules
GOALS	Weight Loss Fat Loss General Health

USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes
DIFF	Intermediate
TAGS	PCOS, diabetes, thyroid, medical, hormones

DEFINITION

Coaching rules for: Hormonal and Medical Rules. 14 IF→THEN rules.

WHY IT MATTERS

- Consistent rule application ensures evidence-based coaching.
- Rules are designed for RAG retrieval on user-specific conditions.

KEY FACTS

- IF: PCOS confirmed → THEN: Apply WL-SP-001 protocol. Low GI, high protein, strength training, inositol supplement.
- IF: Hypothyroidism confirmed (treated) → THEN: Reduce TDEE estimate by 200 kcal vs standard. Extra resistance training. Check ferritin.
- IF: Hypothyroidism uncontrolled (high TSH) → THEN: Refer to endocrinologist. Weight loss is very difficult with uncontrolled hypothyroid.
- IF: Type 2 Diabetes (diet-controlled) → THEN: Apply WL-SP-002 protocol. Walk after every meal. Low GI diet strictly.
- IF: Type 2 Diabetes on medication → THEN: Coordinate with treating doctor. Weight loss may reduce medication needs — dangerous without doctor involvement.
- IF: Prediabetes → THEN: Apply WL-SP-002 protocol. Up to 100% reversal possible with 7-10% body weight loss.
- IF: Postpartum (<8 weeks) → THEN: Do NOT prescribe deficit. Rest, healing, adequate nutrition. No formal exercise.
- IF: Postpartum (8-24 weeks, breastfeeding) → THEN: Minimum 1800 kcal. Very gentle activity only. No aggressive deficit.
- IF: Senior (60+) → THEN: Apply WL-SP-005. Higher protein. Low-impact exercise. Never below 1600 kcal.
- IF: Teen (13-18) → THEN: Never below 1600 kcal. Activity-first approach. No restriction. Nutrient density focus.
- IF: Suspected eating disorder → THEN: STOP coaching. Refer to eating disorder specialist immediately.
- IF: User on steroids/antidepressants/antipsychotics → THEN: These medications cause weight gain. Coordinate with prescribing doctor. Expect harder journey.
- IF: User reports fatigue + cold + hair loss + constipation → THEN: Thyroid screening before proceeding. These are classic hypothyroid symptoms.
- IF: User reports extreme fatigue on diet → THEN: Rule out: anaemia (iron, B12), hypothyroid, vitamin D deficiency. Test first, adjust after.

WL-DE-006	MOD 12 — Decision Engine · Emotional and Psychological Rules
GOALS	Weight Loss Fat Loss General Health
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes
DIFF	Intermediate
TAGS	emotional eating, binge, mindset, psychology

DEFINITION

Coaching rules for: Emotional and Psychological Rules. 10 IF→THEN rules.

WHY IT MATTERS

- Consistent rule application ensures evidence-based coaching.
- Rules are designed for RAG retrieval on user-specific conditions.

KEY FACTS

- IF: User reports emotional eating >2x/week → THEN: WL-BH-005 protocol. Consider referral to therapist if severe.
- IF: User shows binge-restrict pattern → THEN: Break cycle: increase daily intake to reduce binge drive. No compensation rules.
- IF: User is highly self-critical about food → THEN: Compassionate reframe. Shame worsens eating behaviour. Never reinforce guilt.
- IF: User shows signs of orthorexia (fear of 'bad' foods) → THEN: Refer to dietitian specializing in eating disorders. Broaden food acceptance.

- IF: User expresses suicidal ideation or self-harm → THEN: STOP all coaching. Immediately provide crisis resources. This is beyond coaching scope.
- IF: User sets unrealistic timeline (5 kg/month) → THEN: Educate: 2-3 kg/month is excellent. Unrealistic targets cause all-or-nothing failure.
- IF: User compares to Instagram/celebrity transformations → THEN: Reality check: sponsored transformations, Photoshop, and drugs. Focus on personal journey.
- IF: User says 'I'm not losing fast enough' → THEN: Show: 0.5 kg/week = 26 kg/year. Slow is fast compounded.
- IF: User wants to go below 1000 kcal → THEN: Refuse. Educate on muscle loss and metabolic damage. Not an option.
- IF: User is obsessively tracking calories → THEN: Assess for disordered eating. Calorie tracking can cause harm if anxiety-driven.

WL-DE-007	MOD 12 — Decision Engine · Lifestyle Rules
GOALS	Weight Loss Fat Loss General Health
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes
DIFF	Intermediate
TAGS	hostel, night shift, travel, stress, sleep, lifestyle

DEFINITION

Coaching rules for: Lifestyle Rules. 9 IF→THEN rules.

WHY IT MATTERS

- Consistent rule application ensures evidence-based coaching.
- Rules are designed for RAG retrieval on user-specific conditions.

KEY FACTS

- IF: Hostel student with mess food → THEN: Apply WL-SP-008. Carry emergency snacks. Skip mess fried items.
- IF: Night shift worker → THEN: Apply WL-SP-007. Eat main meal before shift. Light eating during night hours.
- IF: Frequent traveler → THEN: Apply WL-PL-005. Carry protein sachets. Damage control mode when traveling.
- IF: User reports high stress + no sleep → THEN: Address sleep and stress BEFORE tightening diet. These are primary blockers.
- IF: User has <6 hours sleep consistently → THEN: Sleep optimization is the intervention. No point tightening diet if sleep is disrupted.
- IF: User reports no time to cook → THEN: Sunday meal prep strategy. 5-minute emergency foods list. Not an excuse — a logistics problem to solve.
- IF: User eating at desk while working → THEN: Mindful eating module. Remove screens from mealtimes. 10-min lunch break minimum.
- IF: User drinks alcohol frequently (4+ times/week) → THEN: Alcohol is 7 kcal/g plus lowers inhibitions for eating. Reduce first before diet changes.
- IF: User's social circle pressures unhealthy eating → THEN: Social eating strategy WL-BH-008. Polite decline scripts. Eat before events.

WL-DE-008	MOD 12 — Decision Engine · Supplement Rules
GOALS	Weight Loss Fat Loss General Health
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes
DIFF	Intermediate
TAGS	supplements, protein powder, fat burner, creatine, vitamins

DEFINITION

Coaching rules for: Supplement Rules. 9 IF→THEN rules.

WHY IT MATTERS

- Consistent rule application ensures evidence-based coaching.
- Rules are designed for RAG retrieval on user-specific conditions.

KEY FACTS

- IF: User buying 'fat burner' products → THEN: Educate: no fat burner works meaningfully. Save money. Use for real food or protein powder.
 - IF: User asking about detox tea → THEN: Educate: laxative effect only. No fat loss. Potential electrolyte imbalance. Do not use.
 - IF: Vegetarian user with low protein (<80g) → THEN: Recommend: whey protein supplement as primary fix. Paneer and dal as secondary.
 - IF: Vegan user with low protein → THEN: Recommend: pea + rice protein blend. Soya chunks. Edamame.
 - IF: All vegetarian/vegan users → THEN: Auto-recommend: B12 supplement (non-negotiable). Vitamin D (universal). Omega-3 (algae oil).
 - IF: User doing resistance training → THEN: Consider: creatine 3-5g/day. Whey protein if under protein target. Safe and effective.
 - IF: User has poor sleep → THEN: Consider: magnesium glycinate 200-300mg. Not a drug. Improves sleep depth.
 - IF: User has strong sugar cravings → THEN: Check magnesium and chromium. Both improve insulin sensitivity and reduce cravings.
 - IF: User asks about protein powder brand → THEN: Reputable Indian brands: MuscleBlaze, Nakpro, MyProtein. International: Optimum Nutrition, Dymatize.
-

Master FAQ Database

WL-FAQ-000	MOD 13 — Master FAQ · Master FAQ Database Overview
GOALS	Weight Loss Fat Loss General Health Weight Maintenance
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes
DIFF	Beginner
TAGS	FAQ, questions, weight loss FAQ, common questions, coaching Q&A;

DEFINITION

Comprehensive FAQ database with 261+ Q&As; across 24 categories.

WHY IT MATTERS

- 80% of weight-loss confusion is answered by the same 50 questions.
- Pre-answered FAQs enable fast, consistent AI coaching responses.

KEY FACTS

- Total FAQs in database: 261+
- Categories: Calories, Protein, Carbs, Fat, Fiber, Rice/Roti, Fruits, Dairy, Eggs, Chicken, Supplements, Protein Powder, Creatine, Walking, Gym, PCOS, Diabetes, Sleep, Stress, Cravings, Hostel, Office, Plateaus, Maintenance

ACTION STEPS

1. Search this database by category when users ask common questions.
2. Use direct answers verbatim or paraphrase for conversational tone.
3. Cross-reference with relevant Module for deeper protocol.

WL-FAQ-001	MOD 13 — Master FAQ · FAQs: Calories
GOALS	Weight Loss Fat Loss General Health Weight Maintenance
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes
DIFF	Beginner
TAGS	calories, calorie counting, calorie deficit, TDEE, calorie target

DEFINITION

Comprehensive FAQ answers for: Calories. 30 questions answered.

WHY IT MATTERS

- Category Calories is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Calories FAQs: 30.

FAQ

Q: How many calories should I eat to lose weight?

A: Eat at TDEE minus 300-500 kcal. Calculate TDEE first (Module 01). Typical targets: sedentary Indian female 1,400-1,600 kcal; male 1,700-2,000 kcal. Never guess.

Q: What is a calorie?

A: A unit of energy. Food provides calories; your body burns calories for all functions. 1 kg fat ≈ 7,700 kcal.

Q: How accurate are calorie counts on food labels?

A: ±20% accuracy is typical. Restaurant food is often 25-50% more than stated. Always estimate slightly higher for restaurant meals.

Q: Does cooking method change calories?

A: Yes significantly. Deep frying adds 8-12g fat per 100g food (72-108 extra kcal). Air frying, steaming, boiling add minimal calories.

Q: Can I eat any food if I'm within my calorie budget?

A: Technically yes for weight loss. Practically no — nutrition quality, satiety, and health matter. 1,500 kcal of junk food and 1,500 kcal of whole foods produce different hunger, energy, and health outcomes.

Q: Why am I gaining weight eating only 1,200 kcal?

A: Most likely: tracking errors (oil, sauces, drinks), hormonal issues (hypothyroid), water retention, or inaccurate calorie data for Indian foods. Verified 7-day food diary with kitchen scale first.

Q: Is it bad to eat below 1,000 kcal?

A: Yes. Below 1,200 kcal (female) and 1,500 kcal (male) risks: muscle loss, nutrient deficiency, metabolic adaptation, hair loss, fatigue, and rebound weight gain.

Q: Do calories matter more than food quality?

A: For weight loss: calories matter more. For health: quality matters more. Both matter — the goal is quality foods within a calorie budget.

Q: Should I count net carbs or total carbs?

A: Total carbs for general use. Net carbs (total minus fiber) for keto-style diets. For most Indians eating normal food, total carbs tracking is sufficient.

Q: How do I count calories in Indian food without a database?

A: Use HealthifyMe (best Indian food database), Cronometer, or MyFitnessPal India database. When in doubt: weigh the main components (oil, dal, rice) separately and sum.

Q: Are there negative-calorie foods?

A: No. Some foods (celery, cucumber) have very few calories but claim more energy to digest — the net is never negative. They are 'near zero' calorie, not negative.

Q: How do I count calories in homemade dishes?

A: Log each ingredient separately with quantities before cooking. Use the raw ingredient weights. The cooking process doesn't reduce total calories significantly.

Q: Does eating less slow my metabolism?

A: If below 1,200 kcal (female) / 1,500 kcal (male) for weeks: yes, metabolic adaptation occurs. Moderate deficit (300-500 kcal) with adequate protein: minimal adaptation.

Q: How many meals should I eat per day?

A: 3-4 meals is optimal for most people for hunger control and muscle protein synthesis. Meal frequency does not directly affect metabolism.

Q: What is calorie cycling?

A: Eating more calories on training days (higher TDEE) and fewer on rest days. Net weekly deficit is what matters. Useful for some; not necessary for most.

Q: Should I eat more when I exercise?

A: Depends on how you calculated TDEE. If you used sedentary multiplier: add 50-70% of exercise calories. If you used an active multiplier: no additional eating needed.

Q: Can I drink alcohol and still lose weight?

A: Yes if it fits calories. Alcohol is 7 kcal/g. 1 large beer (500ml) = 200-250 kcal. It also lowers inhibitions leading to more eating. Limit to 1-2 drinks max.

Q: What are empty calories?

A: Calories with little to no nutritional value: sugary drinks, refined alcohol, candy, most fried snacks. They fill your budget without contributing protein, fiber, vitamins, or minerals.

Q: How do I know if my calorie target is right?

A: After 3-4 weeks of accurate tracking: if losing 0.3-1 kg/week, target is right. If losing too fast, increase by 200. If not losing, decrease by 150-200 or increase movement.

Q: Is intermittent fasting just about calorie restriction?

A: Yes. IF works by creating a shorter eating window, which naturally reduces total intake for most people. It does not have metabolic magic beyond the calorie deficit it creates.

Q: What is the best way to reduce calorie intake?

A: Priority order: (1) Cut liquid calories (sweet chai, cold drinks, juice), (2) Reduce cooking oil by 1-2 tbsp, (3) Reduce rice/roti by 25%, (4) Replace high-calorie snacks with low-calorie ones.

Q: Can I lose weight without counting calories?

A: Yes. Portion control, eliminating junk food, eating mostly whole foods, and mindful eating can create a deficit without counting. Counting improves precision if progress stalls.

Q: Does meal timing affect calorie burning?

A: Minimally. Total daily intake matters most. Slight advantage to eating more earlier in the day vs late at night, but the difference is small vs overall deficit.

Q: What is TDEE?

A: Total Daily Energy Expenditure — all calories your body burns in a day. BMR + NEAT + exercise + TEF. The benchmark your calorie intake is measured against.

Q: How often should I recalculate my calorie target?

A: Every 5 kg of weight change (your TDEE drops as you lose weight) and after major lifestyle changes (new job, injury, activity change).

Q: Are all calories the same?

A: Thermodynamically yes. Metabolically no — 100 kcal of protein vs 100 kcal of sugar produce different hormonal, satiety, and metabolic responses. Source matters even within a calorie budget.

Q: What happens if I eat too little for too long?

A: Metabolic adaptation (TDEE drops by 10-25%), muscle loss, nutrient deficiencies, hair loss, fatigue, hormonal disruption, and increased likelihood of binge eating and yo-yo cycling.

Q: Can stress cause me to burn fewer calories?

A: Yes. Chronic stress increases cortisol, which slightly reduces metabolic efficiency and promotes fat storage. Also causes NEAT reduction (sitting more, less fidgeting). Significant indirect effect.

Q: How much does exercise increase calorie burn?

A: Walking 30 min: 150-200 kcal. Gym session: 250-400 kcal. HIIT 30 min: 300-450 kcal. Less than most people think — diet remains primary lever.

Q: What is TEF and does it matter?

A: Thermic Effect of Food — calories burned digesting food. Protein: 20-30%, carbs: 5-10%, fat: 0-3%. High protein diet effectively burns 50-100 more kcal daily from TEF. Useful but not transformative.

WL-FAQ-002 MOD 13 — Master FAQ · FAQs: Protein

GOALS Weight Loss | Fat Loss | General Health | Weight Maintenance

USERS Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes

DIFF Beginner

TAGS protein, protein intake, vegetarian protein, egg protein, paneer protein

DEFINITION

Comprehensive FAQ answers for: Protein. 30 questions answered.

WHY IT MATTERS

- Category Protein is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Protein FAQs: 30.

FAQ

Q: How much protein do I need per day?

A: For weight loss: 1.6-2.0g per kg body weight. Example: 65 kg person → 104-130g protein daily. Higher end if doing resistance training.

Q: What are the best vegetarian protein sources in India?

A: Paneer 18g/100g, Greek yogurt/hung curd 10-15g/100g, soya chunks 52g/100g dry, moong dal 9g/100g cooked, rajma 15g/100g cooked, tofu 8g/100g, sprouts 4g/100g.

Q: Can I get enough protein without eating meat?

A: Yes, but requires planning. A vegetarian eating eggs, paneer, curd, dal, soya, and legumes can reach 80-100g protein daily. Add protein powder if falling short.

Q: Why is protein important for weight loss?

A: Three reasons: (1) highest satiety per calorie, (2) preserves muscle during fat loss, (3) highest thermic effect (burns 20-30% of its calories during digestion).

Q: Is protein powder safe?

A: Yes. Whey protein is concentrated milk protein. No hormones, no steroids. Safe for healthy adults. Choose FSSAI-certified brands. Not needed if getting adequate protein from food.

Q: How much protein is in one egg?

A: 6-6.5g per whole egg. Egg white: 3.5-4g per white. Yolk: 2.5g protein plus fat, vitamins, choline.

Q: How much protein is in paneer?

A: Regular paneer: 18g/100g. Low-fat paneer: 20-22g/100g. Greek yogurt equivalent protein at lower fat.

Q: Can I eat too much protein?

A: For healthy individuals: no harm at 2-3g/kg body weight. Above this provides no additional benefit. Only concern: people with pre-existing kidney disease (need medical guidance).

Q: Is dal a good protein source?

A: Yes but in portions. 1 cup cooked dal = 9-16g protein depending on type (masoor/moong vs rajma). Good source but need 3-4 cups daily to hit protein targets from dal alone.

Q: What is a complete protein?

A: Contains all 9 essential amino acids. Complete sources: eggs, meat, fish, dairy, quinoa, soy. Plant sources combined: rice + dal, roti + dal, corn + legumes.

Q: Do I need to eat protein right after workout?

A: Within 2 hours is ideal. The '30-minute window' is outdated. If you ate protein 2 hours before training, you're covered. Don't stress about exact timing.

Q: Does eating protein at night cause fat gain?

A: No. Calories, not timing, determine fat gain/loss. A pre-sleep protein-rich meal (curd, milk, casein) actually supports overnight muscle protein synthesis.

Q: What is casein protein?

A: Slow-digesting protein from milk. Ideal before sleep. Found naturally in paneer, curd, and milk. Provides a slow amino acid release over 5-7 hours while sleeping.

Q: How do I add more protein to Indian breakfast?

A: Options: eggs (boiled, scrambled, omelette), moong dal chilla, besan chilla, paneer bhurji, high-protein oats with protein powder, Greek yogurt parfait, sprouts bowl.

Q: Is tofu a good substitute for paneer?

A: Yes. Tofu: 8g protein/100g at 76 kcal. Paneer: 18g protein/100g at 260 kcal. Tofu is lower calorie, similar or higher calcium, excellent for weight loss and vegan diets.

Q: Can soya chunks replace meat?

A: Yes for protein. Dry soya chunks: 52g protein/100g (highest plant protein). Prepare with proper marination as texture differs from meat. Excellent budget protein source.

Q: What is the best protein powder in India?

A: MuscleBlaze Raw Whey (budget, reliable), MyProtein Impact Whey (quality), Optimum Nutrition Gold Standard Whey (premium). For vegans: MuscleBlaze Biozyme Plant Protein.

Q: How do I hit 100g protein on a vegetarian diet?

A: Example day: Greek yogurt 200g (20g) + eggs x2 (13g) + 100g paneer (18g) + 1.5 cups dal (20g) + protein shake 1 scoop (24g) = 95g. Replace eggs with extra soya/sprouts for egg-free.

Q: Does dal+rice give complete protein?

A: Yes. Dal provides lysine (limiting in grains); rice provides methionine (limiting in legumes). Together they form a complementary complete protein. Traditional wisdom was nutritionally correct.

Q: Is chicken the best protein source?

A: Among the best: 31g protein/100g cooked breast at 165 kcal. Zero carbs. Very high protein:calorie ratio. Excellent for fat loss. Comparable to: fish, egg whites, tuna.

Q: What happens if I don't eat enough protein while dieting?

A: Muscle loss (up to 40-60% of weight lost comes from muscle). Lower metabolic rate. Increased hunger. Fatigue and weakness. 'Skinny fat' appearance rather than toned.

Q: When should I drink a protein shake?

A: Whenever you'd otherwise under-eat protein. Good timings: breakfast (quick preparation), post-workout (within 2 hours), as an afternoon snack. No 'wrong' time.

Q: Does cooking destroy protein?

A: No. Heat denatures protein structure (unfolds it) but does not destroy its amino acid content. Cooked chicken, boiled eggs, and heated dal provide same protein as raw.

Q: Is sprouts a good protein source?

A: Moderate. Moong sprouts: 4g protein/100g. Chickpea sprouts: 9g/100g. Better than most vegetables but not as dense as paneer or dal. Excellent for living enzymes and bioavailability.

Q: How much protein in 1 cup milk?

A: Whole milk 240ml: 8g protein. Skimmed milk 240ml: 8.4g protein. Good source — 2-3 glasses daily contributes 16-24g protein.

Q: Should I count protein from vegetables?

A: Minimal. Spinach 2.9g/100g, broccoli 2.8g/100g — these are too low to significantly impact totals. Focus on high-density sources: eggs, dairy, legumes, meat.

Q: What is whey protein isolate vs concentrate?

A: Concentrate: 70-80% protein, small amount of lactose. Isolate: 90%+ protein, virtually lactose-free, slightly higher cost. Both effective. Isolate better for lactose-intolerant.

Q: Can diabetics eat high protein diet?

A: Yes — protein is actually beneficial for blood sugar control. Avoid very high protein if CKD (kidney disease). Otherwise: 1.2-1.6g/kg is safe and beneficial for T2D management.

Q: Is egg protein better than plant protein?

A: Egg has PDCAAS score of 1.0 (perfect) and excellent bioavailability. Plant proteins vary but well-planned combinations achieve similar overall protein quality over the course of a day.

Q: Does hunger mean I need more protein?

A: Often yes. Protein is the most satiating macro. If consistently hungry: check if protein target is being met. Add 20-30g protein and reassess hunger within 3-7 days.

WL-FAQ-003 MOD 13 — Master FAQ · FAQs: Carbohydrates

GOALS Weight Loss | Fat Loss | General Health | Weight Maintenance

USERS Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes

DIFF Beginner

TAGS carbs, rice, roti, oats, carbohydrates, low carb, GI

DEFINITION

Comprehensive FAQ answers for: Carbohydrates. 20 questions answered.

WHY IT MATTERS

- Category Carbohydrates is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Carbohydrates FAQs: 20.

FAQ

Q: Is rice bad for weight loss?

A: No. 1 cup cooked rice (200g) = 260 kcal. The issue is portion size and accompaniments (heavy gravies, ghee). Rice in a calorie-controlled plan with protein and vegetables is fine.

Q: Should I cut carbs to lose weight?

A: Not necessarily. Reduce quantity and improve quality. Cutting carbs works because it usually creates a calorie deficit and reduces insulin spikes — not because of carb magic.

Q: Is brown rice better than white for weight loss?

A: Slightly. Brown rice has more fiber (3.5g vs 0.6g per cup) and lower GI (68 vs 72). Better nutrition profile. If choosing, yes. But white rice in controlled portions is not a barrier to weight loss.

Q: Can I eat roti while losing weight?

A: Yes. 1 whole wheat roti (no oil) = 85 kcal, 3g protein, 2.5g fiber. 2 rotis + dal + salad = excellent weight-loss meal. Problem is when accompanied by excess ghee or paired with rice.

Q: How many rotis should I eat per day?

A: Depends on calorie target. At 1400 kcal: 4-5 rotis (no ghee). At 1600 kcal: 5-6. Never >2 at one meal for best blood sugar response.

Q: What is the best time to eat carbs?

A: Earlier in the day (breakfast and lunch) when activity is higher. Minimize carbs at dinner when activity is lower and insulin sensitivity is slightly reduced.

Q: Are millets better than wheat?

A: Generally yes. Jowar, bajra, ragi have lower GI, more fiber, and more micronutrients than wheat. Excellent for PCOS and diabetes. Taste is different — takes getting used to.

Q: What are good low-carb Indian foods?

A: Paneer, tofu, eggs, chicken, fish, leafy vegetables, broccoli, mushrooms, cucumber, cauliflower, bell peppers. Most non-starchy vegetables are low carb.

Q: Can I eat fruit on a weight loss diet?

A: Yes. Fruit provides fiber, vitamins, and natural sugar. 2-3 fruits/day fits in most plans. Prefer: guava, apple, pear, papaya, berries. Limit: mango, litchi, grapes, banana (higher sugar).

Q: Is oats good for weight loss?

A: One of the best grains for weight loss. High beta-glucan fiber (reduces appetite by 30-40%). Low GI (55). High satiety. Affordable. Easily adapted to Indian taste (masala oats, oats khichdi).

Q: What happens when I stop eating carbs?

A: First 1-3 days: glycogen depletion, water loss of 1-3 kg. Fatigue, 'keto flu' if very low carb. After adaptation (1-2 weeks): energy normalizes. Long-term very low carb: thyroid conversion may decrease.

Q: Is poha healthy?

A: Moderate. 1 cup cooked (200g) = 150-200 kcal, 3-4g protein, 33-40g carbs. Light, easy to digest. Healthy if made with 1 tsp oil and vegetables. Restaurant poha uses more oil.

Q: Should I eat carbs before workout?

A: 30-60 min before workout: 20-40g carbs improves performance. Banana, 1-2 dates, 1 roti. If exercising first thing in morning and prefer fasted: that's fine too — individual preference.

Q: Are root vegetables (potatoes, sweet potatoes) ok?

A: Boiled/baked potato: 77 kcal/100g, good fiber when skin kept, moderately high GI (80). Fine in moderation in any form except fried. Sweet potato: better GI, more nutrients.

Q: What is glycemic index and should I care?

A: GI measures how fast a carb raises blood sugar. Matters most for PCOS and diabetes. For general weight loss: total calories > GI. For metabolic health: lower GI is better.

Q: Is bread ok to eat on a diet?

A: Whole wheat bread: 75 kcal/slice, 6g protein, 13g carbs. Ok in moderation. 2 slices = 150 kcal. Better than white bread. Worse than oats or dal as a carb source.

Q: Can I eat pasta on a weight loss diet?

A: Yes if calories are managed. Whole wheat pasta is better. 1 cup cooked (140g) = 175 kcal, 7.5g protein, 37g carbs. Portion control is key.

Q: What are complex carbs vs simple carbs?

A: Complex: whole grains, legumes, vegetables — digest slowly, sustained energy, high fiber. Simple: white sugar, white bread, fruit juice — digest rapidly, blood sugar spike.

Q: Is jowar/sorghum better than wheat?

A: For weight loss and health: yes. Lower GI (55), more fiber, more micronutrients, gluten-free. Good for people with wheat sensitivity.

Q: Can I eat rice every day?

A: Yes, in controlled portions with protein and fiber. 1 cup cooked rice (not 3 cups) as part of a balanced meal is compatible with weight loss for most people.

WL-FAQ-004 MOD 13 — Master FAQ · FAQs: Dietary Fat

GOALS Weight Loss | Fat Loss | General Health | Weight Maintenance

USERS Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes

DIFF Beginner

TAGS fat, ghee, oil, cooking oil, omega 3, nuts, fatty food

DEFINITION

Comprehensive FAQ answers for: Dietary Fat. 10 questions answered.

WHY IT MATTERS

- Category Dietary Fat is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Dietary Fat FAQs: 10.

FAQ

Q: Is ghee good for weight loss?

A: Not helpful but not forbidden. 1 tsp ghee (5g) = 45 kcal, 5g fat. Benefits: fat-soluble vitamins, CLA, medium-chain fats. During weight loss: limit to 1 tsp/day. Not a 'superfood'.

Q: Which cooking oil is best for weight loss?

A: All oils are ~120 kcal/tbsp. Quantity matters more than type. For health: mustard (good omega-3/6 ratio), groundnut (MUFA), coconut (small amounts). Avoid vanaspati (trans fat).

Q: How much fat should I eat per day?

A: 20-35% of total calories. At 1600 kcal: 36-62g fat/day. Include essential fats (omega-3 and omega-6). Never go below 30g/day — required for hormonal health and fat-soluble vitamins.

Q: Do dietary fats make you fat?

A: Excess calories make you fat. Dietary fat (9 kcal/g) is dense and easy to overconsume. But fat itself is essential — cutting all fat is harmful. Moderate fat intake in calorie budget is fine.

Q: Is coconut oil good for weight loss?

A: No special fat-loss properties. Medium-chain triglycerides (MCTs) in coconut oil have a slight metabolic advantage, but it's still 121 kcal/tbsp. Same calories as other oils.

Q: Are nuts good for weight loss?

A: Yes — in measured portions. 30g almonds (15 nuts) = 174 kcal, 6g protein, 15g fat, 2g fiber. High satiety, good nutrition. Easy to overeat — always pre-measure.

Q: What is trans fat and why avoid it?

A: Artificially hydrogenated fat (vanaspati, many packaged biscuits/snacks). Raises LDL, lowers HDL. Increases cardiovascular risk. No safe amount. Found in dalda, commercial baked goods, cheap fried foods.

Q: Is avocado available in India and worth eating?

A: Available in metros (₹60-120 each). Nutritionally excellent: healthy MUFA, fiber, potassium. 200g avocado = 320 kcal — significant calorie cost. Good if fits budget and calorie plan.

Q: What are omega-3 foods in India?

A: Affordable: flaxseed (chia), mustard oil, walnuts, fatty fish (mackerel, sardines, hilsa). Supplement: algae oil (vegan) or fish oil 1-2g EPA+DHA/day.

Q: Can I eat peanut butter while losing weight?

A: Yes. 1 tbsp (16g) = 94 kcal, 4g protein, 8g fat. High satiety, good protein. Choose natural PB (only peanuts). Measure — it's easy to eat 4 tbsp without noticing.

WL-FAQ-005	MOD 13 — Master FAQ · FAQs: Fiber
GOALS	Weight Loss Fat Loss General Health Weight Maintenance
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes
DIFF	Beginner
TAGS	fiber, dietary fiber, constipation, gut health, psyllium, isabgol

DEFINITION

Comprehensive FAQ answers for: Fiber. 10 questions answered.

WHY IT MATTERS

- Category Fiber is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Fiber FAQs: 10.

FAQ

Q: How much fiber should I eat daily?

A: Women: 25g/day. Men: 38g/day. Most Indians eat only 10-15g/day. Focus on increasing to 25g minimum.

Q: What are high-fiber Indian foods?

A: Rajma 15g/100g cooked, chana 12.5g/100g, guava 8.9g/100g, oats 4g/100g cooked, pear 5.5g, whole wheat roti 2.5g each.

Q: Does fiber have calories?

A: Practically no. Fermentable fiber yields ~2 kcal/g vs 4 kcal/g for digestible carbs. Psyllium husk and non-starchy vegetables are near-zero calorie.

Q: What is psyllium husk and should I take it?

A: Isabgol (psyllium husk): 7g fiber per 2 tbsp. Affordable, safe, effective. Take with 300ml water before meals. Reduces hunger, improves blood sugar, reduces cholesterol.

Q: Does fiber help weight loss?

A: Yes significantly. High fiber foods: slow digestion (sustained fullness), stabilize blood sugar (reduce cravings), feed beneficial gut bacteria (improve metabolism), add volume without calories.

Q: Can too much fiber cause problems?

A: If increased too quickly: gas, bloating, discomfort. Increase by 5g/week. Always drink water with fiber (dehydrated fiber causes constipation, not relief).

Q: Is brown bread high in fiber?

A: Depends on brand. True whole wheat bread: 2-3g fiber/slice. Many 'brown breads' are colored white flour — check ingredient list. First ingredient should be 'whole wheat flour'.

Q: Do vegetables have a lot of fiber?

A: Yes. Most non-starchy vegetables: 2-4g fiber/100g. Spinach, broccoli, bhindi, cabbage, French beans are excellent. Eat 300-500g vegetables daily for 8-15g fiber from vegetables alone.

Q: What is the difference between soluble and insoluble fiber?

A: Soluble (oats, dal, psyllium): forms gel, slows absorption, reduces cholesterol, blood sugar. Insoluble (vegetables, whole grains): adds bulk, speeds gut transit, prevents constipation. Both needed.

Q: Does fiber reduce belly fat?

A: Indirectly yes. Soluble fiber (dal, oats) reduces visceral fat by improving insulin sensitivity. Studies: 10g more soluble fiber daily = 3.7% reduction in visceral fat over 5 years.

WL-FAQ-006 MOD 13 — Master FAQ · FAQs: Rice and Roti**GOALS** Weight Loss | Fat Loss | General Health | Weight Maintenance**USERS** Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes**DIFF** Beginner**TAGS** rice, roti, chapati, can I eat rice, bajra, jowar, millet**DEFINITION**

Comprehensive FAQ answers for: Rice and Roti. 12 questions answered.

WHY IT MATTERS

- Category Rice and Roti is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Rice and Roti FAQs: 12.

FAQ**Q: Can I eat rice and still lose weight?**

A: Yes. Control portion (1 cup cooked = 200g), combine with protein (dal, fish, chicken) and fiber (vegetables), eat earlier in day. Rice itself is not fattening — excess calories are.

Q: Rice vs roti: which is better for weight loss?

A: Similar calories per serving. Roti has more protein and fiber per 100g. Rice is easier to overeat in large portions. Both can be part of a weight-loss diet.

Q: How much rice can I eat while losing weight?

A: 1 cup cooked (200g, 260 kcal) as one serving. Maximum 1.5 cups per meal within your calorie budget. Pair always with protein and fiber.

Q: Is basmati rice better than regular white rice?

A: Yes. Lower GI (58 vs 72), longer grains, more aromatic. Better blood sugar response. Good upgrade for PCOS and diabetes.

Q: How many rotis per day for weight loss?

A: At 1400 kcal: 4-5 rotis (no ghee). At 1600 kcal: 5-6 rotis. Distribute across meals — not all at once.

Q: Is jowar roti better than wheat roti?

A: Yes for PCOS and diabetes. Jowar: lower GI (55 vs 62), more fiber, gluten-free. Nutritionally superior. Slightly denser texture.

Q: Can I eat both rice and roti in the same meal?

A: Not recommended for weight loss — doubles the carb load without doubling nutrition. Choose one per meal.

Q: Is cold rice better for weight loss?

A: Yes. Cooling rice after cooking increases resistant starch by 2-4x. Resistant starch behaves like fiber — feeds gut bacteria, lower calorie absorption, better blood sugar response.

Q: What is the glycemic index of common Indian roti types?

A: Whole wheat: 62, Maida: 70, Jowar: 55, Bajra: 54, Ragi: 68, Multigrain: 55-60.

Q: How to make rice meals healthier?

A: Add dal to the plate first. Use small bowl for rice. Load the rest with vegetables and protein. Walk 15 min after eating.

Q: Is millet roti better for weight loss?

A: Yes. Bajra (pearl millet): GI 54, high fiber, high protein (8g/cup). Jowar: GI 55, high fiber. Both superior to wheat for blood sugar management.

Q: Can diabetics eat rice?

A: Yes, in small portions (1/2 cup cooked) combined with protein and fiber. Choose basmati, eat it last (protein and vegetables first), walk after. Monitor blood sugar response individually.

WL-FAQ-007 MOD 13 — Master FAQ · FAQs: Fruits

GOALS	Weight Loss Fat Loss General Health Weight Maintenance
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes
DIFF	Beginner
TAGS	fruits, banana, mango, fruit sugar, diabetic fruit, best fruit

DEFINITION

Comprehensive FAQ answers for: Fruits. 12 questions answered.

WHY IT MATTERS

- Category Fruits is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Fruits FAQs: 12.

FAQ

Q: Is fruit sugar bad?

A: No. Whole fruit sugar comes with fiber that slows absorption. Fruit sugar in juice (no fiber) behaves like added sugar. Eat the whole fruit, not the juice.

Q: How many fruits can I eat per day on a diet?

A: 2-3 portions per day is appropriate for most people. Choose lower-sugar fruits: guava, apple, papaya, berries, pear. Limit very sweet fruits: mango, banana, litchi, grapes.

Q: Is banana fattening?

A: No. 1 medium banana = 105 kcal, 27g carbs, 3g fiber. Good for energy and potassium. Not fattening in itself — only in excess of calorie budget.

Q: Can diabetics eat fruit?

A: Yes — whole fruit, not juice. Choose lower-GI fruits: apple, pear, guava, papaya, berries, jamun. Limit mango, banana, watermelon (higher GI). 1 fruit per serving, not 3 at once.

Q: Is mango ok to eat while dieting?

A: Yes, in season and in moderation. 1 cup mango (150g) = 99 kcal. The issue: most people eat 2-3 cups plus 2 times daily. 1 cup/day fits in most plans. Don't avoid it — manage portions.

Q: Which fruits are best for weight loss?

A: Guava (high fiber, low calorie), papaya (digestive enzymes, low calorie), watermelon (low calorie, high satiety from water), pear (highest fiber), apple (fiber, satiety).

Q: Is fresh juice a healthy choice?

A: No for weight loss. Juicing removes fiber. 1 glass orange juice (250ml) = 115 kcal, 0g fiber. 2 whole oranges = 120 kcal, 6g fiber. Always eat the fruit whole.

Q: Does fruit make you fat?

A: Extremely unlikely if eating whole fruit. Fructolipogenesis (conversion of fruit sugar to fat) requires significant overconsumption — 8-10+ fruits daily minimum. Normal 2-3 fruits/day: no fat gain.

Q: Is dry fruit a good snack?

A: In small portions. 3 dates = 80 kcal, 21g sugar, 2g fiber. 10g raisins = 30 kcal, 8g sugar. Good nutrition but very calorie-dense because water is removed. 20-30g portion maximum.

Q: Can fruits replace vegetables in a diet?

A: No. Fruits and vegetables have different micronutrient profiles. Fruits are higher in sugar and lower in certain minerals. Need both — 2-3 fruits and 400-500g vegetables daily.

Q: Is coconut water good for weight loss?

A: Yes. 250ml = 46 kcal, good electrolytes, very low sugar. Excellent hydration drink. Better than packaged coconut water (often added sugar). Natural tender coconut water is ideal.

Q: Which fruit is highest in protein?

A: Guava (2.6g/100g), avocado (2g/100g), jackfruit (1.5g/100g). No fruit is a good protein source — protein needs come from other foods.

WL-FAQ-008 MOD 13 — Master FAQ · FAQs: Dairy**GOALS** Weight Loss | Fat Loss | General Health | Weight Maintenance**USERS** Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes**DIFF** Beginner**TAGS** dairy, milk, curd, paneer, ghee, yogurt, lassi, buttermilk**DEFINITION**

Comprehensive FAQ answers for: Dairy. 10 questions answered.

WHY IT MATTERS

- Category Dairy is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Dairy FAQs: 10.

FAQ**Q: Is milk good for weight loss?**

A: Yes in appropriate amounts. 1 cup toned milk (240ml) = 130 kcal, 7.5g protein. Good protein source. 1-2 cups/day is appropriate. Don't drink 4-5 cups while wondering why calories are high.

Q: Is curd better than milk for weight loss?

A: Curd has probiotics, similar protein, and is more satiating due to thickness. 200g low-fat curd = 90-110 kcal, 8-9g protein. Excellent daily food. Greek yogurt/hung curd even better.

Q: Does dairy cause weight gain?

A: Excess calories from any food cause weight gain. Dairy itself is not inherently fattening. Indian overweight is more commonly linked to cooking oil, fried food, and sweets than dairy.

Q: Is paneer good for weight loss?

A: Good protein source but calorie-dense (260 kcal/100g, 20g fat). Limit to 100g serving. Low-fat paneer (165 kcal/100g) is a better option for weight loss.

Q: How to make low-fat paneer at home?

A: Use low-fat milk (1.5-2% fat). Add lemon juice or vinegar at near-boiling temperature. Strain. The lower fat milk produces lower-fat paneer.

Q: Is Greek yogurt available in India?

A: Yes but expensive. Alternative: hung curd — strain regular curd through muslin 3-4 hours. Result: 10-15g protein/100g, similar to Greek yogurt at 1/5th the price.

Q: Does milk increase blood sugar?

A: Moderately. Milk has lactose (milk sugar) with GI ~30-40. 1 cup causes modest blood sugar rise. Toned milk better than full fat. Adding sugar to milk (turmeric milk, flavored milk) increases GI significantly.

Q: Is lassi healthy?

A: Plain lassi: moderate. 1 glass sweet lassi (300ml) = 200-250 kcal and significant sugar. Plain/salted lassi: much better. Switch to chaas (buttermilk) for similar probiotics at 50-60 kcal.

Q: Can lactose-intolerant people eat dairy?

A: Sometimes. Curd is usually tolerated even with lactose intolerance (bacteria consume most lactose). Hard cheeses also low lactose. Milk and soft cheese cause more symptoms. Experiment individually.

Q: Is butter or ghee better?

A: Different fats. Ghee: clarified butter, slightly higher smoke point, pure fat. Butter: 80% fat, 20% water, some milk solids. Neither is a weight-loss food. Ghee marginally better (no milk solids). Limit both.

WL-FAQ-009 MOD 13 — Master FAQ · FAQs: Eggs**GOALS** Weight Loss | Fat Loss | General Health | Weight Maintenance**USERS** Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes**DIFF** Beginner**TAGS** eggs, egg protein, how many eggs, egg yolk, boiled egg

DEFINITION

Comprehensive FAQ answers for: Eggs. 10 questions answered.

WHY IT MATTERS

- Category Eggs is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Eggs FAQs: 10.

FAQ

Q: How many eggs can I eat per day?

A: 3-4 whole eggs daily is safe for most healthy adults (AHA, recent research). Higher LDL sensitivity varies individually — those with familial hypercholesterolaemia should consult doctor. No blanket limit for healthy people.

Q: Is egg yolk unhealthy?

A: No. Yolk contains choline (brain health), vitamins D, B12, A, K, omega-3, and yes — cholesterol and saturated fat. Dietary cholesterol has minimal effect on blood cholesterol for most people.

Q: Are eggs good for weight loss?

A: Excellent. Whole egg (78 kcal) with 6g protein and full nutrient profile. High satiety — people who eat eggs at breakfast consume 400 fewer calories over the day vs toast breakfast.

Q: How should I cook eggs for weight loss?

A: Boiled or poached: lowest calorie. Scrambled with non-stick and no butter: good. Omelette with 1/4 tsp oil: good. Fried with oil: adds 40-80 kcal. Deep fried: avoid.

Q: Is egg white only better than whole egg?

A: Only if calorie-restricted and already meeting all other micronutrient needs. Yolk contains choline, vitamins, and healthy fat. 1-2 whole eggs + additional whites is optimal.

Q: Are eggs vegetarian in India?

A: Culturally complex. Most Indian vegetarians exclude eggs ('eggetarians' eat them). Biologically, unfertilized eggs are animal products. Dietary classification varies by individual.

Q: Can diabetics eat eggs?

A: Yes. Eggs have minimal effect on blood glucose. High protein helps satiety without raising blood sugar. 2-3 eggs daily is safe and beneficial for most T2D patients.

Q: Are brown eggs healthier than white eggs?

A: No. Shell color is determined by the hen's breed — no nutritional difference. Both have identical nutrition.

Q: How long can boiled eggs last in the fridge?

A: Unpeeled: 1 week. Peeled: 5 days in airtight container with water. Boil 6 eggs Sunday for the week.

Q: What is the protein content of an omelette?

A: 2-egg omelette: 12-13g protein, 150-180 kcal depending on added oil. Add egg whites to boost further.

WL-FAQ-010 MOD 13 — Master FAQ · FAQs: Chicken

GOALS Weight Loss | Fat Loss | General Health | Weight Maintenance

USERS Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes

DIFF Beginner

TAGS chicken, chicken breast, tandoori, non veg, chicken protein

DEFINITION

Comprehensive FAQ answers for: Chicken. 10 questions answered.

WHY IT MATTERS

- Category Chicken is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Chicken FAQs: 10.

FAQ

Q: How much protein is in chicken breast?

A: 100g cooked chicken breast: 31g protein, 165 kcal. One of the highest protein-to-calorie ratios of any food.

Q: Is chicken breast or thigh better for weight loss?

A: Breast: 165 kcal, 31g protein, 3.6g fat per 100g. Thigh: 209 kcal, 26g protein, 11g fat per 100g. Breast is leaner. Thigh has more flavour — both are good protein sources.

Q: How often can I eat chicken?

A: 3-5 times/week is common and healthy. Vary with eggs, fish, and legumes for nutritional diversity. No strict upper limit for healthy adults.

Q: Is tandoori chicken healthy?

A: One of the best restaurant choices. Marinated and grilled = minimal added fat. 100g tandoori chicken = 150 kcal, 24g protein. Avoid creamy gravies.

Q: How to cook chicken for weight loss?

A: Best methods: grilled, baked, air-fried, poached. Avoid: deep fried, butter-basted, creamy gravies. Marinate with yogurt and spices for flavour without calories.

Q: Is boiled chicken healthy?

A: Yes for weight loss. Removes most fat during boiling. Bland but functional. Shred for rolls, salads, rice bowls, sandwiches. Add spices after cooking.

Q: Can I eat chicken every day?

A: Yes if you enjoy it. Dietary variety is ideal long-term but daily chicken is nutritionally fine. Rotate with fish for omega-3 benefits 2x/week.

Q: Is chicken good for PCOS?

A: Yes. High protein, low carb, doesn't spike insulin. Pair with vegetables and legumes for balanced PCOS-friendly meal.

Q: How to include chicken in a hostel setting?

A: Order grilled chicken from delivery (avoid gravy). Canned tuna is an accessible alternative. Boiled eggs are the most practical hostel protein.

Q: Does chicken have good fat?

A: Some. Skinless chicken: mostly unsaturated fat. Skin adds significant saturated fat — remove skin before cooking or after. 100g chicken breast with skin: 195 kcal vs 165 kcal skinless.

WL-FAQ-011 MOD 13 — Master FAQ · FAQs: Supplements

GOALS Weight Loss | Fat Loss | General Health | Weight Maintenance

USERS Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes

DIFF Beginner

TAGS supplements, fat burner, green tea, ashwagandha, ACV

DEFINITION

Comprehensive FAQ answers for: Supplements. 10 questions answered.

WHY IT MATTERS

- Category Supplements is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Supplements FAQs: 10.

FAQ

Q: What supplements should I take for weight loss?

A: Priority: Vitamin D + B12 (everyone). Omega-3 (if no fatty fish 2x/week). Whey protein (if under protein target). No supplement burns fat.

Q: Do fat burners work?

A: No meaningful evidence. Most contain caffeine (get from coffee). Some have dangerous stimulants. Spend money on food and protein instead.

Q: Is green tea good for weight loss?

A: Marginally. Green tea increases metabolism by 3-5% = 40-80 kcal/day. Good antioxidants. Beneficial habit but not transformative for weight loss.

Q: Is apple cider vinegar good for weight loss?

A: Very modest effect. ACV (1-2 tbsp before meals) may reduce post-meal blood sugar by 5-10% and slightly reduce meal size. Not a weight-loss supplement — more a digestive aid.

Q: What is ashwagandha and does it help with weight loss?

A: Adaptogen that reduces cortisol by 27% (KSM-66 extract, 300-600mg/day). Indirectly helps weight loss by reducing stress-driven eating and improving sleep. Not a fat burner.

Q: Should I take a multivitamin?

A: Can fill gaps but doesn't replace varied diet. For most Indians: specific supplements (D, B12, iron) are more targeted and effective than generic multivitamin.

Q: Is collagen good for weight loss?

A: No direct fat-loss effect. Collagen is a protein source — can contribute to protein target. Some benefits for joint health and skin. Not a weight-loss supplement.

Q: What is inositol and is it good for PCOS?

A: Myo-inositol (2g) + D-chiro-inositol (50mg) combination reduces insulin resistance, androgen levels, and improves ovulation in PCOS. Good evidence. Available in India (VitaBalance).

Q: Does caffeine help weight loss?

A: Yes modestly. 100-200mg caffeine (1-2 coffees) increases metabolism 4-11% and improves exercise performance. Pre-workout caffeine is the most supported ergogenic aid. Not a substitute for diet.

Q: Is Triphala good for weight loss?

A: Traditional Ayurvedic formula. Some evidence for gut health and mild laxative effect. No direct fat-loss evidence. Not harmful but not a weight-loss supplement.

WL-FAQ-012 MOD 13 — Master FAQ · FAQs: Protein Powder

GOALS Weight Loss | Fat Loss | General Health | Weight Maintenance

USERS Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes

DIFF Beginner

TAGS protein powder, whey, plant protein, MuscleBlaze, protein shake

DEFINITION

Comprehensive FAQ answers for: Protein Powder. 10 questions answered.

WHY IT MATTERS

- Category Protein Powder is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Protein Powder FAQs: 10.

FAQ

Q: Is protein powder necessary for weight loss?

A: No. Food first. Protein powder is convenient when struggling to hit protein targets from food. Especially useful for vegetarians.

Q: Which protein powder is best for vegetarians in India?

A: Whey (from milk): MuscleBlaze Raw Whey, Nakpro. For vegans: MuscleBlaze Biozyme Plant Protein, Fast&Up; Plant Protein (pea + rice blend).

Q: Can protein powder cause kidney damage?

A: No, in healthy individuals. Evidence of harm exists only with pre-existing CKD. Extensive research at 2-3g/kg protein shows no kidney harm in healthy adults.

Q: How do I use protein powder in Indian cooking?

A: Add to: oats porridge, curd, lassi, smoothies, roti dough, besan chilla batter, kheer. Unflavored protein works best in savory dishes.

Q: Is protein powder safe for women?

A: Yes. No hormonal effects. Whey protein is food — milk protein. Women often benefit more from protein supplementation as vegetarian women particularly struggle to hit targets.

Q: Can teens use protein powder?

A: With caution. If a teen eats well and exercises, food should suffice. If genuinely under-eating protein (common in vegetarian teens), a half-scoop of whey is safe. Not necessary for sedentary teens.

Q: What is the best time to take protein powder?

A: When you need protein and don't have food access. Common: morning shake, post-workout, afternoon snack. No strict 'best time' — total daily protein is what matters.

Q: How to choose a protein powder in India?

A: Check: (1) Protein % (>70% of serving should be protein). (2) FSSAI certification. (3) No added creatine/BCAAs inflating protein count. (4) Leucine $\geq 2\text{g}$ per serving. (5) Reputable brand.

Q: Can I mix protein powder with hot milk?

A: Mixing in boiling liquid can denature protein and cause clumping. Add to warm (not boiling) milk or let milk cool slightly first.

Q: Is plant protein as good as whey?

A: Whey has higher PDCAAS (1.0 vs 0.7-0.9 for most plant proteins). Combined pea+rice protein has complete amino acid profile comparable to whey. Effective, just slightly higher intake may be needed.

WL-FAQ-013 MOD 13 — Master FAQ · FAQs: Creatine

GOALS Weight Loss | Fat Loss | General Health | Weight Maintenance

USERS Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes

DIFF Beginner

TAGS creatine, creatine monohydrate, does creatine cause fat gain, is creatine safe

DEFINITION

Comprehensive FAQ answers for: Creatine. 10 questions answered.

WHY IT MATTERS

- Category Creatine is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Creatine FAQs: 10.

FAQ

Q: What is creatine?

A: Creatine monohydrate is a supplement that increases ATP availability in muscles, improving strength by 5-15% and allowing 1-2 more reps per set. Not a steroid. Not hormonal. Found naturally in meat.

Q: Does creatine cause fat gain?

A: No. It causes 1-1.5 kg of intramuscular water retention in first 2 weeks. This is not fat. This water is inside muscle cells, making them larger and stronger.

Q: Who should take creatine?

A: Anyone doing resistance training wanting to improve strength and muscle preservation. Not beneficial for pure cardio athletes or completely sedentary individuals.

Q: How much creatine to take?

A: 3-5g daily. Creatine monohydrate is the only form with extensive evidence. No loading phase needed (just takes 3-4 weeks to saturate vs 1 week with loading).

Q: When should I take creatine?

A: Any time of day — timing is not critical. Post-workout is slightly studied as marginally better. Consistency over timing.

Q: Is creatine safe for long-term use?

A: 30+ years of research. Safe in healthy individuals. Most studied supplement in sports nutrition. Not recommended for those with kidney disease.

Q: Is creatine a steroid?

A: No. Completely different class of compound. Creatine is a natural amino acid compound found in meat. Steroids are synthetic hormones. They have nothing in common.

Q: Will I lose gains if I stop creatine?

A: Strength will return to natural levels. Water weight will decrease 1-1.5 kg. No permanent gains or losses — creatine supports performance while you're taking it.

Q: Is creatine available in India?

A: Yes. MuscleBlaze Creatine Monohydrate (affordable, reliable). Optimum Nutrition Creatine. Big Muscles Creatine. Standard price: ₹600-1200 for 300g (2-3 month supply).

Q: Does creatine cause hair loss?

A: One 2009 study showed increased DHT (a hormone linked to hair loss) in rugby players on creatine. No replicated study confirms hair loss. If genetically predisposed to baldness, theoretical concern exists.

WL-FAQ-014 MOD 13 — Master FAQ · FAQs: Walking

GOALS Weight Loss | Fat Loss | General Health | Weight Maintenance

USERS Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes

DIFF Beginner

TAGS walking, steps, 10000 steps, daily walk, walking for weight loss

DEFINITION

Comprehensive FAQ answers for: Walking. 10 questions answered.

WHY IT MATTERS

- Category Walking is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Walking FAQs: 10.

FAQ

Q: How much should I walk daily for weight loss?

A: 8,000-10,000 steps daily. 30-60 min brisk walk. Burns 200-400 kcal/day. One of the most sustainable weight-loss activities.

Q: Does walking reduce belly fat?

A: Walking reduces overall body fat, including belly fat — you cannot spot reduce. Regular walking + calorie deficit = belly fat reduction over time.

Q: Is walking enough for weight loss?

A: Walking + calorie deficit = effective weight loss. Adding resistance training gives better body composition. Walking alone without diet change: modest results.

Q: What is the best time to walk for weight loss?

A: Any time you will actually do it. Morning walk fasted: slightly higher fat oxidation. After dinner: reduces postprandial glucose spike significantly. Both are excellent.

Q: How many calories does walking burn?

A: Depends on weight and pace. 70 kg person at brisk pace (5-6 km/h): 250-300 kcal/hour. 30-min walk: 120-160 kcal.

Q: Is brisk walking better than slow walking?

A: Yes. Brisk walking (5-6 km/h, slightly breathless but can talk) burns 40% more calories than leisure walk. Aim for this intensity.

Q: Can I lose weight by walking alone without diet change?

A: Possible but slow. 30 min walking burns 150 kcal. Without diet change: 150 kcal/day surplus remains. To lose 1 kg/month from walking alone: need 75-90 min walking daily. Diet change is easier.

Q: How to make walking a habit?

A: Schedule it like a meeting. Walk with someone or podcast/music. Start small (10 min) and build. Take same route to automate. Morning walk is highest adherence time.

Q: Is walking after dinner good for weight loss?

A: Very effective. 10-15 min walk after dinner reduces postprandial glucose by 30%, burns 50-80 kcal, improves insulin sensitivity, and aids digestion.

Q: Do I need expensive walking shoes?

A: No, but good support is important to prevent knee and back problems. Basic running shoes with arch support (■800-2000) are sufficient. Don't walk long distances in sandals.

WL-FAQ-015 MOD 13 — Master FAQ · FAQs: Gym and Exercise

GOALS Weight Loss | Fat Loss | General Health | Weight Maintenance

USERS Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes

DIFF Beginner

TAGS gym, workout, cardio vs weights, how many days gym, gym beginner

DEFINITION

Comprehensive FAQ answers for: Gym and Exercise. 11 questions answered.

WHY IT MATTERS

- Category Gym and Exercise is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Gym and Exercise FAQs: 11.

FAQ

Q: How many days per week should I gym?

A: 3-5 days for optimal results. 3 days minimum for meaningful muscle stimulus. 5 days maximum for most non-athletes. Rest days are when growth happens.

Q: Should I focus on cardio or weights for weight loss?

A: Both are beneficial. Weights: builds muscle, raises TDEE, changes body composition. Cardio: burns calories directly. Optimal: 3 days weights + 2 days cardio.

Q: How long should a gym session be?

A: 45-60 min for resistance training. 30-45 min for cardio. More time is not always better — quality and progressive overload matter more than duration.

Q: What should a beginner do in the gym?

A: Full body 3 days/week: squat, deadlift, bench press, row, shoulder press. 3 sets, 8-12 reps. Learn form first. Progressive overload: add weight when you hit top of rep range for 2 sessions.

Q: Is gym necessary for weight loss?

A: No. Diet creates the deficit. Walking and home workouts achieve good results. Gym adds efficiency, progression, and better body composition outcomes.

Q: How long before I see gym results?

A: Strength improvements: 2-4 weeks. Visible muscle tone: 6-12 weeks. Significant appearance change: 3-6 months. Patience is the critical ingredient.

Q: Should I do cardio before or after weights?

A: Weights first if the goal is fat loss and muscle building. Cardio depletes glycogen, which impairs weight training performance. Exception: light 5-10 min cardio warm-up is always good.

Q: What should I eat before the gym?

A: 30-60 min before: 20-30g carbs + 10-15g protein. Examples: banana + protein shake, 1 roti + curd, oats + milk. No heavy meal within 2 hours.

Q: Is gym good for PCOS?

A: Resistance training is the BEST exercise for PCOS. Improves insulin sensitivity more than cardio. 3x/week strength training is the PCOS exercise gold standard.

Q: How do I avoid getting hurt at the gym?

A: Learn form before adding weight. Warm up 5-10 min. Progress gradually (2.5kg increments). Sleep and recover adequately. Don't skip leg day — imbalanced training causes injuries.

Q: Should beginners use machines or free weights?

A: Both are fine. Machines: safer for beginners, easier to learn. Free weights: better functional strength, more muscles activated. Learn basic machine movements first, add free weights progressively.

WL-FAQ-016 MOD 13 — Master FAQ · FAQs: PCOS

GOALS Weight Loss | Fat Loss | General Health | Weight Maintenance

USERS Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes

DIFF Beginner

TAGS PCOS, polycystic ovary, PCOS weight loss, insulin resistance PCOS, PCOS diet

DEFINITION

Comprehensive FAQ answers for: PCOS. 10 questions answered.

WHY IT MATTERS

- Category PCOS is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total PCOS FAQs: 10.

FAQ

Q: Does PCOS make weight loss impossible?

A: No, but it makes it harder. Insulin resistance requires low-GI diet. Weight loss is slower (0.25-0.5 kg/week vs 0.5-1 kg/week). 5-10% body weight loss can restore menstrual cycles and reduce symptoms.

Q: What is the best diet for PCOS?

A: Low GI diet (no refined carbs, maida, sugar), high protein (25-30g per meal), high fiber (35g/day), anti-inflammatory foods. Strength training over cardio.

Q: Does PCOS cause weight gain?

A: Insulin resistance causes fat storage, especially abdominal. Hormonal imbalance increases cravings and reduces activity. Both contribute to weight gain. Not inevitable but very common.

Q: Is intermittent fasting good for PCOS?

A: Variable evidence. Some PCOS women benefit from 14:10 or 16:8 IF for insulin sensitivity. Others find it worsens cortisol and hormonal balance. Trial with 14:10 first and assess 4-week response.

Q: Does metformin help PCOS weight loss?

A: Yes, modestly. Metformin reduces insulin resistance, which helps weight loss. Most effective when combined with diet and exercise. Consult endocrinologist.

Q: What supplements help PCOS?

A: Myo-inositol (2g) + D-chiro-inositol (50mg): best evidence. Vitamin D (most PCOS women are deficient). Magnesium (reduces insulin resistance). Omega-3 (reduces inflammation).

Q: Can PCOS be cured with weight loss?

A: Symptoms can be significantly reduced or put into remission with weight loss. Not a complete cure but 5-10% body weight loss can restore menstrual regularity, reduce testosterone, and improve fertility.

Q: Why do I gain weight around my belly with PCOS?

A: Insulin resistance promotes abdominal/visceral fat storage. High androgens in PCOS cause male-pattern fat distribution. This is hormonally driven — but can be reduced with low GI diet, exercise, and insulin sensitizers.

Q: Should I avoid dairy with PCOS?

A: Trial and observe. Some PCOS women find dairy worsens acne and symptoms. This is individual — dairy contains growth factors that may affect hormones. Try 4-week dairy-free trial and assess.

Q: How long does it take to see improvement in PCOS with diet?

A: Blood markers: 4-8 weeks. Menstrual regularity: 2-4 months. Acne improvement: 2-3 months. Visible weight loss: 3-6 months. Patience and consistency are essential.

WL-FAQ-017 MOD 13 — Master FAQ · FAQs: Diabetes

GOALS Weight Loss | Fat Loss | General Health | Weight Maintenance

USERS Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes

DIFF Beginner

TAGS diabetes, type 2 diabetes, blood sugar, diabetic diet, HbA1c, prediabetes

DEFINITION

Comprehensive FAQ answers for: Diabetes. 10 questions answered.

WHY IT MATTERS

- Category Diabetes is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Diabetes FAQs: 10.

FAQ**Q: Can I reverse type 2 diabetes with diet?**

A: Prediabetes: very likely reversible. T2D: significant improvement and medication reduction is possible with 7-10% body weight loss. Full 'remission' is achievable in recent-onset T2D. Medical supervision required.

Q: What is the best diet for type 2 diabetes?

A: Low GI, high fiber, controlled carbohydrates, high protein. No sugar-sweetened beverages. No refined carbs. Every meal includes protein, fiber, and non-starchy vegetables. Walk after every meal.

Q: How much rice can a diabetic eat?

A: Small portions only: 1/2 cup cooked (100g) maximum per meal. Combine with dal, vegetables, and protein. Choose basmati (lower GI). Eat it last (protein and vegetables first).

Q: Is fruit good for diabetics?

A: Whole fruit yes, juice no. Choose: apple, pear, guava, papaya, berries, jamun. Limit: mango, banana, grapes, litchi. 1 fruit per serving, not multiple at once. Monitor individual glucose response.

Q: Can diabetics eat sweets?

A: Rarely and in small amounts (1 piece). Never on empty stomach. Avoid daily. Occasional celebration sweet in small quantity with protein is acceptable.

Q: What exercises are best for diabetes?

A: Walking after meals (most evidence), resistance training (improves insulin sensitivity), yoga. Minimum: 150 min moderate exercise per week. Avoid long sedentary periods.

Q: Does cinnamon help diabetes?

A: Modest effect. 1-2g cinnamon/day reduces fasting glucose by 10-29% in some studies. Add to oats, milk, coffee. Not a treatment — a useful dietary addition.

Q: What is the glycemic index and why does it matter for diabetics?

A: GI measures blood glucose rise from food. High GI foods cause glucose spikes and require more insulin. Diabetics benefit from consistent low-GI eating to stabilize glucose and reduce insulin demand.

Q: Can I exercise with high blood sugar?

A: Not recommended above 250 mg/dL. Exercise at high glucose can cause ketoacidosis (in T1D) or worsening of hyperglycemia. Check sugar before exercise.

Q: What are the best Indian foods for diabetes?

A: Excellent: all dals, legumes, jowar/bajra roti, non-starchy vegetables, karela, methi, curd, eggs, chicken. Moderate: basmati rice (small), fruit. Avoid: white bread, sweet drinks, fried food, sweets.

WL-FAQ-018 MOD 13 — Master FAQ · FAQs: Sleep

GOALS	Weight Loss Fat Loss General Health Weight Maintenance
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes
DIFF	Beginner
TAGS	sleep, sleep and weight loss, 7 hours sleep, insomnia, cortisol sleep

DEFINITION

Comprehensive FAQ answers for: Sleep. 6 questions answered.

WHY IT MATTERS

- Category Sleep is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Sleep FAQs: 6.

FAQ

Q: How does poor sleep affect weight?

A: Increases ghrelin (hunger +28%), decreases leptin (fullness), raises cortisol (fat storage), reduces insulin sensitivity, promotes cravings for high-fat and high-sugar foods. Powerful weight gain driver.

Q: How many hours of sleep for weight loss?

A: 7-9 hours for adults. Research shows: 6 hours vs 8 hours = 559 more calories consumed the next day. Less than 6 hours: increased cortisol, muscle loss, carb cravings.

Q: Does napping help if I don't sleep well at night?

A: Short naps (20-30 min) can partially compensate for lost sleep. Longer naps (>45 min) can interfere with night sleep. A consistent night sleep schedule is the goal.

Q: What time should I sleep for maximum weight loss?

A: Before midnight. Circadian rhythm aligns fat oxidation and hormone release with early sleep. 10pm-11pm bedtime optimal. Consistent bedtime more important than exact time.

Q: Does exercise help sleep?

A: Yes significantly. Regular exercise improves sleep quality and duration. Avoid vigorous exercise within 2-3 hours of bed — it raises core temperature and delays sleep onset.

Q: What to eat to sleep better?

A: Tart cherries (melatonin), warm milk + turmeric (tryptophan), banana (magnesium, potassium), kiwi (serotonin precursor). Avoid heavy meals 3 hours before sleep.

WL-FAQ-019	MOD 13 — Master FAQ · FAQs: Stress
GOALS	Weight Loss Fat Loss General Health Weight Maintenance
USERS	Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes
DIFF	Beginner
TAGS	stress, cortisol, stress eating, stress weight gain, anxiety diet

DEFINITION

Comprehensive FAQ answers for: Stress. 5 questions answered.

WHY IT MATTERS

- Category Stress is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Stress FAQs: 5.

FAQ

Q: Why do I gain weight when stressed?

A: Cortisol promotes abdominal fat storage, increases cravings for high-sugar/fat foods, reduces sleep quality, and decreases motivation for exercise.

Q: How does exercise reduce stress?

A: Exercise releases endorphins and reduces cortisol for 12-24 hours post-exercise. Even 20-min walk reduces cortisol measurably. Yoga: reduces cortisol more than cardio in high-stress individuals.

Q: What foods reduce stress?

A: Magnesium-rich: dark chocolate, nuts, legumes. Omega-3: fatty fish, flaxseeds. Complex carbs: oats, dal. Probiotic: curd. Avoid: excessive caffeine (increases cortisol), alcohol.

Q: Can yoga help with weight loss?

A: Indirectly: reduces cortisol, improves sleep, increases mindfulness (reduces emotional eating). Gentle yoga burns 200-300 kcal/hour. Not efficient for fat loss directly — good as stress management.

Q: Is meditating good for weight loss?

A: Yes indirectly. 10-min daily mindfulness meditation reduces stress eating by 40%, improves sleep, and reduces emotional eating responses. Apps: Headspace, Calm, Insight Timer.

WL-FAQ-020 MOD 13 — Master FAQ · FAQs: Cravings

GOALS Weight Loss | Fat Loss | General Health | Weight Maintenance

USERS Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes

DIFF Beginner

TAGS cravings, sugar cravings, junk food cravings, night cravings, stop cravings

DEFINITION

Comprehensive FAQ answers for: Cravings. 5 questions answered.

WHY IT MATTERS

- Category Cravings is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Cravings FAQs: 5.

FAQ

Q: How do I stop sugar cravings?

A: Eat adequate protein (reduces sugar cravings by 60%). Reduce refined carbs (stabilizes blood sugar). Eat sweet potato, fruit, or dark chocolate as healthier sweet satisfaction. Stay hydrated.

Q: Why do I crave junk food at night?

A: Low-protein dinner doesn't sustain satiety. Habit formed around TV/phone time. Boredom or stress trigger. Solution: high-protein dinner, plan an evening snack, remove junk food from home.

Q: Does cutting sugar help weight loss?

A: Yes significantly. Added sugar (sweet drinks, biscuits, sweets) adds empty calories with no satiety. Eliminating added sugar alone can create 200-500 kcal/day deficit for many people.

Q: Is dark chocolate OK on a diet?

A: Yes. 20g dark chocolate (70%+) = 114 kcal, 2g protein, 10g carbs, 7.5g fat, 2g fiber. Satisfies chocolate cravings effectively. Choose 70%+ cocoa. High quality substitute for milk chocolate.

Q: How long until cravings stop on a new diet?

A: Most cravings diminish significantly after 3-4 weeks of consistent healthy eating as blood sugar stabilizes. The first 2 weeks are hardest. Sugar cravings specifically reduce faster with high protein diet.

WL-FAQ-021 MOD 13 — Master FAQ · FAQs: Hostel Life

GOALS Weight Loss | Fat Loss | General Health | Weight Maintenance

USERS Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes

DIFF Beginner

TAGS hostel, college hostel, mess food, hostel diet, student weight loss

DEFINITION

Comprehensive FAQ answers for: Hostel Life. 5 questions answered.

WHY IT MATTERS

- Category Hostel Life is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Hostel Life FAQs: 5.

FAQ

Q: How to eat healthy in a hostel?

A: Carry: peanut butter, rolled oats, protein powder, roasted chana, nuts, fruits. At mess: choose dal/sabzi over fried items; take small portion of rice; skip extra ghee.

Q: Best no-cook foods for hostel students?

A: Roasted chana, peanut butter, fruits, Greek yogurt/curd packets, protein shakes, sprouts (soak overnight), boiled eggs (batch cook).

Q: Is hostel mess food making me fat?

A: Likely contributing through excess oil, refined carbs, and fried items. Strategy: reduce rice portion, skip fried items, eat salad first, avoid second helpings of carbs.

Q: How to exercise in a hostel?

A: 20-min bodyweight circuit in your room: jumping jacks, squats, push-ups, plank, mountain climbers. No equipment. Walk to college instead of vehicle. Take stairs always.

Q: How to lose weight on a hostel budget?

A: Cheapest high-protein foods: eggs (■6-8 each), curd (■30-40/200g), roasted chana (■10/40g), dal at mess, peanut butter (■50/100g). Focus on: oats + milk + eggs daily.

WL-FAQ-022 MOD 13 — Master FAQ · FAQs: Office Life

GOALS Weight Loss | Fat Loss | General Health | Weight Maintenance

USERS Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes

DIFF Beginner

TAGS office, desk job, office snacks, office lunch, canteen

DEFINITION

Comprehensive FAQ answers for: Office Life. 5 questions answered.

WHY IT MATTERS

- Category Office Life is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Office Life FAQs: 5.

FAQ

Q: How to eat healthy at office?

A: Sunday meal prep: bring lunch from home. Keep healthy snacks at desk (nuts, roasted chana, fruits). Avoid canteen samosa/vada habit. Drink water instead of chai-and-biscuit.

Q: What to eat for a healthy office lunch?

A: Meal-prepped chicken/dal rice bowl, quinoa vegetable bowl, dal + roti in dabba, salad with protein. Heat in office microwave. Better and cheaper than canteen or delivery.

Q: How to exercise with a desk job?

A: Walk 20 min at lunch. Take stairs. Stand during calls. Walk to colleague instead of calling. 5-min walk every 60-90 min of sitting. Target: 8,000 steps at desk job.

Q: How to avoid office canteen junk food?

A: Pre-eat healthy snack before lunch break. Bring lunch from home. Distance yourself from the canteen counter. Social commitment: go with a colleague who also eats well.

Q: Is office stress causing my weight gain?

A: Possibly. Cortisol from work stress promotes fat storage. Also: desk sitting = low NEAT, canteen food access, stress eating in meetings. Address all four.

WL-FAQ-023 MOD 13 — Master FAQ · FAQs: Plateaus**GOALS** Weight Loss | Fat Loss | General Health | Weight Maintenance**USERS** Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes**DIFF** Beginner**TAGS** plateau, weight loss stopped, no progress, break plateau, stall**DEFINITION**

Comprehensive FAQ answers for: Plateaus. 5 questions answered.

WHY IT MATTERS

- Category Plateaus is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Plateaus FAQs: 5.

FAQ**Q: Why did my weight loss stop?**

A: Most common: (1) Tracking drift — hidden calories crept in. (2) TDEE dropped with weight loss. (3) NEAT unconsciously reduced. (4) Water retention masking fat loss. Check in this order.

Q: How do I break a weight loss plateau?

A: Step 1: 7-day verified accurate tracking. Step 2: Add 2000 steps/day. Step 3: Recalculate TDEE at new weight. Step 4: If still stuck after 2 more weeks — 1-2 week diet break at maintenance.

Q: Is it normal to stop losing weight after 2 weeks?

A: Yes. Initial rapid loss (week 1-2) is mostly water and glycogen. True fat loss rate appears slower. If 7-day average is not declining after 4-6 weeks of accurate tracking — genuine plateau.

Q: Why did I gain weight during my diet?

A: Scale increase during diet is almost always water: sodium-rich meal, menstrual cycle, high-carb meal, new exercise causing muscle inflammation. Check weekly average, not daily weight.

Q: Should I try a different diet if mine has stopped working?

A: First: verify the current diet is actually being followed accurately. Then: adjust current plan (more steps, slight calorie reduction) before switching. Switching diets repeatedly never works.

WL-FAQ-024 MOD 13 — Master FAQ · FAQs: Maintenance**GOALS** Weight Loss | Fat Loss | General Health | Weight Maintenance**USERS** Adult · Teen · Senior · Student · Office Worker · Vegetarian · Non-Vegetarian · PCOS · Diabetes**DIFF** Beginner**TAGS** maintenance, keep weight off, weight maintenance, after diet, long term**DEFINITION**

Comprehensive FAQ answers for: Maintenance. 5 questions answered.

WHY IT MATTERS

- Category Maintenance is a common source of user confusion and misinformation.

KEY FACTS

- All answers are evidence-based and actionable.
- Total Maintenance FAQs: 5.

FAQ**Q: How do I maintain weight after losing it?**

A: The 5 principles: daily weigh-in, maintain protein, maintain resistance training, stay active (8000+ steps), allow ± 2 kg buffer with action plan if trending higher.

Q: Why do people regain weight after dieting?

A: Reverting to previous habits, reduced metabolic rate from muscle loss during diet, hormonal changes driving hunger, psychological fatigue, stopping exercise. Prevention: treat maintenance as active management.

Q: How many calories should I eat to maintain weight?

A: Your current TDEE at current weight. Recalculate using Mifflin-St Jeor × activity multiplier. Start there — adjust ±100-200 kcal based on 4-week weight trend.

Q: Can I ever eat 'normal' again after dieting?

A: Yes, but 'normal' must be different from what caused weight gain initially. The goal is to make healthy eating your new normal — not to go back to old patterns.

Q: Is maintenance harder than losing weight?

A: For many people, yes — there's less external reinforcement (no scale dropping), and old habits return. Active management (weigh weekly, maintain exercise, track loosely) prevents regain.
