

Chapter 23

Bipolar disorder and plasticity: a key target for new treatment

Sheng-Min Wang and Chi-Un Pae

Department of Psychiatry, College of Medicine, The Catholic University of Korea, Seoul, South Korea

23.1 Introduction

Despite detrimental effect of bipolar disorder (BD), large proportion of patients with BD still do not experience sufficient treatment response leading to significant impairment in everyday functioning. Countless attempts have been taken for deciphering the neurobiology of BD and developing new therapeutic targets. However, “an oldie but a goodie” lithium is still the most effective pharmacological agent for BD (Won & Kim, 2017). Despite its confirmed treatment effects, lithium’s exact mechanism of action in mood regulation is still yet to be clarified. Along with the direct inhibition on glycogen synthase kinase 3 (GSK-3), increasing studies highlight lithium’s various effects on neurotrophic factors, neurotransmitters, oxidative metabolism, apoptosis, neuronal structures and glia, and second messenger systems which all are important factors contributing to plasticity of brain (Machado-Vieira, 2018; Wada, 2009). The evidence of disturbed plasticity in patients with BD is evident in both cellular and anatomical level. This chapter will elaborate current evidence for the dysfunction plasticity in BD, with a particular focus on the target for treatment.

23.2 Concepts of plasticity and implications in bipolar disorder

Plasticity, defined as ability to undergo and sustain change, is essential for the proper functioning of our nervous system by allowing organisms to maintain homeostasis to external change (de Sousa et al., 2014). Neuroplasticity refers to change of synaptic number and strength alternation, remodeling of axonal and dendritic architecture, growth of neuronal cell body, and generation of new neurons (in some parts of brain) (Sweatt, 2016). Synaptic

plasticity is a more specific term within neuroplasticity which refers to ability of synapses to strengthen or weaken over time, in response to increases or decreases in their activity (Fig. 23.1) (Citri & Malenka, 2008). Impairment of neuro- and synaptic plasticity involving dysregulation of neurotrophic factors and proteins are associated with BD while activation of neuroprotective and neurotrophic pathways are linked to the therapeutic effects of mood stabilizers (Soeiro-de-Souza et al., 2012). Lithium, which is considered as a “gold standard mood stabilizer,” is known to exert its effect via regulating neurotrophic factors, MAPK, Bcl-2, phosphoinositol signaling, intracellular calcium, glutamate activity, and GSK-3 which all lead to enhanced neuroplasticity.

23.3 Cellular studies

23.3.1 Mitochondrial-mediated pathway

In cellular level, studies showed that patients with BD have lower volume, decreased density, and smaller number, and reduced size of neurons and glial cells involving limbic regions including subgenual prefrontal cortex (PFC), orbital cortex, dorso-lateral PFC, amygdala, and basal ganglia (Cotter, Pariante, & Everall, 2001; Manji & Duman, 2001). Consistent studies reported mitochondrial dysfunction and associated dysregulation of calcium cascade as one of the most important biological abnormalities reported in BD (de Sousa et al., 2014). Mitochondria is important not only in regulating cellular energy metabolism but is also involved in modulation of cellular calcium levels, production of free radicals, and regulation of apoptosis. Thus it inevitably plays important role in neural plasticity and cellular resilience (Cheng, Hou, & Mattson, 2010). In the intrinsic mitochondrial-mediated




FIGURE 23.1 General concept of plasticity (de Sousa et al., 2014).

pathway, activation of proapoptotic BCL-2 family and heightened cytoplasmic calcium may enhance neuronal cell death and synaptic damage (Youle & Strasser, 2008). Likewise, BCL-2 family is also involved in facilitation of axonal regeneration, neurite growth, neuronal survival, and neurogenesis (Vinet, Bernier, & Parent, 2002). A polymorphism of BCL-2 antiapoptotic gene (rs956572) was found to be associated with an abnormal BCL-2 expression and dysfunctional calcium homeostasis in lymphoblastoid cells from patients with BD (Machado-Vieira et al., 2011). Another study reported that rs956572 polymorphism was related with increased levels of glutamate in anterior cingulate cortex of patients with BD (Soeiro-de-Souza et al., 2013).

Studies further showed that mitochondrial-mediated pathway involving BCL-2 and calcium regulation might be a promising therapeutic target. Chronic lithium therapy was shown to prevent oxidative stress and loss of mitochondrial membrane by blocking increment of calcium (Tseng & Lin-Shiau, 2002). In addition, lithium might reverse BCL-2 polymorphism (rs956572) associated dysfunctional calcium in BD (Yucel et al., 2007). Moreover, Bcl-2 expression and protein levels in the frontal cortex, hippocampus, and striatum were enhanced after long-term lithium therapy (Chen et al., 1999; Manji, Moore, & Chen, 1999). Likewise, electroconvulsive treatment, which is known to have a rapid mood stabilizing effect, is reported to increase BCL-2 expression leading to increased precursor cell proliferation in the dentate gyrus DG (Perera et al., 2007; Soeiro-de-Souza et al., 2012).

23.3.2 Neurotrophic factors and associated pathway

In terms of neurotrophic signaling cascade, brain-derived neurotrophic factor (BDNF) and its related pathway are known to play important roles in the pathophysiology of BD (Bramham & Messaoudi, 2005). BDNF binds to tyrosine kinase B receptor (TrkB) and activates downstream pathway involving phosphatidylinositol-3 kinase (PI3K)/protein kinase B (Akt) signaling pathway and extracellular regulated kinase (ERK)—mitogen-activated protein kinase (ERK/MAPK) pathway. BDNF itself is important for neurogenesis, neuronal survival, and normal maturation of neural development pathways. Moreover, it is important for synaptic plasticity, dendritic growth, and long-term memory consolidation (Post, 2007). Studies suggested that BDNF serum levels are decreased in depressive and manic episodes, which returned to normal levels in euthymia (Cunha et al., 2006; Machado-Vieira et al., 2007). The BDNF level was also shown to be negatively correlated with severity of manic and depressive symptoms (Fernandes et al., 2009). BDNF Val66val polymorphism was associated with increased risk of rapid cycling (Muller et al., 2006) while patients with BDNF val66met polymorphism

responded better to lithium prophylaxis than other genotype (Rybakowski et al., 2005).

The ERK/MAPK pathway is a major intracellular signaling cascade controlling the biological effects of neurotrophic factors and long-term cell plasticity (Chen & Manji, 2006). In the ERK/MAPK pathway, ribosomal S6 kinase is enhanced which activates cAMP response element-binding (CREB) protein (Sakamoto, Karelina, & Obrietan, 2011). CREB plays an important role in expression of the neuroprotective proteins, and this pathway is activated when BDNF binds to TrkB, which is known to be abnormal in patients with BD (Einat & Manji, 2006). PI3K also plays an important role in regulation of cell apoptosis, metabolism, and survival. The PI3K induces phosphoinositide dependent protein kinase-1 activating Akt serine/threonine kinase (Matsuda et al., 2019). The lowered PI3K/Akt signaling pathway eventually results in abnormalities of neuroplasticity.

Multiple studies suggested that BDNF and its related downstream pathway could be a potential therapeutic target of BD. BDNF is closely associated with the mechanism action of not only mood stabilizers but also antidepressants (Castren & Rantamaki, 2010; Coyle & Duman, 2003). For example, blockage of BDNF signaling with either a TrkB inhibitor or ERK/MAPK pathway inhibitor decreased the effects of antidepressants (Saarelainen et al., 2003). Likewise, long-term therapy with lithium and valproate increased BDNF concentration in hippocampus and PFC (Frey et al., 2006; Grande et al., 2010). Chronic lithium and valproate treatment were also found to activate ERK/MAPK pathway (Yuan et al., 2001) and they were also known to increase the levels of activated ERK in the emotion related brain lesions including anterior cingulate cortex and hippocampus in mice model (Chuang et al., 2002).

23.3.3 Glycogen synthase kinase-3 β

GSK-3 of mammals has two isoforms, GSK-3 α and GSK-3 β , and they are associated with the regulation of glycogen synthesis in response to insulin (Wada, 2009). The difference arises from phosphorylation of tyrosine 279 and 216 for GSK-3 α and GSK-3 β , respectively. Although cerebral GSK-3 α is known to be involved in neurodevelopment, no studies to date has reported importance of GSK-3 α in the pathophysiology of major neuropsychiatric disease including BD (Dandekar et al., 2018). In contrast, GSK-3 β is a key regulator of apoptosis, cellular plasticity, and resilience of neurons (Adli et al., 2007; Dandekar et al., 2018). The relationship between GSK-3 β gene polymorphism and symptom severity of patients with mood disorders including BD and major depressive disorder (MDD) were repeatedly shown.

The role of GSK-3 β in the pathophysiology of BD was discovered rather serendipitously from study results of lithium and valproate showing selective inhibitors of GSK-3 β (Adli et al., 2007; Valvassori et al., 2017). Grimes and

Jope proposed that GSK-3 β acts like a “gatekeeper” over a numerous downstream transcription factors and pathways which all play important role in regulating neuroplasticity (Grimes & Jope, 2001). The advantage of targeting GSK-3 β is that GSK-3 β is closely associated with mitochondrial-mediated pathway involving BCL-2, ERK/MAPK pathway, and PI3K/Akt signaling pathway (Soeiro-de-Souza et al., 2012; Soeiro-de-Souza et al., 2013). Disinhibition or increased activity of GSK-3 β can directly cause decreased level of β -catenin resulting in lowered BDNF and neuroplastic factors (Dandekar et al., 2018). The lowered BDNF and neuroplastic factors in turn will cause lowered activities of ERK/MAPK and PI3K/Akt pathway resulting in loss of neuroplasticity. Similarly, GSK-3 β can also directly inhibit mitochondrial-mediated pathway involving BCL-2 accelerating neuronal death and synaptic loss. Theoretically, multiple cellular pathways associated decreased neuroplasticity involved in pathophysiology of BD can either be blocked or reversed by targeting GSK-3 β . Multiple studies repeatedly showed that lithium and valproate inhibit GSK-3 β activity, enhance β -catenin activities, and increase PI3K mediated Akt (Chalecka-Franaszek & Chuang, 1999; Chen et al., 1999; Stambolic, Ruel, & Woodgett, 1996). Thus GSK-3 β inhibition is acknowledged to be a promising target to enhance neuroplasticity in the treatment of BD. Fig. 23.2 summarizes general concept of neuroplasticity disturbance in BD.




FIGURE 23.2 General concept of neuroplasticity disturbance in bipolar disorder (Dandekar et al., 2018; Soeiro-de-Souza et al., 2012). 1. Mitochondrial-mediated pathway: activation of proapoptotic BCL-2 family leads to excess calcium resulting in neuronal cell death and synaptic damage. 2. BDNF related pathway: decreased BDNF activities cause (a) decrement of ERK/MAPK pathway, which leads to lowered CREB activity; (b) decrement of PI3K/Akt signaling pathway, which leads to lowered Akt. Both (a) and (b) result in decreased neuroplasticity. 3. GSK-3 β is closely associated with both mitochondrial-mediated pathway and BDNF related pathway. Thus, theoretically, by inhibiting GSK-3 β majority of cellular pathway leading decreased neuroplasticity associated with BD can be blocked or prevented. *Akt*, protein kinase B; BDNF, brain-derived neurotrophic factor; *CREB*, cAMP response element-binding; *ERK*, extracellular regulated kinase; *MAPK*, mitogen-activated protein kinase; *PI3K*, phosphatidylinositol-3 kinase.

23.4 Neuroimaging studies

23.4.1 Structural magnetic resonance imaging

With the developmental of neuroimaging techniques, it has become possible to visualize plasticity changes per se and its negative impact of BD in human brain. Thus pathological changes in cellular level are now both visible and detectable in an anatomical level. The first wave of translating cellular level to anatomical level started with growing interest of underpinnings neurobiology of BD using structural magnetic resonance imaging (MRI) studies. As discussed earlier, cellular level studies showed that BD are associated with decrease in volume, density, number, and size of neuron and glial cells mainly in subgenual PFC, orbital cortex, dorso lateral PFC, amygdala, and basal ganglia (Cotter et al., 2001; Manji & Duman, 2001). In parallel, reduced gray matter volumes in areas of the orbital and medial PFC, ventral striatum and hippocampus, and enlargement of third ventricles were repeatedly observed in patients with BD (Selvaraj et al., 2012). Moreover, studies indicated that structural abnormalities are already present in first-episode schizophrenia or BD with significant overlap between the two. However, whole gray matter volume deficits and lateral ventricular enlargement were reported to be more prominent in first-episode schizophrenia whereas white matter volume reduction were more prominent in BD (De Peri et al., 2012). Likewise, evidences indicated that these structural brain abnormalities are already present in first-episode BD, but these do not overlap with those emerging from previous metaanalyses performed in patients with chronic BD (McIntosh et al., 2007). Taken these together, the loss of neuroplasticity observed in cellular or molecular level caused cortical and subcortical damage of brain in patients with BD. These findings also suggest that a distinct pattern of morphological changes occurs during the course of BD (Vita, De Peri, & Sacchetti, 2009). Studies showed that patients with longer illness duration have lower gray matter volume suggesting progression of BD could be associated with an abnormal cellular plasticity, especially of gray matter (Brambilla et al., 2001; Moorhead et al., 2007). An imaging study showed that adolescents with BD who were taking mood stabilizer had less volume loss (Chang et al., 2005). More importantly, patients who received long-term therapy with lithium showed an increase in gray matter volume over time, which suggested possibility of positive morphological and plastic alterations in the brain of treated BD patients (Moore et al., 2000; Sassi et al., 2002).

23.4.2 Diffusion tensor imaging

The structural MRI has advantages of assessing cortical volume loss or general cerebral atrophy. However, it has weakness in studying white matter because of its limited contrast detection in white matter tracts. In contrast, diffusion tensor imaging (DTI) is uniquely sensitive to white matter (WM)

microstructure analysis including axonal coherence, fiber density, and myelin integrity (O'Donnell & Westin, 2011). DTI applies two indices to investigate WM's physical integrity: the fractional anisotropy (FA) and the apparent diffusion coefficient (ADC) (Alexander et al., 2007). It may be used to map and characterize the three-dimensional diffusion of water as a function of spatial location, so it has been used extensively to map white matter tractography in the brain (Alexander et al., 2007). Thus it is more sensitive to white matter microstructure analysis including axonal coherence, fiber density, and myelin integrity than conventional structural MRI (Le Bihan, 2003). Thus it provides more specific information on plasticity-related morphological tissue changes than conventional anatomical MRI (Voss & Schiff, 2009).

Multiple studies using region-of-interest based analysis showed that patients with BD exhibited lower FA compared to healthy controls, mainly in WM tracts of prefrontal areas, anterior cingulum, callosal areas, and limbic-striatal areas (Arnone et al., 2008; Beyer et al., 2005; Wang et al., 2008). Likewise, a metaanalysis of voxel-based analysis showed clusters of FA and mean diffusivity alterations in areas relevant for emotional processing (Marlinge, Bellivier, & Houenou, 2014). The voxel-based analysis-DTI studies repeatedly suggested existence of a disconnected framework within fronto-limbic areas as well as between fronto-limbic and temporal, parietal, and occipital lobes. For example, a recent metaanalysis of the voxel-based analysis-diffusion tensor imaging (VBA-DTI) studies showed significant clusters of decreased FA within right temporo-parietal WM, left cingulum, and left anterior cingulate in patients with BD (Nortje et al., 2013). Thus DTI studies have consistently shown an alteration in prefrontal–limbic anatomical connectivity (Bellani et al., 2016). Bellani et al. suggested that due to disconnection between prefrontal and limbic regions, the prefrontal cortices will be unable to downregulate limbic regions leading to emotional instability of BD.

WM alterations in BD have been regarded as a potential for drug discovery and development (Marlinge et al., 2014). A study showed that compared to controls and MDD patients, BD patients had significantly lower FA and increased ADC in the majority of the WM fiber bundles connecting structures of the anterior limbic network (Benedetti et al., 2011). The study further suggested that WM changes might be secondary to myelin sheaths damage, and lithium therapy might protect against such injury. Thus DTI studies could be reliable biomarkers which can be used for development and discover of new drugs for BD treatment.

23.4.3 Resting-state functional magnetic resonance imaging

Resting-state functional MRI (rs-fMRI), by measuring changes in resting-state blood oxygen level dependence, reflects cyclic modulation of gross

cortical excitability and neuronal synchronization signals (Zuo et al., 2010). Thus it provides a noninvasive and task-free approach that removes some performance-related confounds, so it could reflect measure of “baseline” brain activity, connectivity, and plasticity of brain (Guerra-Carrillo, Mackey, & Bunge, 2014). Among diverse large-scale brain networks, aberrance of default mode network (DMN) in BD is consistently reported. According to a recent systematic by Zovetti et al., a minimum of 23 studies investigated DMN connectivity in patients with BD (Zovetti et al., 2020). Most studies repeatedly confirmed that BD patients show aberrant functional connectivity either in frontal or posterior hubs of DMN. However, studies showed conflicting results in terms of type of aberrance (i.e., both hypo and hypoconnectivity) and involved anatomic regions (PFC, anterior and posterior cingulate cortex, and precuneus). The controversial results might be attributed to the fact that rs-fMRI were different depending on patient’s manic, depressive, mixed, or euthymic phase (Anand et al., 2009; Chai et al., 2011; Chepenik et al., 2010).

Another meta-analysis investigated functional connectivity difference among health controls, BD patients, and major depressive patients (Gong et al., 2020). The study showed that patients with BD and MDD had common increment of amplitude of low-frequency fluctuation (ALFF) in the bilateral insula and right medial prefrontal cortex (mPFC) and decrement of ALFF in the left cerebellum posterior lobe. The study further showed that patients with BD had a different regional intrinsic activity from that of patients with MDD. The neuroplastic change illustrated by fMRI were also known to be associated with symptom severity differently in BD and MDD. In patients with MDD, metaregression showed that higher depressive symptoms measured via Hamilton Depression Symptom Scale (HAM-D) was correlated with greater increases in ALFF in the right anterior cingulate cortex (ACC). However, in patients with BD, higher HAM-D was associated with greater decreases in ALFF in the right posterior cingulate cortex (PCC). A more recent study found that BD patients presented different connectivity pattern depending on its disease state (Wang et al., 2020). In acutely ill state, decreased connectivity within the affective network and DMN were noted, which disappeared in the remitted state. Authors further suggested that rs-fMRI could become an important clinically relevant biomarker in guiding treatment strategies for BD. However, there remain issues such as reproducibility of results and lack of data demonstrating longitudinal changes of fMRI findings in BD before it can widely be used as a surrogate biomarker.

23.4.4 Neuroinflammation

Inflammation, by providing protective response to harmful stimuli, such as pathogens, damaged cells, or irritants, play an important role in plasticity (Medzhitov, 2010). Increased neuroinflammation corresponds to activation

of microglia, the resident macrophages of the brain, which is known to be associated with abnormalities of cerebral plasticity such as lowered hippocampal volume (Beumer et al., 2012). Mounting evidences showed strong association between inflammation and neural plasticity, and recent researches further suggested that neuroinflammation might serve as marker reflecting neuroplasticity (Golia et al., 2019; Singhal & Baune, 2017).

With recent advance imaging technique, microglia activation reflecting neuroinflammation can now be visualized in vivo with the radioactive contrast [11C]-(R)-PK11195 by means of positron emission tomography (PET). In a study comprising 14 patients with BD and 11 healthy controls, patients with BD showed a significantly increased [11C]-(R)-PK11195 binding potential, which is indicative of neuroinflammation, in the right hippocampus than HC (Haarman et al., 2014). A more recent study conducted by the same research group compared hippocampal volume and metabolites in bipolar I disorder (BD-I) patients with HCs using MRI and spectroscopy (Haarman et al., 2016). The study furthered early finding by showing that *N*-acetylaspartate (NAA) + *N*-acetyl-aspartyl-glutamate (NAAG) concentration were decreased in the left hippocampus of BD-I patients which suggested that there was a decreased neuronal integrity or lowered plasticity in this region.

23.5 Drugs development targeting GSK-3 β

As stated earlier in the chapter, by targeting GSK-3 β , multiple detrimental cascades related to neuroplasticity can be blocked simultaneously (Fig. 23.2). Thus majority of molecules targeting neuroplasticity act via GSK-3 β inhibition. Pharmacological inhibitors of GSK-3 β could be classified into three following categories: (1) competition with either adenosine triphosphate (ATP) or metal-binding sites that are involved directly in the catalytic process (ATP-competitive inhibitors); (2) primed phosphorylated Ser 9 binding area necessary for binding of substrate (substrate competitive inhibitors); (3) targeting kinase selectivity (non-ATP-competitive inhibitors) (Dandekar et al., 2018).

An initial study showed that AR-A014418, an ATP-competitive inhibitors of GSK-3 β inhibitors, crossed blood–brain barrier and reduced immobility time in rats exposed to the forced swim test, which is a well-established model for antidepressant efficacy (Gould et al., 2004). The study further showed that AR-A014418 decreased amphetamine-induced activity and suggested that it could be a potential molecule to treat BD, especially for bipolar depression. A more recent study showed that AR-A014418 has an antimanic effect by illustrating that it can reverse behavioral and oxidative stress related to manic symptoms induced ouabain (OUA) (Dal-Pont et al., 2019). OUA is known to mimic manic-like symptoms, such as hyperactivity, risk-taking behavior, and stereotypic behaviors via inhibition of Na + K + ATPase activity (El-Mallakh et al., 2003). L803-mts is another

selective peptide inhibitor GSK-3 β . A study showed that mouse injected intracerebroventricularly with L803-mts showed less immobility time in forced swim test (Kaidanovich-Beilin et al., 2004). The study further showed that L803-mts significantly inhibited purified GSK-3 β , and expression levels of beta-catenin was increased in the hippocampus of L803-mts-treated mouse by 50% compared to control peptide treated mouse. The upregulation of beta-catenin, as stated earlier, is known to be associated with inactivation of GSK-3 β (Sakanaka, Weiss, & Williams, 1998).

Most of the GSK-3 β inhibitors developed so far are ATP-competitive inhibitors. However, ATP binding site is highly conserved among numerous other protein kinases. Thus ATP-competitive inhibitor have a serious limitation regarding its low specificity resulting in frequent off-target effects (Eldar-Finkelman et al., 2010). In contrast, substrate competitive inhibitors are expected to be more specific, but its weak binding interaction with the enzyme is its major limitation. Due to this advantage substrate-competitive inhibitors have often been overlooked. In the other perspective, studies showed that the GSK-3 β over-expression does not exceed twofold to threefold over normal levels even in a pathological state. Therefore moderate-to-weak inhibition of GSK-3 β activity would be sufficient to provide good selectivity with desired treatment effect (Dandekar et al., 2018). Palomo et al. reported substituted 5-imino-1,2,4-thiadiazoles as the first small molecules able to inhibit GSK-3 β as a substrate competitive inhibitor (Palomo et al., 2012). The study showed that it can decrease inflammatory activation and selectively differentiate neural stem cells resulting in protection against neuronal cell death and promote endogenous neurogenesis by blocking the GSK-3 β substrate site. It is also known to cross blood–brain barrier penetration and bind to human serum albumin showing good drug-like properties (Perez et al., 2012).

In line with substrate competitive inhibitors, achieving kinase selectivity is one of the major obstacles in drug discovery and design of non-ATP-competitive inhibitors (Eglen & Reisine, 2009). The major advantage of non-ATP-competitive inhibitors is that only pathologic GSK-3 β can be inhibited without causing effect to GSK-3 α . By doing so, it may be possible to achieve high potency with reduced toxicity. However, discovering such molecules is very difficult because the isoenzymes GSK-3 α and GSK-3 β share more than 98% homology. Tideglusib (NP-12, P031112) is a potent and irreversible non-ATP-competitive inhibitor for GSK-3 β (Dominguez et al., 2012). Animal studies showed that it has neuroprotective, antiinflammatory, and neurogenesis-inducing effects (Wang et al., 2016). More importantly, tideglusib is shown to inhibit tau phosphorylation, amyloid deposition, and neuron loss. Thus most studies were done in Alzheimer's disease and other neurodegenerative disorder such as Progressive Supranuclear Palsy, and they all have entered clinical phase (i.e., Phase II for Alzheimer's disease) (del Ser et al., 2013). Thiadiazolidinone-8 (TDZD-8), another non-

ATP-competitive inhibitor, is an effective thiadiazolidinone derivate that is able to suppress the expression of inflammatory cytokines; it also presents tissue protective actions by GSK-3 β inhibition, promoting thus an antiinflammatory effect. Earlier studies showed that it can decrease immobility time in a tail suspension test (Belmaker, Agam, & Bersudsky, 2008) but more recent studies focused on its pro-cognitive and antiinflammatory effect for diverse neurodegenerative disorder including Alzheimer's disease. Table 23.1 provides summary of investigational drug with GSK-3 β inhibition effect.

23.6 Summary

A plethora of research indicated that disturbance of plasticity, or more specifically neuroplasticity, plays an important role in the pathophysiology of BD. The evidence is repeatedly found in cellular studies which became visible and detectable via various neuroimaging studies. The development of neuroimaging studies enabled us to assess diverse changes of neuroplasticity with greater time and space resolution. Investigating volumetric decrement using structural MRI can reflect neuronal damage and cellular death, white matter tract damage with DTI can represent myelination change, functional connectivity aberration with rs-fMRI can show synaptic dysfunction, and PK11195 PET study can mirror neuroinflammatory change of BD. Thus diverse biomarkers related with neuroplasticity can serve as a key target for the future treatment of BD.

In terms of drug development, studies consistently reported that classic mood stabilizers including lithium and valproate exert their effect by inhibiting GSK-3 β . Moreover, multiple cellular pathways associated decreased neuroplasticity including mitochondrial-mediated pathway, ERK/MAPK pathway, and PI3K/Akt signaling pathway could be targeted simultaneously by inhibiting GSK-3 β . Thus most investigational molecules developed to date work as GSK-3 β inhibitors. Despite promising findings, none of GSK-3 β inhibitors or drugs enhancing neuroplasticity reached the market. Several promising GSK-3 β inhibitors failed in the diverse developmental phase due to their poor selectivity and weak BBB permeability. Future studies should focus on increasing the selectivity and BBB permeability to increase its potential as a viable target for BD.

TABLE 23.1 Investigational drug for bipolar disorder (BD) with GSK-3 β inhibition effect.

| Molecule | Mode of action | Study phase | Findings | Target | Study |
|------------------------------|----------------------------------|-------------|---|---------------------|--|
| AR-A014418 | ATP-competitive inhibitors | Preclinical | Reduced immobility time in rats exposed to forced swim test | BD-depression | Gould et al. (2004) |
| | | Preclinical | Reverse OUA-induced behavior disturbance | BD-manic | Dal-Pont et al. (2019) |
| L803-mts | ATP-competitive inhibitors | Preclinical | Reduced immobility time in forced swim test. Upregulation of beta-catenin | BD-depression | Kaidanovich-Beilin et al. (2004) |
| 5-Imino-1,2,4-thiadiazoles | Substrate competitive inhibitors | Preclinical | Decrease neuronal cell death.Increase neurogenesis | Neuroinflammation | Palomo et al. (2012) |
| Tideglusib (NP-12, NP031112) | Non-ATP competitive inhibitors | Clinical | Neuroprotective,antiinflammatory,neurogenesis. Inhibit tau phosphorylation, amyloid deposition, neuron loss | Alzheimer's disease | Wang et al. (2016) |
| TDZD-8 | Non-ATP competitive inhibitors | Preclinical | Antiinflammatory effect | BD-depression | Belmaker et al. (2008) |

OUA, Ouabain; TDZD-8, Thiadiazolidinone-8.

References

- Adli, M., Hollinde, D. L., Stamm, T., Wiethoff, K., Tsahuridu, M., Kirchheiner, J.,
Bauer, M. (2007). Response to lithium augmentation in depression is associated with the glycogen synthase kinase 3-beta -50T/C single nucleotide polymorphism. *Biological Psychiatry*, 62(11), 1295–1302. Available from <https://doi.org/10.1016/j.biopsych.2007.03.023>.
- Alexander, A. L., Lee, J. E., Lazar, M., & Field, A. S. (2007). Diffusion tensor imaging of the brain. *Neurotherapeutics*, 4(3), 316–329. Available from <https://doi.org/10.1016/j.nurt.2007.05.011>.
- Anand, A., Li, Y., Wang, Y., Lowe, M. J., & Dzemidzic, M. (2009). Resting state corticolimbic connectivity abnormalities in unmedicated bipolar disorder and unipolar depression. *Psychiatry Research*, 171(3), 189–198. Available from <https://doi.org/10.1016/j.psychres.2008.03.012>.
- Arnone, D., McIntosh, A. M., Chandra, P., & Ebmeier, K. P. (2008). Meta-analysis of magnetic resonance imaging studies of the corpus callosum in bipolar disorder. *Acta Psychiatrica Scandinavica*, 118(5), 357–362. Available from <https://doi.org/10.1111/j.1600-0447.2008.01229.x>.
- Bellani, M., Boschello, F., Delvecchio, G., Dusi, N., Altamura, C. A., Ruggeri, M., & Brambilla, P. (2016). DTI and Myelin Plasticity in Bipolar Disorder: Integrating Neuroimaging and Neuropathological Findings. *Frontiers in Psychiatry*, 7, 21. Available from <https://doi.org/10.3389/fpsy.2016.00021>.
- Belmaker, R. H., Agam, G., & Bersudsky, Y. (2008). Role of GSK3beta in behavioral abnormalities induced by serotonin deficiency. *Proc Natl Acad Sci U S A*, 105(20), E23. Available from <https://doi.org/10.1073/pnas.0801168105>, author reply E24.
- Benedetti, F., Absinta, M., Rocca, M. A., Radaelli, D., Poletti, S., Bernasconi, A., ... Filippi, M. (2011). Tract-specific white matter structural disruption in patients with bipolar disorder. *Bipolar Disorders*, 13(4), 414–424. Available from <https://doi.org/10.1111/j.1399-5618.2011.00938.x>.
- Beumer, W., Gibney, S. M., Drexhage, R. C., Pont-Lezica, L., Doorduyn, J., Klein, H. C., ... Drexhage, H. A. (2012). The immune theory of psychiatric diseases: a key role for activated microglia and circulating monocytes. *Journal of Leukocyte Biology*, 92(5), 959–975. Available from <https://doi.org/10.1189/jlb.0212100>.
- Beyer, J. L., Taylor, W. D., MacFall, J. R., Kuchibhatla, M., Payne, M. E., Provenzale, J. M., ... Krishnan, K. R. (2005). Cortical white matter microstructural abnormalities in bipolar disorder. *Neuropsychopharmacology*, 30(12), 2225–2229. Available from <https://doi.org/10.1038/sj.npp.1300802>.
- Brambilla, P., Harenski, K., Nicoletti, M., Mallinger, A. G., Frank, E., Kupfer, D. J., ... Soares, J. C. (2001). Differential effects of age on brain gray matter in bipolar patients and healthy individuals. *Neuropsychobiology*, 43(4), 242–247. Available from <https://doi.org/10.1159/000054897>.
- Bramham, C. R., & Messaoudi, E. (2005). BDNF function in adult synaptic plasticity: the synaptic consolidation hypothesis. *Progress in neurobiology*, 76(2), 99–125. Available from <https://doi.org/10.1016/j.pneurobio.2005.06.003>.
- Castren, E., & Rantamaki, T. (2010). The role of BDNF and its receptors in depression and antidepressant drug action: Reactivation of developmental plasticity. *Developmental neurobiology*, 70(5), 289–297. Available from <https://doi.org/10.1002/dneu.20758>.

- Chai, X. J., Whitfield-Gabrieli, S., Shinn, A. K., Gabrieli, J. D., Nieto Castanon, A., McCarthy, J. M., ... Ongur, D. (2011). Abnormal medial prefrontal cortex resting-state connectivity in bipolar disorder and schizophrenia. *Neuropsychopharmacology*, 36(10), 2009–2017. Available from <https://doi.org/10.1038/npp.2011.88>.
- Chalecka-Franaszek, E., & Chuang, D. M. (1999). Lithium activates the serine/threonine kinase Akt-1 and suppresses glutamate-induced inhibition of Akt-1 activity in neurons. *Proc Natl Acad Sci U S A*, 96(15), 8745–8750. Available from <https://doi.org/10.1073/pnas.96.15.8745>.
- Chang, K., Karchemskiy, A., Barnea-Goraly, N., Garrett, A., Simeonova, D. I., & Reiss, A. (2005). Reduced amygdalar gray matter volume in familial pediatric bipolar disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*, 44(6), 565–573. Available from <https://doi.org/10.1097/01.chi.0000159948.75136.0d>.
- Chen, G., Huang, L. D., Jiang, Y. M., & Manji, H. K. (1999). The mood-stabilizing agent valproate inhibits the activity of glycogen synthase kinase-3. *Journal of Neurochemistry*, 72(3), 1327–1330. Available from <https://doi.org/10.1046/j.1471-4159.2000.0721327.x>.
- Chen, G., & Manji, H. K. (2006). The extracellular signal-regulated kinase pathway: an emerging promising target for mood stabilizers. *Current Opinion in Psychiatry*, 19(3), 313–323. Available from <https://doi.org/10.1097/01.yco.0000218604.63463.cd>.
- Chen, G., Zeng, W. Z., Yuan, P. X., Huang, L. D., Jiang, Y. M., Zhao, Z. H., & Manji, H. K. (1999). The mood-stabilizing agents lithium and valproate robustly increase the levels of the neuroprotective protein bcl-2 in the CNS. *Journal of Neurochemistry*, 72(2), 879–882. Available from <https://doi.org/10.1046/j.1471-4159.1999.720879.x>.
- Cheng, A., Hou, Y., & Mattson, M. P. (2010). Mitochondria and neuroplasticity. *ASN NEURO*, 2(5), e00045. Available from <https://doi.org/10.1042/AN20100019>.
- Chepenik, L. G., Raffo, M., Hampson, M., Lacadie, C., Wang, F., Jones, M. M., ... Blumberg, H. P. (2010). Functional connectivity between ventral prefrontal cortex and amygdala at low frequency in the resting state in bipolar disorder. *Psychiatry Research*, 182(3), 207–210. Available from <https://doi.org/10.1016/j.psychresns.2010.04.002>.
- Chuang, D. M., Chen, R. W., Chalecka-Franaszek, E., Ren, M., Hashimoto, R., Senatorov, V., ... Leeds, P. (2002). Neuroprotective effects of lithium in cultured cells and animal models of diseases. *Bipolar Disorders*, 4(2), 129–136. Available from <https://doi.org/10.1034/j.1399-5618.2002.01179.x>.
- Citri, A., & Malenka, R. C. (2008). Synaptic plasticity: multiple forms, functions, and mechanisms. *Neuropsychopharmacology*, 33(1), 18–41. Available from <https://doi.org/10.1038/sj.npp.1301559>.
- Cotter, D. R., Pariante, C. M., & Everall, I. P. (2001). Glial cell abnormalities in major psychiatric disorders: the evidence and implications. *Brain Research Bulletin*, 55(5), 585–595. Available from [https://doi.org/10.1016/s0361-9230\(01\)00527-5](https://doi.org/10.1016/s0361-9230(01)00527-5).
- Coyle, J. T., & Duman, R. S. (2003). Finding the intracellular signaling pathways affected by mood disorder treatments. *Neuron*, 38(2), 157–160. Available from [https://doi.org/10.1016/s0896-6273\(03\)00195-8](https://doi.org/10.1016/s0896-6273(03)00195-8).
- Cunha, A. B., Frey, B. N., Andreazza, A. C., Goi, J. D., Rosa, A. R., Goncalves, C. A., ... Kapczinski, F. (2006). Serum brain-derived neurotrophic factor is decreased in bipolar disorder during depressive and manic episodes. *Neuroscience Letters*, 398(3), 215–219. Available from <https://doi.org/10.1016/j.neulet.2005.12.085>.
- Dal-Pont, G. C., Resende, W. R., Varela, R. B., Menegas, S., Trajano, K. S., Peterle, B. R., ... Valvassori, S. S. (2019). Inhibition of GSK-3beta on Behavioral Changes and Oxidative Stress in an Animal Model of Mania. *Molecular Neurobiology*, 56(4), 2379–2393. Available from <https://doi.org/10.1007/s12035-018-1226-2>.

- Dandekar, M. P., Valvassori, S. S., Dal-Pont, G. C., & Quevedo, J. (2018). Glycogen Synthase Kinase-3beta as a Putative Therapeutic Target for Bipolar Disorder. *Current Drug Metabolism*, 19(8), 663–673. Available from <https://doi.org/10.2174/1389200219666171227203737>.
- De Peri, L., Crescini, A., Deste, G., Fusar-Poli, P., Sacchetti, E., & Vita, A. (2012). Brain structural abnormalities at the onset of schizophrenia and bipolar disorder: a meta-analysis of controlled magnetic resonance imaging studies. *Current Pharmaceutical Design*, 18(4), 486–494. Available from <https://doi.org/10.2174/138161212799316253>.
- de Sousa, R. T., Machado-Vieira, R., Zarate, C. A., Jr., & Manji, H. K. (2014). Targeting mitochondrially mediated plasticity to develop improved therapeutics for bipolar disorder. *Expert opinion on therapeutic targets*, 18(10), 1131–1147. Available from <https://doi.org/10.1517/14728222.2014.940893>.
- del Ser, T., Steinwachs, K. C., Gertz, H. J., Andres, M. V., Gomez-Carrillo, B., Medina, M., ... Leon, T. (2013). Treatment of Alzheimer's disease with the GSK-3 inhibitor tideglusib: a pilot study. *Journal of Alzheimer's disease : JAD*, 33(1), 205–215. Available from <https://doi.org/10.3233/JAD-2012-120805>.
- Dominguez, J. M., Fuertes, A., Orozco, L., del Monte-Millan, M., Delgado, E., & Medina, M. (2012). Evidence for irreversible inhibition of glycogen synthase kinase-3beta by tideglusib. *The Journal of Biological Chemistry*, 287(2), 893–904. Available from <https://doi.org/10.1074/jbc.M111.306472>.
- Eglen, R. M., & Reisine, T. (2009). The current status of drug discovery against the human kinome. *Assay and drug development technologies*, 7(1), 22–43. Available from <https://doi.org/10.1089/adt.2008.164>.
- Einat, H., & Manji, H. K. (2006). Cellular plasticity cascades: genes-to-behavior pathways in animal models of bipolar disorder. *Biological Psychiatry*, 59(12), 1160–1171. Available from <https://doi.org/10.1016/j.biopsych.2005.11.004>.
- El-Mallakh, R. S., El-Masri, M. A., Huff, M. O., Li, X. P., Decker, S., & Levy, R. S. (2003). Intracerebroventricular administration of ouabain as a model of mania in rats. *Bipolar Disorders*, 5(5), 362–365. Available from <https://doi.org/10.1034/j.1399-5618.2003.00053.x>.
- Eldar-Finkelman, H., Licht-Murava, A., Pietrokovski, S., & Eisenstein, M. (2010). Substrate competitive GSK-3 inhibitors - strategy and implications. *Biochimica et Biophysica Acta*, 1804(3), 598–603. Available from <https://doi.org/10.1016/j.bbapap.2009.09.010>.
- Fernandes, B. S., Gama, C. S., Kauer-Sant'Anna, M., Lobato, M. I., Belmonte-de-Abreu, P., & Kapczinski, F. (2009). Serum brain-derived neurotrophic factor in bipolar and unipolar depression: a potential adjunctive tool for differential diagnosis. *Journal of Psychiatric Research*, 43(15), 1200–1204. Available from <https://doi.org/10.1016/j.jpsychires.2009.04.010>.
- Frey, B. N., Andreazza, A. C., Cereser, K. M., Martins, M. R., Valvassori, S. S., Reus, G. Z., ... Kapczinski, F. (2006). Effects of mood stabilizers on hippocampus BDNF levels in an animal model of mania. *Life Sciences*, 79(3), 281–286. Available from <https://doi.org/10.1016/j.lfs.2006.01.002>.
- Golia, M. T., Poggini, S., Alboni, S., Garofalo, S., Ciano Albanese, N., Viglione, A., ... Maggi, L. (2019). Interplay between inflammation and neural plasticity: Both immune activation and suppression impair LTP and BDNF expression. *Brain, Behavior, and Immunity*, 81, 484–494. Available from <https://doi.org/10.1016/j.bbi.2019.07.003>.
- Gong, J., Wang, J., Qiu, S., Chen, P., Luo, Z., Wang, J., ... Wang, Y. (2020). Common and distinct patterns of intrinsic brain activity alterations in major depression and bipolar disorder: voxel-based meta-analysis. *Translational Psychiatry*, 10(1), 353. Available from <https://doi.org/10.1038/s41398-020-01036-5>.

- Gould, T. D., Einat, H., Bhat, R., & Manji, H. K. (2004). AR-A014418, a selective GSK-3 inhibitor, produces antidepressant-like effects in the forced swim test. *The international journal of neuropsychopharmacology / official scientific journal of the Collegium Internationale Neuropsychopharmacologicum (CINP)*, 7(4), 387–390. Available from <https://doi.org/10.1017/S1461145704004535>.
- Grande, I., Fries, G. R., Kunz, M., & Kapczinski, F. (2010). The role of BDNF as a mediator of neuroplasticity in bipolar disorder. *Psychiatry Investigation*, 7(4), 243–250. Available from <https://doi.org/10.4306/pi.2010.7.4.243>.
- Grimes, C. A., & Jope, R. S. (2001). The multifaceted roles of glycogen synthase kinase 3 β in cellular signaling. *Progress in neurobiology*, 65(4), 391–426. Available from [https://doi.org/10.1016/s0301-0082\(01\)00011-9](https://doi.org/10.1016/s0301-0082(01)00011-9).
- Guerra-Carrillo, B., Mackey, A. P., & Bunge, S. A. (2014). Resting-state fMRI: a window into human brain plasticity. *The Neuroscientist*, 20(5), 522–533. Available from <https://doi.org/10.1177/1073858414524442>.
- Haarman, B. C., Burger, H., Doorduyn, J., Renken, R. J., Sibeijn-Kuiper, A. J., Marsman, J. B., ... Riemersma-Van, & der Lek, R. F. (2016). Volume, metabolites and neuroinflammation of the hippocampus in bipolar disorder - A combined magnetic resonance imaging and positron emission tomography study. *Brain, Behavior, and Immunity*, 56, 21–33. Available from <https://doi.org/10.1016/j.bbi.2015.09.004>.
- Haarman, B. C., Riemersma-Van der Lek, R. F., de Groot, J. C., Ruhe, H. G., Klein, H. C., Zandstra, T. E., ... Doorduyn, J. (2014). Neuroinflammation in bipolar disorder - A [(11)C]-(R)-PK11195 positron emission tomography study. *Brain, Behavior, and Immunity*, 40, 219–225. Available from <https://doi.org/10.1016/j.bbi.2014.03.016>.
- Kaidanovich-Beilin, O., Milman, A., Weizman, A., Pick, C. G., & Eldar-Finkelman, H. (2004). Rapid antidepressive-like activity of specific glycogen synthase kinase-3 inhibitor and its effect on beta-catenin in mouse hippocampus. *Biological Psychiatry*, 55(8), 781–784. Available from <https://doi.org/10.1016/j.biopsych.2004.01.008>.
- Le Bihan, D. (2003). Looking into the functional architecture of the brain with diffusion MRI. *Nature Reviews Neuroscience*, 4(6), 469–480. Available from <https://doi.org/10.1038/nrn1119>.
- Machado-Vieira, R. (2018). Lithium, Stress, and Resilience in Bipolar Disorder: Deciphering this key homeostatic synaptic plasticity regulator. *Journal of Affective Disorders*, 233, 92–99. Available from <https://doi.org/10.1016/j.jad.2017.12.026>.
- Machado-Vieira, R., Dietrich, M. O., Leke, R., Cereser, V. H., Zanatto, V., Kapczinski, F., ... Gentil, V. (2007). Decreased plasma brain derived neurotrophic factor levels in unmedicated bipolar patients during manic episode. *Biological Psychiatry*, 61(2), 142–144. Available from <https://doi.org/10.1016/j.biopsych.2006.03.070>.
- Machado-Vieira, R., Pivovarov, N. B., Stanika, R. I., Yuan, P., Wang, Y., Zhou, R., ... Andrews, S. B. (2011). The Bcl-2 gene polymorphism rs956572AA increases inositol 1,4,5-trisphosphate receptor-mediated endoplasmic reticulum calcium release in subjects with bipolar disorder. *Biological Psychiatry*, 69(4), 344–352. Available from <https://doi.org/10.1016/j.biopsych.2010.10.019>.
- Manji, H. K., & Duman, R. S. (2001).). Impairments of neuroplasticity and cellular resilience in severe mood disorders: implications for the development of novel therapeutics. *Psychopharmacology Bulletin*, 35(2), 5–49, Retrieved from. Available from <https://www.ncbi.nlm.nih.gov/pubmed/12397885>.
- Manji, H. K., Moore, G. J., & Chen, G. (1999). Lithium at 50: have the neuroprotective effects of this unique cation been overlooked? *Biological Psychiatry*, 46(7), 929–940. Available from [https://doi.org/10.1016/s0006-3223\(99\)00165-1](https://doi.org/10.1016/s0006-3223(99)00165-1).

- Marlinge, E., Bellivier, F., & Houenou, J. (2014). White matter alterations in bipolar disorder: potential for drug discovery and development. *Bipolar Disorders*, 16(2), 97–112. Available from <https://doi.org/10.1111/bdi.12135>.
- Matsuda, S., Ikeda, Y., Murakami, M., Nakagawa, Y., Tsuji, A., & Kitagishi, Y. (2019). Roles of PI3K/AKT/GSK3 Pathway Involved in Psychiatric Illnesses. *Diseases*, 7(1). Available from <https://doi.org/10.3390/diseases7010022>.
- McIntosh, A. M., Moorhead, T. W., McKirdy, J., Sussmann, J. E., Hall, J., Johnstone, E. C., & Lawrie, S. M. (2007). Temporal grey matter reductions in bipolar disorder are associated with the BDNF Val66Met polymorphism. *Molecular Psychiatry*, 12(10), 902–903. Available from <https://doi.org/10.1038/sj.mp.4002044>.
- Medzhitov, R. (2010). Inflammation 2010: new adventures of an old flame. *Cell*, 140(6), 771–776. Available from <https://doi.org/10.1016/j.cell.2010.03.006>.
- Moore, G. J., Bebchuk, J. M., Wilds, I. B., Chen, G., & Manji, H. K. (2000). Lithium-induced increase in human brain grey matter. *Lancet*, 356(9237), 1241–1242. Available from [https://doi.org/10.1016/s0140-6736\(00\)02793-8](https://doi.org/10.1016/s0140-6736(00)02793-8).
- Moorhead, T. W., McKirdy, J., Sussmann, J. E., Hall, J., Lawrie, S. M., Johnstone, E. C., & McIntosh, A. M. (2007). Progressive gray matter loss in patients with bipolar disorder. *Biological Psychiatry*, 62(8), 894–900. Available from <https://doi.org/10.1016/j.biopsych.2007.03.005>.
- Muller, D. J., de Luca, V., Sicard, T., King, N., Strauss, J., & Kennedy, J. L. (2006). Brain-derived neurotrophic factor (BDNF) gene and rapid-cycling bipolar disorder: family-based association study. *The British Journal of Psychiatry*, 189, 317–323. Available from <https://doi.org/10.1192/bjp.bp.105.010587>.
- Nortje, G., Stein, D. J., Radua, J., Mataix-Cols, D., & Horn, N. (2013). Systematic review and voxel-based meta-analysis of diffusion tensor imaging studies in bipolar disorder. *Journal of Affective Disorders*, 150(2), 192–200. Available from <https://doi.org/10.1016/j.jad.2013.05.034>.
- O'Donnell, L. J., & Westin, C. F. (2011). An introduction to diffusion tensor image analysis. *Neurosurgery Clinics of North America*, 22(2), 185–196. Available from <https://doi.org/10.1016/j.nec.2010.12.004>, viii.
- Palomo, V., Perez, D. I., Perez, C., Morales-Garcia, J. A., Soteras, I., Alonso-Gil, S., ... Martinez, A. (2012). 5-imino-1,2,4-thiadiazoles: first small molecules as substrate competitive inhibitors of glycogen synthase kinase 3. *Journal of Medicinal Chemistry*, 55(4), 1645–1661. Available from <https://doi.org/10.1021/jm201463v>.
- Perera, T. D., Coplan, J. D., Lisanby, S. H., Lipira, C. M., Arif, M., Carpio, C., ... Dwork, A. J. (2007). Antidepressant-induced neurogenesis in the hippocampus of adult nonhuman primates. *The Journal of Neuroscience*, 27(18), 4894–4901. Available from <https://doi.org/10.1523/JNEUROSCI.0237-07.2007>.
- Perez, D. I., Pistolozzi, M., Palomo, V., Redondo, M., Fortugno, C., Gil, C., ... Bertucci, C. (2012). 5-Imino-1,2,4-thiadiazoles and quinazolines derivatives as glycogen synthase kinase 3beta (GSK-3beta) and phosphodiesterase 7 (PDE7) inhibitors: determination of blood-brain barrier penetration and binding to human serum albumin. *European Journal of Pharmaceutical Sciences*, 45(5), 677–684. Available from <https://doi.org/10.1016/j.ejps.2012.01.007>.
- Post, R. M. (2007). Role of BDNF in bipolar and unipolar disorder: clinical and theoretical implications. *Journal of Psychiatric Research*, 41(12), 979–990. Available from <https://doi.org/10.1016/j.jpsychires.2006.09.009>.
- Rybakowski, J. K., Suwalska, A., Skibinska, M., Szczepankiewicz, A., Leszczynska-Rodziewicz, A., Permoda, A., ... Hauser, J. (2005). Prophylactic lithium response and polymorphism of

- the brain-derived neurotrophic factor gene. *Pharmacopsychiatry*, 38(4), 166–170. Available from <https://doi.org/10.1055/s-2005-871239>.
- Saarelainen, T., Hendolin, P., Lucas, G., Koponen, E., Sairanen, M., MacDonald, E., . . . Castren, E. (2003). Activation of the TrkB neurotrophin receptor is induced by antidepressant drugs and is required for antidepressant-induced behavioral effects. *The Journal of Neuroscience*, 23(1), 349–357. Retrieved from. Available from <https://www.ncbi.nlm.nih.gov/pubmed/12514234>.
- Sakamoto, K., Karelina, K., & Obrietan, K. (2011). CREB: a multifaceted regulator of neuronal plasticity and protection. *Journal of Neurochemistry*, 116(1), 1–9. Available from <https://doi.org/10.1111/j.1471-4159.2010.07080.x>.
- Sakanaka, C., Weiss, J. B., & Williams, L. T. (1998). Bridging of beta-catenin and glycogen synthase kinase-3 β by axin and inhibition of beta-catenin-mediated transcription. *Proc Natl Acad Sci U S A*, 95(6), 3020–3023. Available from <https://doi.org/10.1073/pnas.95.6.3020>.
- Sassi, R. B., Nicoletti, M., Brambilla, P., Mallinger, A. G., Frank, E., Kupfer, D. J., . . . Soares, J. C. (2002). Increased gray matter volume in lithium-treated bipolar disorder patients. *Neuroscience Letters*, 329(2), 243–245. Available from [https://doi.org/10.1016/s0304-3940\(02\)00615-8](https://doi.org/10.1016/s0304-3940(02)00615-8).
- Selvaraj, S., Arnone, D., Job, D., Stanfield, A., Farrow, T. F., Nugent, A. C., . . . McIntosh, A. M. (2012). Grey matter differences in bipolar disorder: a meta-analysis of voxel-based morphometry studies. *Bipolar Disorders*, 14(2), 135–145. Available from <https://doi.org/10.1111/j.1399-5618.2012.01000.x>.
- Singhal, G., & Baune, B. T. (2017). Microglia: An Interface between the Loss of Neuroplasticity and Depression. *Frontiers in Cellular Neuroscience*, 11, 270. Available from <https://doi.org/10.3389/fncel.2017.00270>.
- Soeiro-de-Souza, M. G., Dias, V. V., Figueira, M. L., Forlenza, O. V., Gattaz, W. F., Zarate, C. A., Jr., & Machado-Vieira, R. (2012). Translating neurotrophic and cellular plasticity: from pathophysiology to improved therapeutics for bipolar disorder. *Acta Psychiatrica Scandinavica*, 126(5), 332–341. Available from <https://doi.org/10.1111/j.1600-0447.2012.01889.x>.
- Soeiro-de-Souza, M. G., Salvatore, G., Moreno, R. A., Otaduy, M. C., Chaim, K. T., Gattaz, W. F., . . . Machado-Vieira, R. (2013). Bcl-2 rs956572 polymorphism is associated with increased anterior cingulate cortical glutamate in euthymic bipolar I disorder. *Neuropsychopharmacology*, 38(3), 468–475. Available from <https://doi.org/10.1038/npp.2012.203>.
- Stambolic, V., Ruel, L., & Woodgett, J. R. (1996). Lithium inhibits glycogen synthase kinase-3 activity and mimics wingless signalling in intact cells. *Current Biology*, 6(12), 1664–1668. Available from [https://doi.org/10.1016/s0960-9822\(02\)70790-2](https://doi.org/10.1016/s0960-9822(02)70790-2).
- Sweatt, J. D. (2016). Neural plasticity and behavior - sixty years of conceptual advances. *Journal of Neurochemistry*, 139(Suppl 2), 179–199. Available from <https://doi.org/10.1111/jnc.13580>.
- Tseng, W. P., & Lin-Shiau, S. Y. (2002). Long-term lithium treatment prevents neurotoxic effects of beta-bungarotoxin in primary cultured neurons. *Journal of Neuroscience Research*, 69(5), 633–641. Available from <https://doi.org/10.1002/jnr.10318>.
- Valvassori, S. S., Dal-Pont, G. C., Resende, W. R., Jornada, L. K., Peterle, B. R., Machado, A. G., . . . Quevedo, J. (2017). Lithium and valproate act on the GSK-3 β signaling pathway to reverse manic-like behavior in an animal model of mania induced by ouabain. *Neuropharmacology*, 117, 447–459. Available from <https://doi.org/10.1016/j.neuropharm.2016.10.015>.

- Vinet, J., Bernier, P. J., & Parent, A. (2002). Bcl-2 expression in thalamus, brainstem, cerebellum and visual cortex of adult primate. *Neuroscience Research*, 42(4), 269–277. Available from [https://doi.org/10.1016/s0168-0102\(02\)00004-4](https://doi.org/10.1016/s0168-0102(02)00004-4).
- Vita, A., De Peri, L., & Sacchetti, E. (2009). Gray matter, white matter, brain, and intracranial volumes in first-episode bipolar disorder: a meta-analysis of magnetic resonance imaging studies. *Bipolar Disorders*, 11(8), 807–814. Available from <https://doi.org/10.1111/j.1399-5618.2009.00759.x>.
- Voss, H. U., & Schiff, N. D. (2009). MRI of neuronal network structure, function, and plasticity. *Progress in Brain Research*, 175, 483–496. Available from [https://doi.org/10.1016/S0079-6123\(09\)17532-5](https://doi.org/10.1016/S0079-6123(09)17532-5).
- Wada, A. (2009a). GSK-3 inhibitors and insulin receptor signaling in health, disease, and therapeutics. *Front Biosci (Landmark Ed)*, 14, 1558–1570. Available from <https://doi.org/10.2741/3324>.
- Wada, A. (2009b). Lithium and neuropsychiatric therapeutics: neuroplasticity via glycogen synthase kinase-3beta, beta-catenin, and neurotrophin cascades. *Journal of pharmacological sciences*, 110(1), 14–28. Available from <https://doi.org/10.1254/jphs.09r02cr>.
- Wang, F., Jackowski, M., Kalmar, J. H., Chepenik, L. G., Tie, K., Qiu, M., . . . Blumberg, H. P. (2008). Abnormal anterior cingulum integrity in bipolar disorder determined through diffusion tensor imaging. *The British Journal of Psychiatry*, 193(2), 126–129. Available from <https://doi.org/10.1192/bjp.bp.107.048793>.
- Wang, H., Huang, S., Yan, K., Fang, X., Abussaud, A., Martinez, A., . . . Feng, Z. P. (2016). Tideglusib, a chemical inhibitor of GSK3beta, attenuates hypoxic-ischemic brain injury in neonatal mice. *Biochimica et Biophysica Acta*, 1860(10), 2076–2085. Available from <https://doi.org/10.1016/j.bbagen.2016.06.027>.
- Wang, Y., Gao, Y., Tang, S., Lu, L., Zhang, L., Bu, X., . . . Huang, X. (2020). Large-scale network dysfunction in the acute state compared to the remitted state of bipolar disorder: A meta-analysis of resting-state functional connectivity. *EBioMedicine*, 54, 102742. Available from <https://doi.org/10.1016/j.ebiom.2020.102742>.
- Won, E., & Kim, Y. K. (2017). An Oldie but Goodie: Lithium in the Treatment of Bipolar Disorder through Neuroprotective and Neurotrophic Mechanisms. *International Journal of Molecular Sciences*, 18(12). Available from <https://doi.org/10.3390/ijms18122679>.
- Youle, R. J., & Strasser, A. (2008). The BCL-2 protein family: opposing activities that mediate cell death. *Nature Reviews Molecular Cell Biology*, 9(1), 47–59. Available from <https://doi.org/10.1038/nrm2308>.
- Yuan, P. X., Huang, L. D., Jiang, Y. M., Gutkind, J. S., Manji, H. K., & Chen, G. (2001). The mood stabilizer valproic acid activates mitogen-activated protein kinases and promotes neurite growth. *The Journal of Biological Chemistry*, 276(34), 31674–31683. Available from <https://doi.org/10.1074/jbc.M104309200>.
- Yucel, K., McKinnon, M. C., Taylor, V. H., Macdonald, K., Alda, M., Young, L. T., & MacQueen, G. M. (2007). Bilateral hippocampal volume increases after long-term lithium treatment in patients with bipolar disorder: a longitudinal MRI study. *Psychopharmacology (Berl)*, 195(3), 357–367. Available from <https://doi.org/10.1007/s00213-007-0906-9>.
- Zovetti, N., Rossetti, M. G., Perlini, C., Maggioni, E., Bontempi, P., Bellani, M., & Brambilla, P. (2020). Default mode network activity in bipolar disorder. *Epidemiol Psychiatr Sci*, 29, e166. Available from <https://doi.org/10.1017/S2045796020000803>.
- Zuo, X. N., Di Martino, A., Kelly, C., Shehzad, Z. E., Gee, D. G., Klein, D. F., . . . Milham, M. P. (2010). The oscillating brain: complex and reliable. *NeuroImage*, 49(2), 1432–1445. Available from <https://doi.org/10.1016/j.neuroimage.2009.09.037>.