

List of contributors

- Brandon Bacalzo** Department of Psychiatry and Iowa Neuroscience Institute, University of Iowa Health Care, Iowa City, IA, United States
- Izabela G. Barbosa** Izabela Departamento of Mental Health, School of Medicine, Federal University of Minas Gerais (UFMG), Belo Horizonte, Brazil
- Moises E. Bauer** Moises Laboratory of Immunobiology, School of Health and Life Sciences, Pontifical Catholic University of Rio Grande do Sul (PUCRS), Porto Alegre, Brazil
- Kyle J. Burghardt** Eugene Applebaum College of Pharmacy and Health Sciences, Wayne State University, Detroit, MI, United States
- Mehmet Akif Camkurt** Center of Excellence on Mood Disorders, Faillace Department of Psychiatry and Behavioral Sciences, McGovern Medical School, The University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States
- Deniz Ceylan** Department of Psychiatry, Faculty of Medicine, Koç University, Istanbul, Turkey
- Audrey C. Chong** Department of Psychiatry and Behavioral Sciences, University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States; McGovern Medical School, University of Texas Health Science Center, Houston, TX, United States
- Anthony J. Cleare** Centre for Affective Disorders, Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, United Kingdom
- Gustavo C. Dal-Pont** Translational Psychiatry Laboratory, Graduate Program in Health Sciences, University of Southern Santa Catarina (UNESC), Criciúma, SC, Brazil
- Amanda Decker** Department of Psychiatry and Iowa Neuroscience Institute, University of Iowa Health Care, Iowa City, IA, United States
- Alexandre Paim Diaz** Center of Excellence on Mood Disorders, Faillace Department of Psychiatry and Behavioral Sciences, McGovern Medical School, The University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States; Translational Psychiatry Program, Faillace Department of Psychiatry and Behavioral Sciences, McGovern Medical School, The University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States

Breno Satler Diniz Department of Psychiatry, University of Connecticut Health Center, CT, United States

Brisa S. Fernandes Center of Excellence on Mood Disorders, Faillace Department of Psychiatry and Behavioral Sciences, McGovern Medical School, The University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States; iMPACT, The Institute for Mental and Physical Health and Clinical Translation, School of Medicine, Barwon Health, Deakin University, Geelong, VIC, Australia; Translational Psychiatry Program, Faillace Department of Psychiatry and Behavioral Sciences, McGovern Medical School, The University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States

Gabriel R. Fries Translational Psychiatry Program, Faillace Department of Psychiatry and Behavioral Sciences, McGovern Medical School, The University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States; Neuroscience Program, MD Anderson Cancer Center UTHealth Graduate School of Biomedical Sciences, Houston, TX, United States; Center for Precision Health, School of Biomedical Informatics, The University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States

Fernando S. Goes Department of Psychiatry and Behavioral Sciences, Johns Hopkins School of Medicine, Baltimore, MD, United States

Iria Grande Bipolar and Depressive Disorders Unit, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM, Barcelona, Spain

Bradley H. Howlett Eugene Applebaum College of Pharmacy and Health Sciences, Wayne State University, Detroit, MI, United States

Fabrice Jollant Health Faculty, Department of Psychiatry, Université de Paris, France; GHU Paris Psychiatrie et Neuroscience, Sainte-Anne Hospital, Paris, France; Department of Psychiatry, Academic Hospital (CHU) Nîmes, France; Department of Psychiatry, McGill University and McGill Group for Suicide Studies, Douglas Hospital Research Centre, Verdun, QC, Canada; Department of Psychiatry and Psychotherapy, Jena University Hospital, Jena, Germany; Moods Team, INSERM UMR-1178, CESP, Le Kremlin-Bicêtre, France

Gregory Jones Department of Psychiatry and Behavioral Sciences, University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States

Mario F. Juruena Centre for Affective Disorders, Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, United Kingdom

Marissa A. Kassa Eugene Applebaum College of Pharmacy and Health Sciences, Wayne State University, Detroit, MI, United States

Tadafumi Kato Department of Psychiatry & Behavioral Science, Juntendo University Graduate School of Medicine, Tokyo, Japan

Camila Nayane de Carvalho Lima Translational Psychiatry Program, Faillace Department of Psychiatry and Behavioral Sciences, McGovern Medical School, The University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States

- Rodrigo Machado-Vieira** Experimental Therapeutics and Molecular Pathophysiology Program, Louis Faillace, M.D Department of Psychiatry and Behavioral Sciences, McGovern Medical School, UTHealth Science Center, Houston, TX, United States
- Naguib Mechawar** Department of Psychiatry, McGill University and McGill Group for Suicide Studies, Douglas Hospital Research Centre, Verdun, QC, Canada
- Gustavo C. Medeiros** Department of Psychiatry and Behavioral Sciences, Johns Hopkins School of Medicine, Baltimore, MD, United States
- Kathleen Ries Merikangas** Genetic Epidemiology Research Branch, Intramural Research Program, National Institute of Mental Health, Bethesda, Maryland, United States
- Jose Alberto Moreno** Bipolar and Depressive Disorders Unit, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM, Barcelona, Spain; Department of Psychiatry, Hospital de la Vega Baja, Orihuela, Spain
- Ather Muneer** Department of Psychiatry, Rawal Institute of Health Sciences, Islamabad, Pakistan
- Benson Mwangi** Center of Excellence on Mood Disorders, Faillace Department of Psychiatry and Behavioral Sciences, McGovern Medical School, The University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States
- Tanya T. Nguyen** Department of Psychiatry, University of California San Diego, San Diego, CA, United States; Sam and Rose Stein Institute for Research on Aging, University of California San Diego, San Diego, CA, United States; Veterans Affairs San Diego Healthcare System, San Diego, CA, United States
- Mark J. Niciu** Department of Psychiatry and Iowa Neuroscience Institute, University of Iowa Health Care, Iowa City, IA, United States
- Ayşegül Özerdem** Department of Psychiatry and Psychology, Mayo Clinic, Rochester, MN, United States
- Chi-Un Pae** Department of Psychiatry, College of Medicine, The Catholic University of Korea, Seoul, South Korea
- Ives Cavalcante Passos** Laboratory of Molecular Psychiatry, Centro de Pesquisa Experimental (CPE) e Centro de Pesquisa Clínica (CPC), Hospital de Clínicas de Porto Alegre (HCPA), Porto Alegre, Brazil; Faculty of Medicine, Graduate Program in Psychiatry and Behavioral Sciences, Department of Psychiatry, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil
- João Quevedo** Center of Excellence on Mood Disorders, Faillace Department of Psychiatry and Behavioral Sciences, McGovern Medical School, The University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States; Translational Psychiatry Program, Faillace Department of Psychiatry and Behavioral Sciences, McGovern Medical School, The University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States; Neuroscience Graduate Program, The University of Texas MD Anderson Cancer Center UTHealth Graduate School of Biomedical Sciences, Houston, TX, United States; Translational Psychiatry Laboratory, Graduate Program in Health Sciences, University of Southern Santa Catarina (UNESC), Criciúma, Brazil

Carola Rong Department of Psychiatry and Behavioral Sciences, University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States

Thiago Henrique Roza Laboratory of Molecular Psychiatry, Centro de Pesquisa Experimental (CPE) e Centro de Pesquisa Clínica (CPC), Hospital de Clínicas de Porto Alegre (HCPA), Porto Alegre, Brazil; Faculty of Medicine, Graduate Program in Psychiatry and Behavioral Sciences, Department of Psychiatry, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil

Ana C. Ruiz Department of Psychiatry and Behavioral Sciences, University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States

Janusz K. Rybakowski Department of Adult Psychiatry, Poznan University of Medical Sciences, Poznan, Poland

Estela Salagre Bipolar and Depressive Disorders Unit, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM, Barcelona, Spain

Thyago Antonelli Salgado Laboratory of Molecular Psychiatry, Centro de Pesquisa Experimental (CPE) e Centro de Pesquisa Clínica (CPC), Hospital de Clínicas de Porto Alegre (HCPA), Porto Alegre, Brazil; Faculty of Medicine, Graduate Program in Psychiatry and Behavioral Sciences, Department of Psychiatry, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil

Marsal Sanches Center of Excellence on Mood Disorders, Faillace Department of Psychiatry and Behavioral Sciences, McGovern Medical School, The University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States

Lokesh Shahani Center of Excellence on Mood Disorders, Department of Psychiatry and Behavioral Sciences, McGovern Medical School, The University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States

Jair C. Soares UT Center of Excellence on Mood Disorders, Louis Faillace, M.D. Department of Psychiatry and Behavioral Sciences, McGovern Medical School, UTHealth Science Center, Houston, TX, United States

Ananya Swaminathan Genetic Epidemiology Research Branch, Intramural Research Program, National Institute of Mental Health, Bethesda, Maryland, United States; Department of Biomedical Engineering, Johns Hopkins University, Baltimore, MD, United States

Antonio L. Teixeira Antonio Department of Psychiatry and Behavioral Sciences, McGovern Medical School, University of Texas Health Science Center, Houston. Faculdade Santa Casa BH, Belo Horizonte, Brazil

Bárbara Tietbohl-Santos Laboratory of Molecular Psychiatry, Centro de Pesquisa Experimental (CPE) e Centro de Pesquisa Clínica (CPC), Hospital de Clínicas de Porto Alegre (HCPA), Porto Alegre, Brazil; Faculty of Medicine, Graduate Program in Psychiatry and Behavioral Sciences, Department of Psychiatry, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil

Massimo Tusconi Department of Medical Sciences and Public Health, Unit of Psychiatry, University of Cagliari, Cagliari, Italy

Eleonora D. Vaccariello Department of Psychiatry, University of California San Diego, San Diego, CA, United States; Department of Psychology and Neuroscience, Maastricht University, Maastricht, The Netherlands

Samira S. Valvassori Translational Psychiatry Laboratory, Graduate Program in Health Sciences, University of Southern Santa Catarina (UNESC), Criciúma, SC, Brazil

Roger B. Varela Queensland Brain Institute, Queensland University, St Lucia, QLD, Australia

Courtney M. Vecera Department of Psychiatry and Behavioral Sciences, University of Texas Health Science Center at Houston (UTHealth), Houston, TX, United States

Eduard Vieta Bipolar and Depressive Disorders Unit, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM, Barcelona, Spain

Sheng-Min Wang Department of Psychiatry, College of Medicine, The Catholic University of Korea, Seoul, South Korea

Allan H. Young Centre for Affective Disorders, Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, United Kingdom