From Published Evidence to Analyzable Biomedical Cohorts with Scientific Agents

Author details pending confirmation
Affiliation pending confirmation

Abstract

Researchers may need paired biomedical measurements that are available only as published group summaries or regression coefficients. We study whether a scientific agent can use these findings to construct complete virtual records for a specified population. The researcher declares the population, variables, analysis and permitted evidence. The agent writes a stochastic program, specifying how reported statistics constrain the model and how it will generate relationships the studies leave open. Numerical tools sample complete tables from the program. In two glycemia cases, supplying external associations reduced the mean absolute discrepancy for five of six regression coefficients. In an iron-status study, two agents used the same source threshold table, which implies an odds ratio of 6.65. Their different assumptions for the target population implied odds ratios of 4.65 and 6.65, compared with 17.79 in the target sample. One generator came closer on the proportion with both biomarkers low, the other on their continuous rank correlation. Both cases in an earlier prospective evaluation fell short of their prespecified overall criterion. The frozen models make it possible to trace an analysis result to a published finding or an explicit assumption and examine how much its interpretation depends on that choice.

CCS Concepts

• **Computing methodologies** → **Modeling and simulation**; • **Applied computing** → *Life and medical sciences.*

Keywords

Synthetic cohorts, biomedical evidence, scientific agents, epidemiology

1 Introduction

Suppose a researcher wants to study iron status among young adult women. She can find a paper giving ferritin concentrations within hemoglobin bands and counts of women jointly classified by both thresholds, but she cannot access the paired measurements [33]. She wants to know how often both values are low, how strongly low ferritin is associated with low hemoglobin, and how the two concentrations vary together over their full ranges. With the original measurements, she could answer all three questions on the same rows. With a published table, she can recover the first two quantities for the source sample; she cannot recover the third exactly, or assume that any of them will carry over unchanged to another population.

This is a common use for synthetic biomedical data: the records needed for an analysis are inaccessible, incomplete, or were never measured together. The researcher may have no individual records at all. Studies nevertheless report means, subgroup frequencies and regression coefficients that constrain a possible cohort. The challenge is deciding what each result constrains. A crude odds ratio for raw ferritin and an odds ratio for inflammation-corrected ferritin describe different measurements [33,34]. Even two regression coefficients measured in the same units need not belong in the same multivariable equation. A system that treats these findings as interchangeable can generate plausible-looking records that answer the wrong question.

The researcher specifies the population, variables and analyses and what information the agent may use. Depending on the study, the agent receives a reviewed evidence packet or searches for publications. It authors a joint probability model and the code to sample from it. A numerical worker calls this program with a seed and a row count. Researchers can use the resulting table in ordinary analysis software and inspect the program to see what supplied the associations in the table. The research questions are visible during construction. The study protocol separately determines when the measured reference may be accessed.

We organize the experiments around three questions. **RQ1:** When the agent receives external relationship evidence in addition to marginal information, which target regression estimates improve? **RQ2:** Which analyses can the same continuous virtual cohort support, including analyses not numerically specified by a source? **RQ3:** How does that pattern change across biomedical requests, and what survives when measured reference results are first revealed after generation? These questions separate agreement with an input statistic, the usefulness of a completed joint model, and transfer to a new reference.

2 Related work and problem definition

2.1 From available information to analyzable records

Filshtein et al. share fitted study models so a recipient can simulate complete cohorts for epidemiological analyses [1]. SynthIPD uses published survival curves and trial summaries to reconstruct time-to-event records with covariates [2]. SynDI combines internal individual data with external regression information across populations to fit a target regression [29]. Multilevel network meta-regression likewise relates aggregate evidence to individual-level models under population assumptions [3]. These approaches differ in what the recipient knows. Filshtein et al. supply the comparison for our HRS experiment because the published exports and the scientific analysis can be matched. In our external-evidence experiments, the agent must also choose the joint dependence structure needed to perform the requested analyses.

G-Sim combines LLM-authored simulators with numerical calibration and structural refinement [4]. LLMSynthor uses aggregate discrepancies to guide proposals for new records and evaluates mobility, e-commerce and household population synthesis [5]. SynthCraft studies healthcare synthetic-data workflows [6], and SPADA samples from a dependency graph fitted with observed tabular data

[15]. Our methodological question concerns the preceding biomedical decision: how findings with different populations, scales and adjustment sets constrain one generator, and which analyses the resulting data support.

2.2 Evidence-conditioned cohort construction

A request specifies the target population, eligibility and measurements, desired variables, permitted observations and intended analyses. Each evidence record identifies its source population, measurement scale, statistical operator, adjustment set, estimate, available uncertainty and provenance. The agent returns an executable generator, the numerical uses of that evidence and the assumptions needed to complete the model:

```
(generator, evidence uses, completion assumptions)

    = agent(scientific request, evidence, optional
     observations)

rng = RNG(seed)

virtual data = generator(row count, rng, context)

scientific results = requested analyses(virtual data).
```

When individual records are unavailable, the program generates every requested variable. For enrichment, it conditions on supplied observations to generate an additional variable. For partial completion, it can preserve observed entries and fill the designated missing cells. Our principal experiments evaluate complete-cohort generation. In the iron case the population definition limits eligibility to nonpregnant women aged 20–44. The request calls for two continuous output columns, hemoglobin in g/dL and raw ferritin in µg/L.

A finding may constrain the law numerically, provide a check on its implications, or remain contextual. Algebraic consequences of accepted findings are derived evidence. Unreported dependence and transfer from a source population to the request are completion assumptions. Several laws can satisfy the available information while implying different conditional results, so the output includes both the table and its construction specification.

We evaluate usefulness at the level of scientific analysis. For an adjusted coefficient, correlation or dispersion statistic, the generated estimate is compared with the relevant sample reference in its native units. Reproducing a supplied summary measures reconstruction; agreement with a target result not supplied to the author addresses a different question. A table can support several analyses with different degrees of evidential support. These distinctions guide both construction and evaluation.

3 Agent-controlled evidence-to-cohort construction

3.1 Specify the request and qualify the evidence

The investigator supplies the target population, eligibility, requested columns and analyses, the observations available and any restrictions on sources. In Q1 and the iron study, researchers prepared the evidence packets. In Q4, each author chose searches and original papers through Europe PMC/PMC, within a budget of eight query credits and four original articles. Q5 supplied reviewed articles and permitted additional search. Each author constructed its own generator. The number of authors in an experiment counts separate constructions.

For each finding, the construction records the study population, definition and units of the measurement, the statistic and adjustment set, and its numerical role. For example, a crude threshold association for raw ferritin and one using inflammation-corrected ferritin are different quantities. In the iron study, the packet already marked the Cuban result as contextual. The agents then decided how to model continuous measurements within the hemoglobin bands and how much of the source dependence to carry into the US target. Table 1 identifies these separate contributions.

3.2 Translate findings into numerical model constraints

The agent writes both the sampling program and a record linking its choices to the publications. A reported value can set a model parameter directly, enter a calculation that determines several parameters, or serve as a check. The record identifies assumptions needed to complete the model, including dependence between variables or transfer across populations. A reviewer can inspect where a number enters the program or why it was used only as context.

In a glycemia task the packet explicitly identified the BMI and glucose coefficients reported by Jansen et al. as separate age- and sex-adjusted projections [9]. The constructor still needed coefficients for a single joint response model. For residualized BMI and glucose with SDs `sB` and `sG` and correlation `r`, the joint coefficients `thetaB` and `thetaG` satisfy

```
cB = thetaB + thetaG * r * sG/sB

cG = thetaG + thetaB * r * sB/sG.
```

CHNS E1 constructor A01, selected by ordinal identifier, used reported slopes `cB = 0.13` and `cG = 2.29`, residual SDs 4.07611 and 0.49694, and the declared assumption `r = 0.25`. Solving the equations gave 0.0642173 and 2.1583154 in IFCC response units. Multiplication by 0.09148 yielded 0.00587460 and 0.19744269 in HbA1c percentage-point units [11]. The source projections therefore informed the joint law through their mathematical relationship, with residual dependence and population transfer remaining explicit assumptions. The worked trace connects the derivation to frozen code, an unfitted correlation check and the resulting target BMI discrepancy.

3.3 Complete an executable joint law

The author chooses the probability family, sampling order, parameters and dependence assumptions needed for the requested columns. It writes a program that takes a row count and a random number generator. Where the protocol provides a population context, the program can use different rules for the source and target. In the iron case both implementations return a table of paired hemoglobin




Figure 1: A completed evidence-to-cohort construction. The agent converts the Hainan male glucose–HbA1c Spearman coefficient 0.44 into latent Gaussian correlation 0.457. Selected entries of the sex-stratified copula matrix distinguish the derived glucose–HbA1c relation from the assumed BMI–glucose value 0.25. Standard-normal noise, Cholesky sampling and shifted-lognormal transforms generate BMI, glucose and HbA1c; age has a bounded-integer transform. Table cells are schematic. The scientific law, assumptions, execution checks and requested regressions form one inspectable path. This worked case is separate from the information-contrast and cross-task experiments below.

Table 1: From an iron-status request to two continuous columns. Both constructors received the same task and sources. Each delivered three tables from its own frozen law.

Who or what	What it supplies in this case
Investigator	US eligibility, two biomarkers, three intended analyses and an audited evidence packet. The packet identifies the Cuban corrected-ferritin result as contextual.
Source study	Counts jointly classified by low hemoglobin and low raw ferritin, plus ferritin geometric means within three hemoglobin bands [33].
Agent A	Authors the continuous sampling law. For transfer it chooses a relative weight of 2/3 for low ferritin, then an independent hemoglobin draw for 20% of rows.
Agent B	Authors a copula linking ranks within each band and carries its source law into the target population.
Numerical worker and evaluator	Run each program at the prescribed seeds. Analyze the same two columns for the threshold odds ratio, joint proportion and rank correlation.

and ferritin measurements. The threshold indicators and rank correlation are computed later from those two columns.

Figure 1 presents a completed construction using a sex-stratified Gaussian copula and positive biomarker margins. The agent converted a reported glucose–HbA1c Spearman relationship into a latent Gaussian correlation, declared the missing BMI–glucose dependence and implemented sampling and measurement transforms. Other requests led to bounded conditional laws, gamma innovations, lognormal mixtures or a recursive longitudinal family. The scientific contribution lies in choosing and connecting these components to the evidence, rather than introducing a new distribution family.

3.4 Execute and inspect the resulting analyses

The numerical worker runs the program in isolation. Where source replay is available, it draws from the source context and applies the published statistic to those rows. The author can retain or revise the model within the protocol budget. In the iron study each author received one source-only review, using 100,000 source draws at seed 4101. A revised the upper tail of ferritin to obtain finite variance. B

Table 2: What the constructor could know, and what the evaluation can establish. Q1–Q5 are the identifiers of the frozen experiments. The iron study and the earlier prospective tests have separate protocols. All listed tasks generate complete cohorts.

Study	Construction inputs	Constructions and final deliveries	Analysis
Q1: CHNS and ENSANUT	Basic evidence (E0); E0 plus external relations (E1)	20 agent + 4 prescribed; 72 tables of 3,000	Six adjusted glycemia coefficients
Iron status	External evidence; target eligibility; no target biomarkers	2 agent; 6 tables of 5,000	Threshold association, joint burden and continuous ranks
Q5: glycemia and lipids	Reviewed external papers, population description, optional retrieval	6 agent; 18 tables of 3,000	Six requested effects or correlations
Earlier prospective wave	External summaries and population description; exact target results withheld	5 agent + 1 prescribed per case; 36 tables of 5,000	HAALSI glucose and Turkana lipid analyses
Q3: longitudinal reconstruction	Published HRS fitted-model artifacts	1 agent + 1 reconstructed; 6 valid tables, 11,509 subjects × 4 visits each	Age–HbA1c interactions in memory trajectories
Q2 / Q4: supporting conditions	Q2: with or without 30 target summaries; Q4: open paper retrieval without bibliographic seeds	Q2: 6 agent laws / 18 distinct tables; Q4: 4 / 12	Reconstruction of supplied summaries; repeated open retrieval

retained its program. In Q5 all 132 proposed source comparisons were unavailable under their declared definitions. The authors could still inspect summaries of their own generated outputs.

For the iron target, the worker ran each final program at seeds 5101–5103 for 5,000 rows per seed. The six output tables were sealed before the target biomarkers were accessed. The evaluator applied the three requested analyses to each table and separately computed the survey-weighted reference. The stored program, seeds and analysis code specify how to repeat this numerical stage. Asking a fresh agent to interpret the same evidence may produce another model. The supplement records the construction inputs and the limits of the surviving model-service metadata.

4 Evaluation design

For each request we fixed the scientific analysis and applied it to the generated records and the appropriate reference (Table 2). A construction is a separately authored probability model. A seed samples that fixed model and measures variation across its draws. We report effects in their scientific units and preserve each study's original decision rule.

CHNS (6,631 adults, ages 35–74) uses unweighted estimates for the selected sample. ENSANUT (6,931 adults, ages 25–79) assigns equal mass to eight sex-by-age cells. In Q1, each case and information condition has five fresh agent authors and one prescribed statistical law. We average absolute discrepancy over three draws within each law, then across laws within the arm. E0 and E1 authors are not paired. The prescribed law illustrates a statistical completion from the same information; it is not a competitive search over statistical methods.

The reference conditions differ. Q1–Q5 reused project-level references where applicable. For iron status we sealed the protocol and the six deliveries before acquiring the target biomarkers; NHANES was already known to the project. In the earlier prospective wave, a custodian withheld the exact HAALSI and Turkana results, although investigators knew related Turkana findings. We distinguish target sampling intervals, resampling stability bands and model-conditional intervals in the figures. None makes a sample estimate the true population value.

5 Results

5.1 RQ1: What does adding external relationship evidence change?

Across CHNS and ENSANUT, the agent arms given external relationships (E1) had smaller mean absolute discrepancies than the basic-evidence arms (E0) for five of six coefficients (Figure 2). The clearest change was the CHNS glucose slope: error fell from 0.134019 to 0.024079 HbA1c percentage points per mmol/L. In contrast, CHNS BMI error rose from 0.020380 to 0.031557 percentage points per kg/m². All three ENSANUT discrepancies decreased. The same-information statistical construction was closer on some E1 endpoints, including CHNS glucose. The result identifies analyses for which additional evidence helped in these cases; it does not establish that agent authors outperform statistical generators.

5.2 RQ2: Which uses survive in the same iron-status cohort?

Zhu et al. studied 1,671 parous women who used IUDs in northern China, two to four years postpartum [33]. The source counts imply a crude odds ratio of 6.65. The agents' source replays gave 6.58 and 6.73. Both programs used the hemoglobin bands and their ferritin summaries. The Cuban study used inflammation-corrected ferritin [34]. As specified in the shared packet, its odds ratio supplied context for the transfer assumptions (Table 1).

Figure 3 traces the implications of the frozen code. In A, giving low-ferritin observations a relative weight of two thirds preserves the threshold odds ratio, since the factor cancels in the cross-product of counts. It reduces the joint-low proportion from 10.35% to 7.67%. For one fifth of rows A then draws hemoglobin independently from the same target hemoglobin distribution. The resulting model implies a joint-low proportion of 6.73% and an odds ratio of 4.65. B




Figure 2: Relational evidence and scientific analysis agreement. CHNS panels show BMI adjusted for age/sex (a), glucose adjusted for age/sex (b), and age adjusted for sex/BMI/glucose (c). ENSANUT panels show age adjusted for sex/BMI/glucose (d), BMI adjusted for age/sex (e), and BMI additionally adjusted for glucose (f). All six analyses retain native units. E0 contains basic/marginal evidence; E1 adds relations. Each agent arm has five independently authored laws and each statistical arm one prescribed law, with all three final draws retained: 24 constructions, 72 tables and 216 coefficient estimates. Dots show seeds, open diamonds constructor means and thin segments seed ranges; dashed lines mark sample references. Reference bands are CHNS community-cluster CR1-t intervals or ENSANUT cluster-resampling stability bands. Author indices do not pair E0 with E1. Table S1 gives every endpoint's mean absolute discrepancy and E1–E0 contrast.

retains the source joint-low proportion of 10.35% and odds ratio of 6.65. These values follow from the two authored models. The colored markers in panels c to e show estimates from the six previously sealed tables.

The survey reference included 979 eligible women with both measurements [35]. Comparison on the log-odds-ratio scale was primary. All six delivered odds ratios (A 4.835–4.866, B 6.011–6.565) fell below the survey's lower confidence limit of 9.28 (estimate 17.79, 95% CI 9.28–34.11). A came closer on the joint-low proportion (6.66–6.88% versus 7.63%), while B came closer on the continuous Spearman correlation (0.391–0.430 versus 0.398). For B the joint-low proportion was 10.02–10.44%, and for A the correlation was 0.156–0.195. The sources reported no continuous rank coefficient. Their ferritin summaries by hemoglobin band constrain ranks indirectly, leaving within-band dependence to the author. The same two columns in each delivery supplied all three estimates.

5.3 RQ3: Which findings carry across requests and reference conditions?

For glycemia and lipids, three authors per request generated 18 tables in total. All 54 primary estimates agreed in point direction with the corresponding sample estimates across six distinct analyses (Figure 4). The tables supported age- and sex-adjusted BMI models and further adjustment for glucose. The three constructor-mean correlations of total and HDL cholesterol after adjustment for age, sex and BMI were 0.108–0.141, against 0.512 in the measured sample. The total-cholesterol slope's reference interval crossed zero. A post hoc check of the sealed lipid tables found HDL above total cholesterol in 69 of 27,000 synthetic rows (0.256%). Both concentrations were expressed in mmol/L. This check measures the biological consistency of individual pairs separately from the prespecified association results.




Figure 3: How two transfer choices affect analyses on the same iron-status records. Panels a and b are a post hoc calculation from the source counts and frozen rules. Panels c to e show three sealed tables of 5,000 rows per law and the NHANES estimate with its design-based 95% interval. Low hemoglobin is below 12 g/dL and low raw ferritin below 15 µg/L. The two secondary analyses had no prespecified equivalence margin.

The earlier prospective wave tested a stricter question. In HAALSI adults aged 40 or older [30,31], all 15 agent draws recovered the positive directions of the adjusted BMI and age associations with capillary glucose; only eight recovered the negative difference between female and male BMI slopes. The rule required four of five laws per case to meet all prespecified margins on every seed. Three passed in HAALSI and none in Turkana [32], so the planned expansion stopped. Even a zero-effect rule met the HAALSI margin for all three endpoints. The earlier method and its gate remain a separate prospective result, not a fresh evaluation of the later workflow on the same Turkana participants.

5.4 What changes when the scientific relationship is already supplied?

In the HRS longitudinal benchmark both methods received the published model exports [1]. Below the 6.5% HbA1c knot, the age–HbA1c interaction in memory trajectories averaged −0.04464 for agent tables and −0.04050 for the reconstructed published generator, against the supplied −0.04684. Above the knot the mean signs differed; five of six intervals included zero (Figure S6). This is a matched reconstruction check, not transfer from partial evidence. Similarly, supplying 30 target summaries in Q2 allowed joint reconstruction of the five highlighted regression and correlation analyses (Table S2). In Q4, open retrieval yielded twelve distinct original studies across four authors and twelve complete tables (Tables S3–S4); the design did not isolate the effect of retrieval. HRS and Q2 test reconstruction when target-model information is supplied; Q4 instead demonstrates a way to obtain external evidence without supplying more information about the target.

6 Discussion

A researcher with published threshold counts can reconstruct a source association directly. A virtual table adds a way to run analyses on continuous measurements and to inspect the assumptions that connect them. In the iron case, the odds ratio can be traced through each agent's transfer decision, while the rank correlation depends partly on how the model pairs values within a hemoglobin band. A researcher deciding whether to use the table for that correlation would benefit from paired measurements within those bands in a comparable population. Paired lipid measurements under the specified adjustment would likewise address the dependence left uncertain in the lipid case. Across these studies the frozen records let us explain particular outcomes after comparison with the reference. Establishing which analyses can be trusted before seeing a reference would require a separate evaluation.

7 Limitations and Ethical Considerations

The evidence cannot identify every joint relationship. Sources differ in eligibility, age, geography, specimen handling and adjustment; reference samples have sampling uncertainty. A small number of




Figure 4: Analysis-specific behavior across glycemia and lipid requests. HbA1c panels in the left column show age adjusted for sex/BMI/glucose (a), BMI adjusted for age/sex (c), and BMI additionally adjusted for glucose (e). The right column shows age/sex-adjusted HDL and TC BMI slopes (b,d) and residual Pearson correlation after age/sex/BMI adjustment (f). All six laws and 54 estimates from 18 tables are retained. Circles show seeds, open diamonds means and thin segments seed ranges; dark diamonds and dashed lines mark sample references. HbA1c units are percentage points per 10 years or 5 kg/m²; lipid slopes are mg/dL per 5 kg/m². Bands are ENSANUT empirical cluster-resampling stability bands and lipid HC3 assumption-conditional sensitivity intervals. The TC band crosses zero; no residual-correlation interval is available. Seed ranges describe sampling under each frozen law. Table S5 supplies all numerical summaries and Figure S3 the dispersion results.

agent constructions share a model service, and repeated draws do not represent independent clinical cohorts. The protocols differ across studies. No held-out trial shows that review improves outcomes or that the workflow saves a measured amount of human time. For research planning, the source-based and assumed parts should travel with the records. Values must be labeled synthetic and must not be presented as measured individual biomarkers, clinical diagnoses or substitutes for independent patient evidence. Original cohort access and reuse conditions continue to apply; the public artifacts contain code and aggregate results rather than participant-level tables.

8 Generative AI Usage

Language models participated in research planning and manuscript editing and were an object of the experiments. In the scientific

workflow, agent authors interpreted evidence, selected or received studies according to protocol, proposed statistical models, and wrote or revised generator code. Deterministic tools sampled the records and computed the reported analyses; separate implementations checked the numerical results where specified. Study protocols and source records document these roles. The final author list, study-specific declarations, and applicable oversight and data-use statements must be confirmed before submission.

Supporting materials and reproducibility

The public research repository contains protocols, executable code and aggregate audits. The iron protocol seal and delivery seal precede target evaluation. Supplements retain all prespecified outcomes, unfavorable cases and implementation incidents. No new cohort was generated for this revision.

References

(1) Filshtein TJ, Li X, Zimmerman SC, Ackley SF, Glymour MM, Power MC. Proof of concept example for use of simulation to allow data pooling despite privacy restrictions. *Epidemiology*. 2021;32(5):638–647. DOI and full text.

(2) Zhao Z et al. SynthIPD: training-free synthetic individual patient data generation. arXiv:2509.16466v2, 24 July 2026. Versioned full text. Publication beyond this preprint was not established in the positioning audit.

(3) Phillippo DM et al. Multilevel network meta-regression for population-adjusted treatment comparisons. *Journal of the Royal Statistical Society: Series A*. 2020;183(3):1189–1210. DOI/full text.

(4) Holt S, Ruiz Luyten M, Berthon A, van der Schaar M. G-Sim: Generative Simulations with Large Language Models and Gradient-Free Calibration. *Proceedings of ICML*. PMLR 267;2025:23513–23561. Published record.

(5) Tang Y, Kong M, He J, Nie T, Ma W, Sun L. LLMSynthor: Macro-Aligned Micro-Records Synthesis with Large Language Models. KDD 2026. DOI; full text v4.

(6) Callender T, Boyd A, Davis R, Ruhrberg Estevez S, Lavista Ferres JM, van der Schaar M. SynthCraft: An AI partner for synthetic data generation to support data access and augmentation in healthcare. *PLOS Digital Health*. 2026;5(3):e0001290. Published article.

(7) Bao Y, Ma X, Li H, Zhou M, Hu C, Wu H, Tang J, Hou X, Xiang K, Jia W. Glycated haemoglobin A1c for diagnosing diabetes in Chinese population: cross sectional epidemiological survey. *BMJ*. 2010;340:c2249. DOI; bibliographic record.

(8) Wang P, Li Y, Wang M, Song L, Dong Y, Han X, Tuomilehto J, Wang Y, Du Y, Qiu C. Comparing glycemic traits in defining diabetes among rural Chinese older adults. *PLOS ONE*. 2024;19(1):e0296694. Published article.

(9) Jansen H, Stolk RP, Nolte IM, Kema IP, Wolffenbuttel BHR, Snieder H. Determinants of HbA1c in nondiabetic Dutch adults: genetic loci and clinical and lifestyle parameters, and their interactions in the lifelines cohort study. *Journal of Internal Medicine*. 2013;273(3):283–293. Published article.

(10) Pani LN, Korenda L, Meigs JB, Driver C, Chamany S, Fox CS, Sullivan L, D'Agostino RB, Nathan DM. Effect of aging on A1C levels in individuals without diabetes: evidence from the Framingham Offspring Study and the National Health and Nutrition Examination Survey 2001-2004. *Diabetes Care*. 2008;31(10):1991–1996. DOI; bibliographic record.

(11) NGSP. The IFCC and NGSP, Table 2. Official measurement-scale guidance.

(12) Li T, Quan H, Zhang H, Lin L, Lin L, Ou Q, Chen K. Type 2 diabetes is more predictable in women than men by multiple anthropometric and biochemical measures. *Scientific Reports*. 2021;11:6062. DOI; original full text.

(13) Sakane N, Sato J, Tsushita K, Tsujii S, Kotani K, Tominaga M, Kawazu S, Sato Y, Usui T, Kamae I, Yoshida T, Kiyohara Y, Sato S, Tsuzaki K, Nirengi S, Takahashi K, Kuzuya H, JDPP Research Group. Determinants of Glycated Hemoglobin in Subjects With Impaired Glucose Tolerance: Subanalysis of the Japan Diabetes Prevention Program. *Journal of Clinical Medicine Research*. 2017;9(4):360–365. DOI; original full text.

(14) Wisgerhof W, Ruijgrok C, den Braver NR, Borgonjen—van den Berg KJ, van der Heijden AAWA, Elders PJM, Beulens JWJ, Alssema M. Phenotypic and lifestyle determinants of HbA1c in the general population–The Hoorn Study. *PLOS ONE*. 2020;15(6):e0233769. DOI; original full text.

(15) Yang S, Zhang Z, Prenkaj B, Kasneci G. Doubling Your Data in Minutes: Ultra-fast Tabular Data Generation via LLM-Induced Dependency Graphs. *Proceedings of the 2025 Conference on Empirical Methods in Natural Language Processing*. 2025:10337–10358. Official article and PDF; DOI.

(16) Wu et al. Effect of age on the diagnostic efficiency of HbA1c for diabetes in a Chinese middle-aged and elderly population: The Shanghai Changfeng Study. *PLoS ONE*. 2017. Admitted original.

(17) Skogberg et al. The association between anthropometric measures and glycated haemoglobin (HbA1c) is different in Russian, Somali and Kurdish origin migrants compared with the general population in Finland: a cross-sectional population-based study. *BMC Public Health*. 2019. Admitted original.

(18) Zhang et al. The Association of Retinopathy and Plasma Glucose and HbA1c: A Validation of Diabetes Diagnostic Criteria in a Chinese Population. *Journal of Diabetes Research*. 2016. Admitted original.

(19) Venkataraman et al. Ethnicity modifies the relation between fasting plasma glucose and HbA1c in Indians, Malays and Chinese. *Diabetic Medicine*. 2012. Admitted original.

(20) Liu et al. Ideal glycated hemoglobin cut-off points for screening diabetes and prediabetes in a Chinese population. *Journal of Diabetes Investigation*. 2016. Admitted original.

(21) Huang et al. Hemoglobin A1c Levels Associated with Age and Gender in Taiwanese Adults without Prior Diagnosis with Diabetes. *International Journal of Environmental Research and Public Health*. 2021. Admitted original.

(22) Ghazanfari et al. A Comparison of HbA1c and Fasting Blood Sugar Tests in General Population. *International Journal of Preventive Medicine*. 2010. Admitted original.

(23) Masuch et al. Preventing misdiagnosis of diabetes in the elderly: age-dependent HbA1c reference intervals derived from two population-based study cohorts. *BMC Endocrine Disorders*. 2019. Admitted original.

(24) Marini et al. Association between hemoglobin glycation index with insulin resistance and carotid atherosclerosis in non-diabetic individuals. *PLoS ONE*. 2017. Admitted original.

(25) Greiner et al. Associations between low HDL, sex and cardiovascular risk markers are substantially different in sub-Saharan Africa and the UK: analysis of four population studies. *BMJ Global Health*. 2021;6:e005222. DOI; published correction to author names.

(26) Obirikorang et al. Obesity and Cardio-Metabolic Risk Factors in an Urban and Rural Population in the Ashanti Region-Ghana: A Comparative Cross-Sectional Study. *PLOS ONE*. 2015;10:e0129494. DOI; admitted original.

(27) Agongo et al. The burden of dyslipidaemia and factors associated with lipid levels among adults in rural northern Ghana: An AWI-Gen sub-study. *PLOS ONE*. 2018;13:e0206326. DOI; admitted original.

(28) Agongo et al. Correction: The burden of dyslipidaemia and factors associated with lipid levels among adults in rural northern Ghana: An AWI-Gen sub-study. *PLOS ONE*. 2019;14:e0213233. DOI; correction. The correction concerns hip/waist entries and a citation; those entries were not used in the lipid constructions.

(29) Gu T, Taylor JMG, Mukherjee B. A synthetic data integration framework to leverage external summary-level information from heterogeneous populations. *Biometrics*. 2023;79:3831–3845. DOI.

(30) Gómez-Olivé FX et al. Cohort Profile: Health and Ageing in Africa: A Longitudinal Study of an INDEPTH Community in South Africa (HAALSI). *International Journal of Epidemiology*. 2018;47:689–690j. Cohort profile.

(31) Harvard Center for Population and Development Studies. HAALSI Baseline Survey. Harvard Dataverse, 2016, version 3.0. Dataset.

(32) Lea AJ, Martins D, Kamau J, Gurven M, Ayroles JF. Urbanization and market integration have strong, nonlinear effects on cardiometabolic health in the Turkana. *Science Advances*. 2020;6:eabb1430. Article. Dataset.

(33) Zhu JH et al. Iron, folate, and B12 deficiencies and their associations with anemia among women of childbearing age in a rural area in northern China. International Journal for Vitamin and Nutrition Research. 2010;80:144–154. Article.

(34) Pita-Rodríguez GM et al. Prevalence of Anemia and Iron Deficiency in Women of Reproductive Age in Cuba and Associated Factors. International Journal of Environmental Research and Public Health. 2023;20:5110. Article.

(35) National Center for Health Statistics. NHANES, August 2021–August 2023: Ferritin (FERTIN_L); Complete Blood Count (CBC_L); Demographic Variables and Sample Weights (DEMO_L). Ferritin and phlebotomy weights; complete blood count; demographics.

Supplementary results

The supplement retains the complete numerical comparisons and supporting displays under their original study identities. Figure and table numbers follow their new placement; no construction, primary result or unavailable comparison is removed by the shorter main narrative. Q1–Q5 remain distinct experiments with their original information conditions.

S1. Complete external-evidence contrast

All 24 Q1 constructions produced their three scheduled final tables. The 72 final tables were distinct and supported every requested regression and diagnostic; seven exploratory tables brought physical executions to 79. Additional relational evidence reduced constructor-averaged mean absolute discrepancy for five of the six designated coefficients (Table S1). CHNS glucose discrepancy decreased from 0.134019 to 0.024079 HbA1c percentage points per mmol/L. All three ENSANUT contrasts improved. CHNS BMI discrepancy increased from 0.020380 to 0.031557.

The effect displays explain what these changes mean. CHNS E1 glucose constructor means were 0.363363–0.414189 around the 0.402093 reference. Its BMI means were positive but lower, 0.012400–0.045757 versus 0.054946. Conditional-age means ranged from −0.000509 to 0.007393 versus 0.008107; one author retained a negative mean and two negative seeds. All ENSANUT E1 focal coefficients were positive, although its age means of 0.097769–0.102452 per decade exceeded 0.075132 and its BMI means remained below their references. Preserving a direction and recovering its magnitude are distinct aspects of usefulness.

Every E1 author used at least one relationship as a numerical constraint. The complete uptake records include constraints, unfitted analytic checks and contextual uses, including the projection calculation in Section 3.2. Statistical references also achieved useful results: under E1, the prescribed constructions had lower seed-averaged discrepancies than every corresponding agent for CHNS glucose and ENSANUT conditional BMI. Other comparisons favored agent constructions. The evidence therefore supports scientific utility and the value of additional information, rather than a universal algorithmic ranking.

S2. Complete supplied-summary reconstruction

The separate CHNS experiment gave three authors external literature (L) and three authors that literature plus 30 target summaries and their metadata (R). Each author delivered three tables and had one additional diagnostic-review opportunity. All six retained their models. There were 42 physical executions: six trials and 36 initial/final deliveries. Final bytes duplicated their matching initial outputs, leaving 24 distinct tables overall and 18 distinct delivery tables. The identical stages are displayed once.

R supported close reconstruction of all five highlighted scientific analyses (Table S2; Figure S1). These values were legitimate construction inputs; their agreement measures reconstruction rather than prediction of hidden coefficients. The L/R packet and authors differ from Q1, so those two information comparisons are kept separate.

Agreement extended beyond central values. R constructor-mean BMI SDs were 3.38965–3.40243 versus 3.40632, glucose SDs were 1.48817–1.58968 versus 1.54455, and HbA1c SDs were 0.95208–0.96554 versus 0.95357. A separate three-table statistical calibration supplement also closely reconstructed the regression summaries, with mean coefficients 0.055726, 0.412568 and 0.007679. That purpose-built family was developed with coding-agent assistance and is not an agent-removal ablation. Both approaches demonstrated the usefulness of constructing a joint table from accessible aggregate information.

S3. Open retrieval across population requests

The unified workflow allocated two fresh scientific authors to each of the existing CHNS and ENSANUT requests. Each author received the intended analyses, population/measurement definitions, task interface and retrieval budget, but no target coefficients, desired signs, individual records or prior construction. CHNS- and ENSANUT-derived studies were excluded as generating evidence. Across 24 query credits, the authors selected sixteen paper uses representing twelve distinct originals. Independent review checked the original statistical definitions, numerical derivations, source applicability and required finite moments before execution. Sources could overlap across authors; four constructions do not imply four independent bodies of clinical evidence.

The agents chose materially different laws. C01 used an age-conditioned multivariate-lognormal mixture informed by Chinese population summaries [7,16] and contextual age/body-size findings [10,17]. C02 combined copula predictors with an affine HbA1c conditional mean using Pinggu summaries and Singapore coefficients [18,19], with other populations as applicability context [14,20]. E01 used bounded beta predictors and a conditional gamma response [10,17,21,22]. E02 combined lognormal BMI, conditional truncated-normal glucose and a positive conditional-mean lognormal response [10,21,23,24]. Their unreported dependencies, population transfers and residual choices remained declared assumptions. The shared controller did not prescribe these families or insert target coefficients.

Each author completed one 3,000-row trial, inspected generated-only diagnostics and source feedback, then selected its final model before three fixed draws. All four retained the initial law. The sixteen physical executions comprised four trials and twelve unique final tables, or 36,000 final virtual rows. Feedback contained 29 qualified approximate comparisons and 39 unavailable checks across the four trials, not source-population replays or 29 passed constraints. After all sessions closed, the evaluator reused the fixed sample references. A separate numerical implementation reproduced all 366 final coefficient/diagnostic values; no clinical data were downloaded or refitted.

All 36 primary seed estimates had the same positive directions as their corresponding references (Figure S2). These are repeated measurements of six analysis definitions across four laws, not 36 independent confirmations. Magnitude recovery varied by analysis. CHNS C01/C02 means were 0.03992/0.05338 for BMI against 0.05495, 0.37754/0.49241 for glucose against 0.40209, and 0.00311/0.00563 for age against 0.00811. Units are respectively HbA1c percentage points per kg/m², mmol/L and year. C02 was closer for BMI and age, while C01 was closer for glucose; no construction dominated all analyses.

Table S1: Complete main-study information contrast. Discrepancies retain the units of each row; they are not pooled. The reference response is HbA1c percentage points. E1–E0 uses unrounded quantities before display rounding.

Case / focal analysis	Predictor increment	E0 mean absolute discrepancy	E1 mean absolute discrepancy	E1 − E0
CHNS BMI, adjusted age/sex	1 kg/m²	0.020380	0.031557	+0.011178
CHNS glucose, adjusted age/sex	1 mmol/L	0.134019	0.024079	−0.109939
CHNS age, adjusted sex/BMI/glucose	1 year	0.007281	0.004854	−0.002426
ENSANUT age, adjusted sex/BMI/glucose	10 years	0.063944	0.024670	−0.039274
ENSANUT BMI, adjusted age/sex	5 kg/m²	0.056883	0.023732	−0.033151
ENSANUT BMI, adjusted age/sex/glucose	5 kg/m²	0.048797	0.025649	−0.023147




Figure S1: Supplied summaries become jointly analyzable data. Five panels show BMI, glucose and age regression coefficients plus glucose–HbA1c Pearson and Spearman correlations. L01–L03 use literature; R01–R03 additionally receive 30 target summaries and metadata. Points show all three final draws, diamonds constructor means and segments seed ranges. Dashed lines mark reference summaries available to R; regression bands are CHNS community-cluster CR1-t intervals, with no supplied correlation interval. All six constructors, 18 distinct delivery tables, 30 means and 90 seed statistics appear once. Initial and final deliveries were identical after retention; Table S2 gives the complete five-analysis numerical comparison.

Table S2: Joint-statistic reconstruction with available summaries. Ranges span three constructor means over three fixed seeds, not confidence intervals. Regression units are HbA1c percentage points per kg/m², mmol/L and year, respectively.

Analysis	Supplied reference	External literature L	Literature + summaries R
BMI coefficient, age/sex adjusted	0.054946	0.013001–0.034287	0.052821–0.055789
Glucose coefficient, age/sex adjusted	0.402093	0.356064–0.487815	0.396941–0.413715
Age coefficient, sex/BMI/glucose adjusted	0.008107	0.003552–0.006910	0.007569–0.007947
Glucose–HbA1c Pearson correlation	0.660968	0.633764–0.813170	0.650561–0.666438
Glucose–HbA1c Spearman correlation	0.361457	0.501339–0.566477	0.357571–0.360983




Figure S2: Repeated open-retrieval construction across two biomedical requests. All six prespecified primary analyses are shown for four agent-authored laws and twelve final 3,000-row virtual tables. Blue and purple identify the two constructions within each request. Open circles show seeds 8101–8103, ordered top to bottom within each author row; vertical spacing is solely for legibility. Diamonds and horizontal segments below the circles show the three-seed mean and minimum–maximum range. Coral squares in separate reference rows show previously exposed selected-sample estimates. Solid coral intervals are CHNS community-cluster CR1-t 95% confidence intervals; dashed coral intervals are ENSANUT empirical cluster-resampling stability bands from 399 attempts. These reference intervals have different meanings and neither is an acceptance or equivalence threshold. Axes retain native effect units and zero; pp denotes HbA1c percentage points. All generated coefficients are positive, while both CHNS age estimates are weaker and both ENSANUT age estimates stronger than their references. Seed ranges describe sampling under each fixed law, not confidence intervals or uncertainty over evidence, model choice or transport. The four constructions provide developmental workflow replication on two existing references, not independent clinical replication.

For ENSANUT, E01/E02 means were 0.10308/0.09772 for adjusted age against 0.07513 pp/decade, 0.09165/0.07140 for age/sex-adjusted BMI against 0.08364 pp per 5 kg/m², and 0.07509/0.04733 after additionally adjusting for glucose against 0.05664. Every draw retained the reduction of the BMI coefficient after glucose adjustment, an association pattern rather than causal mediation. Both authors built an age coefficient near 0.10 pp/decade and shared some sources; their agreement does not establish independent evidence convergence.

The complete diagnostics reveal remaining limits. CHNS C01 glucose–HbA1c Pearson correlation was approximately 0.666 versus 0.661, but Spearman correlation was 0.635 versus 0.361. ENSANUT residual HbA1c RMS was approximately 0.295/0.324 versus 0.488, and total SD was 0.362/0.406 versus 0.527. Correct directions and a close Pearson coefficient therefore coexisted with rank and dispersion mismatches. The complete replication report, every diagnostic and Supplementary Table S3 retain the full results. A scalar-type interface defect in the original evaluator was corrected additively on unchanged frozen tables, with its failed output and independent audit preserved; no scientific model was adjusted after comparison.

CHNS BMI/glucose analyses adjust for age and sex; CHNS age adjusts for sex, BMI and glucose. ENSANUT age adjusts for sex, BMI and glucose; its BMI analyses adjust for age/sex, with or without glucose. All Q4 diagnostic values remain in the complete diagnostic table.

The reviewed evidence/specification files distinguish reported statistics, algebraic derivations and assumed relations. The public experiment record includes per-source eligibility, measurement, adjustment, overlap and reporting-quality notes; full third-party originals and participant-like rows remain local.

S4. Longitudinal reconstruction from published artifacts

The HRS recipient-stage example [1] tested a different data product. A fresh agent received published model exports, a variable specification and the intended mixed-model analysis. It authored the recursive conditional generating family with parameter provenance. A separate implementation reconstructed the published law. Each generated 11,509 subjects at 0, 2, 4 and 6 years, producing 46,036 rows per table. No HRS individual records were used, and the source analysis coefficients were construction inputs.

All three agent and three corrected-law tables supported the native R/lme4 model with 16 fixed effects, individual intercepts/slopes and piecewise HbA1c terms around 6.5%, with age centered at 60 and scaled per decade. The below-threshold HbA1c-by-age interaction averaged −0.044635 in agent data, with range −0.051059 to −0.038247, against the supplied −0.046837. The reconstructed law averaged −0.040500. Both preserved its negative direction in all three draws (Figure S6).

Figure S6 shows the estimates and their per-table Wald intervals. The longitudinal analysis record gives all 16 fixed effects and the comparator correction.

The weaker above-threshold interaction was less stable: agent estimates averaged 0.026485 against 0.012867, while corrected-law estimates averaged −0.011691. Overall agent HbA1c mean/SD was 5.8725/0.9676 versus the original published 5.9/1.0. The high-HbA1c subgroup had mean/SD 7.1515/0.5630, compared with original 7.7/1.4 and published synthetic 7.2/0.6. Thus useful source-model behavior coexisted with inadequate upper-group spread.

A CRLF-sensitive parser initially overwrote seven coefficient/SE pairs in our comparator preparation. A separately reviewed source-bound repair produced the three corrected comparator tables without changing the frozen agent outputs. The full record contains nine physical generations, six valid comparison tables and three invalid initial comparator attempts. This was our implementation defect, not a failure of the published method. The incident record and complete effects retain the correction and all outcomes.

S5. Cross-task construction, dispersion and source uses

A post hoc biological consistency check reports the HDL and total cholesterol ordering counts by frozen law and seed. The comparison uses generated pairs in their original common units.

The next study applied the common workflow to HbA1c analyses in the ENSANUT2018 analytical design and BMI-related TC/HDL analyses in the Turkana study context. Three fresh agent contexts constructed a law for each request, using reviewed external articles, population/measurement metadata and optional literature tools. No participant inputs, target coefficients, target means or prior generators were supplied. The six contexts used the same model service and shared some source evidence. This study therefore tests repeated construction across two requests, with the law chosen by each author, rather than independent clinical replication or a controlled comparison of workflow components.

The HbA1c analyses retained the existing age and BMI estimands and equal allocation to eight sex-by-age cells. The lipid analyses used unweighted age/sex-adjusted BMI coefficients for TC and HDL, plus the Pearson correlation of their residuals after age, sex and BMI adjustment. Lipid eligibility required adults with recognized recorded sex and finite, positive BMI and lipid values. Generated lipid concentrations were converted from mmol/L to mg/dL using the frozen factor 38.67. The reference represents a selected Turkana/Laikipia study sample; fasting, assay transfer and population composition remained qualified in the evidence record. No new participant records were read, downloaded or refitted for this study.

The agents made different scientific choices within the common interface. H01 combined bounded predictors, conditionally truncated glucose and a shifted gamma HbA1c innovation; H02 used bounded beta conditional laws. H03 solved cross-source moment equations under declared dependence and dispersion assumptions, then used conditional beta laws. Their source records included overlapping age, body-size and glycemia evidence [10,17,21,22], with additional check-only/contextual evidence for H01 [14]. The lipid authors combined Gaussian copulas and lognormal margins using geometric summaries from Greiner et al. [25], rural Ashanti rank correlations [26] and arithmetic moments from Agongo et al. [27,28]. Choices about arithmetic versus geometric location, transfer of coefficients of variation, sex strata and age dependence differed among L01–L03. These are applications of established probability families whose numerical use was selected and explained by the agent.

Table S3: Complete Q4 open-retrieval primary summaries. Means and ranges summarize three final seeds under each fixed construction. Units are HbA1c percentage points per stated predictor increment. No row is omitted; ranges are not confidence intervals.

Request / analysis	Author	Increment	Reference	Generated mean	Seed min–max
CHNS BMI	C01	1 kg/m²	0.054946	0.039923	0.034342–0.045708
CHNS glucose	C01	1 mmol/L	0.402093	0.377538	0.357560–0.398892
CHNS adjusted age	C01	1 year	0.008107	0.003109	0.002461–0.003755
CHNS BMI	C02	1 kg/m²	0.054946	0.053375	0.047088–0.059055
CHNS glucose	C02	1 mmol/L	0.402093	0.492412	0.488649–0.494875
CHNS adjusted age	C02	1 year	0.008107	0.005628	0.005137–0.006301
ENSANUT adjusted age	E01	10 years	0.075132	0.103084	0.101546–0.104315
ENSANUT BMI	E01	5 kg/m²	0.083643	0.091651	0.089940–0.094413
ENSANUT glucose-adjusted BMI	E01	5 kg/m²	0.056638	0.075089	0.072005–0.076661
ENSANUT adjusted age	E02	10 years	0.075132	0.097724	0.094947–0.099135
ENSANUT BMI	E02	5 kg/m²	0.083643	0.071399	0.071232–0.071563
ENSANUT glucose-adjusted BMI	E02	5 kg/m²	0.056638	0.047325	0.045022–0.048592

Table S4: Original-source inventory for Q4 open retrieval. Twelve unique originals supplied sixteen author-specific paper uses. Sharing a source is not independent clinical evidence; source populations are not the target populations.

Original	Reference	Constructing authors
Wu et al., 2017	[16]; PMC5591004	C01
Bao et al., 2010	[7]; PMC2871988	C01
Pani et al., 2008	[10]; PMC2551641	C01, E01, E02
Skogberg et al., 2019	[17]; PMC6458679	C01, E01
Zhang et al., 2016	[18]; PMC5078665	C02
Venkataraman et al., 2012	[19]; PMC3504343	C02
Liu et al., 2016	[20]; PMC5009131	C02
Wisgerhof et al., 2020	[14]; PMC7272077	C02
Huang et al., 2021	[21]; PMC8038122	E01, E02
Ghazanfari et al., 2010	[22]; PMC3075530	E01
Masuch et al., 2019	[23]; PMC6371438	E02
Marini et al., 2017	[24]; PMC5398507	E02

Independent source and implementation reviews checked the statistical meanings, specification/code consistency and finite moments before isolated execution. Each constructor completed one scheduled trial, inspected generated diagnostics and source feedback, and could revise once. All six retained their initial laws. Three fixed final seeds, 9101–9103, produced 18 distinct 3,000-row tables: 54,000 fictional rows from 24 successful physical attempts, including the six trials. No final execution failed and no seed was replaced. Reference comparison began after all six author sessions closed and reused previously exposed aggregate results.

All 54 primary point estimates had the same direction as their corresponding sample estimates, with quantitative agreement varying by analysis (Figure 4; Table S5). HbA1c age effects averaged 0.09954–0.10091 percentage points per decade versus 0.07513 in the reference; age/sex-adjusted BMI effects averaged 0.04892–0.06030 versus 0.08364 per 5 kg/m². With glucose additionally adjusted, H03 averaged 0.05098 versus 0.05664, whereas H01 and H02 averaged 0.03047 and 0.01936. Thus the shared age direction and attenuation after glucose adjustment did not imply recovery of each effect magnitude.

For lipids, mean HDL BMI coefficients ranged from −2.1865 to −1.9868 mg/dL per 5 kg/m² versus −4.6230 in the reference, while TC coefficients ranged from 10.0151 to 11.2609 versus 5.6831. The TC reference sensitivity interval was −3.7535 to 15.1197, so matching its positive point estimate does not establish replication of a definite positive population effect. Residual TC–HDL correlations averaged 0.10828–0.14134 versus 0.51207. The generators preserved the directions of these focal point estimates while substantially underrepresenting the lipid dependence remaining after the specified adjustment.

Dispersion changes the assessment of usefulness (Figure S3). H03's closer glucose-adjusted BMI coefficient accompanied lower HbA1c marginal SD and full-model residual RMS, 0.3523 and 0.3006 percentage points versus 0.5274 and 0.4885. H02 retained more variation, with corresponding values 0.4825 and 0.4036, but smaller BMI coefficients. L02's TC SD was close to the reference, 43.6778 versus 42.8466 mg/dL, although its residual TC–HDL correlation remained weak. All three lipid laws had HDL SDs below the reference. No construction uniformly dominated across focal effects, dependence and dispersion.

The evaluation retained all 513 prespecified estimates and diagnostics, including the 54 primary outputs. Independent recomputation verified every value and all 324 matched reference comparisons; the maximum numerical difference was 5.33×10^{-14}. The remaining 189 diagnostics lacked matching cached reference estimators and remained generated-only results. Across authors, eight original studies supplied 22 author–paper uses, including context/check-only uses; optional retrieval consumed seven query credits and one original download. All 132 declared source-check entries were unavailable under the implemented semantic rules because weighting or essential operator definitions were unresolved or unsupported (Table S6). Numerical source values still informed reviewed parameter derivations, but this batch provides neither measured numerical source-fit feedback nor a before/after revision benefit. The complete results and reviews retain every endpoint, unavailable comparison and implementation incident, including correction of the independent auditor's identifier parser without changing any generated table.

HbA1c age adjusts for sex, BMI and glucose. HbA1c BMI adjusts for age/sex, with glucose added in the third analysis. Lipid BMI slopes adjust for age/sex; the residual correlation removes age/sex/BMI projections from both lipids on the same rows. H01–H03 belong only to this Q5 HbA1c request, and L01–L03 only to this Q5 lipid request; the L labels do not refer to the earlier Q2 literature arm. All 54 individual primary outputs and all 513 prespecified estimates/diagnostics remain in the complete endpoint table. The 324 matched reference comparisons and 189 generated-only diagnostics retain their original availability labels.

The evidence-use inventory and source/code review retain the source IDs, numerical roles, unresolved definitions and actual retrieval activity. Unresolved weighting and unsupported or incompletely specified operators were not replaced with different statistics. Parameter derivations can therefore use reviewed source information while the corresponding runtime source comparison remains unavailable.

S6. Dependence sensitivity and the supporting construction

The preceding open-search example acquired five seriously reviewed studies [9,10,12,13,14] without bibliographic seeds or desired signs. A fresh constructor selected a sex-stratified Gaussian copula with positive biomarker margins and generated one trial plus three complete 3,000-person tables. It retained the model after eleven approximate and 23 unavailable source comparisons per execution. Secondary BMI coefficients were 0.04217–0.05176 versus 0.05495 in the reused CHNS reference, secondary age was 0.01613–0.01771 versus 0.01530, and primary age was 0.00842–0.01084 versus 0.00811. Primary BMI/glucose were attenuated, and the female point direction and distributional tails remained imperfect. Supplementary Figure S4 retains all seven nonintercept effects and their uncertainty labels.

Before generating data or comparing targets, the source review identified an unreported BMI–glucose dependence. Two valid latent correlations, 0.10 and 0.40, implied conditional BMI coefficients of +0.03422 and −0.01095 while retaining the same margins, other bivariate laws and secondary projection. The delivered model used the declared assumption 0.25. This is an analytical sensitivity, not additional generated cohorts or a causal intervention. It demonstrates why an explicit data-generating law can identify which missing relationship matters to an intended use.

A separate analysis contextualized the same three tables with 100 community-disjoint resplits of the existing CHNS sample, drawing 3,000 people per side and retaining all 36 metrics. Secondary BMI and age discrepancies were on scales comparable with within-sample variation, while primary glucose/female and several joint/tail discrepancies were larger. The complete table preserves the nonexchangeable real-real and generated comparisons. These are descriptive interpretation aids, not a new population or retrospectively selected equivalence cutoff.

S7. Prospective evaluation under the earlier method

In HAALSI, eligible ages started at 40 years and the analysis used the released capillary point-of-care glucose measurement. Fasting was not an eligibility requirement. The fixed BMI and age contrasts were taken from the same additive glucose model with recorded sex. A second model estimated the difference in the BMI slope between women and men, adjusting for age. In Turkana, the two BMI slopes were adjusted for age and recorded sex. The correlation uses the residuals after adjusting both lipid measurements for age, sex and BMI.

Each case had five separately authored agent laws and one prespecified statistical generator (S01). The experiment produced 48 source-context and 36 target-context tables of 5,000 rows each. All authors retained their initial laws after source-only feedback. Initial and final labels therefore refer to the same 36 target tables. The public checkpoint preceding disclosure records work in progress. The final evaluation record documents the complete inventory, one reveal and independent recomputation of 6,704 checks. HAALSI had a prospectively withheld target. Turkana's exact estimates were withheld, with prior investigator exposure to adjacent findings. Later Q5 reused the same 397-person Turkana reference.

The primary rule divided the absolute difference in each slope by the measured outcome's sample SD and used the absolute correlation difference directly. A construction passed when all nine comparisons (three analyses by three seeds) were at most 0.10. The agent counts at thresholds 0.05, 0.10, 0.20 and 0.30 were 1, 3, 5 and 5 of five for HAALSI and 0, 0, 1 and 4 for Turkana. Both S01 laws missed the primary criterion. For HAALSI, the discrepancies for three zero-valued effects were 0.0990, 0.0733 and 0.0770. The frozen decision closed the expansion. The prospective study predates the later interface for documenting numerical evidence use




Figure S3: Dispersion across all cross-task constructions. Panels show marginal HbA1c SD, HbA1c full-model residual RMS, and marginal TC and HDL SDs for each construction and its three final draws, retaining all 36 diagnostic values from the same 18 tables. HbA1c uses the population-SD diagnostic; lipids use sample SD. These quantities measure individual or residual spread, not uncertainty in a regression coefficient. H03's relatively close adjusted BMI effect in Figure 4 coexists with reduced HbA1c spread, and L02's close TC SD coexists with weak adjusted lipid dependence. Seed ranges reflect Monte Carlo variation conditional on the selected law. Gray HbA1c bands are empirical cluster-resampling stability bands; no reference interval is supplied for lipid SDs.

and supplies a separate evaluation of the underlying construction approach.

S8. Full records and retained implementation incidents

The public repository provides original code, protocols, aggregate results and independent checks. The complete v0.9 extended record preserves all six original numerical tables and full methods/results. Its Table 2 supplies every main endpoint's adjustment set, units and reference interval; Tables 3–4 retain the main and summary-assisted values; Table 5 retains all latest seven effects; Table 6 retains the real-real comparison values. These archived tables preserve the earlier analyses alongside the separately sealed iron study.

The main study, summary-assisted study, statistical supplement, HRS example, preceding open-search construction, unified workflow replication and cross-task evidence-use study preserve separate frozen identities and complete endpoint outputs. The cross-task protocol, complete endpoint table, independent numerical review and scientific interpretation document the new unit. Earlier live A/B results, including the conditioned BMI mismatch and source-replay unavailability, remain in the extended record. Figure updates and added tables consume reviewed aggregates only; manuscript integration creates no cohorts or clinical fits. In the main study, seven authors inspected exploratory outputs and kept their laws, and thirteen finalized without exploration. The two earlier live authors both kept their models after the available review.

Individual real/generated tables, third-party originals and credentials remain local; clinical reruns require lawful access and the appropriate environment. Historical model identifiers and token costs are incompletely metered. Author/affiliation/funding/conflict/ethics/access/AI declarations and final submission formatting remain to be completed. No manuscript has been submitted.

The historical supplement retains earlier enrichment and direct-row construction, the divergent-moment repair, failed transport criteria, tail defects and revision trade-offs. The v0.16 reading draft preserves the earlier layout.

Table S5: Complete Q5 cross-task primary summaries. Each row summarizes three fixed final draws from one law. All 18 author–analysis combinations are retained. HbA1c entries are percentage points (pp); lipid slopes are mg/dL per 5 kg/m². Ranges are seed minima and maxima, not confidence intervals or evidence/model uncertainty.

Request / analysis	Author	Units / increment	Reference	Generated mean	Seed min–max
HbA1c adjusted age	H01	pp / 10 years	0.075132	0.099545	0.094075–0.103606
HbA1c BMI	H01	pp / 5 kg/m²	0.083643	0.060295	0.055769–0.063938
HbA1c glucose-adjusted BMI	H01	pp / 5 kg/m²	0.056638	0.030469	0.027956–0.034291
HbA1c adjusted age	H02	pp / 10 years	0.075132	0.099539	0.098162–0.100396
HbA1c BMI	H02	pp / 5 kg/m²	0.083643	0.048923	0.029090–0.065499
HbA1c glucose-adjusted BMI	H02	pp / 5 kg/m²	0.056638	0.019363	0.002308–0.030777
HbA1c adjusted age	H03	pp / 10 years	0.075132	0.100912	0.096356–0.105015
HbA1c BMI	H03	pp / 5 kg/m²	0.083643	0.056649	0.048449–0.063758
HbA1c glucose-adjusted BMI	H03	pp / 5 kg/m²	0.056638	0.050978	0.044994–0.056901
HDL BMI	L01	mg/dL / 5 kg/m²	−4.623004	−2.186526	−2.286805–−2.118624
TC BMI	L01	mg/dL / 5 kg/m²	5.683108	10.393977	9.992186–10.676486
TC–HDL residual correlation	L01	dimensionless	0.512070	0.111234	0.108517–0.113140
HDL BMI	L02	mg/dL / 5 kg/m²	−4.623004	−2.100390	−2.194806–−2.035712
TC BMI	L02	mg/dL / 5 kg/m²	5.683108	11.260864	10.811415–11.565687
TC–HDL residual correlation	L02	dimensionless	0.512070	0.108283	0.105395–0.110194
HDL BMI	L03	mg/dL / 5 kg/m²	−4.623004	−1.986781	−2.122835–−1.913541
TC BMI	L03	mg/dL / 5 kg/m²	5.683108	10.015112	8.617345–11.698752
TC–HDL residual correlation	L03	dimensionless	0.512070	0.141343	0.121087–0.155368

Table S6: Q5 original-source uses and unavailable feedback. Eight distinct originals supplied 22 author–paper uses, including context/check-only evidence. All six authors retained their initial laws. The 132 source-check entries were unavailable, not successful numerical comparisons or failed generators. The inventory counts the three lipid originals, with the Agongo correction [28] attached to its source rather than counted as a separate original study.

Author	Original-source references	Paper uses	Unavailable source checks
H01	Pani [10]; Wisgerhof [14]; Skogberg [17]; Huang [21]; Ghazanfari [22]	5	13
H02	Pani [10]; Skogberg [17]; Huang [21]; Ghazanfari [22]	4	18
H03	Pani [10]; Skogberg [17]; Huang [21]; Ghazanfari [22]	4	32
L01	Greiner [25]; Obirikorang [26]; Agongo [27,28]	3	29
L02	Greiner [25]; Obirikorang [26]; Agongo [27,28]	3	19
L03	Greiner [25]; Obirikorang [26]; Agongo [27,28]	3	21
Total	8 distinct original studies	22	132

S9. Iron-status analysis and independent verification

All six deliveries used two continuous columns, with three target analyses per delivery. The target survey comparison used the phlebotomy weight and 15 design degrees of freedom; 979 of 1,065 eligible positive-weight women had both measurements. A separate implementation reproduced all 21 point estimates (three reference, eighteen generated), the six target interval endpoints and the four unweighted cell counts. The study report gives all seed values and




Figure S4: Analysis-specific usefulness and an unresolved joint dependence. Panels (a–d) show all primary-model nonintercept effects and (e–g) all secondary-model effects. Blue points are three fixed 3,000-person draws with conditional 95% HC3-t intervals. Gray dashed lines and bands are exposed CHNS reference estimates and 95% community-cluster CR1-t intervals; solid lines mark zero. All effects retain native units (pp: HbA1c percentage points), with shared limits for matching predictors. Generated intervals omit source, structural and transport uncertainty; overlap is not equivalence. Panel (h) displays two pre-generation analytical scenarios: latent BMI–glucose correlations 0.10 and 0.40 imply conditional BMI coefficients +0.03422 and −0.01095 while leaving marginal laws, other bivariate laws and the secondary projection unchanged. The delivered model used 0.25. These are not new draws, a fitted curve or a causal intervention. Retained initial/final stages alias the same three tables.

Table S7: Six prespecified analyses under the earlier workflow. The columns report the measured-sample estimate, the minimum and maximum across 15 agent draws, the number agreeing in point sign, and the range of the three draws of the prescribed statistical law. The ranges describe draws under the candidate models. HAALSI glucose contrasts are in mmol/L, lipid slopes in mg/dL and the adjusted correlation is unitless.

Analysis	Measured	Agent range	Same sign	S01 range
HAALSI glucose, BMI per 5 kg/m²	0.3138	0.3285 to 0.5698	15/15	−0.0174 to 0.0180
HAALSI glucose, age per 10 years	0.2322	0.1136 to 0.3005	15/15	−0.0170 to 0.0077
HAALSI female-minus-male BMI slope per 5 kg/m²	−0.2440	−0.2487 to 0.1903	8/15	−0.0524 to 0.0046
Turkana HDL, BMI per 5 kg/m²	−4.6230	−3.5509 to −1.8173	15/15	−0.1120 to 0.0710
Turkana TC, BMI per 5 kg/m²	5.6831	10.2822 to 15.0413	15/15	−0.4657 to 0.0334
Turkana residual TC–HDL correlation	0.5121	0.1162 to 0.5361	15/15	0.3625 to 0.3840




Figure S5: Earlier prospective evaluation of whole-cohort generation. Blue circles show the three draws of each agent law and purple squares the three draws of the prescribed statistical law (S01). Short segments span draws. Dashed lines mark measured-sample estimates, and shaded bands give approximate 95% HC3 sensitivity intervals for the five slopes. The residual correlation has no reference interval. HAALSI glucose main effects were directionally stable across agent draws. The sex difference in the BMI slope varied by construction. Turkana's total cholesterol point estimate had a reference interval that included zero. Its quantitative estimates were withheld before generation despite prior exposure to related published findings.

the evaluation protocol specifies the source replay, primary log-odds-ratio comparison and secondary analyses.

S10. Reproducing a program and tracing an agent decision

In the iron task each fresh author received the same target description, source packet and clarification about a source summary. Only external source materials were allowed. The protocol specified numpy and pandas, a supplied random generator, the separate source and target contexts, and one source-only retain-or-revise opportunity. The task, evidence packet, A's final program and B's final program document the handoff. Run the programs with the recorded seeds and worker. The published code and aggregates support computational verification. Historical author-model snapshots and per-construction token counts are incompletely recorded, which limits reproduction of the original model-authoring sessions.

For the analytic part of Figure 3, let the four source counts be 173 (both low), 85 (low hemoglobin only), 331 (low ferritin only) and 1,082 (neither). Weight the first and third cells by two thirds and renormalize. Under A the final joint-low probability equals 0.8 times the reweighted joint-low probability plus 0.2 times the product of the two reweighted marginal probabilities. B keeps the original probabilities. This calculation describes the selected models and can be reproduced from published counts without sampling. The figure retains the original sealed deliveries and reference analysis alongside these model implications.




Figure S6: Two prespecified age–HbA1c interactions in a longitudinal memory model. Diamonds show the supplied coefficients; circles and squares show estimates from three draws per law (seeds 6101–6103, top to bottom). Bars show approximate 95% Wald intervals calculated from the exported source standard errors or from the generated-table fits. Units are standardized memory-score change per decade per percentage point of HbA1c within the indicated branch. Each table contains 11,509 synthetic subjects and four visits per subject. The upper-branch signs differed between constructions. The six tables describe sampling under two frozen generating laws.