

Derive marginalised probability

Ultimately want to maximise $P(\theta | \{\log L_i\})$. Bayes' rule to invert in terms of known quantities

$$P(\{\log L_i\} | \theta) = \frac{P(\{\log L_i\} | \theta) \times P(\theta)}{P(\{\log L_i\})} \quad (1)$$

Assume an uninformative constant prior $P(\theta)$, and disregard the total evidence $P(\{\log L_i\})$ which is independent of θ . The remaining quantity is the evidence for θ , which is calculated by marginalising over \mathbf{X}

$$P(\{\log L_i\} | \theta) \propto P(\{\log L_i\} | \theta) = \int P(\{\log L_i\} | \theta, \mathbf{X}) P(\mathbf{X}) d\mathbf{X} \quad (2)$$

$$P(\mathbf{X}) = \frac{1}{2\pi|\Sigma|} \exp \left[-\frac{1}{2}(\mathbf{X} - \boldsymbol{\mu})^T \Sigma^{-1}(\mathbf{X} - \boldsymbol{\mu}) \right] \quad (3)$$

$$P(\{\log L_i\} | \mathbf{X}, \theta) = \prod_i \delta(\log L_i - f(X_i, \theta)) \quad (4)$$

$$\therefore P(\theta | \{\log L_i\}) = \int \frac{1}{2\pi|\Sigma|} \exp \left[-\frac{1}{2}(\mathbf{X} - \boldsymbol{\mu})^T \Sigma^{-1}(\mathbf{X} - \boldsymbol{\mu}) \right] \left(\prod_i \delta(\log L_i - f(X_i, \theta)) \right) dX_1 \cdots dX_n \quad (5)$$

$$f(X_i, \theta) = \log L_{\max} - \frac{X_i^{2/d}}{2\sigma^2} \quad (6)$$

Evaluate integral

Consider the identity

$$\delta(g(x)) = \sum_{\text{roots } j} \frac{\delta(x - x_j)}{|g'(x_j)|} \quad (7)$$

For the above case we expect only one root for each X_i , namely at $\log L_i = f(X_i^*, \theta)$ (assuming that f is one-to-one, which it is for the form we are using). (4) reduces to

$$\prod_i \frac{\delta(X_i - X_i^*)}{|f'(X_i^*, \theta)|} \quad (8)$$

Putting this together, the integral evaluates to

$$P(\theta | \log \mathbf{L}) = \frac{1}{2\pi|\Sigma|} \left(\prod_i \frac{1}{|f'(X_i^*, \theta)|} \right) \exp \left[-\frac{1}{2}(\mathbf{X}^* - \boldsymbol{\mu})^T \Sigma^{-1}(\mathbf{X}^* - \boldsymbol{\mu}) \right] \quad (9)$$

where the \mathbf{X}^* are given by the inverse of f . Discard normalising term:

$$\log P(\theta | \log \mathbf{L}) = - \left(\sum_i \log |f'(X_i^*, \theta)| \right) - \frac{1}{2}(\mathbf{X}^* - \boldsymbol{\mu})^T \Sigma^{-1}(\mathbf{X}^* - \boldsymbol{\mu}) \quad (10)$$

Substitute specific form of f

Invert f to find X_i^* and $|f'(X_i^*)|$

$$f(X_i, \theta) = f(X_i, d) = -X_i^{2/d} \quad (11)$$

$$X_i^* = (-\log L_i)^{d/2} \quad (12)$$

$$f'(X_i, d) = -\frac{2}{d} X_i^{2/d-1} \quad (13)$$

$$\therefore \log P(\theta \mid \log \mathbf{L}) = - \left(\sum_i \log \left| \frac{2}{d} X_i^{*2/d-1} \right| \right) - \frac{1}{2} (\mathbf{X}^* - \boldsymbol{\mu})^\top \Sigma^{-1} (\mathbf{X}^* - \boldsymbol{\mu}) \quad (14)$$