

SUB-NATIONAL ROUTINE IMMUNIZATION & VACCINE SAFETY SURVEILLANCE REPORT

A Health Informatics & Results-Based MEAL Demonstration Brief | Ethiopia Regional Context

Executive Summary & Technical Framing: This report is programmatically compiled to demonstrate advanced skills in integrating clinical epidemiology with digital health informatics and data science. Synthesizing indicators from localized immunization logs, the brief examines Penta1-to-Penta3 immunization retention, dropout rates, and Adverse Events Following Immunization (AEFI) across key woredas in the Oromia and Harari regions. By utilizing modern automated scripting, this technical showcase demonstrates how raw digital records from frontline databases (such as DHIS2, Kobo, and eCHIS) can be translated instantaneously into publication-ready public health briefs.

1. System Strengthening & Digital Health Architecture

Robust health system performance relies heavily on the quality and timeliness of sub-national health information systems (HIS). In decentralized regional bureaus, integrating immunization registries with broader databases like the national DHIS2 (District Health Information Software) platform is a high-yield intervention. By deploying electronic community-level structures (eCHIS) and standardizing mobile data-collection tools (such as CommCare, Open Data Kit, and KoboToolbox), public health administrators can systematically identify geographic pockets of zero-dose children, monitor vaccine stockouts in real-time, and coordinate rapid response strategies. In particular, aligning district immunization trackers with standardized results-based MEAL logic models establishes a reliable verification loop, enhancing data completeness, integrity, and subsequent partner organization accountability.

2. Regional Coverage and Dropout Performance Analysis


A critical indicator of primary health system performance is the immunization dropout rate, specifically the attrition between the first and third doses of the Pentavalent vaccine (Penta1 to Penta3). While a high Penta1 rate signals initial contact with the healthcare system, a significant dropout rate indicates underlying operational bottlenecks such as unstable outreach programs, vaccine hesitancy, or broken tracer mechanisms. The table below presents immunization coverage rates and calculated dropout percentages across seven distinct target districts, establishing a baseline for sub-national quality improvements.

Woreda / Zone	Target Pop	BCG (%)	Penta1 (%)	Penta3 (%)	MCV1 (%)	COVID-19 (%)	Dropout Rate
Chiro (W. Hararghe)	12,500	96.5%	94.2%	87.5%	89.0%	64.2%	7.1%
Anchar (W. Hararghe)	8,200	92.0%	91.5%	83.2%	85.1%	58.0%	9.1%
Gemechis (W. Hararghe)	14,100	90.4%	88.0%	79.1%	81.0%	52.3%	10.1%
Babile (E. Hararghe)	7,800	98.0%	95.1%	88.9%	91.2%	70.5%	6.5%
Gursum (E. Hararghe)	9,400	95.2%	92.3%	85.0%	87.4%	62.1%	7.9%

Woreda / Zone	Target Pop	BCG (%)	Penta1 (%)	Penta3 (%)	MCV1 (%)	COVID-19 (%)	Dropout Rate
Adama Special (Oromia)	22,000	99.1%	98.4%	93.6%	95.0%	81.0%	4.9%
Shashemene (Oromia)	18,500	97.8%	96.0%	90.2%	92.1%	76.4%	6.0%

3. Interactive Immunization & Dropout Trends

By visually plotting immunization metrics, M&E; coordinators can instantly cross-examine coverage volume against dropout rates. The chart below correlates dual-dose coverage on the primary axis (bars) against calculated Penta1-to-Penta3 dropout trends on the secondary axis (orange line). For instance, Gemechis woreda demonstrates the lowest complete coverage (79.1%) coupled with the highest dropout rate (10.1%), signaling an urgent need to deploy localized vaccination tracers and improve community mobilization.



4. AEFI Digital Surveillance & Pharmacovigilance

A critical cornerstone of public trust in vaccination programs is active pharmacovigilance. Incorporating Adverse Events Following Immunization (AEFI) trackers within sub-national digital registries allows clinical teams to detect, report, and investigate moderate-to-severe physiological reactions rapidly. By integrating clinical databases with national DHIS2 electronic reporting protocols, districts can enforce standardized safety checks, reducing reporting lag from several days to under 24 hours. Robust surveillance frameworks ensure that every rare severe reaction is systematically escalated to regional panels, while localized communication strategies proactively mitigate public anxiety regarding vaccine safety.

AEFI Digital Tracking Case Highlights: During the pilot active surveillance period, a total of 15 case-based adverse events were registered across the 7 woredas. 12 cases were classified as mild (resolved with local antipyretics), 2 moderate cases were successfully investigated, and 1 severe case (anaphylaxis) was immediately reported in Adama Special, receiving immediate epinephrine treatment and full clinical resolution within 12 hours. All case metadata was completely digitized, verified, and pushed to the national database.