

WINTER 2017/18 TERM 2 STAT 302: ASSIGNMENT 3

Due: 2pm on Tuesday March 27, 2018

- Please remember to **include a cover sheet** when you submit your assignment. Please hand in your assignment in the STAT 302 assignment box on the ground floor of the Earth Sciences Building (ESB) by the due time.
- When answering the questions, writing down the final answer will not be sufficient to receive full marks. Please show all calculations unless otherwise specified. Also define any events and notation that you use in your solutions.

Question 1: We are studying a collection of molluscs living on a large beach in northern BC. The molluscs live near the high-tide line and it is known that their location is uniformly distributed with minimum -1 and maximum +2 metres from this line. Their daily energy intake (in kilocalories) turns out to be 1 plus 25% of their squared location.

(a) What is the cumulative distribution function of the daily energy intake?

Let X denote the location of the molluscs. $X \sim U(-1, 2)$

Let Y denote the daily energy intake, $Y = 1 + 25\%X^2$

$$f(x) = \begin{cases} 1/3, & x \in [-1, 2] \\ 0, & \text{otherwise} \end{cases}$$

$$Pr(X \leq x) = F(x) = \begin{cases} 0, & x < -1 \\ \frac{x+1}{3}, & x \in [-1, 2] \\ 1, & x > 2 \end{cases}$$

$$Pr(Y \leq y) = Pr(1 + 0.25X^2 \leq y) = Pr(X^2 \leq 4(y-1)) = Pr(X \geq -2\sqrt{y-1}, X \leq 2\sqrt{y-1})$$

Compute the domain of y :

$$-1 \leq -2\sqrt{y-1} \leq 0 \implies 1 \leq y \leq \frac{5}{4}$$

$$0 \leq 2\sqrt{y-1} \leq 2 \implies 1 \leq y \leq 2$$

$$\text{Therefore, when } 1 \leq y \leq \frac{5}{4}, F(y) = \frac{2\sqrt{y-1}+1}{3} - \frac{-2\sqrt{y-1}+1}{3} = \frac{4\sqrt{y-1}}{3}$$

So the cumulative distribution function of daily energy intake is:

$$F(y) = \begin{cases} 0, & y < 1 \\ \frac{4\sqrt{y-1}}{3}, & 1 \leq y \leq \frac{5}{4} \\ \frac{2\sqrt{y-1}+1}{3}, & \frac{5}{4} \leq y \leq 2 \\ 1, & y > 2 \end{cases}$$

(b) What is the probability density function of the daily energy intake?

$$f(y) = F'(y) = \begin{cases} 0, & y < 1 \text{ or } y > 2 \\ \frac{2}{3\sqrt{y-1}}, & 1 \leq y \leq \frac{5}{4} \\ \frac{1}{3\sqrt{y-1}}, & \frac{5}{4} \leq y \leq 2 \end{cases}$$

(c) What is the expected daily energy intake?

$$\mathbb{E}(Y) = \int_{-\infty}^{\infty} yf(y)dy = \int_1^{5/4} \frac{2y}{3\sqrt{y-1}}dy + \int_{5/4}^2 \frac{y}{3\sqrt{y-1}}dy$$

$$\mathbb{E}(Y) = \int_1^{5/4} \frac{2}{3}\sqrt{y-1}dy + \int_1^{5/4} \frac{2}{3\sqrt{y-1}}dy + \int_{5/4}^2 \frac{1}{3}\sqrt{y-1}dy + \int_{5/4}^2 \frac{1}{3\sqrt{y-1}}dy$$

$$\begin{aligned}\mathbb{E}(Y) &= \frac{4}{9}(y-1)^{3/2}|_1^{5/4} + \frac{4}{3}\sqrt{y-1}|_1^{5/4} + \frac{2}{9}(y-1)^{3/2}|_{5/4}^2 + \frac{2}{3}\sqrt{y-1}|_{5/4}^2 \\ \mathbb{E}(Y) &= \frac{4}{9} \times \frac{1}{8} + \frac{4}{3} \times \frac{1}{2} + \frac{2}{9} \times \frac{7}{8} + \frac{2}{3} \times \frac{1}{2}\end{aligned}$$

$$\mathbb{E}(Y) = \frac{5}{4}$$

Question 2: Let X and Y be two independent Bernoulli(0.5) random variables and define $U = X + Y$ and $V = X - Y$.

- Find the joint and marginal probability mass functions for U and V . [It is sufficient to construct a table to describe these mass functions.]
- Are U and V independent? Why or why not?
- Find the conditional probability mass functions $p_{U|V=v}(u)$ and $p_{V|U=u}(v)$. [Again, you can construct a table to describe these mass functions.]

Question 3: This question will provide an intriguing contrast to Question 2. Recall that if we have a continuous random variable X defined by a pdf $f_X(x)$, and we define a new random variable $Y = g(X)$ where g is a bijective (i.e. one-to-one) transformation, then the inverse of g is well-defined everywhere, g^{-1} , and the density of Y is given by

$$f_Y(y) = f_X(g^{-1}(y)) \cdot \left| \frac{d}{dy} g^{-1}(y) \right|.$$

We can generalize this to the bivariate setting as follows. Suppose X_1 and X_2 are continuous random variables with joint pdf $f_{X_1, X_2}(x_1, x_2)$ and suppose that both $u = g_1(X_1, X_2)$ and $v = g_2(X_1, X_2)$ are bijective (i.e. one-to-one) transformations with inverses $g_1^{-1}(u, v)$ and $g_2^{-1}(u, v)$. If these inverse functions have continuous partial derivatives and nonzero *Jacobian*

$$J = \det \begin{bmatrix} \frac{\partial g_1^{-1}}{\partial u} & \frac{\partial g_1^{-1}}{\partial v} \\ \frac{\partial g_2^{-1}}{\partial u} & \frac{\partial g_2^{-1}}{\partial v} \end{bmatrix} = \frac{\partial g_1^{-1}}{\partial u} \frac{\partial g_2^{-1}}{\partial v} - \frac{\partial g_2^{-1}}{\partial u} \frac{\partial g_1^{-1}}{\partial v} \neq 0,$$

then the joint density of u and v is

$$f_{u,v}(u, v) = f_{X_1, X_2}(g_1^{-1}(u, v), g_2^{-1}(u, v)) |J|,$$

where $|J|$ is the absolute value of the Jacobian.

- Let X_1 and X_2 be independent standard normal random variables. Write down the joint probability density function of X_1 and X_2 . Moreover, compute $\Pr(X_1 < 1, X_2 < 1)$.
- Define the transformations $u = g_1(x_1, x_2) = x_1 + x_2$ and $v = g_2(x_1, x_2) = x_1 - x_2$. Find the inverse functions $g_1^{-1}(u, v)$ and $g_2^{-1}(u, v)$ and compute the Jacobian of this bivariate transformation of variables.
- Write down the joint pdf of $U = X_1 + X_2$ and $V = X_1 - X_2$ and show that this density separates over variables; i.e. show $f_{U,V}(u, v) = a(u)b(v)$ for some real functions $a(u)$ and $b(v)$. Recall from class that this implies that U and V are actually *independent*.

- (d) Let X_1, \dots, X_n be a random sample from a normal distribution with mean μ and variance σ^2 . If $n = 2$, show that the sample mean, \bar{X} , and the sample variance, S_X^2 , are independent random variables. [Hint: first write \bar{X} and S_X^2 in terms of U and V as above, remembering that U and V are linear combinations of *standard* normal random variables.]

Question 4: Let W be a Gamma random variable with parameters $(\alpha = 2, \lambda = 2)$. Conditional on the value $W = w$, X is an exponential random variable with rate parameter w .

- (a) What is the conditional density function for X given $W = w$? Be sure to indicate any restrictions on the values of x and w . If $W = 2$, what is the probability that $X \leq 2$?
- (b) What is the probability that W is greater than its expected value? Do *not* use an online applet to find this probability; calculate by hand.
- (c) Show that the conditional distribution of W given $X = 2$ is Gamma distributed with parameters $(\alpha = 3, \lambda = 4)$. [Recall that $\Gamma(\alpha) = (\alpha - 1)!$, and use the fact that $f_X(2) = 1/8$.]

Question 5: Let X be the amount of time that a student spends walking from the Earth Sciences Building to the Mathematics Building, and let Y be the amount of time that a student spends walking from the Earth Sciences Building to the Pharmaceutical Sciences Building. Suppose that the joint density of X and Y is given by the following function

$$f_{X,Y}(x, y) = \begin{cases} \frac{kx}{y} & \text{if } 0 < x < y < 10 \\ 0 & \text{otherwise} \end{cases}$$

for some fixed constant k .

- (a) What value of k makes $f_{X,Y}(x, y)$ an honest probability density function?
- (b) Find the marginal probability density function for X . [Don't forget to specify the support of the function!]
- (c) Find the probability $\Pr(Y + X \leq 10)$.