Subjective Annotation Studies: Valid Inference and Annotation Design

Click here for the latest version

Abstract

We develop a statistical framework for annotation studies of inherently subjective features of unstructured data. In the absence of a conventional ground-truth measurement, we define document-level measurements relative to a reference population of evaluators whose responses are relevant to the underlying economic question. Our goal is valid inference for weighted corpus averages of these measurements and parameters derived from them. We specify an annotation design under which inexpensive machine predictions for the full corpus can be combined with relatively scarce human evaluations of sampled documents. Under this design, we establish asymptotically valid inference and characterize the variance arising from document sampling, limited evaluations per document, and evaluator reuse. For weighted corpus averages, we use this variance to derive an optimal allocation of the annotation budget. The allocation increases a document's sampling probability before increasing its number of evaluations above the required minimum and directs more resources toward documents with larger absolute weights in the corpus average, greater prediction errors, or more variable human responses. We illustrate the method with simulations calibrated to Wikipedia toxicity annotations and an empirical application based on Shapiro and Wilson (2022a).

1 Introduction

Economists increasingly use unstructured data, such as text, audio, images, and video, to study questions for which conventional datasets lack suitable measures (Gentzkow et al., 2019; Dell, 2025; Ash and Hansen, 2023). To use these records in econometric analysis, however, researchers must first construct numerical measurements with clear interpretations, a process in which human annotation often plays a central role. Much of the existing literature studies applications in which a ground-truth measurement exists but is costly to obtain. Human labels are then treated as expensive, high-quality validation data that directly reveal the truth or provide information that helps identify it. Examples include determining whether a vacancy permits remote work (Hansen et al., 2023) or transcribing information from an archival record (Dell, 2025), among other applications (see also Burchfield et al., 2006; Donaldson and Hornbeck, 2016; Henderson et al., 2021). There is, however, a wide class of applications where a ground-truth measurement is conceptually difficult to establish. Dupas et al. (2026), for example, investigate whether women presenting research in economics seminars encounter more negative interactions than men. What constitutes a negative interaction, however, is inherently subjective and can mean different things to different people; two careful labelers may therefore evaluate the same interaction differently without either being wrong. In this setting, the absence of a traditional ground-truth target makes the construction of an interpretable measurement a problem in its own right and creates corresponding difficulties for the interpretation and inference of downstream parameters.

Such settings are not uncommon in empirical work. Researchers have constructed measures of the ideological slant of news coverage (Gentzkow and Shapiro, 2010), the emotional tone conveyed by central-bank communication (Gorodnichenko et al., 2023), and the perceived trustworthiness of borrowers from their photographs (Duarte et al., 2012). Although they differ in subject matter, these applications concern subjective judgments on which careful evaluators may disagree without error. Recent work in machine learning has similarly argued that disagreement in subjective annotations can be systematic and informative and that the resulting labels depend on whose responses are represented (Plank, 2022; Gordon et al., 2022). We give this insight a statistical formulation. The researcher first specifies a reference population of potential annotators whose assessments are directly relevant to the research question. We use "document" to refer to an individual text, audio recording, image, or video, and "corpus" to refer to the collection under study. For each document, the researcher then defines measurements that summarize the distribution of responses in this population. Observed annotations therefore record responses from sampled members of the reference population and carry information about these document-level measurements, rather than directly revealing a ground-truth value. Within this framework, we first specify a class of annotation designs describing how documents are sampled, how many evaluations each receives, and how evaluators are assigned to documents. Under these designs, we use established estimators that combine a limited set of human annotations with machine predictions for the full corpus. We establish valid inference for corpus-level parameters and characterize the asymptotic variance. We then use this variance characterization to derive optimal allocations of annotation resources.

In the absence of a conventional ground-truth target, we first define the document-level measurement by fixing a substantively relevant reference population. The reference population should be treated as part of the economics research question. A population may be relevant because its members possess the interpretive context needed to evaluate the object of interest, because their perceptions are implicated by the empirical question, or both. Members of the economics profession, for example, possess the contextual knowledge needed to interpret seminar interactions, while their perceptions of professional conduct are also substantively relevant in the seminar setting discussed above. In the absence of a conventional ground-truth target, different reference populations therefore define different measurements.

Under a fixed annotation protocol, each member of the reference population has a potential response to document j. A natural measurement is the mean of these responses, μ_j, which we use as our primary document-level measure. Disagreement among evaluators can also be economically informative. For example, differences in responses to written descriptions of macroeconomic shocks reveal differences in how people understand the economy (**?**). For binary responses, the mean determines the response distribution and hence the extent of disagreement. With richer response scales, however, documents with the same mean can exhibit different levels of disagreement. We therefore also develop theory for measurements based on pairs of independently drawn responses to the same document. These include response variance and the probability that two categorical responses differ, both of which measure disagreement directly. The same framework also provides unbiased estimates of the squared means μ_j^2 needed for regressions that use μ_j as an explanatory variable.

We study estimation and inference for corpus level weighted averages of these document-level measurements and smooth functions of those averages. We take the J-document corpus and its observed characteristics as given, with weights determined by these characteristics and the economic question. This class and its applications include corpus means (Gentzkow and Shapiro, 2010), comparisons across subcorpora (Dupas et al., 2026), and regression coefficients when the document-level measurement is the outcome (Shapiro and Wilson, 2022a). The analysis extends naturally to fixed-dimensional vectors of such averages and smooth transformations of these vectors. These extensions cover regression coefficients

when a subjective measurement is used as a regressor (Möbius and Rosenblat, 2006) and parameters from nonlinear specifications (Kessler et al., 2019).

To estimate these corpus-level parameters, researchers must choose which documents to annotate, how many evaluations each receives, and how evaluators are assigned to documents. With unlimited resources, one could repeatedly draw independent evaluators from the reference population for every document, allowing each document-level measurement to be estimated arbitrarily precisely. In realistic annotation studies, however, limited budgets make this infeasible. Researchers must therefore decide how to distribute the available evaluations across documents and evaluators, and these choices affect the precision of the resulting corpus-level estimates. We specify a class of designs in which documents are sampled independently using probabilities chosen in advance, a procedure known as Poisson sampling (Särndal et al., 1992). The number of evaluations each document receives if sampled is also chosen in advance. We impose a fixed cap on each evaluator's workload (Derczynski et al., 2016; Potash et al., 2019; Haagsma et al., 2020). To limit the recruitment burden, we use the smallest feasible workforce under this cap and fill all but possibly one evaluator to capacity. We assign evaluators to documents at random, requiring each sampled document to receive its prescribed number of evaluations and each evaluator to complete their assigned workload. No evaluator evaluates the same document twice, and all assignments satisfying these requirements are equally likely. Within this class, researchers can vary both the probability that a document is sampled and the number of evaluations it receives if sampled. This leads to a natural question: how should researchers choose these quantities under an annotation budget to estimate the aforementioned corpus-level parameters as precisely as possible?

Our estimator builds on established methods for combining machine predictions with observed measurements, but human annotations in our setting cannot be treated as simple ground-truth document level measurements. Inference must therefore account for how these measurements are estimated from evaluator responses. Human evaluations are costly to collect, whereas machine generated outputs can be obtained inexpensively for the full corpus. Our estimator follows an established solution in these type of situations Särndal et al. (1992); Angelopoulos et al. (2023); Zrnic and Candès (2024): it first computes the relevant corpus averages using machine predictions for all documents, then adds a correction based on human measurements from sampled documents. This correction averages the differences between those measurements and the corresponding predictions, accounting for unequal sampling probabilities. In our setting, each sampled document's measurement must itself be estimated from a small number of evaluator responses. Reusing evaluators can therefore correlate the resulting estimation errors across documents, with the dependence pattern determined by how evaluators are assigned to documents.

This complicates statistical inference, which must account for the random selection of documents, the random allocation of their evaluations across workers, and the dependence created when workers evaluate multiple documents. A Gaussian approximation for the estimator may be straightforward to establish conditional on the documents selected for annotation and their assignment to evaluators. To quantify uncertainty about the corpus-level parameter, however, we must also account for variation across possible document selections and document allocation. This requires averaging the conditional distributions over these possibilities, and the resulting distribution need not be Gaussian under arbitrary allocation mechanism. We show that under our proposed annotation design, the estimator has a Gaussian limit under standard regularity conditions. We also construct a consistent estimator of the resulting asymptotic variance, yielding standard feasible Gaussian inference.

The variance characterization obtained from our asymptotic analysis also allows us to study how annotation resources should be allocated. Choosing a sampling design to improve precision under a budget is a classical problem in survey sampling (Stuart, 1954; Särndal et al., 1992). More recent work on active

statistical inference and stratified prediction-powered inference also started studying how the allocation of labels can improve precision (Zrnic and Candès, 2024; Fisch et al., 2024). Our reference-population setting introduces a distinct allocation decision because each document-level measurement must itself be estimated from evaluator responses. Collecting more evaluations thus improves the precision of that measurement but leaves fewer resources for annotating other documents. We therefore jointly choose document inclusion probabilities and annotation depths to minimize the asymptotic variance of the resulting estimator under an annotation budget.

We study this problem for weighted averages of the reference population mean, referring to the weights in these averages as target weights. Importantly, within our design class, the first-order variance contribution from worker reuse is common across allocation policies, so worker reuse affects attainable precision but not how resources should be distributed across documents. We also show that, for any fixed expected expenditure on a document, it is weakly preferable to increase its inclusion probability before increasing the labels it receives. The optimal allocation therefore follows a coverage-before-depth rule, increasing labels per document only after the document is sampled with certainty. Profiling out this choice yields a convex problem with a generalized Neyman solution (Neyman, 1934). At a specified annotation depth, the solution assigns expected expenditure according to the document's absolute target weight and the square root of a document-specific difficulty measure. This measure depends on how much evaluators' responses vary for a document, how far the machine prediction is from the reference-population mean, and how many evaluations the document receives if sampled. Thus, more resources are directed toward documents with larger absolute target weights, noisier human responses, or less accurate machine predictions.

In practice, the machine-prediction errors and evaluator-response variation needed by the optimal allocation rules are unknown before annotation. A pilot with repeated annotations, or comparable annotated data, can be used to predict them from observed document characteristics. When reliable information for ranking document difficulty is unavailable, we propose a transparent default that treats documents as equally difficult in expectation for planning purposes and divides the budget using only the fixed weights in the weighted corpus average. When these weights are not too concentrated and the budget is small relative to the corpus, expected expenditure on each document is proportional to the absolute value of its weight, and every sampled document receives the minimum annotation depth. Two examples of this allocation rule are particularly interpretable. For a corpus mean, this rule samples all documents with the same probability. For a contrast between two group means, it assigns half of the expected budget to each group and therefore samples documents in the smaller group with higher probability.

To illustrate the performance of our estimator in a setting grounded in real annotation data, we conduct a calibrated simulation using the Wikipedia Talk Labels: Toxicity corpus (Wulczyn et al., 2017; Thain et al., 2017). We use the observed comments and human judgments to estimate comment-level toxicity probabilities and calibrate a synthetic population of workers. We apply a pretrained Cardiff offensive-language classifier to each comment to generate the machine predictions used by the estimator (Barbieri et al., 2020). For the corpus mean, nominal 95% confidence intervals achieve 93.2–94.5% coverage, compared with 54.8–54.9% when each comment's average label is treated as error-free. In a stress test with approximately 30 or 60 workers, coverage ranges from 93.5% to 94.7%. For the difference in mean toxicity between unregistered and registered contributors, allocation based on target weights reduces empirical variance by 25.4% relative to uniform sampling. The full-information optimal allocation and its pilot-estimated version reduce variance by 36.2% and 29.4%, respectively.

We apply our framework to the setting studied by Shapiro and Wilson (2022b), who relate sentiment in Federal Open Market Committee (FOMC) deliberations to economic conditions to investigate central-

bank objectives. They measure sentiment using a dictionary-based score: the share of negative words minus the share of positive words. They then fit a quadratic relationship between this score and inflation to estimate the inflation rate associated with the least negative sentiment (the inflation target). We conduct an annotation study with Prolific participants selected for relevant educational, professional, or investment experience, replacing dictionary-based sentiment with the mean assessment of this reference population. We combine these human annotations with either predictions from GPT-6 Astra or a rescaled version of the authors' original dictionary score. The resulting turning-point estimates are 1.68% and 1.79%, respectively. Both are close to the authors' original estimate of approximately 1.69%, supporting the effectiveness of their dictionary-based measure for this economic task.

The remainder of the paper is organized as follows. Section 2 formalizes the reference-population measurement framework and defines the corpus-level targets, and Section 3 introduces the annotation design. Section 4 develops the estimator and establishes valid inference under evaluator reuse, while Section 5 derives the optimal allocation and its practical implementation. Section 6 uses a simulation calibrated to the Wikipedia toxicity data to evaluate inference under worker reuse, finite-sample performance, and the precision gains from targeted annotation design. Proofs and supporting results are collected in the appendices.

2 Model

2.1 Annotation outcomes

Let $W_{1:J} := (W_1, \ldots, W_J)$ denote a fixed but large corpus of J document. In the main text, we treat corpus construction as completed before the annotation study begins, and $W_{1:J}$ is held fixed throughout the main-text analysis. The parameters below therefore concern this fixed corpus and require no probability model for document generation. When a superpopulation model for document generation can be credibly imposed, it can be combined with the main analysis. Online Appendix OA.4 provides such an example for the model-based extensions.

Let P_θ denote the distribution of worker characteristics in a prespecified reference population. Potential workers are indexed by $i \in \mathbb{N}$, and θ_i collects all latent characteristics that may persist across evaluations, such as background, prior beliefs, and interpretive tendencies. For each worker–document pair, let ε_{ij} represent influences on worker i's response to document j that are not captured by the document W_j or the worker's persistent characteristics θ_i. Let Y_{ij}^* denote the response that worker i would provide if assigned to evaluate document j. Notice, this potential response is defined whether or not that assignment is made. We assume the responses are bounded, that is $|Y_{ij}^*| \le C_Y$ for all i, j, where $C_Y < \infty$ is a fixed constant independent of i, j, and J. We write $\mathcal{Y} \subseteq [-C_Y, C_Y]$ for the set of possible response values. We impose the following assumptions.

(S1) Stable potential responses. There is a measurable function h, common to all pairs and otherwise unrestricted, such that

$$Y_{ij}^* = h(W_j, \theta_i, \varepsilon_{ij}), \qquad i \ge 1, \quad j \le J.$$

The response is invariant to the realized document sample, worker assignment, evaluation order, and workload.

(S2) Pair-specific shocks. The shocks $\{\varepsilon_{ij} : i \ge 1, j \le J\}$ are mutually i.i.d. and independent of $\{\theta_i : i \ge 1\}$.

(RW0) Reference-population sampling. The worker types $\{\theta_i\}_{i \ge 1}$ are i.i.d. draws from P_θ.

Assumption **(S1)** ensures that the potential-response notation is well defined: the response of worker i to document j is a fixed function of the pair alone, invariant to the realized annotation design. It rules out fatigue, order, and workload effects, under which a worker's response to a document would depend on the rest of their assignment.

Assumption **(S2)** makes ε_{ij} a genuinely pair-specific shock. It rules out response dependence across worker–document pairs beyond that induced by persistent worker characteristics, including communication or herding among annotators. Responses from the same worker may nevertheless remain dependent across documents through θ_i. Importantly, θ_i need not represent only an additive worker effect: it may encode the worker's entire response rule, allowing different worker types to map the same document into very different response distributions. The common function h therefore does not impose a common interpretation rule across workers.

A well-designed annotation exercise should seek to limit violations of Assumptions **(S1)**–**(S2)**. Preventing communication among annotators can limit dependence across workers, while limiting the number of documents per session and requiring regular breaks can help reduce fatigue. Attention checks can also help detect inattention during the study. These practices are increasingly practical to implement as researchers gain access to tools for managing annotation workloads, scheduling breaks, and monitoring attention both in online and offline settings.

Assumption **(RW0)** permits the reference population and its eligibility rules to reflect the application and corpus domain. Such choices define the population being averaged over and therefore form part of the measurement itself. Conditional on the workforce size required by the design, the recruited annotators must represent the announced population; their identities or composition cannot be selected using the sampled documents or any final-study annotation outcomes.

2.2 Document-level measurements, characteristics, and pre-trained predictions

Let θ denote a generic draw from P_θ, and let ε denote an independent draw from the distribution of the pair-specific shocks. Our primary document-level measurement is the reference-population mean response

$$\mu_j := \mathbb{E}_{\theta,\varepsilon}[h(W_j, \theta, \varepsilon)] \,.$$

The expectation averages over worker characteristics and pair-specific shocks, with the document W_j held fixed.

To interpret μ_j, consider Dupas et al. (2026), who study, in one specification, the number of seminar interruptions that fellow economists rate as negative in tenor. In our notation, let W_j contain the interactions in seminar j, and let Y_{ij}^* be the number that economist i would rate as negative. Their in-person seminar sample illustrates how such a measure can be constructed by asking economists to assess the interactions directly.

If annotation resources were unlimited, the researcher could draw M economists independently from a prespecified population of economics researchers, ask each to evaluate seminar j once, and average their responses. Under Assumptions **(S1)**–**(S2)** and **(RW0)**,

$$\frac{1}{M}\sum_{i=1}^{M} Y_{ij}^* \xrightarrow{\text{a.s.}} \mu_j \qquad \text{as } M \to \infty.$$

The mean provides a natural measure of response level and is the primary document level measurement we consider. However, In some applications, heterogeneity in reference-population responses is itself substantively important. To some extent, the mean framework itself can accommodate questions about response heterogeneity by changing the outcome being averaged. For example, if Y_{ij}^* is an ordered categorical response, the researcher may instead take the outcome to be the indicator that the response lies at or below a prespecified cutoff c. The resulting document-level mean is

$$\mu_j(c) := \mathbb{E}\big[\mathbf{1}\{Y_{ij}^* \leq c\}\big] = \Pr(Y_{ij}^* \leq c),$$

the probability that a randomly drawn member of the reference population returns a response at or below that cutoff. Different cutoffs reveal different features of the response distribution.

In many applications, however, researchers may prefer a measurement that directly captures the extent of disagreement or contestedness. Let Y_{1j}^* and Y_{2j}^* denote the responses of two independently drawn workers, including independent pair-specific shocks. A measure of response dispersion can be defined as

$$\nu_j := \frac{1}{2}\mathbb{E}\big[(Y_{1j}^* - Y_{2j}^*)^2\big] = \mathrm{Var}(Y_{ij}^*).$$

Whereas a recoded mean describes a particular feature of the response distribution, ν_j directly summarizes how differently two randomly drawn members of the reference population are expected to respond.

Pairs of responses are also useful when μ_j enters a linear regression as a regressor. Such regressions require corpus averages of μ_j^2. However, simply squaring a document's average annotation overestimates μ_j^2 on average by the variance of that average. The resulting measurement-error bias need not disappear as the corpus grows if each document receives only a few evaluations. Products of responses from two independently drawn evaluators instead have expectation μ_j^2, allowing us to estimate the required corpus moment consistently.

More generally, we extend our analysis to pairwise response measurements of the form

$$\vartheta_j(\ell) := \mathbb{E}\big[\ell_j(Y_{1j}^*, Y_{2j}^*)\big],$$

where ℓ_j is a prespecified symmetric comparison. Their natural sample analogues compare pairs of annotations within a document and are order-two U-statistics.

Before annotation begins, the researcher may already have some information for every document in the corpus. Because W_j is an unstructured object, it enters the econometric analysis only after being mapped into numerical variables or scores. Let $X_j \in \mathbb{R}^d$, for fixed d, collect prespecified document characteristics and numerical representations of its content, such as its source and date, characteristics of its author or speaker, pretrained text, image, or audio embeddings. Researchers may use these characteristics as regressors in downstream analyses or to guide the choice of document sampling probabilities and the number of evaluations per sampled document.

Before documents are selected for human annotation, the researcher may also construct a scalar prediction g_j of μ_j from W_j, X_j, or both. The prediction procedure may be an off-the-shelf classifier, a model trained on auxiliary data, or an LLM. In the LLM case, the prespecified procedure includes the model version, prompt, conversion of the response into a numerical score, number of calls, and any rule used to combine outputs across calls.

Once the procedure has been applied to every document, the resulting vector $g_{1:J}$ is held fixed: it is not regenerated, re-trained, or revised after the annotation sample is drawn or in response to the final-

study annotations. We call g_j a *frozen prediction* precisely in this timing sense. If the procedure is stochastic, the subsequent analysis conditions on the realized predictions. Thus, both a single realized call and a prespecified aggregation of repeated calls are permitted.[1] The frozen prediction may inform the annotation design and enter the prediction-assisted estimator. It need not be unbiased, calibrated, or consistent. As we show below, its predictive quality affects efficiency rather than validity. However, we do require it to be bounded, given the original annotation outcome is bounded.

(RW2) Bounded frozen predictions. We assume that $|g_j| \leq \bar{g}$ for every document j, where $\bar{g} < \infty$ is a fixed constant independent of j and J.

2.3 Economic parameters generated by corpus moments

The preceding subsection defines document-level measurements, including the reference-population mean μ_j and the pairwise measurements $\vartheta_j(\ell)$. Economic applications, however, generally concern corpus-level parameters. We consider parameters built from weighted averages of these document-level measurements of the form

$$\Phi_J(a) = \frac{1}{J}\sum_{j=1}^{J} a_j \mu_j, \qquad \Phi_J^{\ell}(a) = \frac{1}{J}\sum_{j=1}^{J} a_j \vartheta_j(\ell),$$

where the weights a_j may depend on observed corpus characteristics, such as the X_j's, but not on the potential responses Y_{ij}^*.

Let $\mathbf{\Phi}_J$ denote a fixed-dimensional vector whose components take the form $\Phi_J(a)$ or $\Phi_J^{\ell}(a)$. We study corpus-level parameters that can be written as

$$\boldsymbol{\beta}_J = H_J(\mathbf{\Phi}_J), \tag{2.1}$$

where H_J is a known, corpus-specific map that is continuously differentiable in a neighborhood of $\mathbf{\Phi}_J$. The map may depend on fixed observed attributes of the entire corpus.

The following applications fall in this class.

Reader-perceived media slant (Gentzkow and Shapiro, 2010). The authors study political slant in U.S. newspapers and construct an automated measure based on partisan phrase use. In our framework, W_j is the text of article j, and Y_{ij}^* is reader i's assessment of the article's political position. A natural reference population is U.S. newspaper readers. Thus μ_j is the mean perceived political position of article j in this population. For a newspaper n with a corpus C_n of J_n articles, the newspaper-level target is

$$\beta_{J_n,n} = \frac{1}{J_n}\sum_{j \in C_n} \mu_j = \Phi_{J_n}(\mathbf{1}).$$

An article-level adaptation of the authors' phrase-scoring rule could provide a natural g_j.

Negative interruptions in economics seminars (Dupas et al., 2026). The authors study whether female presenters are treated more harshly than male presenters in economics seminars. They combine human coding with audio-processing algorithms. In our framework, W_j is the available audio or audiovisual record of seminar j, X_j includes whether the seminar was held online or in person, and D_j indicates a female presenter. The authors recruited human coders from the economics research community, mainly graduate students. We therefore take the reference population to be economics researchers. Let Y_{ij}^* be

[1] Generative-model outputs may vary across calls. Han (2025) discusses the use of separate LLM sessions, while Gao et al. (2026) treats a pretrained LLM and its prompting protocol as a fixed prediction rule.

the number of audience interactions in recording j that researcher i rates as negative in tenor. Thus μ_j is the mean number of audience interactions rated as negative in seminar j, averaging over the population of economics researchers. With $J_1 := \sum_{j=1}^{J} D_j$ and $J_0 := J - J_1$, the female–male contrast is

$$\beta_J = \frac{1}{J_1} \sum_{j:D_j=1} \mu_j - \frac{1}{J_0} \sum_{j:D_j=0} \mu_j = \Phi_J(a^{\Delta}), \qquad a_j^{\Delta} = J \left\{ \frac{D_j}{J_1} - \frac{1 - D_j}{J_0} \right\}.$$

This unadjusted contrast is a weighted corpus moment. If $\widetilde{D}_j$ residualizes presenter gender on prespecified seminar-series effects and fixed controls, the corresponding finite-corpus regression coefficient is also a weighted mean, with $a_j = J\widetilde{D}_j / \sum_{k=1}^{J} \widetilde{D}_k^2$. A natural g_j is the seminar-level count of interruptions classified by the authors' pretrained audio model as having negative tone.

More generally, linear regressions with μ_j as the outcome and observed corpus characteristics as regressors also fall into this class. For the seminar example, let $\widetilde{X}_j = (1, D_j, X_j')'$, where X_j contains the prespecified controls. Provided the regressor second-moment matrix is nonsingular, the least-squares coefficients are

$$\boldsymbol{\beta}_J = \arg\min_{\boldsymbol{b}} \frac{1}{J} \sum_{j=1}^{J} (\mu_j - \widetilde{X}_j' \boldsymbol{b})^2 = \left(\frac{1}{J} \sum_{j=1}^{J} \widetilde{X}_j \widetilde{X}_j' \right)^{-1} \left(\frac{1}{J} \sum_{j=1}^{J} \widetilde{X}_j \mu_j \right).$$

Perceived attractiveness as a regressor (Möbius and Rosenblat, 2006). The authors study the beauty premium in an experimental labor market. They have photographs of all participants rated on a five-point attractiveness scale by 50 high-school students from Tucumán and use the resulting beauty measure in regressions for worker confidence and productivity and for employers' wage assessments. To illustrate how our framework could be applied to a worker-level regression, let the corpus consist of photographs of the workers included in that regression. Then W_j is worker j's photograph and Y_{ij}^* is the attractiveness rating that evaluator i would assign to it. One natural reference population, matching the original evaluator pool, is high-school students in Tucumán. Thus $\mu_j = \mathbb{E}[Y_{ij}^*]$ is worker j's mean perceived attractiveness in this population.

Let R_j be a fixed worker outcome, such as reported confidence or realized productivity, let X_j contain the remaining observed regressors, and write $\widetilde{X}_j = (1, X_j', \mu_j)'$. The regression target under our framework is

$$\boldsymbol{\beta}_J = \arg\min_{\boldsymbol{b}} \frac{1}{J} \sum_{j=1}^{J} (R_j - \widetilde{X}_j' \boldsymbol{b})^2.$$

The document-level measurements enter these coefficients through corpus averages of μ_j, $X_j\mu_j$, $R_j\mu_j$, and μ_j^2; all remaining quantities are observed for the full corpus. The first three are weighted averages of μ_j. The last can be expressed using a pairwise measurement, since the product of responses from two independently drawn evaluators has expectation μ_j^2. If the regressor second-moment matrix is nonsingular, the coefficients are a smooth function of these corpus averages, as in equation (2.1).

A natural g_j is a frozen vision-model prediction of the reference-population mean rating under the same rating protocol. Its square, g_j^2, can serve as an auxiliary prediction for μ_j^2.

Employer evaluations of résumés (Kessler et al., 2019). The authors ask employers to rate hypothetical résumés on an ordered hiring-interest scale; their appendix also dichotomizes the rating at several cutoffs and estimates logit specifications for the resulting counterfactual-callback outcomes. Let W_j be the complete résumé, let Y_{ij}^* be employer i's hiring-interest rating, and, for a prespecified cutoff

c, define

$$B^*_{ij}(c) = \mathbf{1}\{Y^*_{ij} \geq c\}, \qquad \mu_j(c) = \mathbb{E}[B^*_{ij}(c)].$$

The reference population can be defined as employers recruiting Penn seniors in the relevant field, with separate populations for STEM and business/humanities/social-science recruiting when their evaluation criteria differ. Let Z_j collect the fixed résumé characteristics in the economic specification. Let $\Lambda(t) := \{1 + \exp(-t)\}^{-1}$ denote the logistic link. We define the logistic regression coefficients by

$$\boldsymbol{\beta}_J(c) = \arg\min_{\boldsymbol{b}} \frac{1}{J} \sum_{j=1}^{J} \big[-\mu_j(c) \log \Lambda(Z_j'\boldsymbol{b}) - \{1 - \mu_j(c)\} \log\{1 - \Lambda(Z_j'\boldsymbol{b})\} \big]. \tag{2.2}$$

Define

$$\boldsymbol{\Phi}_J(c) = \frac{1}{J} \sum_{j=1}^{J} Z_j \mu_j(c).$$

Each coordinate of $\boldsymbol{\Phi}_J(c)$ is a weighted first-order corpus moment. For a finite, locally unique minimizer of (2.2), the score equation is

$$\boldsymbol{\Phi}_J(c) - \frac{1}{J} \sum_{j=1}^{J} Z_j \Lambda\big(Z_j'\boldsymbol{\beta}_J(c)\big) = 0.$$

If its Jacobian with respect to $\boldsymbol{\beta}$ is nonsingular, the implicit-function theorem gives a locally continuously differentiable map H_J such that

$$\boldsymbol{\beta}_J(c) = H_J\{\boldsymbol{\Phi}_J(c)\}.$$

Although these applications report different economic parameters, their dependence on the reference-population response distributions is captured by a fixed-dimensional collection of weighted first-order and order-two corpus moments. These moments are therefore the common population objects developed in the sections that follow.

3 Annotation design

Machine predictions can be generated inexpensively for the full corpus, whereas human evaluations are costly and time-consuming. Recruiting fresh annotators for every annotation may also be impractical. Under a limited annotation budget, the researcher must therefore choose which documents to annotate, how many evaluations each receives, and how evaluators are assigned to documents.

3.1 Document sampling and the annotation budget

Throughout the analysis, annotation expenditure is measured by the number of requested labels. The document characteristics $X_{1:J}$, machine predictions $g_{1:J}$, and target weights $a_{1:J}$ are fixed. Using this information, the researcher specifies in advance the probability that each document j is sampled (its inclusion probability, $\pi_j \in (0, 1]$). The researcher also specifies how many evaluations it receives if sampled (its annotation depth, $m_j \in \{1, \ldots, \bar{m}\}$), where $\bar{m}$ is a fixed finite integer. These are the two document-level design choices. The inclusion probabilities govern the extensive margin: how much of the corpus is expected to be covered and which documents are more likely to be annotated. The annotation depths govern the intensive margin: how precisely each sampled document's measurement can be estimated.

Let $I_j = 1$ if document j is sampled and $I_j = 0$ otherwise.

(D1) Poisson document sampling. The indicators $\{I_j\}_{j=1}^{J}$ are mutually independent, with

$$I_j \sim \text{Bernoulli}(\pi_j), \qquad j = 1, \ldots, J.$$

Poisson sampling with unequal inclusion probabilities is standard in survey sampling (Särndal et al., 1992; Tillé, 2006). It is particularly useful here because it gives the researcher direct control over each document's marginal inclusion probability while preserving independence across document-selection decisions. This independence simplifies the variance analysis. The expected budget constraint $\sum_j \pi_j m_j = \bar{B}$ is also additive across documents, making it easier to optimize inclusion probabilities and annotation depths for later analysis.

This convenience comes with a limitation: the realized annotation expenditure B is random. A constraint on expected expenditure does not guarantee that realized expenditure stays within that budget. Such situation also arise in optimal experimental design and treatment-policy choice (Hahn et al., 2011; Sun, 2026). In our framework, an strict upper bound can nevertheless be enforced with asymptotically negligible distortion. Let $B_{\max}$ be the largest feasible number of labels and choose the expected budget below this cap so that, for example,

$$B_{\max} - \bar{B} \geq \sqrt{\bar{B} \log \bar{B}}.$$

Under the bounded-depth condition, Lemma 5 implies

$$\Pr(B > B_{\max}) \longrightarrow 0.$$

If the cap must never be exceeded, samples with $B > B_{\max}$ can be rejected and redrawn before annotation begins. The accepted-sample law has total variation distance $\Pr(B > B_{\max})$ from the original Poisson law, so the rejection step does not alter the asymptotic conclusions.

When the annotation budget must be fixed exactly, Online Appendix OA.2 develops a stratified simple-random-sampling alternative for these applications. A fixed number n_h of documents is sampled without replacement from each prespecified stratum h of size J_h, and every selected document in that stratum receives a common annotation depth m_h. Consequently,

$$B = \sum_{h=1}^{H} n_h m_h$$

is fixed by design. We refer to this as the stratified fixed-budget design, Design **(SFB)**. It sacrifices fully document-specific control of π_j and m_j in exchange for stratum-level control and an exact finite-sample annotation budget. Poisson sampling will remain the main sampling regime in this paper, while Design **(SFB)** provides a fixed-budget alternative for applications that require strict budget control.

3.2 Annotator allocation

After document sampling, the researcher must assign the B requested evaluations to workers. Each selected document j is evaluated by m_j distinct workers. We allow workers to evaluate multiple documents, but impose a fixed workload cap K.

Recruiting and managing annotators is itself costly, so it is desirable to use recruited workers up to their

workload cap. For $B > 0$, define

$$n_0 := \left\lceil \frac{B}{K} \right\rceil, \qquad r_B := B - K(n_0 - 1) \in \{1, \dots, K\}.$$

Our capacity-filled default uses n_0 workers: $n_0 - 1$ receive K documents and the remaining worker receives r_B. Thus n_0 is the smallest workforce compatible with the cap, and all but possibly one active worker are used to capacity. The cap K limits the number of documents evaluated by the same worker and hence the amount of dependence created by worker reuse.

Once the document and worker workloads have been specified, the remaining design choice is how to match documents to workers. We represent such a matching by a simple bipartite graph A_J, whose two vertex sets are the selected documents and the active workers. An edge (j, i) means that worker i evaluates document j. The document degrees therefore record the requested annotation depths, while the worker degrees record worker workloads. We impose simplicity on the graph, that is, the same worker does not evaluate a document more than once. Let M_j denote the set of workers assigned to evaluate document j, with $M_j = \varnothing$ when $I_j = 0$.

In the baseline design, workers are ex ante exchangeable: they are recruited from the same reference population, and the allocation rule does not use worker types or potential responses. Conditional on the prescribed document and worker workloads, there is, therefore, no reason to prefer one feasible matching over another. We implement this symmetry by drawing the allocation uniformly from all feasible matchings.

Formally, let

$$\boldsymbol{d}_J := (m_j)_{j \in \mathcal{S}_J}, \qquad \boldsymbol{k}_J^{\mathrm{cf}} := (\underbrace{K, \dots, K}_{n_0 - 1}, r_B)$$

denote the document- and worker-degree sequences. Both have degree sum B. For degree vectors $\boldsymbol{d}$ and $\boldsymbol{k}$, let

$$\mathfrak{A}_J(\boldsymbol{d}, \boldsymbol{k})$$

denote the class of labelled simple bipartite graphs on document vertices $\mathcal{S}_J$ and worker vertices $\{1, \dots, n_0\}$ having those degree sequences.

(RW1) Random-worker allocation. If $B > 0$ and the prescribed workloads are feasible,[2] then, conditional on $\mathcal{S}_J$,

$$A_J \sim \mathrm{Unif}\Big(\mathfrak{A}_J(\boldsymbol{d}_J, \boldsymbol{k}_J^{\mathrm{cf}})\Big).$$

The allocation can be generated using established sampling methods. A simple approach uniformly matches the requested evaluations to worker slots with the prescribed workloads, repeating the entire draw if any worker is assigned the same document twice. This produces an exact uniform allocation (**?**). When rejection becomes computationally costly, established swap algorithms provide an alternative: they approximate the uniform distribution by repeatedly proposing exchanges of assignments between workers while preserving workloads and preventing repeated worker–document assignments (Erdős et al., 2018). Appendix E.1 describes this implementation and shows that its approximation error can be made negligible for our asymptotic results.

Let $\mathcal{G}_J$ denote the information contained in the realized document sample and worker allocation, and let

[2] If $B > 0$ and these workloads are infeasible, we recruit B workers and assign each requested evaluation to a different worker. The probability that this alternative is needed tends to zero as $\bar{B} \to \infty$; see Appendix C.3. If $B = 0$, no workers are recruited and A_J is the empty graph.

$\mathcal{L}_J$ denote the information contained in all worker types and worker–document response shocks:

$$\mathcal{G}_J := \sigma(\mathcal{S}_J, A_J),$$
$$\mathcal{L}_J := \sigma(\{\theta_i\}_{i\geq 1}, \{\varepsilon_{ij} : i \geq 1,\ j \leq J\}).$$

(D2) Design exogeneity. The sigma-fields $\mathcal{G}_J$ and $\mathcal{L}_J$ are independent.

Assumption **(D2)** permits the design to use document-side information fixed before the current annotation study, but rules out allocation based on worker types, potential responses, or pair-specific shocks from the study.

Allocation under the exact-budget design Under Design **(SFB)**, the same capacity-filled allocation is conducted separately within each stratum. Stratum h contributes the fixed workload $B_h = n_h m_h$; conditional on its document sample, the allocation is drawn uniformly from the feasible graphs with document degree m_h and the corresponding capacity-filled worker degrees. Worker pools are disjoint and graph draws are independent across strata. Thus each stratum fills its assigned budget, all but possibly one active worker per stratum are used to capacity, and no worker is shared across strata.

4 Estimation and inference

Subsection 2.3 expressed the economic parameters of interest as smooth functions of weighted averages at the corpus level. The main text develops estimation and inference for the building block of such parameters:

$$\Phi_J(a) := \frac{1}{J}\sum_{j=1}^{J} a_j \mu_j,$$

The weighted averages built from the primary document-level measurements μ_j. Online Appendix OA.1 develops the corresponding results for two-response document measurements, while Online Appendix OA.3 combines a fixed number of mean and two-response moments and extend their joint limit theory to smooth and implicitly defined economic parameters using delta-method and implicit-function arguments.

The framework targets applications with reasonably large annotation budgets, typically comprising hundreds or thousands of labels, but much larger corpora containing thousands, tens of thousands, or more documents. Formally, we use the framework of finite-population asymptotics (Abadie et al., 2020, Section 3.1), under which

$$J \to \infty \qquad \text{and} \qquad \bar{B} \to \infty.$$

No limit is imposed on $\bar{B}/J$. In particular, the analysis permits the sparse regime $\bar{B}/J \to 0$, at any rate consistent with $\bar{B} \to \infty$, as well as regimes in which the annotation budget remains nonnegligible relative to the corpus size.

The researcher observes X_j, g_j, and a_j for every document in the corpus, but observes human evaluations only for documents selected for annotation. This section investigates the commonly used prediction-assisted estimator for $\Phi_J(a)$ and investigates its' statistical properties.

4.1 Prediction-assisted estimation and unbiasedness

For each selected document $j \in \mathcal{S}_J$, let

$$\bar{Y}_j := \frac{1}{m_j} \sum_{i \in M_j} Y_{ij}^*$$

denote its average observed evaluation. Set $\bar{Y}_j = 0$ when $I_j = 0$. Assumptions **(D2)** and **(RW0)** imply

$$\mathbb{E}\big[\bar{Y}_j \mid \mathcal{G}_J\big] = \mu_j \qquad \text{on } \{I_j = 1\}.$$

Thus, conditional on the realized sample and worker allocation, $\bar{Y}_j$ is an unbiased measurement of μ_j.

We estimate $\Phi_J(a) = J^{-1} \sum_{j=1}^J a_j \mu_j$ following a standard approach in survey sampling (Cassel et al., 1976; Särndal et al., 1992) that is also used in active statistical inference (Zrnic and Candès, 2024):

$$\widehat{\Phi}_J(a) = \frac{1}{J} \sum_{j=1}^J a_j g_j + \frac{1}{J} \sum_{j \in \mathcal{S}_J} \frac{a_j}{\pi_j} \big(\bar{Y}_j - g_j\big). \tag{4.1}$$

The first term applies the machine prediction to every document in the corpus. The second uses the annotated sample to estimate the corresponding full-corpus weighted prediction error, with inverse-probability weighting correcting for document selection. The predictions therefore provide an initial full-corpus estimate, while the human evaluations estimate the correction required to recover the target. If $g_j = 0$ for every j, equation (4.1) reduces to the ordinary Horvitz–Thompson estimator based on the observed document means.

Proposition 1 (Unbiasedness)**.** *Suppose Assumptions **(RW0)**, **(S1)**–**(S2)**, and **(D2)** hold, and the allocation is simple, so that each selected document is evaluated by m_j distinct workers. Under any document-sampling design with known positive first-order inclusion probabilities*

$$\Pr(I_j = 1) = \pi_j > 0, \qquad j = 1, \ldots, J,$$

for every finite J,

$$\mathbb{E}\Big[\widehat{\Phi}_J(a)\Big] = \Phi_J(a).$$

Proof. Design exogeneity and reference-population sampling give

$$\mathbb{E}\big[\bar{Y}_j - g_j \,\big|\, I_j = 1\big] = \mu_j - g_j.$$

Consequently,

$$\mathbb{E}\left[\frac{I_j}{\pi_j} a_j (\bar{Y}_j - g_j)\right] = \frac{a_j}{\pi_j} \Pr(I_j = 1)(\mu_j - g_j) = a_j(\mu_j - g_j).$$

Summing across documents and adding $J^{-1} \sum_{j=1}^J a_j g_j$ proves the result. □

The proof uses only the first-order inclusion probabilities and does not require independent document selection. Under the baseline Poisson design, these probabilities are the π_j specified in Assumption **(D1)**. Under Design **(SFB)**, a document in stratum h has inclusion probability $\pi_h = n_h / J_h$. Since $\rho_h = J_h / J$,

$$\frac{1}{J\pi_h} = \frac{\rho_h}{n_h},$$

so equation (4.1) reduces exactly to the stratified version in Online Appendix OA.2. Notice, The sampling law affects only the variance and subsequent inference, but not unbiasedness.

Equation (4.1) is also the fixed-corpus, unequal-probability analogue of the mean estimator in prediction-powered inference (Angelopoulos et al., 2023). The channel through which predictions can improve efficiency is the same. As discussed by Angelopoulos et al. (2023), predictions available for the full corpus allow the annotated sample to estimate only the prediction error. When g_j tracks μ_j well in the design-weighted sense, the residual correction is less variable, reducing document-sampling uncertainty. The predictions need not be unbiased, calibrated, or consistent; poor predictions may indeed increase this component of uncertainty, but cannot introduce bias.

The inferential challenge here, however, is substantially different. In standard model-assisted survey sampling and prediction-powered inference, the outcome is observed once a unit enters the labeled sample. Here, selecting document j does not reveal μ_j: its correction must itself be estimated from finitely many worker evaluations, and worker reuse can correlate these measurement errors across documents. Valid inference must therefore account jointly for document sampling, finite annotation depth, and the dependence induced by the allocation graph.

4.2 Asymptotic normality under worker reuse

The asymptotic argument begins with an decomposition of the estimation error. Let

$$r_j := \mu_j - g_j$$

denote the prediction residual for document j. Adding and subtracting the sampled residual correction gives

$$\widehat{\Phi}_J(a) - \Phi_J(a) = \underbrace{\frac{1}{J}\sum_{j=1}^{J}\left(\frac{I_j}{\pi_j} - 1\right) a_j r_j}_{T_J^s(a)} + \underbrace{\frac{1}{J}\sum_{j\in\mathcal{S}_J}\frac{a_j}{\pi_j}\left(\bar{Y}_j - \mu_j\right)}_{T_J^m(a)}. \tag{4.2}$$

The first term is the usual Horvitz–Thompson error applied to the prediction residual. As in model-assisted survey sampling and prediction-powered inference, it arises because the residual correction is observed only for the sampled documents. The second term is annotation measurement error and marks the main departure from those settings. Even for a sampled document, μ_j is not directly observed: the collected labels are finitely many draws from the reference population whose mean is μ_j. We analyze the two components separately before combining them.

As with other estimators built from inverse-probability weights, asymptotic normality requires some control over extreme design weights. In our setting, the inclusion probability π_j and annotation depth m_j determine expected expenditure $\pi_j m_j$ on document j. Dividing the expected annotation budget equally across documents would assign $\bar{B}/J$ labels to each document in expectation. Define

$$\lambda_j := \frac{J\pi_j m_j}{\bar{B}},$$

which compares document j's expected expenditure with the expenditure under this equal allocation.

Assumption 1 (Balanced spending)**.** *For some constant $c_\lambda \in (0, 1]$, independent of J,*

$$\lambda_j \geq c_\lambda, \qquad j = 1, \ldots, J. \tag{4.3}$$

Assumption 1 imposes a one-sided balance requirement: expected spending on each document must be at least a fixed fraction c_λ of the expenditure under equal allocation. Individual expected expenditures may be small, provided they satisfy this relative lower bound. The condition permits heterogeneous inclusion probabilities and annotation depths, including sparse sampling with $\bar{B}/J \to 0$, provided that $\bar{B} \to \infty$. Together with the maintained boundedness conditions and fixed worker capacity, this lower bound controls the inverse-probability weights and ensures that no individual document or worker dominates the normalized estimation error. These properties support the central limit theorem established below.

4.2.1 Document-sampling uncertainty

We first consider the sampling term, $T_J^s(a)$. Under Poisson sampling, the scaled variance of the first term in (4.2) is

$$v_{r,J}(a) := \bar{B}\,\mathrm{Var}\{T_J^s(a)\} = \frac{\bar{B}}{J^2}\sum_{j=1}^{J}\frac{1-\pi_j}{\pi_j}a_j^2 r_j^2. \tag{4.4}$$

The document-sampling CLT requires that no document receive enough inverse-probability weight to affect the limit on its own. Under uniformly bounded target weights, $|a_j| \le \bar{a}$, and annotation depths, $m_j \le \bar{m}$, balanced spending gives

$$\delta_J^{\mathrm{lev}}(a) := \max_{1\le j\le J}\frac{\sqrt{\bar{B}}}{J\pi_j}|a_j| = \max_j \frac{m_j|a_j|}{\lambda_j\sqrt{\bar{B}}} \le \frac{\bar{m}\bar{a}}{c_\lambda\sqrt{\bar{B}}} \longrightarrow 0. \tag{4.5}$$

The quantity $\delta_J^{\mathrm{lev}}(a)$ measures the largest document-level leverage after the $\sqrt{\bar{B}}$ normalization. With uniformly bounded prediction residuals, (4.5) implies the Lindeberg condition for the document-sampling term: no individual document makes a first-order contribution to its asymptotic distribution.

Lemma 1 (Document-sampling central limit theorem)**.** *Suppose documents are selected independently by Poisson sampling, $\bar{B} \to \infty$, Assumption 1 holds, and the target weights, annotation depths, and prediction residuals are uniformly bounded. If*

$$v_{r,J}(a) \longrightarrow v_r(a), \tag{4.6}$$

then

$$\sqrt{\bar{B}}\,T_J^s(a) \xrightarrow{d} \mathcal{N}(0, v_r(a))\,.$$

Equation (4.6) is Condition **(RW4-S)**. Conditions of this form, requiring a normalized design variance or covariance matrix to converge, are standard in finite-population design-based asymptotics; see, for example, Francisco and Fuller (1991, Condition 3), Deville and Särndal (1992, p. 379), and Boistard et al. (2017, Condition **(HT2)**). In our setting, the condition requires only that the design-weighted second moment of the prediction residuals converge to a finite limit $v_r(a) \ge 0$. The condition follows, for example, from an ordinary law of large numbers under a well-behaved document superpopulation. It can equally hold for a deterministic sequence of fixed corpora, for example when the corpus has finitely many document strata whose proportions and within-stratum weighted second moments stabilize. The lemma is the standard triangular-array Lindeberg–Feller result; its proof is given in Appendix C.1.

4.2.2 Measurement uncertainty and the high-level limit

The measurement term requires more care. Reusing a worker can induce correlation across the documents evaluated by that worker. Following the usual remedy in clustered-data settings (Hansen and Lee, 2019), we rewrite the measurement term as a sum of worker contributions. Let $\mathcal{J}_i$ denote the documents

assigned to worker i, and define

$$\zeta_i(a) := \frac{\sqrt{\bar{B}}}{J} \sum_{j \in \mathcal{J}_i} \frac{a_j}{\pi_j m_j} \big(Y_{ij}^* - \mu_j\big), \qquad \sqrt{\bar{B}}\, T_J^m(a) = \sum_i \zeta_i(a).$$

Conditional on $\mathcal{G}_J$, the worker contributions are independent and have mean zero. Balanced spending, together with bounded worker capacity and the maintained boundedness conditions, ensures that no worker dominates the sum. Appendix C.1 therefore gives a conditional Gaussian approximation for the complete measurement term.

However, this is not quite enough. The remaining issue is most directly illustrated by the conditional variance. Define

$$\mu_j^\theta(\theta) := \mathbb{E}\big[Y_{ij}^* \mid \theta_i = \theta\big], \qquad \tau_{jk} := \mathrm{Cov}_\theta\big\{\mu_j^\theta(\theta), \mu_k^\theta(\theta)\big\},$$

and let $N_{jk} := |M_j \cap M_k|$. As shown in the appendix, the graph-specific measurement variance is

$$\begin{aligned} v_{\varepsilon,J}(a) &:= \bar{B}\, \mathrm{Var}\{T_J^m(a) \mid \mathcal{G}_J\} \\ &= \frac{\bar{B}}{J^2} \sum_{j=1}^{J} \frac{I_j a_j^2 \nu_j}{\pi_j^2 m_j} + \frac{\bar{B}}{J^2} \sum_{j \neq k} \frac{I_j I_k N_{jk}}{\pi_j \pi_k m_j m_k} a_j a_k \tau_{jk}, \end{aligned} \tag{4.7}$$

where the second sum is over ordered document pairs. The first component captures variation across fresh evaluations of the same document. The second is the overlap component generated by worker reuse. In general, this variance—especially its overlap component—depends on the realized sample and allocation graph through quantities such as N_{jk}.

Conditional on $\mathcal{G}_J$, the worker-level argument gives a centered Gaussian approximation for $\sqrt{\bar{B}}\, T_J^m(a)$ with variance $v_{\varepsilon,J}(a)$. The sampling term $T_J^s(a)$, however, is then a fixed and generally nonzero shift. Integrating over $\mathcal{G}_J$ therefore averages Gaussian approximations with different locations and variances. The resulting limit need not be Gaussian, even though the sampling and measurement terms can be shown to be uncorrelated.

This resembles the random-environment problem in Andrews (2005), where random conditional variances generate mixed-normal limits. The additional complication here is that the environment also contains the document-selection realization and hence fixes the separate sampling shift. Standardizing by the graph-specific measurement variance alone therefore does not remove the random location.

Martingale central limit theory suggests a way forward. A standard condition in this literature is that the conditional variance converges to a nonrandom limit (Hall and Heyde, 1980, Chapter 3). We impose a similar variance-stabilization condition **(RW4-M)**:

$$v_{\varepsilon,J}(a) \xrightarrow{p} v_\varepsilon(a), \tag{4.8}$$

where $v_\varepsilon(a)$ is a finite, nonrandom, nonnegative constant. To understand its role here, recall that the normalized measurement term is approximately Gaussian conditional on the document sample and worker assignment, with conditional mean zero. Since a Gaussian distribution is determined by its mean and variance, this approximation depends on the realized sample and assignment only through its' conditional variance. Variance stabilization makes the approximating distribution asymptotically the same across realized document samples and worker assignments, so the measurement term becomes asymptotically independent of both. Since the document-sampling term is determined by the selected documents, the two terms have independent Gaussian limits. Their sum therefore has a Gaussian limit with variance

$v_r(a) + v_\varepsilon(a)$. Appendix C.1 formalizes this argument.

Theorem 1 (Asymptotic normality under worker reuse)**.** *Maintain Assumptions **(RW0)**, **(RW2)**, **(S1)**–**(S2)**, and **(D1)**–**(D2)**. Suppose $J \to \infty$, $\bar{B} \to \infty$, the target weights and annotation depths are uniformly bounded, and the allocation is simple with every worker assigned at most a fixed number K of documents. Suppose also that Assumption 1 holds. If* (4.6) *and* (4.8) *hold and*

$$v_r(a) + v_\varepsilon(a) > 0,$$

then

$$\sqrt{\bar{B}}\left\{\widehat{\Phi}_J(a) - \Phi_J(a)\right\} \xrightarrow{d} \mathcal{N}(0, v_r(a) + v_\varepsilon(a)). \tag{4.9}$$

Because $B/\bar{B} \xrightarrow{p} 1$, the same conclusion holds with $\sqrt{B}$ in place of $\sqrt{\bar{B}}$.

It should be noted, although similar, condition **(RW4-M)** is fundamentally different from Condition **(RW4-S)**. Condition **(RW4-S)** concerns convergence of the deterministic document-sampling variance along the fixed corpus sequence, whereas Condition **(RW4-M)** requires the random conditional measurement variance to converge to a nonrandom limit despite variation in the sampled documents and worker assignments. Its validity therefore depends on the allocation rule as well as document sampling. At this stage, we impose this variance behavior as a high-level condition. The next subsection shows that, under additional regularity conditions, our uniform allocation scheme ensures variance stabilization.

4.2.3 Variance stabilization under uniform allocation

We begin with two document-side stability conditions analogous to Condition **(RW4-S)**. Suppose, for $v_{\text{diag}}(a; \pi, m) \geq 0$ and $\tau(a) \geq 0$, that

$$\frac{1}{J}\sum_{j=1}^{J} \frac{a_j^2 \nu_j}{\lambda_j} \longrightarrow v_{\text{diag}}(a; \pi, m), \qquad \frac{1}{J^2}\sum_{j \neq k} a_j a_k \tau_{jk} \longrightarrow \tau(a), \tag{4.10}$$

where the second sum is over ordered document pairs. As with Condition **(RW4-S)**, these limits may hold along a regular deterministic sequence of fixed corpora or follow from ordinary and U-statistic laws of large numbers under a well-behaved optional document superpopulation. Although the finite-J off-diagonal average can be negative, any limit $\tau(a)$ is nonnegative: the covariance matrix $(\tau_{jk})_{j,k \leq J}$ is positive semidefinite, and its omitted diagonal contribution is $O(J^{-1})$ under the maintained bounds.

Conditions (4.10) alone do not ensure variance stabilization because they do not determine which document pairs share workers. Even with bounded worker loads, an allocation rule can systematically concentrate overlaps on particular document classes and make the conditional measurement variance depend persistently on the realized design. Appendix C.2 gives such a counterexample.

Uniform sampling over all feasible bipartite graphs provides this much-needed allocation regularity. The proof relies on two averaging arguments. First, capacity filling fixes the overall amount of worker reuse, while uniform allocation randomizes which document pairs share workers. On the high-probability event that the capacity-filled allocation is feasible, the fixed-degree graph result of McKay (2010, Theorem 2.2) gives, uniformly for selected $j \neq k$,

$$\mathbb{E}[N_{jk} \mid \mathcal{S}_J] = (K-1)\frac{m_j m_k}{B} + O\Big(\frac{m_j m_k}{B^2}\Big). \tag{4.11}$$

Thus pairs receiving more annotations are proportionally more likely to share a worker. The proof also shows that the resulting weighted aggregate of worker overlaps concentrates around these expected

values.

Second, the factors $m_j m_k$ in (4.11) cancel the corresponding annotation-depth adjustments in (4.7). The remaining average over selected document pairs converges under Poisson sampling to the corpus-wide limit $\tau(a)$, while the diagonal component converges to $v_{\text{diag}}(a;\pi,m)$ following standard Horvitz–Thompson concentration arguments.

Theorem 2 (Variance stabilization under uniform allocation)**.** *Maintain Assumptions **(RW0)**–**(RW2)**, **(S1)**–**(S2)**, and **(D1)**–**(D2)**. Suppose documents are selected independently by Poisson sampling, $J \to \infty$, $\bar{B} \to \infty$, the target weights and annotation depths are uniformly bounded, worker capacity K is fixed, and Assumption 1 holds. Suppose also that allocation uses the uniform capacity-filled mechanism of Subsection 3.2 and that* (4.10) *holds. Then*

$$v_{\varepsilon,J}(a) \xrightarrow{p} v_{\text{diag}}(a;\pi,m) + (K-1)\tau(a). \tag{4.12}$$

Combining the theorem with the high-level Gaussian result gives the following operational conclusion.

Corollary 1 (Operational Gaussian limit)**.** *Suppose Condition **(RW4-S)** and the conditions of Theorem 2 hold. The conclusion of Theorem 1 then holds with*

$$v_\varepsilon(a) = v_{\text{diag}}(a;\pi,m) + (K-1)\tau(a),$$

provided

$$v_r(a) + v_{\text{diag}}(a;\pi,m) + (K-1)\tau(a) > 0.$$

It should be noted that uniform allocation is one transparent sufficient mechanism, not a necessary condition. Other allocation schemes can be treated case by case when their overlap structure and the corresponding, potentially allocation-dependent, fixed-array limits jointly imply Condition **(RW4-M)**. Under an optional document-superpopulation model, those limits may often be verified using ordinary or U-statistic laws of large numbers, but regularity of the allocation mechanism must still be established case by case.

4.3 Variance estimation and feasible inference

Throughout this subsection, we restrict attention to designs with $m_j \geq 2$ for every document (we explain this restriction later in the subsection). If the document-level means $\{\mu_j\}_{j=1}^J$ were known, the prediction residuals $r_j = \mu_j - g_j$ would also be known, and the document-sampling variance $v_{r,J}(a)$ could be calculated directly from (4.4). Conditional on the sampled documents and worker assignment, the measurement variance $v_{\varepsilon,J}(a)$ could be estimated by centering responses using the known document means and summing squared worker contributions. This worker-level "cluster robust" type calculation accounts for worker reuse. The difficulty is that the document means needed for these calculations are unknown and must themselves be estimated from a small number of responses.

Although $\bar{Y}_j$ is unbiased for μ_j, substituting it for μ_j in squared residuals and covariance products can introduce bias. Because the number of labels per document remains bounded, this bias may not disappear as more documents are annotated. To address this problem, we follow the leave-out logic of Kline et al. (2020) and construct unbiased estimators of the moments needed for variance estimation. The remaining mean-zero estimation errors can then average out under the maintained regularity conditions.

4.3.1 Document-sampling variance

For the document-sampling component, the finite-label bias takes a particularly simple form. For a selected document,

$$\mathbb{E}\big[(\bar{Y}_j - g_j)^2 \,\big|\, \mathcal{G}_J\big] = r_j^2 + \frac{\nu_j}{m_j}. \tag{4.13}$$

The additional term is the finite-label variance of the observed document mean. Because m_j remains bounded, this term need not vanish as the annotation budget grows, and its weighted aggregate can generate first-order bias in the estimated document-sampling variance.

This expression suggests a direct correction. For $j \in \mathcal{S}_J$, let

$$\widehat{\sigma}_j^2 := \frac{1}{m_j - 1} \sum_{i \in M_j} \big(Y_{ij}^* - \bar{Y}_j\big)^2$$

denote the usual within-document sample variance, and define

$$\widehat{r}_j^2 := (\bar{Y}_j - g_j)^2 - \frac{\widehat{\sigma}_j^2}{m_j}. \tag{4.14}$$

Since $\widehat{\sigma}_j^2$ is unbiased for ν_j, $\widehat{r}_j^2$ is unbiased for r_j^2. Equivalently, $\widehat{r}_j^2$ can be written as the average product of each worker's label and the mean of the remaining workers' labels, after both have been centered at g_j. Thus, (4.14) is a document-level leave-out correction of the type used to estimate quadratic objects in Kline et al. (2020).

Substituting the corrected residual square into the Horvitz–Thompson variance formula gives

$$\widehat{v}_{r,J}(a) := \frac{\bar{B}}{J^2} \sum_{j \in \mathcal{S}_J} \frac{1 - \pi_j}{\pi_j^2} a_j^2 \widehat{r}_j^2. \tag{4.15}$$

4.3.2 Measurement variance

The diagonal term in the measurement variance is already handled by $\widehat{\sigma}_j^2$. It remains to estimate the overlap contribution $N_{jk}\tau_{jk}$ for document pairs that share workers. For a worker i shared by documents j and k, the corresponding oracle covariance product is

$$\big(Y_{ij}^* - \mu_j\big)\big(Y_{ik}^* - \mu_k\big),$$

whose conditional expectation is τ_{jk}. Because μ_j and μ_k are unknown, centering instead by the full document means would use worker i's own labels in both centering terms. We remove this self-centering by defining

$$\bar{Y}_{j,-i} := \frac{1}{m_j - 1} \sum_{\ell \in M_j \setminus \{i\}} Y_{\ell j}^*.$$

When documents j and k share more than one worker, however, their leave-out means can still contain the other $N_{jk} - 1$ shared workers. The following factor removes the resulting additional covariance:

$$\kappa_{jk}^{\mathrm{loo}} := \begin{cases} \dfrac{(m_j - 1)(m_k - 1)}{(m_j - 1)(m_k - 1) + N_{jk} - 1}, & N_{jk} \geq 1, \\ 0, & N_{jk} = 0. \end{cases}$$

For distinct selected documents j and k, define

$$\widehat{\Gamma}_{jk} := \kappa_{jk}^{\text{loo}} \sum_{i \in M_j \cap M_k} \big(Y_{ij}^* - \bar{Y}_{j,-i}\big)\big(Y_{ik}^* - \bar{Y}_{k,-i}\big).$$

With this adjustment, as shown in the appendix,

$$\mathbb{E}[\widehat{\Gamma}_{jk} \mid \mathcal{G}_J] = N_{jk}\tau_{jk}.$$

Combining the diagonal and overlap terms gives the measurement-variance estimator

$$\widehat{v}_{\varepsilon,J}(a) := \frac{\bar{B}}{J^2}\left[\sum_{j \in \mathcal{S}_J} \frac{a_j^2 \widehat{\sigma}_j^2}{\pi_j^2 m_j} + \sum_{\substack{j,k \in \mathcal{S}_J \\ j \neq k}} \frac{a_j a_k \widehat{\Gamma}_{jk}}{\pi_j \pi_k m_j m_k}\right], \tag{4.16}$$

where the second sum is over ordered document pairs, as in (4.7). The estimator of the total asymptotic variance is

$$\widehat{v}_J(a) := \widehat{v}_{r,J}(a) + \widehat{v}_{\varepsilon,J}(a). \tag{4.17}$$

Under the maintained response, design-exogeneity, Poisson-sampling, and simple-allocation assumptions, the two corrections are unbiased $m_j \geq 2$:

$$\mathbb{E}[\widehat{v}_{r,J}(a)] = v_{r,J}(a), \qquad \mathbb{E}[\widehat{v}_{\varepsilon,J}(a) \,|\, \mathcal{G}_J] = v_{\varepsilon,J}(a). \tag{4.18}$$

Define

$$\bar{v}_{\varepsilon,J}(a) := \mathbb{E}[v_{\varepsilon,J}(a)].$$

Applying the law of iterated expectations to average over the document sample and worker assignment therefore gives

$$\mathbb{E}[\widehat{v}_J(a)] = v_{r,J}(a) + \bar{v}_{\varepsilon,J}(a) = \bar{B}\ \mathrm{Var}\Big\{\widehat{\Phi}_J(a) - \Phi_J(a)\Big\}. \tag{4.19}$$

Unbiasedness alone does not imply consistency. Conditional on the document sample and worker assignment, responses from different workers are independent. Bounded workloads and annotation depths ensure that changing one worker's responses affects only a limited number of terms in the variance estimators. Balanced spending makes these individual contributions small as the budget grows. The Efron–Stein inequality translates this limited influence into an $O(\bar{B}^{-1})$ mean-square bound for the error due to worker responses. Independent Poisson sampling controls the remaining error due to document selection in the sampling-variance estimator. Appendix D gives the formal bounds.

Neither finite-sample unbiasedness nor this concentration argument requires uniform allocation; graph simplicity and bounded worker load are the only allocation-specific requirements. Uniform allocation enters through variance stabilization, which makes $v_{\varepsilon,J}(a)$ converge to the nonrandom limit used for studentization. Consequently, $\widehat{v}_J(a)$ consistently estimates the asymptotic variance in Corollary 1. This retains the crucial modular structure of the results: if researchers desire, or are compelled, to use a different type of allocation mechanism, then, as long as variance stabilization can be established, the variance estimator facilitates feasible inference.

Because the leave-out corrections involve subtraction, the raw estimator $\widehat{v}_J(a)$ is not constrained to be nonnegative in finite samples. This is a familiar possibility for dependence-robust covariance estimators involving subtractive corrections, such as multiway-clustered covariance estimators (Cameron et al.,

2011). For any deterministic sequence $\eta_J > 0$ with $\eta_J \to 0$, define

$$\widehat{v}_J^+(a) := \max\{\widehat{v}_J(a), \eta_J\}.$$

This floor is only a numerical finite-sample safeguard.

Theorem 3 (Feasible inference)**.** *Suppose the conditions of Corollary 1 hold and $m_j \geq 2$ for every document. Then*

$$\widehat{v}_{r,J}(a) - v_{r,J}(a) \xrightarrow{p} 0, \qquad \widehat{v}_{\varepsilon,J}(a) - v_{\varepsilon,J}(a) \xrightarrow{p} 0,$$

and hence

$$\widehat{v}_J(a) \xrightarrow{p} v_r(a) + v_{\text{diag}}(a; \pi, m) + (K-1)\tau(a).$$

If

$$v_r(a) + v_{\text{diag}}(a; \pi, m) + (K-1)\tau(a) > 0,$$

then

$$\frac{\sqrt{\bar{B}}\,\{\widehat{\Phi}_J(a) - \Phi_J(a)\}}{\sqrt{\widehat{v}_J^+(a)}} \Rightarrow \mathcal{N}(0,1).$$

The same conclusion holds with $\sqrt{B}$ in place of $\sqrt{\bar{B}}$.

Proof. Proposition 4 gives componentwise consistency under the stated conditions. Corollary 1, variance stabilization, and Slutsky's theorem then give the studentized limit; the replacement of $\sqrt{\bar{B}}$ by $\sqrt{B}$ follows from Lemma 5. □

Relative to Corollary 1, feasible inference requires only that each document receive at least two labels, so that the within-document variance and leave-out means are well defined. Strict positivity is the usual nondegeneracy condition: it holds whenever document sampling or annotation contributes nonvanishing first-order uncertainty in the target direction. Under this condition, the probability that the numerical safeguard binds converges to zero.

4.4 Important extensions

4.4.1 Pairwise response measurements

Online Appendix OA.1 extends the analysis to the order-two measurements $\vartheta_j(\ell)$ introduced in Subsection 2.2. For a selected document, the natural estimator is the within-document U-statistic

$$\widehat{\vartheta}_j = \binom{m_j}{2}^{-1} \sum_{\substack{i,r \in M_j \\ i<r}} \ell_j(Y_{ij}^*, Y_{rj}^*).$$

Substituting $\widehat{\vartheta}_j$ for the document mean in the same prediction-assisted Horvitz–Thompson construction gives an unbiased estimator of the corresponding corpus target. Its document-sampling component is handled exactly as in the mean case.

The new issue is measurement uncertainty. The Hoeffding decomposition separates the error in $\widehat{\vartheta}_j$ into a linear component indexed by individual workers and a canonical component indexed by pairs of workers. The linear component has the same structure as that of the mean, so the same overlap-averaging argument applies.

The canonical component requires additional care. Because annotation depth remains bounded, its contribution to the within-document measurement variance cannot be discarded. Canonical degeneracy means that each pairwise term has mean zero conditional on either worker's responses. Together with independence across workers, this implies that, conditional on the sample and assignment, canonical terms from different documents have zero covariance unless the same pair of workers evaluates both. Under our allocation scheme, such overlaps are sufficiently rare that their combined contribution to the normalized measurement variance is asymptotically negligible.

One additional complication is that the measurement error can no longer be written as a sum of independent worker contributions, so the worker-block central limit theorem used for the mean is unavailable. Nevertheless, balanced spending and bounded document and worker degrees permit an application of the dependency-graph central limit theorem of Janson (1988, Theorem 2), which provides a conditional Gaussian approximation for the measurement term. With variance stabilization established, the measurement and document-sampling components are then combined using the same conditioning argument as in the mean case.

The appendix consequently establishes Gaussian and feasible inference for bounded symmetric pairwise kernels, including response variance and categorical disagreement. Point estimation and the central limit theorem require $m_j \geq 2$. The feasible variance estimator uses products formed from disjoint worker pairs and therefore requires $m_j \geq 4$.

4.4.2 Stratified exact-budget inference

Online Appendix OA.2 develops the corresponding inference theory under Design **(SFB)**, which fixes the number of sampled documents and their common annotation depth within each stratum, so that total expenditure is fixed by design.

The estimator retains the same decomposition into document-sampling and measurement components. The sampling component is now a stratified finite-population average, so the Poisson-sampling central limit theorem is replaced by a stratumwise finite-population central limit theorem under the usual Hájek no-dominance condition (Li and Ding, 2017, Theorem 1). Independence across the fixed number of strata then combines the stratum-specific limits. Conditional on the samples and allocation graphs, the measurement component remains a sum of independent worker contributions, so the conditional worker-block argument used for the mean continues to apply.

The remaining step is variance stabilization. Within each stratum, simple random sampling makes the sampled document and document-pair moments concentrate around their finite-population counterparts. Common annotation depth makes selected documents exchangeable under uniform allocation, which spreads the capacity-fixed overlap opportunities symmetrically across document pairs. Separate worker pools eliminate cross-stratum measurement dependence. These properties make the allocation dependent measurement variance converge to its stratum-level limit. The sampling and measurement components can then be combined using the same conditioning argument as under the baseline design.

The appendix also constructs a consistent feasible variance estimator using the same leave-out moments, together with a within-stratum correction for the finite-label noise in observed prediction residuals. Pairwise measurements are also covered.

4.4.3 Joint and smooth-functional inference

Online Appendix OA.3 extends the scalar results to a fixed-dimensional vector containing both mean and order-two corpus moments. A mean coordinate based on a bounded single-response transform h_j

can be represented exactly by the symmetric kernel $\{h_j(y_1) + h_j(y_2)\}/2$, allowing mean and genuinely pairwise coordinates, including their mixed covariances, to be treated within one construction. Every fixed linear combination of the moment vector is then a scalar target covered by the preceding results. Applying the scalar result to every fixed linear combination and invoking the Cramér–Wold device, we obtain a joint Gaussian limit. The covariance matrix is then constructed from the difference between scalar variance estimators for the coordinate and coordinate-sum directions using the variance-of-a-sum identity. Standard delta-method and implicit-function arguments consequently provide feasible inference.

4.4.4 Inference beyond the realized corpus

The fixed-corpus perspective does not preclude generalization to a document population when a credible document model is available. Let $\boldsymbol{\beta}_J$ denote the conditional-mean regression coefficient for the realized corpus and let $\boldsymbol{\beta}_0$ denote its document-population counterpart. Writing $B_J^\circ = \bar{B}$ under the Poisson design and $B_J^\circ = B$ under the exact-budget design,

$$\sqrt{B_J^\circ}(\widehat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}_0) = \sqrt{B_J^\circ}(\widehat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}_J) + \sqrt{\frac{B_J^\circ}{J}}\,\sqrt{J}(\boldsymbol{\beta}_J - \boldsymbol{\beta}_0).$$

The first term is uncertainty from the annotation study, while the second reflects variation in the realized corpus. If $\sqrt{J}(\boldsymbol{\beta}_J - \boldsymbol{\beta}_0) = O_p(1)$ and $B_J^\circ/J \to 0$, document-population variation is negligible at the annotation-study scale, so the existing fixed-corpus procedure already provides inference for $\boldsymbol{\beta}_0$. If instead $B_J^\circ/J \to \kappa \in (0, \infty)$, document heterogeneity may be first order. Online Appendix OA.4 then constructs a sandwich covariance estimator combining a standard heteroskedasticity-robust document component with the required sampling and worker-overlap corrections. Under Poisson sampling, this has a particularly simple form: the outer product of the complete prediction-assisted score already incorporates document heterogeneity, document selection, and diagonal annotation noise, so only the off-diagonal worker-overlap correction must be added.

5 Choosing document inclusion probabilities and annotation depths

We now use the preceding variance characterization to choose the document inclusion probabilities $\{\pi_j\}_{j=1}^J$ and annotation depths $\{m_j\}_{j=1}^J$ subject to the expected label budget

$$\sum_{j=1}^{J} \pi_j m_j = \bar{B}.$$

The analysis is restricted to the scaler weighted mean target

$$\Phi_J(a) = \frac{1}{J}\sum_{j=1}^{J} a_j \mu_j$$

under the baseline Poisson document-sampling design. We leave the optimal allocation for pairwise response measurements and stratified exact-budget designs to future research.

All comparisons are restricted to the admissible design class maintained for inference. In particular, design choices are chosen before the final annotation outcomes are observed, documents are selected independently, worker allocation follows the uniform capacity-filled mechanism with fixed capacity K,

and annotation depths are uniformly bounded. We also maintain balanced spending,

$$c_\lambda \le \frac{J\pi_j m_j}{\bar{B}}, \qquad j = 1, \dots, J,$$

together with $\bar{B} \to \infty$ and the corpus-array convergence and nondegeneracy conditions required by the preceding inference results. For feasible mean inference, we impose $2 \le m_{\min} \le m_j \le \bar{m}$. Documents with zero target weight can be removed from the design frame, with the remaining weights rescaled so that the target is unchanged.

5.1 Oracle benchmark

To expose the allocation trade-offs, suppose initially that the prediction residual

$$r_j := \mu_j - g_j$$

and the fresh-worker response variance ν_j are known. For designs satisfying the conditions stated above, our variance-stabilization result gives

$$v_{r,J}(a) + v_{\varepsilon,J}(a) = \frac{\bar{B}}{J^2} \sum_{j=1}^{J} a_j^2 \left\{ \frac{1-\pi_j}{\pi_j} r_j^2 + \frac{\nu_j}{\pi_j m_j} \right\} + v_{\mathrm{ov},J}(a) + o_p(1), \tag{5.1}$$

$$v_{\mathrm{ov},J}(a) := \frac{K-1}{J^2} \sum_{j \ne k} a_j a_k \tau_{jk}.$$

The variance-stabilization result shows that, within our design class and for fixed K, the worker-overlap contribution to the asymptotic variance does not depend on the document inclusion probabilities or annotation depths. Minimizing the asymptotic variance is therefore equivalent to minimizing the remaining terms in (5.1), with the worker-overlap contribution omitted.

We approximate the allocation problem by solving its continuous relaxation, treating annotation depths m_j as continuous variables between $m_{\min}$ and $\bar{m}$. Let

$$s_j := \pi_j m_j$$

denote the expected expenditure on document j. At a fixed value of s_j, that document's contribution to the allocation criterion is

$$a_j^2 \left\{ \frac{\nu_j}{s_j} + r_j^2 \left(\frac{1}{\pi_j} - 1 \right) \right\}. \tag{5.2}$$

The measurement component depends only on total expected expenditure, whereas the document-sampling component decreases as π_j increases. For a fixed expected expenditure s_j, increasing the inclusion probability while reducing annotation depth strictly reduces the document's contribution to the variance whenever $r_j \ne 0$. An optimal implementation of s_j therefore expands coverage before increasing depth:

$$\pi_j^\star(s_j) = \min\left\{1, \frac{s_j}{m_{\min}}\right\}, \qquad m_j^\star(s_j) = \frac{s_j}{\pi_j^\star(s_j)}. \tag{5.3}$$

Thus depth remains at $m_{\min}$ until document j is included with probability one; only then does additional expenditure increase its depth. It should be noted, at this point, this is just a within-document statement. It does not require every document to reach full coverage before depth increases for any document.

Using the coverage-first rule, we profile out π_j and m_j, leaving an optimization problem in expected

expenditures s_j alone. (5.3) into (5.2) gives the contribution $a_j^2\phi_j(s_j)$, where

$$\phi_j(s) = \begin{cases} \dfrac{\nu_j + m_{\min} r_j^2}{s} - r_j^2, & s \le m_{\min}, \\ \dfrac{\nu_j}{s}, & s > m_{\min}. \end{cases} \tag{5.4}$$

The remaining choice is how to divide expected expenditure across documents. The allocation problem therefore becomes

$$\begin{aligned} \min_{\{s_j\}} \quad & \sum_{j=1}^{J} a_j^2 \phi_j(s_j) \\ \text{subject to} \quad & \sum_{j=1}^{J} s_j = \bar{B}, \\ & c_\lambda \frac{\bar{B}}{J} \le s_j \le \bar{m}, \qquad j = 1, \dots, J. \end{aligned} \tag{5.5}$$

The spending bounds preserve the balanced-spending restriction and the maximum annotation depth. The objective is convex, including where its two branches meet at $m_{\min}$. Its Karush–Kuhn–Tucker (KKT) conditions characterize the optimal allocation and yield the rule in the following proposition.

Proposition 2 (Structure of the first-order oracle allocation)**.** *Suppose* $0 < \bar{B} \le J\bar{m}$ *and* $0 < c_\lambda \le 1$. *An optimal solution to* (5.5) *can be implemented using the coverage-first rule in* (5.3). *There is a common multiplier* $\xi \ge 0$ *on the budget constraint. When* $\xi > 0$*, documents whose optimal expenditure is neither at a spending bound nor at the full-coverage threshold* $m_{\min}$ *satisfy*

$$s_j^\star = \begin{cases} |a_j| \sqrt{\dfrac{\nu_j + m_{\min} r_j^2}{\xi}}, & s_j^\star < m_{\min}, \\ |a_j| \sqrt{\dfrac{\nu_j}{\xi}}, & s_j^\star > m_{\min}. \end{cases}$$

The multiplier and any expenditures at the threshold or spending bounds are jointly chosen so that $\sum_j s_j^\star = \bar{B}$.

Appendix F gives the complete constrained characterization, including the profiled convex program, documents placed exactly at full coverage, and binding spend bounds.

The proposition gives a generalized Neyman allocation with two economically distinct regimes. Before full coverage, expected expenditure is proportional to

$$|a_j| \sqrt{\nu_j + m_{\min} r_j^2},$$

because additional expenditure reduces uncertainty from both document selection and finite annotation. Documents receive more resources when they matter more for the target, when the machine prediction is less accurate, or when worker responses are more variable. After full coverage, sampling uncertainty has disappeared, so expenditure is proportional to

$$|a_j| \sqrt{\nu_j},$$

and only annotation uncertainty determines the return to additional depth.

5.2 Practical implementation

The oracle quantities r_j^2 and ν_j are unobserved before annotation. A pilot with repeated annotations, or other comparable annotated data, can be used to estimate these quantities for pilot documents and fit prediction rules based on document characteristics observed throughout the corpus. This parallels two-stage experimental designs that use first-wave outcomes to choose second-wave assignment probabilities or strata (Hahn et al., 2011; Tabord-Meehan, 2023). Applying the fitted rules to the full corpus provides predicted difficulty measures for the allocation problem. Negative fitted variances or squared residuals, if any, are truncated at zero. The researcher then solves the oracle program using these predictions, subject to the same budget, depth, and balanced-spending restrictions imposed above.

Without reliable outcome information for ranking document difficulty, a simple default is to assign every document the same expected response variance and the same expected squared prediction error. Let $\mathcal{I}_J$ denote the information available when the design is chosen, and let $\mathbb{E}_0$ denote the researcher's planning expectation. Our planning convention is

$$\mathbb{E}_0[\nu_j \mid \mathcal{I}_J] = \nu_J^0, \qquad \mathbb{E}_0[r_j^2 \mid \mathcal{I}_J] = q_J^0, \qquad j = 1, \ldots, J,$$

where ν_J^0 and q_J^0 are finite, nonnegative values that are not both zero.[3] It should be stressed that these planning expectations do not assert that the actual document-specific variances or squared prediction errors are equal. They are used only to choose the allocation and are not assumptions required for inference.

Under this convention, suppose that the target weights are not too uneven and that the labeling budget is small relative to the corpus, specifically,

$$c_\lambda \leq \frac{J|a_j|}{\sum_{k=1}^J |a_k|} \quad \text{for every } j, \qquad \frac{\bar{B}}{Jm_{\min}} \frac{J \max_j |a_j|}{\sum_{k=1}^J |a_k|} \leq 1. \tag{5.6}$$

Under this condition, the allocation never assigns more than the minimum admissible annotation depth to any document. The resulting rule is

$$s_j^0 = \bar{B} \frac{|a_j|}{\sum_{k=1}^J |a_k|}, \qquad m_j^0 = m_{\min}, \qquad \pi_j^0 = \frac{s_j^0}{m_{\min}}. \tag{5.7}$$

Thus, every sampled document receives $m_{\min}$ annotations, while documents with larger absolute target weights are sampled with higher probability. Condition (5.6) is fully determined by the target weights, the budget, and the corpus size, and can therefore be checked before annotation begins. Appendix F.1 gives the complete result for this common-expected-difficulty default.

Two especially interpretable examples are the following. For the corpus mean, $a_j = 1$, so the rule spends equally across documents. For a difference between two group means, the absolute target weights sum to the same amount within each group. Each group therefore receives one half of the expected budget, with greater expenditure per document in the smaller group. In both cases, expenditure on a document is first used to increase its inclusion probability while keeping $m_j = m_{\min}$. Annotation depth increases only after that document's inclusion probability reaches one.

Finally, we emphasize again that inaccurate or misspecified document difficulty models may reduce efficiency by leading to suboptimal inclusion probabilities and annotation depths. They do not invalidate inference when the resulting design remains within the maintained admissible class.

[3] Appendix F.1 establishes the allocation result under the weaker condition that $\mathbb{E}_0[\nu_j + m_{\min} r_j^2 \mid \mathcal{I}_J]$ has the same positive finite value across documents. We state the separate conditions for ease of interpretation.

6 Calibrated Simulation: Measuring Toxicity

This section uses an empirically anchored calibrated simulation to examine three features of the proposed framework: inference under worker reuse, finite-sample performance with few sampling units and workers, and the efficiency gains from targeted annotation design. We use the Wikipedia Talk Labels: Toxicity corpus, which contains 159,686 comments and 1,598,289 binary toxicity judgments from 4,301 workers (Wulczyn et al., 2017; Thain et al., 2017). The corpus links worker-identified judgments to comment text and metadata recording contributor registration status, posting year, and talk-page namespace. These data are used to construct fixed comment-level toxicity probabilities and calibrate the synthetic worker population. For each comment, we also attach a fixed machine prediction g_j, defined as the offensive-class probability from the pretrained `cardiffnlp/twitter-roberta-base-offensive` model (Barbieri et al., 2020). The prediction is computed once and held fixed across replications; no model is trained, fine-tuned, or cross-fitted within the simulation.

6.1 Calibrated simulation population

We construct an empirically anchored, semi-synthetic population from the Wikipedia Detox corpus, comprising 159,686 comments and approximately 1.6 million binary annotations from 4,301 workers. For each comment, we retain its estimated toxicity probability μ_j, registration status, and fixed Cardiff prediction. Importantly, the Cardiff prediction is not re-trained or cross-fitted within the simulation. It is used directly off-the-shelf.

Worker responses follow a random-coefficients logit model:

$$Y_{ij} \mid u_i, \beta_i \sim \text{Bernoulli}(p_{ij}), \tag{6.1}$$

$$\text{logit}(p_{ij}) = \alpha_j + u_i + x_j^\top \beta_i. \tag{6.2}$$

The vector x_j contains five comment characteristics. Three are indicators supplied with the dataset: whether the comment appears on a user-talk page, belongs to the blocked-user sample, or was posted by an unregistered contributor. We add two measures constructed from the comment text: Length is measured by $-\log(1 + \text{word count})$, so larger values indicate shorter comments. Abusive-word presence is an indicator for whether the comment contains at least one word classified as abusive in the published dictionary of Wiegand et al. (2018).

We specify the worker intercept and slopes as

$$u_i = \sigma Z_i + \delta(A_i - q), \qquad \beta_{ik} = cZ_i + \tau V_{ik}, \quad k = 1, \ldots, 5,$$

where $Z_i, V_{i1}, \ldots, V_{i5}$ are independent standard normal draws and A_i is an independent Bernoulli draw with probability q.

The worker intercept u_i captures differences in the overall tendency to label comments toxic, while the five slopes in β_i allow workers to respond differently to comment characteristics. Each worker's intercept and slopes are drawn jointly once and retained for every comment assigned to that worker.

To capture the skewness and upper tail of workers' average differences from other reviewers, we allow a share q of workers to belong to a class whose average intercept is higher by δ. Within each class, σ controls variation in workers' overall tendency to label comments toxic.

The random slopes capture differences in responses to particular types of comments that cannot be represented by a worker intercept alone. Through the shared draw Z_i, the parameter c allows workers'

responses to comment characteristics to be associated with their overall tendency to label comments toxic. The separate draws V_{ik}, scaled by τ, allow further variation in each slope.

For each candidate pair (q, δ), we fit (σ, c, τ) to covariance comparisons across different groups of comments reviewed by the same worker. We compare the resulting specifications using the standard deviation, skewness and excess kurtosis of workers' average differences from other reviewers. Throughout calibration, the comment intercepts α_j are recomputed so that each comment's response probability, averaged over workers, remains μ_j. Appendix **??** reports the parameter values, the calibration procedure and the comparison of observed and simulated statistics.

6.2 Annotation uncertainty under worker overlap

We consider two corpus-level parameters. The first is the finite-corpus mean toxicity probability,

$$\theta_M = \frac{1}{J} \sum_{j=1}^{J} \mu_j.$$

Corpus-wide averages of this kind parallel text-based measures such as the economic policy uncertainty index of Baker et al. (2016), which aggregates newspaper content into a period-level measure.

The second parameter is the difference in mean toxicity between comments posted by unregistered and registered contributors,

$$\theta_\Delta = \frac{1}{J_U} \sum_{j:G_j=1} \mu_j - \frac{1}{J_R} \sum_{j:G_j=0} \mu_j,$$

where $G_j = 1$ denotes an unregistered contributor and J_U and J_R are the corresponding group sizes. An analogous text-based group contrast appears in Wu (2020), who compares mean counts of professional and personal terms in EJMR posts about women with those in posts about men. In the present application, the contrast measures whether toxic content is disproportionately concentrated among unregistered contributions, which may itself be of interest, as it could inform the allocation of moderation resources.

Each sampled comment receives $m_j = 2$ labels from distinct workers. Comments are sampled uniformly for the corpus mean and in proportion to the absolute target weights for the registration contrast, thereby dividing the expected annotation budget equally between the two groups as prescribed in Subsection 5.2. We consider expected budgets of $\bar{B} = 7{,}500$ and $15{,}000$, holding worker capacity fixed at $K = 125$. These designs correspond to approximately 60 and 120 workers, respectively; realized worker counts vary with the Poisson sample.

Table 6.1 compares two variance calculations applied to the same Cardiff-assisted point estimate. The full estimator accounts for document sampling, finite annotation depth, and covariance between comments labeled by the same worker. The *collapsed-label* calculation first averages the annotations for each comment and then treats the resulting comment-level outcome as fixed and error-free.

Coverage from the full estimator is close to the nominal level in both designs, whereas the collapsed-label calculation is substantially too optimistic, particularly for the corpus mean. For the mean, same-worker covariances accumulate because all target weights have the same sign. The opposing weights in the registration contrast partially cancel the common worker-harshness component. This explain the disparity between the undcoverage observed in the two cases. Importnatly, Holding worker capacity fixed at $K = 100$, doubling both the expected annotation budget and the approximate workforce leaves collapsed-label undercoverage essentially unchanged.

Table 6.1: Coverage under worker overlap

		Coverage (%)		
$(\bar{B}, K)$	Target	Full	Collapsed label	Full/Collapsed variance
(7,500, 125)	Corpus mean	93.2	54.9	6.77
(7,500, 125)	Registration contrast	94.5	82.4	2.07
(15,000, 125)	Corpus mean	94.5	54.8	6.92
(15,000, 125)	Registration contrast	94.5	82.4	2.16

Notes: Results are based on 5,000 replications and nominal 95% intervals. Each sampled comment receives two labels from distinct workers, and K is the maximum number of comments assigned to one worker. The final column is the ratio of the average full variance estimate to the average collapsed-label variance estimate. Both calculations use the same point estimate.

Table 6.2: Finite-sample stress test with package-level count outcomes

$\bar{B}$	Workers	Target	Full cov.	Collapsed cov.
240	30	Package mean	0.935	0.642
240	30	Registration contrast	0.938	0.839
480	60	Package mean	0.944	0.609
480	60	Registration contrast	0.947	0.793

Notes: Each specification uses 5,000 Monte Carlo replications, two distinct ratings per sampled package, and $K = 8$. Each target receives the full stated budget $\bar{B}$. "Full" uses the proposed variance correction. "Collapsed" treats the aggregated human label as noise-free.

6.3 Finite-sample stress test

We next examine performance when both the number of sampling units and the number of annotators are small. We aggregate the 159,686 Wikipedia Detox comments into a fixed population of 1,004 approximately equal, registration-status-pure packages of 159–160 comments, using an outcome-blind ordering within registration groups. The package is both the sampling and annotation unit, and the observed outcome is the number of its comments labeled toxic. Each sampled package is rated by two distinct workers, and each worker rates at most $K = 8$ packages. We consider expected package-rating budgets of $\bar{B} = 240$ and 480 per target, corresponding to 120 and 240 sampled packages in expectation and approximately 30 and 60 workers. The package mean count target uses uniform Poisson sampling, whereas the unregistered-minus-registered count contrast uses Poisson sampling proportional to inverse group size.

This construction closely mirrors the high-level structure of the online-seminar application in Dupas et al. (2026). Their online sample contains 994 talks, and the outcome is a talk-level count of interruptions aggregated from lower-level speaking events. The mean interruption level provides a good summary statistics, while the female-minus-male presenter contrast is the substantive comparison.

Table 6.2 reports results from 5,000 Monte Carlo replications. The Full procedure exhibits modest undercoverage with approximately 30 workers, with coverage moving closer to nominal with approximately 60 workers. Its estimated variances closely track the empirical Monte Carlo variance. By contrast, treating the collapsed package label as noise-free produces substantial undercoverage: coverage is approximately 61–64% for the package mean and 79–84% for the registration contrast. Increasing the annotation budget does not resolve this undercoverage.

6.4 Efficiency of the annotation design

Table 6.3 compares four allocation designs for the registration contrast at $\bar{B} = 15{,}000$ and $K = 150$. Each design is evaluated using the same expected main-stage annotation budget.

Table 6.3: Efficiency of alternative annotation designs

	Variance ($\times 10^5$)		Relative
Design	Empirical	Estimated	efficiency
Uniform	9.417	9.304	1.000
Target-weighted	7.025	6.816	1.340
Full oracle	6.009	6.092	1.567
Pilot-estimated	6.651	6.520	1.416

Notes: Results are based on 5,000 replications. Estimated variance is the average full variance estimate. Relative efficiency is the empirical variance under uniform sampling divided by that under the indicated design.

The full oracle reduces empirical variance by 40.5% relative to uniform sampling, a substantial precision gain at the same expected annotation budget. The oracle allocation is also informative about how this gain is achieved: it sets $m_j = m_{\min} = 2$ for every sampled comment. The entire improvement therefore comes from reallocating sampling probabilities across comments rather than increasing annotation depth. This provides a concrete numerical illustration of the coverage-before-depth rule in (5.3).

We next consider the feasible allocations introduced in Subsection 5.2. The target-weighted rule uses only the fixed target weights and requires no pilot annotation data. It reduces variance by 25.4% and captures 70.2% of the reduction achieved by the full oracle. We then estimate the oracle difficulty measure using a fixed historical pilot of 1,000 comments, each with two observed labels. The pilot comments are excluded from the analysis population for all four designs, and their 2,000 labels are separate from the main-stage budget. The pilot observations and resulting allocation are held fixed across replications. The pilot-estimated design reduces variance by 29.4% and captures 81.2% of the oracle gain. Across all four designs, average estimated variance closely tracks empirical variance, with estimated-to-empirical ratios between 0.970 and 1.014. This again verified the robustness of our inference result to allocation mechanism.

7 Empirical Application: Taking the Fed at its Word

In this section, we apply our method to the setting studied by Shapiro and Wilson (2022b), who relate the sentiment of Federal Open Market Committee (FOMC) deliberations to economic conditions to investigate central-bank objectives. The corpus contains 1,676 speaker-meeting documents from 42 speakers at 97 meetings during 2000–2011. The authors measure sentiment using a dictionary-based net-negativity score: the share of negative words minus the share of positive words, classified using the Loughran–McDonald dictionaries. Their quadratic specification with the full control set can be written as

$$N_{st} = \alpha_s + \theta\pi_t + \delta\pi_t^2 + z_t'\gamma + \varepsilon_{st},$$

where N_{st} is the net negativity of speaker s at meeting t, α_s denotes a speaker fixed effect, and π_t is the average of current-quarter and one-quarter-ahead Greenbook forecasts of core PCE inflation. The vector z_t collects the unemployment gap and real GDP growth, their squares, and financial controls: the preceding three-month S&P 500 return, the previous month's Shiller price–earnings ratio, VIX and short-term Treasury-yield volatility. Under the authors' interpretation of negativity as a proxy for the central bank's loss, positive curvature implies a loss-minimizing inflation rate $\pi^* = -\theta/(2\delta)$.

A limitation of dictionary-based sentiment measures is that word counts may miss the context in which language is used. For example, a discussion of possible risks need not express the speaker's own negative assessment. Human readers can interpret these distinctions when assessing sentiment. We define the reference population as English-fluent adult Prolific participants who satisfy at least one of three con-

ditions: (i) their highest degree is in accounting, economics or finance; (ii) their primary employment is in finance; or (iii) they have previously invested in the stock market. Index each speaker–meeting document by a single index j, and let $Y_{ij}^* \in \{1, 2, 3, 4, 5\}$ denote the rating that reader i would assign to document j, ranging from very positive (1) to very negative (5), with neutral sentiment coded as 3. We replace the dictionary score with the reference-population mean of the human response, using the regression target described in Section 2, Example 2.

The study was originally planned as a single collection with the larger combined budget. Because the full intended funding was unavailable at the outset, data collection instead proceeded in two waves with separate annotation budgets. The expected budgets were $\bar{B}_1 = 800$ and $\bar{B}_2 = 400$ individual document ratings, with realised budgets of $B_1 = 842$ and $B_2 = 384$, respectively. Each sampled document received two ratings, and each reader was assigned at most eight documents. Within each wave, documents were assigned to readers using the uniform allocation scheme in Section 3. Further details of recruitment, document sampling and inference are provided in the appendix.

7.1 Results

In our preferred specification, we collapse ratings 1 and 2 to 1, retain rating 3, and collapse ratings 4 and 5 to 5. Annotator feedback indicated difficulty distinguishing degrees of positive or negative sentiment, suggesting that differences between ratings 1 and 2, or between ratings 4 and 5, may partly reflect response noise. Results using the original five-point scale, together with alternative nonlinear specifications, are retained as robustness checks in the appendix. Let C denote this recoding. For each document j, we define

$$\mu_j = \mathbb{E}\big[C(Y_{ij}^*)\big],$$

where the expectation is taken over readers drawn from the reference population. Thus, μ_j is the expected collapsed rating of document j.

We use two auxiliary predictors. The first is constructed using the GPT-6 Astra model, which is prompted to predict how eligible readers would rate each document. The model returns a probability for each of the five possible responses; the exact prompt is provided in the appendix. We form its prediction by averaging the response values using these probabilities, applying the same recoding to the response values when using the collapsed outcome. The second predictor is based on the original authors' dictionary score. To put this score on the 1–5 rating scale, we order all $J = 1,676$ documents from the lowest to the highest dictionary score and assign each document its position in this ordering. Documents with the same score receive the average of the positions they occupy. If r_j denotes this position, the transformed score is

$$g_j^{\text{dict}} = 1 + 4\frac{r_j - 0.5}{J}.$$

Thus, documents with higher dictionary negativity receive higher predicted ratings, with the lowest and highest scores mapped close to 1 and 5. Fractional values are retained in both linear specifications.

Table 7.1 reports the prediction-assisted estimates. The two estimators imply minimum-negativity inflation rates of 1.678% and 1.788%, both close to the 1.691% estimate in the full-controls quadratic specification of Shapiro and Wilson (2022b, Table 1, column (4)). Both yield negative linear and positive quadratic inflation coefficients, individually significant at the nominal 10% level. However, Both turning-point confidence intervals include 2%, which contradicts with the original conclusion of the paper. However, it is difficult to comment on meaningfulness of this results, as the precision might increase with more labels collected, evetually support the authors claim.

Table 7.1: Inflation estimates using human-assessed sentiment

Auxiliary predictor	Inflation θ	Inflation squared δ	Assisted turning point	90% delta-method interval	Predictor-only turning point
GPT-6 Astra	−1.313	0.391	1.678	[1.322, 2.033]	1.326
	(0.610)	(0.172)			
Authors' score	−1.261	0.353	1.788	[1.259, 2.317]	1.727
	(0.705)	(0.193)			

Notes: The outcome is the collapsed 1–3–5 rating. Coefficient estimates are prediction-assisted, with standard errors in parentheses. Both specifications include the economic controls and speaker fixed effects described above. Standard errors account for document sampling, annotation variability and reader overlap. Turning points are measured in inflation percentage units. Predictor-only turning points use the same regression specification on full-corpus predictions, without human correction. The authors' score is rescaled using full-corpus midpoint ranks. Intervals are nominal pointwise 90% delta-method confidence intervals.

The GPT-6 Astra predictor alone implies a lower turning point of 1.326%. Correction using human ratings raises this estimate by 0.352 percentage points to 1.678%, illustrating that the predictor-only and reference-population projections can differ even when a predictor is useful for estimation, and human correction is meaningful in this estimation exercise.

Relative to assistance using the authors' transformed dictionary score, GPT-6 Astra reduces the estimated standard errors of the linear and quadratic inflation coefficients by 13.4% and 11.0%, respectively. This pattern is consistent with a precision benefit from a predictor that accounts for linguistic context. The relatively modest gains in coefficient precision also suggest that the dictionary score remains a competitive auxiliary predictor in this application.

The overlap correction is small for the two inflation coefficients in the GPT-6 Astra-assisted specification, accounting for 1.5% and 2.4% of their total estimated variances, respectively. However, this is not the case for every coefficient. Two notable examples are the unemployment gap and the Shiller price–earnings ratio, for which overlap accounts for 22.4% and 35.2% of total estimated variance.

Table 7.2: Contribution of reader overlap to estimated uncertainty

Coefficient	SE without overlap	SE with overlap	Overlap share of total variance	SE increase
Inflation	0.6057	0.6103	1.5%	0.8%
Inflation squared	0.1700	0.1720	2.4%	1.2%
Unemployment gap	0.1575	0.1788	22.4%	13.5%
Shiller P/E	0.0177	0.0219	35.2%	24.2%

Notes: GPT-6 Astra auxiliary predictor and the collapsed 1–3–5 outcome, with the same controls and speaker effects as Table 7.1. Let $\widehat{V}$ denote total estimated variance and $\widehat{O}$ its overlap component. The overlap share is $100\widehat{O}/\widehat{V}$. Standard errors with and without overlap are $\sqrt{\widehat{V}}$ and $\sqrt{\widehat{V}-\widehat{O}}$, respectively. The SE increase is $100\{\sqrt{\widehat{V}/(\widehat{V}-\widehat{O})}-1\}$. Only the overlap component is removed; point estimates and other covariance components are unchanged.

References

Alberto Abadie, Susan Athey, Guido W. Imbens, and Jeffrey M. Wooldridge. Sampling-based versus design-based uncertainty in regression analysis. *Econometrica*, 88(1):265–296, 2020. doi: 10.3982/ECTA12675.

Donald W. K. Andrews. Cross-section regression with common shocks. *Econometrica*, 73(5):1551–1585, 2005. doi: 10.1111/j.1468-0262.2005.00629.x.

Anastasios N. Angelopoulos, Stephen Bates, Clara Fannjiang, Michael I. Jordan, and Tijana Zrnic. Prediction-powered inference. *Science*, 382(6671):669–674, 2023. doi: 10.1126/science.adi6000.

Elliott Ash and Stephen Hansen. Text algorithms in economics. *Annual Review of Economics*, 15: 659–688, 2023. doi: 10.1146/annurev-economics-082222-074352.

Scott R. Baker, Nicholas Bloom, and Steven J. Davis. Measuring economic policy uncertainty. *The Quarterly Journal of Economics*, 131(4):1593–1636, 2016. doi: 10.1093/qje/qjw024.

Francesco Barbieri, Jose Camacho-Collados, Leonardo Neves, and Luis Espinosa Anke. TweetEval: Unified benchmark and comparative evaluation for tweet classification. In *Findings of the Association for Computational Linguistics: EMNLP 2020*, pages 1644–1650. Association for Computational Linguistics, 2020. doi: 10.18653/v1/2020.findings-emnlp.148. URL `https://aclanthology.org/2020.findings-emnlp.148/`.

Patrick Billingsley. *Probability and Measure*. John Wiley & Sons, New York, 3 edition, 1995.

Hélène Boistard, Hendrik P. Lopuhaä, and Anne Ruiz-Gazen. Functional central limit theorems for single-stage sampling designs. *The Annals of Statistics*, 45(4):1728–1758, 2017. doi: 10.1214/16-AOS1507.

Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration Inequalities: A Nonasymptotic Theory of Independence*. Oxford University Press, Oxford, 2013. doi: 10.1093/acprof:oso/9780199535255.001.0001.

Marcy Burchfield, Henry G. Overman, Diego Puga, and Matthew A. Turner. Causes of sprawl: A portrait from space. *The Quarterly Journal of Economics*, 121(2):587–633, 2006. doi: 10.1162/qjec.2006.121.2.587.

A. Colin Cameron, Jonah B. Gelbach, and Douglas L. Miller. Robust inference with multiway clustering. *Journal of Business & Economic Statistics*, 29(2):238–249, 2011. doi: 10.1198/jbes.2010.07136.

Claes-Magnus Cassel, Carl-Erik Särndal, and Jan H. Wretman. Some results on generalized difference estimation and generalized regression estimation for finite populations. *Biometrika*, 63(3):615–620, 1976. doi: 10.1093/biomet/63.3.615.

Melissa Dell. Deep learning for economists. *Journal of Economic Literature*, 63(1):5–58, 2025. doi: 10.1257/jel.20241733.

Leon Derczynski, Kalina Bontcheva, and Ian Roberts. Broad Twitter corpus: A diverse named entity recognition resource. In *Proceedings of COLING 2016, the 26th International Conference on Computational Linguistics: Technical Papers*, pages 1169–1179. The COLING 2016 Organizing Committee, 2016. URL `https://aclanthology.org/C16-1111/`.

Jean-Claude Deville and Carl-Erik Särndal. Calibration estimators in survey sampling. *Journal of the American Statistical Association*, 87(418):376–382, 1992. doi: 10.1080/01621459.1992.10475217.

Dave Donaldson and Richard Hornbeck. Railroads and American economic growth: A "market access" approach. *The Quarterly Journal of Economics*, 131(2):799–858, 2016. doi: 10.1093/qje/qjw002.

Jefferson Duarte, Stephan Siegel, and Lance Young. Trust and credit: The role of appearance in peer-to-peer lending. *The Review of Financial Studies*, 25(8):2455–2484, 2012. doi: 10.1093/rfs/hhs071.

Pascaline Dupas, Amy Handlan, Alicia Sasser Modestino, Muriel Niederle, Mateo Seré, Haoyu Sheng, Justin Wolfers, and Seminar Dynamics Collective. Gender differences in economics seminars. *American Economic Review*, 116(2):749–789, 2026. doi: 10.1257/aer.20241718.

Rick Durrett. *Probability: Theory and Examples*. Cambridge University Press, Cambridge, 5 edition, 2019. doi: 10.1017/9781108591034.

Péter L. Erdős, Tamás Róbert Mezei, István Miklós, and Dániel Soltész. Efficiently sampling the realizations of bounded, irregular degree sequences of bipartite and directed graphs. *PLOS ONE*, 13(8): e0201995, 2018. doi: 10.1371/journal.pone.0201995.

Adam Fisch, Joshua Maynez, R. Alex Hofer, Bhuwan Dhingra, Amir Globerson, and William W. Cohen. Stratified prediction-powered inference for effective hybrid evaluation of language models. In *Advances in Neural Information Processing Systems*, volume 37, 2024. doi: 10.52202/079017-3541.

Carol A. Francisco and Wayne A. Fuller. Quantile estimation with a complex survey design. *The Annals of Statistics*, 19(1):454–469, 1991. doi: 10.1214/aos/1176347993.

Wayne Gao, Sukjin Han, and Annie Liang. How well do LLMs predict human behavior? a measure of their pretrained knowledge, 2026.

Matthew Gentzkow and Jesse M. Shapiro. What drives media slant? evidence from U.S. daily newspapers. *Econometrica*, 78(1):35–71, 2010. doi: 10.3982/ECTA7195.

Matthew Gentzkow, Bryan Kelly, and Matt Taddy. Text as data. *Journal of Economic Literature*, 57 (3):535–574, 2019. doi: 10.1257/jel.20181020.

Mitchell L. Gordon, Michelle S. Lam, Joon Sung Park, Kayur Patel, Jeffrey T. Hancock, Tatsunori Hashimoto, and Michael S. Bernstein. Jury learning: Integrating dissenting voices into machine learning models. In *Proceedings of the 2022 CHI Conference on Human Factors in Computing Systems*, pages 1–19. Association for Computing Machinery, 2022. doi: 10.1145/3491102.3502004.

Yuriy Gorodnichenko, Tho Pham, and Oleksandr Talavera. The voice of monetary policy. *American Economic Review*, 113(2):548–584, 2023. doi: 10.1257/aer.20220129.

Hessel Haagsma, Johan Bos, and Malvina Nissim. MAGPIE: A large corpus of potentially idiomatic expressions. In *Proceedings of the Twelfth Language Resources and Evaluation Conference*, pages 279–287. European Language Resources Association, 2020. URL `https://aclanthology.org/2020.lrec-1.35/`.

Jinyong Hahn, Keisuke Hirano, and Dean Karlan. Adaptive experimental design using the propensity score. *Journal of Business & Economic Statistics*, 29(1):96–108, 2011. doi: 10.1198/jbes.2009.08161.

Peter Hall and C. C. Heyde. *Martingale Limit Theory and Its Application*. Academic Press, New York, 1980.

Sukjin Han. Mining causality: AI-assisted search for instrumental variables, 2025.

Bruce E. Hansen and Seojeong Lee. Asymptotic theory for clustered samples. *Journal of Econometrics*, 210(2):268–290, 2019. doi: 10.1016/j.jeconom.2019.02.001.

Stephen Hansen, Peter John Lambert, Nicholas Bloom, Steven J. Davis, Raffaella Sadun, and Bledi Taska. Remote work across jobs, companies, and space. NBER Working Paper 31007, National Bureau of Economic Research, 2023. URL https://www.nber.org/papers/w31007.

J. Vernon Henderson, Tanner Regan, and Anthony J. Venables. Building the city: From slums to a modern metropolis. *The Review of Economic Studies*, 88(3):1157–1192, 2021. doi: 10.1093/restud/rdaa042.

Svante Janson. Normal convergence by higher semiinvariants with applications to sums of dependent random variables and random graphs. *The Annals of Probability*, 16(1):305–312, 1988. doi: 10.1214/aop/1176991903.

Judd B. Kessler, Corinne Low, and Colin D. Sullivan. Incentivized resume rating: Eliciting employer preferences without deception. *American Economic Review*, 109(11):3713–3744, 2019. doi: 10.1257/aer.20181714.

Patrick Kline, Raffaele Saggio, and Mikkel Sølvsten. Leave-out estimation of variance components. *Econometrica*, 88(5):1859–1898, 2020. doi: 10.3982/ECTA16410.

Xinran Li and Peng Ding. General forms of finite population central limit theorems with applications to causal inference. *Journal of the American Statistical Association*, 112(520):1759–1769, 2017. doi: 10.1080/01621459.2017.1295865.

Brendan D. McKay. Subgraphs of random graphs with specified degrees. In *Proceedings of the International Congress of Mathematicians 2010*, volume IV, pages 2489–2501, New Delhi, 2010. Hindustan Book Agency. doi: 10.1142/9789814324359_0155.

Markus M. Möbius and Tanya S. Rosenblat. Why beauty matters. *American Economic Review*, 96(1):222–235, 2006. doi: 10.1257/000282806776157515.

Jerzy Neyman. On the two different aspects of the representative method: The method of stratified sampling and the method of purposive selection. *Journal of the Royal Statistical Society*, 97(4):558–625, 1934. doi: 10.1111/j.2397-2335.1934.tb04184.x.

Michael Patriksson. A survey on the continuous nonlinear resource allocation problem. *European Journal of Operational Research*, 185(1):1–46, 2008. doi: 10.1016/j.ejor.2006.12.006.

Barbara Plank. The "problem" of human label variation: On ground truth in data, modeling and evaluation. In *Proceedings of the 2022 Conference on Empirical Methods in Natural Language Processing*, pages 10671–10682. Association for Computational Linguistics, 2022. doi: 10.18653/v1/2022.emnlp-main.731. URL https://aclanthology.org/2022.emnlp-main.731/.

Peter Potash, Adam Ferguson, and Timothy J. Hazen. Ranking passages for argument convincingness. In *Proceedings of the 6th Workshop on Argument Mining*, pages 146–155. Association for Computational Linguistics, 2019. doi: 10.18653/v1/W19-4517.

Carl-Erik Särndal, Bengt Swensson, and Jan Wretman. *Model Assisted Survey Sampling*. Springer, New York, 1992.

Adam Hale Shapiro and Daniel J. Wilson. Taking the Fed at its word: A new approach to estimating central bank objectives using text analysis. *The Review of Economic Studies*, 89(5):2768–2805, 2022a. doi: 10.1093/restud/rdab094.

Adam Hale Shapiro and Daniel J. Wilson. Taking the Fed at its word: A new approach to estimating

central bank objectives using text analysis. *The Review of Economic Studies*, 89(5):2768–2805, 2022b. doi: 10.1093/restud/rdab094. URL https://doi.org/10.1093/restud/rdab094.

Alan Stuart. A simple presentation of optimum sampling results. *Journal of the Royal Statistical Society: Series B (Methodological)*, 16(2):239–241, 1954. doi: 10.1111/j.2517-6161.1954.tb00165.x.

Liyang Sun. Empirical welfare maximization with constraints. *Journal of Econometrics*, 253:106169, 2026. doi: 10.1016/j.jeconom.2025.106169.

Max Tabord-Meehan. Stratification trees for adaptive randomisation in randomised controlled trials. *The Review of Economic Studies*, 90(5):2646–2673, 2023. doi: 10.1093/restud/rdac083.

Nithum Thain, Lucas Dixon, and Ellery Wulczyn. Wikipedia talk labels: Toxicity. figshare dataset, version 2, 2017. URL https://figshare.com/articles/dataset/Wikipedia_Talk_Labels_Toxicity/4563973.

Yves Tillé. *Sampling Algorithms*. Springer, New York, 2006. doi: 10.1007/0-387-34240-0.

Michael Wiegand, Josef Ruppenhofer, Anna Schmidt, and Clayton Greenberg. Inducing a lexicon of abusive words: A feature-based approach. In *Proceedings of the 2018 Conference of the North American Chapter of the Association for Computational Linguistics: Human Language Technologies, Volume 1 (Long Papers)*, pages 1046–1056, New Orleans, Louisiana, 2018. Association for Computational Linguistics. doi: 10.18653/v1/N18-1095. URL https://aclanthology.org/N18-1095/.

Alice H. Wu. Gender bias among professionals: An identity-based interpretation. *The Review of Economics and Statistics*, 102(5):867–880, 2020. doi: 10.1162/rest_a_00877.

Ellery Wulczyn, Nithum Thain, and Lucas Dixon. Ex machina: Personal attacks seen at scale. In *Proceedings of the 26th International Conference on World Wide Web*, pages 1391–1399. International World Wide Web Conferences Steering Committee, 2017. doi: 10.1145/3038912.3052591.

Tijana Zrnic and Emmanuel J. Candès. Active statistical inference. In *Proceedings of the 41st International Conference on Machine Learning*, volume 235 of *Proceedings of Machine Learning Research*, pages 62993–63010. PMLR, 2024. URL https://proceedings.mlr.press/v235/zrnic24a.html.

Appendix

A Conditioning and probability conventions

We adopt a fixed-corpus asymptotic framework, analogous to the fixed-design and finite-population conventions in Kline et al. (2020, Section 1) and Abadie et al. (2020, Section 3.1). For each J, the corpus $W_{J,1:J}$ and all quantities determined before document sampling, including $\{X_{Jj}, a_{Jj}, g_{Jj}, \pi_{Jj}, m_{Jj}\}_{j=1}^{J}$, are treated as nonrandom. For different J, they form a deterministic sequence of finite arrays satisfying the assumptions made in the paper. For tractability, we suppress dependence on J in the notation when there is no ambiguity. Quantities derived from the fixed corpus and the maintained worker-response law, such as μ_j, ν_j, and τ_{jk}, are also nonrandom.

For each J, let $(\Omega_J, \mathscr{F}_J, \mathbb{P}_J)$ support all final-study random quantities, including document selection, worker allocation, and annotation outcomes. Let $\mathbb{E}_J$ and Var_J denote expectation and variance under $\mathbb{P}_J$. Unless stated otherwise, stochastic orders, probability limits, and weak-convergence statements refer to the corresponding row laws as $J \to \infty$. For this construction, no coupling across rows needs to be assumed.

Equivalently, all probabilities and expectations can be understood as conditional on the corpus and on the pre-study information used to construct the fixed quantities above. In particular, the predictions $\{g_j\}_{j=1}^{J}$ and the design rules may be estimated or constructed using auxiliary or pilot data available before the final study. Throughout the analysis, we condition on this information and treat the resulting values and rules as nonrandom components of the row-J array.

B Key decomposition and budget control

We maintain Assumptions **(RW0)**, **(S1)**–**(S2)**, and **(D2)**. Document inclusion probabilities satisfy $\Pr(I_j = 1) = \pi_j > 0$, and each selected document j is assigned to m_j distinct workers. Let n denote the number of active workers, indexed by $1, \ldots, n$, with $n = 0$ if $B = 0$.

Lemma 2 (Decomposition and worker representation)**.** *For $T_J^s(a)$, $T_J^m(a)$, and $\zeta_i(a)$ defined in Subsection 4.2, we have*

$$\widehat{\Phi}_J(a) - \Phi_J(a) = T_J^s(a) + T_J^m(a), \tag{B.1}$$

$$\sqrt{\bar{B}}\, T_J^m(a) = \sum_{i=1}^{n} \zeta_i(a). \tag{B.2}$$

Conditional on $\mathcal{G}_J$, the worker response vectors $\{(Y_{ij}^*)_{j=1}^{J}\}_{i=1}^{n}$ are mutually independent, with $\mathbb{E}[Y_{ij}^* \mid \mathcal{G}_J] = \mu_j$. Consequently, the contributions $\{\zeta_i(a)\}_{i=1}^{n}$ are conditionally independent and satisfy

$$\mathbb{E}[\zeta_i(a) \mid \mathcal{G}_J] = 0, \qquad \mathbb{E}[T_J^m(a) \mid \mathcal{G}_J] = 0. \tag{B.3}$$

Proof. Using $r_j = \mu_j - g_j$, we obtain

$$\begin{aligned}
\widehat{\Phi}_J(a) - \Phi_J(a) &= \frac{1}{J}\sum_{j=1}^{J} a_j(g_j - \mu_j) + \frac{1}{J}\sum_{j\in\mathcal{S}_J} \frac{a_j}{\pi_j}(\bar{Y}_j - g_j) \\
&= -\frac{1}{J}\sum_{j=1}^{J} a_j r_j + \frac{1}{J}\sum_{j\in\mathcal{S}_J} \frac{a_j}{\pi_j}\big[r_j + (\bar{Y}_j - \mu_j)\big] \\
&= \frac{1}{J}\sum_{j=1}^{J} \left(\frac{I_j}{\pi_j} - 1\right) a_j r_j + \frac{1}{J}\sum_{j\in\mathcal{S}_J} \frac{a_j}{\pi_j}(\bar{Y}_j - \mu_j) \\
&= T_J^s(a) + T_J^m(a).
\end{aligned}$$

For the worker representation, the definition of $\bar{Y}_j$ and interchanging the two finite sums give

$$\begin{aligned}
\sqrt{\bar{B}}\, T_J^m(a) &= \frac{\sqrt{\bar{B}}}{J}\sum_{j\in\mathcal{S}_J}\sum_{i\in M_j} \frac{a_j}{\pi_j m_j}(Y_{ij}^* - \mu_j) \\
&= \frac{\sqrt{\bar{B}}}{J}\sum_{i=1}^{n}\sum_{j\in\mathcal{J}_i} \frac{a_j}{\pi_j m_j}(Y_{ij}^* - \mu_j) \\
&= \sum_{i=1}^{n} \zeta_i(a).
\end{aligned}$$

To establish the conditional claims, write $L_i = (\theta_i, (\varepsilon_{ij})_{j\le J})$. Assumptions **(RW0)** and **(S2)** make the blocks $\{L_i\}_{i\ge 1}$ mutually independent, while Assumption **(D2)** makes $\mathcal{G}_J$ independent of their joint sigma-field. Thus, for every positive integer q and every bounded measurable f,

$$\mathbb{E}[f(L_1, \ldots, L_q) \mid \mathcal{G}_J] = \mathbb{E}[f(L_1, \ldots, L_q)]. \tag{B.4}$$

By Assumption **(S1)**, each worker's response vector is a measurable function of that worker's block and the fixed corpus. The response vectors therefore remain independent conditional on $\mathcal{G}_J$, with conditional means μ_j. Since $\mathcal{G}_J$ fixes the number of active workers, their assignments, and the coefficients in $\zeta_i(a)$, conditional independence and centering of the worker contributions follow. Equation (B.2) then gives $\mathbb{E}[T_J^m(a) \mid \mathcal{G}_J] = 0$. □

Lemma 3 (Conditional measurement variance)**.** *The conditional measurement variance is*

$$\begin{aligned}
v_{\varepsilon,J}(a) &:= \bar{B}\ \mathrm{Var}(T_J^m(a) \mid \mathcal{G}_J) \\
&= \frac{\bar{B}}{J^2}\sum_{j=1}^{J} \frac{I_j a_j^2 \nu_j}{\pi_j^2 m_j} + \frac{\bar{B}}{J^2}\sum_{\substack{1\le j,k\le J \\ j\ne k}} \frac{I_j I_k N_{jk} a_j a_k \tau_{jk}}{\pi_j \pi_k m_j m_k},
\end{aligned} \tag{B.5}$$

where the second sum is over ordered document pairs. Equivalently,

$$v_{\varepsilon,J}(a) = \sum_{i=1}^{n} \mathrm{Var}(\zeta_i(a) \mid \mathcal{G}_J). \tag{B.6}$$

Proof. Each selected document j receives m_j labels from distinct workers. By (B.4), these labels are conditionally independent, each with variance ν_j. Hence

$$\mathrm{Var}(\bar{Y}_j - \mu_j \mid \mathcal{G}_J) = \frac{\nu_j}{m_j}.$$

For distinct documents $j \neq k$, design exogeneity and Assumptions **(S1)**–**(S2)** give

$$\mathrm{Cov}(Y_{ij}^*, Y_{\ell k}^* \mid \mathcal{G}_J) = \mathbf{1}\{i = \ell\}\tau_{jk}.$$

responses from different workers are independent. Note:

$$\begin{aligned}\mathrm{Cov}(Y_{ij}^*, Y_{ik}^* \mid \mathcal{G}_J) &= \mathrm{Cov}(Y_{ij}^*, Y_{ik}^*) \\ &= \underbrace{\mathbb{E}\big[\mathrm{Cov}(Y_{ij}^*, Y_{ik}^* \mid \theta_i)\big]}_{=0} + \mathrm{Cov}\big(\mathbb{E}[Y_{ij}^* \mid \theta_i], \mathbb{E}[Y_{ik}^* \mid \theta_i]\big) \\ &= \tau_{jk}, \qquad j \neq k.\end{aligned}$$

Thus only the N_{jk} shared workers contribute to the covariance between the means of two selected documents:

$$\mathrm{Cov}(\bar{Y}_j - \mu_j, \bar{Y}_k - \mu_k \mid \mathcal{G}_J) = \frac{N_{jk}}{m_j m_k}\tau_{jk}.$$

Using the preceding variance and covariance identities, we obtain

$$\begin{aligned}\bar{B}\ \mathrm{Var}(T_J^m(a) \mid \mathcal{G}_J) &= \frac{\bar{B}}{J^2}\sum_{j,k\in\mathcal{S}_J}\frac{a_j a_k}{\pi_j\pi_k}\,\mathrm{Cov}(\bar{Y}_j - \mu_j, \bar{Y}_k - \mu_k \mid \mathcal{G}_J) \\ &= \frac{\bar{B}}{J^2}\sum_{j=1}^{J}\frac{I_j a_j^2 \nu_j}{\pi_j^2 m_j} + \frac{\bar{B}}{J^2}\sum_{\substack{1\le j,k\le J\\ j\neq k}}\frac{I_j I_k N_{jk} a_j a_k \tau_{jk}}{\pi_j\pi_k m_j m_k},\end{aligned}$$

which proves (B.5). The worker representation and conditional independence in Lemma 2 give (B.6). □

Lemma 4 (Orthogonality and unconditional variance)**.** *The sampling and measurement components satisfy*

$$\mathrm{Cov}(T_J^s(a), T_J^m(a)) = 0. \tag{B.7}$$

The total variance satisfies

$$\bar{B}\ \mathrm{Var}(\widehat{\Phi}_J(a) - \Phi_J(a)) = v_{r,J}(a) + \bar{v}_{\varepsilon,J}(a), \tag{B.8}$$

where $v_{r,J}(a) := \bar{B}\ \mathrm{Var}(T_J^s(a))$ *and* $\bar{v}_{\varepsilon,J}(a) := \mathbb{E}[v_{\varepsilon,J}(a)]$*. Under Assumption* ***(D1)*** *(Poisson sampling),*

$$v_{r,J}(a) = \frac{\bar{B}}{J^2}\sum_{j=1}^{J}\frac{1-\pi_j}{\pi_j}a_j^2 r_j^2. \tag{B.9}$$

Proof. By Lemma 2, $\mathbb{E}[T_J^m(a) \mid \mathcal{G}_J] = 0$. Since $T_J^s(a)$ is $\mathcal{G}_J$-measurable,

$$\begin{aligned}\mathrm{Cov}(T_J^s(a), T_J^m(a)) &= \mathbb{E}[T_J^s(a) T_J^m(a)] \\ &= \mathbb{E}[T_J^s(a)\mathbb{E}[T_J^m(a) \mid \mathcal{G}_J]] = 0.\end{aligned}$$

This proves (B.7).

Conditional centering and the law of total variance give

$$\begin{aligned}\bar{B}\ \mathrm{Var}(T_J^m(a)) &= \mathbb{E}\big[\bar{B}\ \mathrm{Var}(T_J^m(a) \mid \mathcal{G}_J)\big] \\ &= \bar{v}_{\varepsilon,J}(a).\end{aligned}$$

Combining this identity with the decomposition and orthogonality proves (B.8).

Finally, under Assumption **(D1)**, the indicators I_j are independent Bernoulli variables with variances $\pi_j(1-\pi_j)$. Therefore,

$$v_{r,J}(a) = \frac{\bar{B}}{J^2}\sum_{j=1}^{J}\frac{a_j^2 r_j^2}{\pi_j^2}\operatorname{Var}(I_j) = \frac{\bar{B}}{J^2}\sum_{j=1}^{J}\frac{1-\pi_j}{\pi_j}a_j^2 r_j^2,$$

which proves (B.9). □

Lemma 5 (Bernstein control of the realized label count)**.** *Suppose Assumption **(D1)** holds and* $1 \le m_j \le \bar{m} < \infty$ *for every* j*, where* $\bar{m}$ *does not depend on* J*. Then* $\mathbb{E}[B] = \bar{B}$ *and*

$$V_{B,J} := \operatorname{Var}(B) = \sum_{j=1}^{J} m_j^2 \pi_j(1-\pi_j) \le \bar{m}\bar{B}. \tag{B.10}$$

For every $t > 0$*,*

$$\mathbb{P}(|B-\bar{B}| \ge t) \le 2\exp\left\{-\frac{t^2}{2(V_{B,J}+\bar{m}t/3)}\right\} \le 2\exp\left\{-\frac{t^2}{2\bar{m}(\bar{B}+t/3)}\right\}. \tag{B.11}$$

Consequently, for $0 < \alpha < 1$*, let* $L_\alpha := \log(2/\alpha)$*. With probability at least* $1-\alpha$*,*

$$|B-\bar{B}| \le \sqrt{2V_{B,J}L_\alpha} + \frac{2\bar{m}}{3}L_\alpha \le \sqrt{2\bar{m}\bar{B}L_\alpha} + \frac{2\bar{m}}{3}L_\alpha. \tag{B.12}$$

If $\bar{B} \to \infty$*, then*

$$\frac{B}{\bar{B}} \xrightarrow{p} 1. \tag{B.13}$$

In particular, $\sqrt{B/\bar{B}} \xrightarrow{p} 1$*,* $\mathbb{P}(B=0) \to 0$*, and* $\mathbb{P}(\bar{B}/2 \le B \le 2\bar{B}) \to 1$*.*

Proof. The mean and variance identities follow from $B = \sum_j m_j I_j$, and $m_j^2 \le \bar{m}m_j$ gives $V_{B,J} \le \bar{m}\sum_j \pi_j m_j = \bar{m}\bar{B}$. The variables $m_j(I_j - \pi_j)$ are independent, centered, bounded in absolute value by $\bar{m}$, and have total variance $V_{B,J}$. Bernstein's inequality applied to these variables and their negatives gives (B.11); see Boucheron et al. (2013, Equation (2.10)).

For $t_\alpha := \sqrt{2V_{B,J}L_\alpha} + (2\bar{m}/3)L_\alpha$,

$$t_\alpha^2 - 2L_\alpha\big(V_{B,J} + \bar{m}t_\alpha/3\big) = \frac{2\bar{m}L_\alpha}{3}\sqrt{2V_{B,J}L_\alpha} \ge 0.$$

This proves the first inequality in (B.12); the variance bound proves the second. Finally, setting $t = \epsilon\bar{B}$ in (B.11) gives

$$\mathbb{P}\left(\left|\frac{B}{\bar{B}} - 1\right| \ge \epsilon\right) \le 2\exp\left\{-\frac{\epsilon^2\bar{B}}{2\bar{m}(1+\epsilon/3)}\right\} \longrightarrow 0.$$

The remaining conclusions follow by continuity and elementary event inclusion. □

C Proofs for asymptotic normality

C.1 The high-level central limit theorem

C.1.1 Document-sampling component

We first establish the Gaussian limit for the document-sampling component.

Proof of Lemma 1. Define

$$Z_{Jj}^s := \frac{\sqrt{\bar{B}}}{J}\left(\frac{I_j}{\pi_j} - 1\right) a_j r_j, \qquad j = 1, \ldots, J.$$

Under Assumption **(D1)**, these variables are independent and centered. By (B.9) and Condition **(RW4-S)**,

$$\sqrt{\bar{B}}\, T_J^s(a) = \sum_{j=1}^{J} Z_{Jj}^s, \qquad \sum_{j=1}^{J} \mathrm{Var}(Z_{Jj}^s) = v_{r,J}(a) \longrightarrow v_r(a).$$

Boundedness of the response scale and Assumption **(RW2)** imply that, for some finite C_r,

$$\max_{j \le J} |r_j| \le C_r.$$

Consequently,

$$\max_{j \le J} |Z_{Jj}^s| \le C_r \max_{j \le J} \frac{\sqrt{\bar{B}}}{J\pi_j}|a_j| = C_r \delta_J^{\mathrm{lev}}(a) \longrightarrow 0$$

by (4.5).

Suppose first that $v_r(a) > 0$, and put $s_{s,J}^2 := v_{r,J}(a)$. Then $s_{s,J} \to \sqrt{v_r(a)} > 0$, so

$$\max_{j \le J} \frac{|Z_{Jj}^s|}{s_{s,J}} \longrightarrow 0.$$

Hence, for every $\epsilon > 0$,

$$\frac{1}{s_{s,J}^2} \sum_{j=1}^{J} \mathbb{E}\big[(Z_{Jj}^s)^2 \mathbf{1}\{|Z_{Jj}^s| > \epsilon s_{s,J}\}\big] = 0$$

for all sufficiently large J. The Lindeberg theorem (Billingsley, 1995, Theorem 27.2) therefore gives

$$\frac{\sum_{j=1}^{J} Z_{Jj}^s}{s_{s,J}} \Rightarrow \mathcal{N}(0,1).$$

Slutsky's theorem then yields

$$\sqrt{\bar{B}}\, T_J^s(a) \Rightarrow \mathcal{N}\big(0, v_r(a)\big).$$

If $v_r(a) = 0$, the same conclusion follows directly from Chebyshev's inequality, since

$$\mathrm{Var}\Big(\sqrt{\bar{B}}\, T_J^s(a)\Big) = v_{r,J}(a) \longrightarrow 0.$$

The exact variance formula was established in Lemma 4. □

The next lemma gives the conditional Gaussian approximation needed for the measurement term. Its approximating variance remains specific to the realized sample and allocation, so Condition **(RW4-M)** is not imposed.

C.1.2 Measurement component

We first bound the conditional Gaussian approximation error in terms of the third absolute moments of the worker contributions. Under the regularity conditions of Theorem 1, we can control their sum using the same constants for every admissible graph and every J. We can then show that the resulting error bound vanishes in probability as $\bar{B} \to \infty$.

Lemma 6 (Gaussian approximation conditional on the realized design)**.** *Suppose the conditions of Lemmas 2 and 3 hold. Define*

$$\Lambda_J(a) := \sum_{i=1}^{n} \mathbb{E}\big[|\zeta_i(a)|^3 \mid \mathcal{G}_J\big] .$$

Then, for every fixed $t \in \mathbb{R}$, the following inequality holds almost surely:

$$\left|\mathbb{E}\Big[\exp\Big\{it\sqrt{\bar{B}}\, T_J^m(a)\Big\} \,\Big|\, \mathcal{G}_J\Big] - \exp\Big\{-\frac{t^2}{2} v_{\varepsilon,J}(a)\Big\}\right| \le \frac{|t|^3}{6}\Lambda_J(a) + \frac{t^4}{8}\Lambda_J(a)^{4/3}. \tag{C.1}$$

Consequently, if $\Lambda_J(a) \xrightarrow{p} 0$,

$$\mathbb{E}\Big[\exp\Big\{it\sqrt{\bar{B}}\, T_J^m(a)\Big\} \,\Big|\, \mathcal{G}_J\Big] = \exp\Big\{-\frac{t^2}{2} v_{\varepsilon,J}(a)\Big\} + o_p(1). \tag{C.2}$$

Proof. Conditional on $\mathcal{G}_J$, the number n of active workers and their assignments are fixed, and

$$\sqrt{\bar{B}}\, T_J^m(a) = \sum_{i=1}^{n} \zeta_i(a)$$

is a sum of independent centered variables. Write

$$v_{Ji}(a) := \mathbb{E}[\zeta_i(a)^2 \mid \mathcal{G}_J], \qquad q_{Ji}(a) := \mathbb{E}[|\zeta_i(a)|^3 \mid \mathcal{G}_J].$$

By Lemma 3 and the definition of $\Lambda_J(a)$,

$$\sum_{i=1}^{n} v_{Ji}(a) = v_{\varepsilon,J}(a), \qquad \sum_{i=1}^{n} q_{Ji}(a) = \Lambda_J(a).$$

When $n = 0$, all sums and products below are interpreted using the usual empty-sum and empty-product conventions.

For each active worker, define

$$\phi_{Ji}(t) := \mathbb{E}\Big[e^{it\zeta_i(a)} \mid \mathcal{G}_J\Big], \qquad g_{Ji}(t) := \exp\Big\{-\frac{t^2}{2} v_{Ji}(a)\Big\}.$$

The second-order Taylor remainder for the complex exponential gives

$$\left|e^{ix} - \left(1 + ix - \frac{x^2}{2}\right)\right| \le \frac{|x|^3}{6}, \qquad x \in \mathbb{R};$$

see Billingsley (1995, equation (26.4)). Using $\mathbb{E}[\zeta_i(a) \mid \mathcal{G}_J] = 0$ and the definition of $v_{Ji}(a)$, we obtain

$$\begin{aligned}
&\left|\phi_{Ji}(t) - \left\{1 - \frac{t^2}{2} v_{Ji}(a)\right\}\right| \\
&\quad = \left|\mathbb{E}\left[e^{it\zeta_i(a)} - 1 - it\zeta_i(a) + \frac{t^2}{2}\zeta_i(a)^2 \,\middle|\, \mathcal{G}_J\right]\right| \\
&\quad \le \mathbb{E}\left[\left|e^{it\zeta_i(a)} - 1 - it\zeta_i(a) + \frac{t^2}{2}\zeta_i(a)^2\right| \,\middle|\, \mathcal{G}_J\right] \\
&\quad \le \frac{|t|^3}{6} q_{Ji}(a),
\end{aligned}$$

where the last inequality applies the Taylor bound with $x = t\zeta_i(a)$.

Moreover, for $x \ge 0$,

$$0 \le e^{-x} - (1 - x) = \int_0^x (1 - e^{-s})\, ds \le \int_0^x s\, ds = \frac{x^2}{2}.$$

Applying this with $x = t^2 v_{Ji}(a)/2$ yields

$$\left|g_{Ji}(t) - \left\{1 - \frac{t^2}{2} v_{Ji}(a)\right\}\right| \le \frac{t^4}{8} v_{Ji}(a)^2.$$

The same factorwise exponential remainder is used in the characteristic-function proof of the Berry–Esseen theorem; see Durrett (2019, proof of Theorem 3.4.17, p. 160). Combining the last two bounds gives

$$|\phi_{Ji}(t) - g_{Ji}(t)| \le \frac{|t|^3}{6} q_{Ji}(a) + \frac{t^4}{8} v_{Ji}(a)^2. \tag{C.3}$$

Conditional independence gives

$$\mathbb{E}\left[e^{it\sum_{i=1}^n \zeta_i(a)} \,\middle|\, \mathcal{G}_J\right] = \prod_{i=1}^n \phi_{Ji}(t), \qquad \exp\left\{-\frac{t^2}{2}\sum_{i=1}^n v_{Ji}(a)\right\} = \prod_{i=1}^n g_{Ji}(t).$$

Replacing the factors one at a time gives the exact identity

$$\prod_{i=1}^n \phi_{Ji}(t) - \prod_{i=1}^n g_{Ji}(t) = \sum_{k=1}^n \left\{\prod_{i<k} \phi_{Ji}(t)\right\} \{\phi_{Jk}(t) - g_{Jk}(t)\} \left\{\prod_{i>k} g_{Ji}(t)\right\}.$$

Because $|\phi_{Ji}(t)| \le 1$ and $0 < g_{Ji}(t) \le 1$, taking absolute values yields

$$\left|\prod_{i=1}^n \phi_{Ji}(t) - \prod_{i=1}^n g_{Ji}(t)\right| \le \sum_{i=1}^n |\phi_{Ji}(t) - g_{Ji}(t)|.$$

This is the product comparison in Billingsley (1995, equation (27.5)). Compare also the proof of Theorem 27.2, especially equations (27.11)–(27.15). Consequently,

$$\begin{aligned}
&\left|\mathbb{E}\left[e^{it\sum_{i=1}^n \zeta_i(a)} \,\middle|\, \mathcal{G}_J\right] - \exp\left\{-\frac{t^2}{2}\sum_{i=1}^n v_{Ji}(a)\right\}\right| \\
&\qquad \le \frac{|t|^3}{6}\sum_{i=1}^n q_{Ji}(a) + \frac{t^4}{8}\sum_{i=1}^n v_{Ji}(a)^2.
\end{aligned} \tag{C.4}$$

The identity and inequality remain valid when $n = 0$ under the stated empty-product convention.

Finally, conditional Lyapunov's moment inequality gives

$$v_{Ji}(a)^{3/2} = \left\{\mathbb{E}[\zeta_i(a)^2 \mid \mathcal{G}_J]\right\}^{3/2} \le \mathbb{E}[|\zeta_i(a)|^3 \mid \mathcal{G}_J] = q_{Ji}(a).$$

Together with monotonicity of finite-dimensional ℓ^p norms, this implies

$$\sum_{i=1}^{n} v_{Ji}(a)^2 \le \left\{\sum_{i=1}^{n} v_{Ji}(a)^{3/2}\right\}^{4/3} \le \left\{\sum_{i=1}^{n} q_{Ji}(a)\right\}^{4/3} = \Lambda_J(a)^{4/3}.$$

Substitution into (C.4), together with the exact identities for the two sums, proves (C.1). If $\Lambda_J(a) \xrightarrow{p} 0$, the right-hand side of (C.1) is $o_p(1)$. This proves (C.2). □

C.1.3 Combining the two terms

We now combine the sampling limit with the conditional measurement approximation. Variance stabilization makes the limiting conditional characteristic function nonrandom, allowing the characteristic function of the sum to factor asymptotically.

Proof of Theorem 1. Write

$$S_J := \sqrt{\bar{B}}\, T_J^s(a), \qquad M_J := \sqrt{\bar{B}}\, T_J^m(a), \qquad \varphi_J(t) := \mathbb{E}[e^{itM_J} \mid \mathcal{G}_J].$$

Let $R_Y := \sup_{y,y' \in \mathcal{Y}} |y - y'|$. Balanced spending and the worker-degree bound give

$$|\zeta_i(a)| \le \frac{K\bar{a}R_Y}{c_\lambda \sqrt{\bar{B}}}.$$

Since the number of active workers is at most B, Lemma 5 implies

$$\Lambda_J(a) \le B\left(\frac{K\bar{a}R_Y}{c_\lambda\sqrt{\bar{B}}}\right)^3 = O_p(\bar{B}^{-1/2}) = o_p(1).$$

Because S_J is $\mathcal{G}_J$-measurable, the tower property gives

$$\mathbb{E}[e^{it(S_J+M_J)}] = \mathbb{E}[e^{itS_J}\varphi_J(t)]. \tag{C.5}$$

Let

$$c_t := \exp\left\{-\frac{t^2}{2}v_\varepsilon(a)\right\}, \qquad D_J(t) := \varphi_J(t) - c_t.$$

Lemma 6 therefore gives

$$\varphi_J(t) - \exp\left\{-\frac{t^2}{2}v_{\varepsilon,J}(a)\right\} \xrightarrow{p} 0,$$

while Condition **(RW4-M)** and continuity of the exponential give

$$\exp\left\{-\frac{t^2}{2}v_{\varepsilon,J}(a)\right\} - c_t \xrightarrow{p} 0.$$

Hence $D_J(t) \xrightarrow{p} 0$. Moreover, $|D_J(t)| \le 2$. Thus, for every $\eta > 0$,

$$\mathbb{E}|D_J(t)| \le \eta + 2\mathbb{P}\{|D_J(t)| > \eta\}.$$

Taking the limit superior and then letting $\eta \downarrow 0$ shows that $\mathbb{E}|D_J(t)| \to 0$. Therefore

$$\left|\mathbb{E}[e^{itS_J} D_J(t)]\right| \leq \mathbb{E}|D_J(t)| \longrightarrow 0.$$

Substitution into (C.5) yields

$$\mathbb{E}[e^{it(S_J+M_J)}] = \exp\left\{-\frac{t^2}{2} v_\varepsilon(a)\right\} \mathbb{E}[e^{itS_J}] + o(1). \tag{C.6}$$

By Lemma 1,

$$\mathbb{E}[e^{itS_J}] \longrightarrow \exp\left\{-\frac{t^2}{2} v_r(a)\right\}.$$

Combining this with (C.6) and applying Lévy's continuity theorem (Billingsley, 1995, Theorem 26.3) gives

$$\sqrt{\bar{B}}\{\widehat{\Phi}_J(a) - \Phi_J(a)\} = S_J + M_J \Rightarrow \mathcal{N}(0, v_r(a) + v_\varepsilon(a)).$$

Finally, if $1 \leq m_j \leq \bar{m}$, Lemma 5 gives $\sqrt{B/\bar{B}} \xrightarrow{p} 1$. Hence

$$\sqrt{B}\{\widehat{\Phi}_J(a) - \Phi_J(a)\} = \sqrt{B/\bar{B}}\, \sqrt{\bar{B}}\{\widehat{\Phi}_J(a) - \Phi_J(a)\}$$

has the same limit by Slutsky's theorem. □

C.2 Why allocation regularity is needed

Bounded degrees and regular document arrays do not by themselves stabilize the graph-specific measurement variance. The following fixed-budget example uses a sample-level imbalance to select between two bounded-load allocation rules. That imbalance also determines the first-order sampling error.

Proposition 3 (Failure under sample-adaptive allocation)**.** *There exists a fixed-corpus sequence under fixed-size simple random sampling with*

$$B = \bar{B} = N_J, \qquad m_j = a_j = \lambda_j = 1, \qquad n = N_J/2, \qquad K = 2,$$

*satisfying the bounded-response model, Assumptions **(RW0)**, **(S1)**–**(S2)**, and **(D2)**, Assumption **(RW2)**, the leverage bound* (4.5)*, the condition $\Lambda_J(a) \xrightarrow{p} 0$, the corresponding document-sampling CLT and variance limit, and the two document-array limits used for variance stabilization, but not Condition **(RW4-M)**. Its normalized estimator has a non-Gaussian limit. Neither exact integrated-variance standardization nor oracle hybrid studentization using the graph-specific measurement variance has a standard-normal limit.*

*Fixed-size SRS replaces Assumption **(D1)**, and the sample-adaptive allocation intentionally falls outside Assumption **(RW1)**.*

Proof. Let J and N_J be even and suppose

$$N_J \to \infty, \qquad J - N_J \to \infty, \qquad f_J := N_J/J \to f \in (0,1).$$

Divide the fixed corpus into two equally sized classes,

$$\chi_j = \begin{cases} 1, & j \le J/2, \\ -1, & j > J/2. \end{cases}$$

Set $\pi_j = f_J$, $m_j = a_j = 1$, and $g_j = c_0$. Thus $\bar{B} = Jf_J = N_J$ and $\lambda_j := J\pi_j m_j/\bar{B} = 1$.

Let $F_i, G_{i,1}, G_{i,-1}$, and U_{ij} be mutually independent Rademacher variables, with the worker blocks i.i.d. across i, and set

$$Y_{ij}^* = c_0 + \delta\chi_j + \tau_0 F_i + \tau_1 G_{i,\chi_j} + \eta U_{ij}, \qquad \delta, \tau_1 > 0.$$

Choose the constants so that the maintained outcome bound holds. This is the maintained response model with

$$\theta_i = (F_i, G_{i,1}, G_{i,-1}), \qquad \varepsilon_{ij} = U_{ij}.$$

Consequently,

$$\mu_j = c_0 + \delta\chi_j, \qquad r_j = \delta\chi_j,$$

and

$$\nu_j = \eta^2 + \tau_0^2 + \tau_1^2, \qquad \tau_{jk} = \tau_0^2 + \tau_1^2 \mathbf{1}\{\chi_j = \chi_k\}. \tag{C.7}$$

Draw exactly N_J documents by SRS without replacement and define

$$N_{+,J} := \sum_{j \in \mathcal{S}_J} \mathbf{1}\{\chi_j = 1\}, \quad N_{-,J} := \sum_{j \in \mathcal{S}_J} \mathbf{1}\{\chi_j = -1\},$$
$$D_J := N_{+,J} - N_{-,J}.$$

Pair the selected documents and assign the $N_J/2$ pairs to distinct workers. If $D_J \ge 0$, draw uniformly from the matchings with as many within-class pairs as parity permits. If $D_J < 0$, draw a uniform perfect matching of all selected documents. Both the SRS draw and all matching randomization are independent of the worker-response blocks, so Assumption **(D2)** holds. Both branches have document degree one and worker degree two.

Write

$$S_J := \sqrt{N_J}\, T_J^s(a), \qquad M_J := \sqrt{N_J}\, T_J^m(a).$$

Because the corpus class counts are equal and the number of selected positive-class documents has a hypergeometric distribution

$$S_J = \delta \frac{D_J}{\sqrt{N_J}}, \qquad \mathrm{Var}(D_J) = \frac{N_J(J - N_J)}{J - 1}.$$

The finite-population central limit theorem (Li and Ding, 2017, Theorem 1) therefore gives

$$S_J \Rightarrow S, \qquad S \sim \mathcal{N}(0, v_r), \qquad v_r := \delta^2(1 - f). \tag{C.8}$$

Let ρ_J denote the fraction of workers whose paired documents have the same class, and set $V_J := v_{\varepsilon,J}(a)$. Lemma **??**(c) and (C.7) give

$$V_J = N_J \,\mathrm{Var}\{T_J^m(a) \mid \mathcal{G}_J\} = \eta^2 + 2\tau_0^2 + \tau_1^2(1 + \rho_J). \tag{C.9}$$

In the within-class branch, at most one pair is cross-class, so

$$0 \le 1 - \rho_J \le \frac{2}{N_J}.$$

In the pooled branch,

$$\mathbb{E}[\rho_J \mid \mathcal{S}_J] = \frac{N_{+,J}(N_{+,J}-1) + N_{-,J}(N_{-,J}-1)}{N_J(N_J-1)} = \frac{1}{2} + \frac{D_J^2 - N_J}{2N_J(N_J-1)} \xrightarrow{p} \frac{1}{2}.$$

For $N_J \ge 4$, concentration in the pooled branch follows by writing $\rho_J = 1 - 2C_J/N_J$, where C_J is the number of cross-class pairs. For each selected + document j, let H_j indicate that it is matched to a − document, so that $C_J = \sum_j H_j$. Conditional on $\mathcal{S}_J$, the one- and two-indicator probabilities are $N_{-,J}/(N_J-1)$ and $N_{-,J}(N_{-,J}-1)/\{(N_J-1)(N_J-3)\}$, respectively. It follows that

$$\mathrm{Var}(\rho_J \mid \mathcal{S}_J) = \frac{8N_{+,J}(N_{+,J}-1)N_{-,J}(N_{-,J}-1)}{N_J^2(N_J-1)^2(N_J-3)} = O(N_J^{-1}).$$

It follows that

$$V_J - \{v_H \mathbf{1}\{D_J \ge 0\} + v_P \mathbf{1}\{D_J < 0\}\} \xrightarrow{p} 0, \qquad \begin{cases} v_H = \eta^2 + 2\tau_0^2 + 2\tau_1^2, \\ v_P = \eta^2 + 2\tau_0^2 + \frac{3}{2}\tau_1^2. \end{cases} \tag{C.10}$$

Because $v_H \neq v_P$ and $\mathbb{P}(D_J \ge 0), \mathbb{P}(D_J < 0) \to 1/2$, V_J does not converge in probability to a constant, so Condition **(RW4-M)** fails.

Each of the $N_J/2$ worker contributions to M_J is bounded by $C/\sqrt{N_J}$. Hence, uniformly in $\mathcal{G}_J$,

$$\Lambda_J(a) = \sum_i \mathbb{E}[|\zeta_i(a)|^3 \mid \mathcal{G}_J] = O(N_J^{-1/2}),$$

so $\Lambda_J(a) \xrightarrow{p} 0$. By Lemma 6,

$$\mathbb{E}(e^{itM_J} \mid \mathcal{G}_J) = e^{-t^2 V_J/2} + o_p(1).$$

The error is bounded and hence is also $o(1)$ in L^1. Therefore, for every $u, t, w \in \mathbb{R}$,

$$\begin{aligned} \mathbb{E}[e^{iuS_J + itM_J + iwV_J}] &= \mathbb{E}\big[e^{iuS_J + iwV_J}\mathbb{E}(e^{itM_J} \mid \mathcal{G}_J)\big] \\ &= \mathbb{E}[e^{iuS_J + iwV_J - t^2V_J/2}] + o(1) \\ &\longrightarrow \mathbb{E}[e^{iuS + iwv(S) - t^2 v(S)/2}], \end{aligned}$$

where

$$v(S) := v_H \mathbf{1}\{S \ge 0\} + v_P \mathbf{1}\{S < 0\}.$$

The final convergence follows from (C.8)–(C.10) and the extended continuous mapping theorem, since $\mathrm{sign}(S_J) = \mathrm{sign}(D_J)$ and $\mathbb{P}(S = 0) = 0$. The limiting characteristic function identifies

$$(S_J, M_J, V_J) \Rightarrow \big(S, \sqrt{v(S)}Z, v(S)\big), \qquad Z \sim \mathcal{N}(0,1), \quad Z \perp S.$$

Consequently,

$$\sqrt{N_J}\{\widehat{\Phi}_J(a) - \Phi_J(a)\} = S_J + M_J \Rightarrow X := S + \sqrt{v(S)}Z. \tag{C.11}$$

Although $\mathbb{E}[X] = 0$, symmetry of S gives

$$\mathbb{E}[X^3] = 3\mathbb{E}[Sv(S)] = 3(v_H - v_P)\sqrt{\frac{v_r}{2\pi}} \neq 0,$$

so X is not Gaussian.

The exact scaled SRS sampling variance is

$$v_{s,J}^{\mathrm{SRS}} := N_J \operatorname{Var}\{T_J^s(a)\} = \delta^2 \frac{J - N_J}{J - 1} \longrightarrow v_r.$$

Conditional centering gives exact orthogonality of the sampling and measurement components. Hence their exact integrated variance is

$$v_{\mathrm{int},J} := N_J \operatorname{Var}\{\widehat{\Phi}_J(a) - \Phi_J(a)\} = v_{s,J}^{\mathrm{SRS}} + \mathbb{E}[V_J].$$

Since V_J is uniformly bounded and each branch has limiting probability $1/2$,

$$v_{\mathrm{int},J} \longrightarrow v_r + \frac{v_H + v_P}{2}.$$

It follows that

$$\frac{\sqrt{N_J}\{\widehat{\Phi}_J(a) - \Phi_J(a)\}}{\sqrt{v_{\mathrm{int},J}}} \Rightarrow \frac{X}{\sqrt{v_r + (v_H + v_P)/2}},$$

which remains non-Gaussian. The preceding joint convergence also yields the oracle hybrid limit

$$\frac{\sqrt{N_J}\{\widehat{\Phi}_J(a) - \Phi_J(a)\}}{\sqrt{v_{s,J}^{\mathrm{SRS}} + V_J}} \Rightarrow \frac{S + \sqrt{v(S)}Z}{\sqrt{v_r + v(S)}}.$$

The mean of the limiting variable is

$$\sqrt{\frac{v_r}{2\pi}} \left\{ \frac{1}{\sqrt{v_r + v_H}} - \frac{1}{\sqrt{v_r + v_P}} \right\} \neq 0,$$

so this limit is not standard normal either.

Finally, the document arrays are regular:

$$\frac{1}{J}\sum_{j=1}^{J} \frac{a_j^2 \nu_j}{\lambda_j} = \eta^2 + \tau_0^2 + \tau_1^2, \qquad \frac{1}{J^2}\sum_{j \neq k} a_j a_k \tau_{jk} \longrightarrow \tau_0^2 + \frac{1}{2}\tau_1^2,$$

while

$$\lambda_j = 1, \qquad \delta_J^{\mathrm{lev}}(a) = \frac{\sqrt{N_J}}{J f_J} = \frac{1}{\sqrt{N_J}} \longrightarrow 0.$$

Thus the failure is caused by the sample-adaptive global allocation switch, not by irregular document moments, excessive degrees, or failure of the conditional worker Gaussian approximation. □

C.3 Primitive conditions for variance stabilization

This subsection proves Theorem 2. Let

$$\mathcal{E}_J^{\mathrm{feas}} := \{B = 0\} \cup \left\{ B > 0,\ \mathfrak{A}_J(\boldsymbol{d}_J, \boldsymbol{k}_J^{\mathrm{cf}}) \neq \varnothing \right\}$$

be the event on which the capacity-filled graph is feasible (with the empty graph used when $B = 0$), and define

$$\mathcal{H}_J := \mathcal{E}_J^{\text{feas}} \cap \left\{ \frac{\bar{B}}{2} \le B \le 2\bar{B} \right\}. \tag{C.12}$$

We first rigorously verify that, whenever $B \ge K\bar{m}$, the requested annotation depths and prescribed worker workloads can be satisfied without assigning the same document to a worker twice. For $B > 0$, let $m_{(1)} \ge \cdots \ge m_{(|\mathcal{S}_J|)}$ be the selected document degrees in decreasing order, and let $k_1, \ldots, k_{n_0}$ be the prescribed capacity-filled worker degrees. Since both degree sequences sum to B, the Gale–Ryser criterion states that a simple bipartite graph with these degrees exists if and only if

$$\sum_{\ell=1}^{t} m_{(\ell)} \le \sum_{i=1}^{n_0} \min(t, k_i), \qquad t = 1, \ldots, |\mathcal{S}_J|.$$

The left-hand side is the evaluation demand of the t most heavily annotated documents; the right-hand side is the total capacity workers can contribute to those documents without repeated evaluations. For $t \ge K$, the right-hand side equals B, so the inequality holds automatically. For $t < K$, capacity filling provides $\lfloor B/K \rfloor$ workers with workload K. If $B \ge K\bar{m}$, then

$$\sum_{\ell=1}^{t} m_{(\ell)} \le t\bar{m} \le t\lfloor B/K \rfloor \le \sum_{i=1}^{n_0} \min(t, k_i).$$

Thus the capacity-filled allocation is feasible whenever $B \ge K\bar{m}$.

Since $K\bar{m}$ is fixed and $\bar{B} \to \infty$, we have $\bar{B}/2 \ge K\bar{m}$ for all sufficiently large J. The preceding feasibility argument therefore gives $\mathcal{H}_J = \{\bar{B}/2 \le B \le 2\bar{B}\}$ for such J. By Lemma 5,

$$\mathbb{P}(\mathcal{H}_J^c) = \mathbb{P}\big(B \notin [\bar{B}/2, 2\bar{B}]\big) \longrightarrow 0. \tag{C.13}$$

The event $\mathcal{H}_J$ is determined by $\mathcal{S}_J$. On this event, $B \asymp \bar{B} \to \infty$, and, conditional on $\mathcal{S}_J$, Assumption **(RW1)** makes A_J uniform over the nonempty class of simple bipartite graphs with the prescribed degrees. We apply this uniform law below only on $\mathcal{H}_J$.

Now, Introducing some necessary notation. For nonnegative integers x and r, write

$$(x)_r := x(x-1)\cdots(x-r+1), \qquad (x)_0 := 1.$$

The next lemma approximates the probability that a particular pattern of worker–document assignments occurs under our allocation scheme. This is directly relevant to N_{jk}: a given worker evaluating both documents j and k forms a two-edge pattern, and N_{jk} counts the workers for whom this pattern occurs. For patterns involving a bounded number of assignments, we show that their probabilities under our allocation scheme can be approximated by those obtained by uniformly matching document slots to worker slots without imposing simplicity.

Lemma 7 (Local patterns in a uniform fixed-degree graph)**.** *Consider a sequence of nonempty classes of simple bipartite graphs with left degrees $\{m_j\}$, right degrees $\{k_i\}$, and common degree sum*

$$B = \sum_j m_j = \sum_i k_i \longrightarrow \infty.$$

Suppose that the maximum degree on both sides is uniformly bounded. For each class, let H be a specified

simple bipartite pattern on its labelled vertex classes, with $|E(H)|$ uniformly bounded, and let $x_j(H)$ and $y_i(H)$ denote its degrees on the two sides. If $\mathcal{A}$ is uniform over the corresponding graph class, then

$$\mathbb{P}(H \subseteq \mathcal{A}) = \frac{\prod_j (m_j)_{x_j(H)} \prod_i (k_i)_{y_i(H)}}{(B)_{|E(H)|}} \{1 + O(B^{-1})\}.$$

The remainder is uniform over the degree sequences and patterns satisfying these bounds.

Proof. If a degree of H exceeds the corresponding prescribed degree, the claim is immediate. Otherwise, let D be a common bound on the prescribed degrees. Each vertex class then contains at least B/D positive-degree vertices, so both classes diverge as required by McKay's asymptotic convention. Apply McKay (2010, Theorem 2.2) with

$$s_j = m_j, \quad t_i = k_i, \quad x_j = x_j(H), \quad y_i = y_i(H), \quad E = B, \quad X = |E(H)|.$$

Writing $s_{\max}, t_{\max}, x_{\max}, y_{\max}$ for the corresponding maximum degrees, McKay's condition is $\Delta' X = o(E)$, and his relative error is $O(\Delta' X/E)$, where

$$\Delta' = (s_{\max} + t_{\max})(s_{\max} + t_{\max} + x_{\max} + y_{\max}).$$

Under the stated bounds, $\Delta' X = O(1) = o(B)$. Since $E = B$, McKay's formula gives the stated $O(B^{-1})$ remainder, uniformly under these bounds. □

We first show that the worker reuse part of the variance concentrates around its conditional mean given the document sample. Thus, for a given sample, variation in the measurement variance across worker allocations becomes asymptotically negligible.

The bounded-response-scale assumption allows us to fix $C_Y < \infty$, uniformly in the array indices, such that

$$|Y_{ij}^* - \mu_j| \le C_Y, \qquad \nu_j \le C_Y, \qquad |\tau_{jk}| \le C_Y.$$

For $j \neq k$, define

$$\beta_{jk}^{\mathrm{ov}} := \frac{\bar{B}}{J^2} \frac{a_j a_k \tau_{jk}}{\pi_j \pi_k m_j m_k} = \frac{a_j a_k \tau_{jk}}{\lambda_j \lambda_k \bar{B}}. \tag{C.14}$$

Lemma 8 (Pairwise mean and concentration of the weighted overlap)**.** *Suppose the conditions of Theorem 2 hold. On $\mathcal{H}_J$, uniformly over distinct selected documents $j \neq k$,*

$$\mathbb{E}[N_{jk} \,|\, \mathcal{S}_J] = (K-1)\frac{m_j m_k}{B} + O\Big(\frac{m_j m_k}{B^2}\Big). \tag{C.15}$$

Define the centered allocation remainder

$$R_J^{\mathrm{alloc}}(a) := \sum_{j \neq k} I_j I_k \beta_{jk}^{\mathrm{ov}} \{N_{jk} - \mathbb{E}[N_{jk} \,|\, \mathcal{S}_J]\}, \tag{C.16}$$

where the sum is over ordered document pairs. Then

$$R_J^{\mathrm{alloc}}(a) \xrightarrow{p} 0.$$

Proof. We first prove the mean statement. Since $B \ge \bar{B}/2 \to \infty$ uniformly over realizations in $\mathcal{H}_J$, the McKay expansion applies to every such realization for all sufficiently large J. Fix one such realization. For distinct selected documents $j \neq k$ and an active worker i, the event $\{i \in M_j \cap M_k\}$ is the containment

event for the two-edge pattern $\{(j,i),(k,i)\}$. Lemma 7 therefore gives

$$\mathbb{P}(i \in M_j \cap M_k \,|\, \mathcal{S}_J) = \frac{m_j m_k \, k_i(k_i-1)}{B(B-1)}\{1 + O(B^{-1})\},$$

uniformly over i, j, k and the stated realizations. If $k_i < 2$, both sides are zero.

On $\mathcal{H}_J$, the worker degrees are fixed conditional on $\mathcal{S}_J$. Capacity filling assigns load K to all but the last worker, whose load is $r_B \in \{1, \dots, K\}$. Hence

$$\sum_{i=1}^{n} k_i(k_i-1) = (K-1)B - r_B(K-r_B) = (K-1)B + O(1),$$

where the remainder is uniformly bounded because K is fixed.

Since $N_{jk} = \sum_{i=1}^{n} \mathbf{1}\{i \in M_j \cap M_k\}$, summing the uniform error bounds gives

$$\begin{aligned}
\mathbb{E}[N_{jk} \,|\, \mathcal{S}_J] &= \frac{m_j m_k}{B(B-1)} \sum_{i=1}^{n} k_i(k_i-1) + O\left(\frac{m_j m_k}{B^2(B-1)} \sum_{i=1}^{n} k_i(k_i-1) \right) \\
&= (K-1)\frac{m_j m_k}{B-1} + O\Big(\frac{m_j m_k}{B^2}\Big) \\
&= (K-1)\frac{m_j m_k}{B} + O\Big(\frac{m_j m_k}{B^2}\Big).
\end{aligned}$$

The second equality uses the capacity-filling identity above and $\sum_i k_i(k_i-1) \le (K-1)B$. The last equality uses $(B-1)^{-1} = B^{-1} + O(B^{-2})$. All remainder bounds are uniform over distinct selected documents $j \neq k$ and sampling realizations in $\mathcal{H}_J$, by Lemma 7, the fixed bound K, and $B \ge \bar{B}/2 \to \infty$. This proves (C.15).

We next establish concentration around the conditional mean. We first rewrite the worker-reuse component of the variance from a sum over document pairs to a sum of worker contributions. We then show that these contributions are small and weakly dependent. Concentration around the conditional mean then follows from an LLN-type argument.

Let

$$M_J^{\mathrm{ov}} := \max_{j \neq k} |\beta_{jk}^{\mathrm{ov}}|.$$

The bounds $\lambda_j \ge c_\lambda$ and $|a_j| \le \bar{a}$ give

$$M_J^{\mathrm{ov}} \le \frac{\bar{a}^2 C_Y}{c_\lambda^2 \bar{B}} = O(\bar{B}^{-1}). \tag{C.17}$$

Here and below, $C_\star < \infty$ may change from line to line but depends only on the fixed bounds.

All sums over $j \neq k$ below are over ordered pairs. For each active worker i, define

$$U_i := \sum_{\substack{j,k \in \mathcal{J}_i \\ j \neq k}} \beta_{jk}^{\mathrm{ov}}.$$

Then $|U_i| \le K(K-1)M_J^{\mathrm{ov}}$. This established that each worker contribution is small.

Counting each worker shared by an ordered document pair once gives

$$R_J^{\mathrm{alloc}}(a) = \sum_{i=1}^{n} \{U_i - \mathbb{E}[U_i \mid \mathcal{S}_J]\}. \tag{C.18}$$

We next establish weak dependence between two worker assigned document set, which we refer to as worker neighborhoods. On $\mathcal{H}_J$, condition on $\mathcal{S}_J$. All patterns below have at most $2K$ edges and degrees bounded in terms of $(\bar{m}, K)$, so the remainder in Lemma 7 is uniform over all workers, neighborhoods, and selection realizations considered below. For $A \subseteq \mathcal{S}_J$ with $|A| = k_i$, containment of the full star from worker i to A is exactly the event $\{\mathcal{J}_i = A\}$, because the star exhausts worker i's prescribed degree. Hence

$$p_i(A) := \mathbb{P}(\mathcal{J}_i = A \mid \mathcal{S}_J) = \frac{k_i! \prod_{j \in A} m_j}{(B)_{k_i}} \{1 + O(B^{-1})\}. \tag{C.19}$$

To bound the conditional covariance between worker contributions, we compare the joint distribution of their assigned document sets with a distribution under which the two sets are independent, while preserving each worker's assignment probabilities given the document sample. Since each contribution is a bounded function of its assigned document set, a bound on the total variation distance between these distributions gives a bound on the conditional covariance.

For distinct active workers $i \neq \ell$, let $P_{i\ell}$ be the conditional joint law of $(\mathcal{J}_i, \mathcal{J}_\ell)$, and let $Q_{i\ell}$ be the product of its exact conditional marginals, so that $Q_{i\ell}(A, C) = p_i(A)p_\ell(C)$. If $|A| = k_i$, $|C| = k_\ell$, and $A \cap C = \varnothing$, the union of the two full stars exhausts both worker degrees. Lemma 7 therefore gives

$$P_{i\ell}(A, C) = \frac{k_i! k_\ell! \prod_{j \in A \cup C} m_j}{(B)_{k_i + k_\ell}} \{1 + O(B^{-1})\}.$$

If $Q_{i\ell}(A, C) = 0$, then $P_{i\ell}(A, C) = 0$ as well, because the joint event is contained in each marginal event. Otherwise, dividing by the two marginal formulas in (C.19), and using disjointness to cancel the document factors, gives For disjoint A, C with $Q_{i\ell}(A, C) > 0$,

$$\begin{aligned}
\frac{P_{i\ell}(A, C)}{Q_{i\ell}(A, C)} &= \frac{(B)_{k_i}(B)_{k_\ell}}{(B)_{k_i + k_\ell}} \frac{1 + O(B^{-1})}{\{1 + O(B^{-1})\}\{1 + O(B^{-1})\}} \\
&= \prod_{r=0}^{k_\ell - 1} \left(1 + \frac{k_i}{B - k_i - r}\right) \{1 + O(B^{-1})\} \\
&= 1 + O(B^{-1}).
\end{aligned}$$

The last equality uses $k_i, k_\ell \leq K$, with K fixed.

Consequently,

$$|P_{i\ell}(A, C) - Q_{i\ell}(A, C)| \leq \frac{C_\star}{B} Q_{i\ell}(A, C), \qquad A \cap C = \varnothing. \tag{C.20}$$

Let $D := \{(A, C) : A \cap C = \varnothing\}$. For each $j \in \mathcal{S}_J$, the one- and two-edge cases of Lemma 7 give

$$\begin{aligned}
\mathbb{P}(j \in \mathcal{J}_i \mid \mathcal{S}_J) &= \frac{k_i m_j}{B} \{1 + O(B^{-1})\}, \\
\mathbb{P}(j \in \mathcal{J}_i \cap \mathcal{J}_\ell \mid \mathcal{S}_J) &= \frac{k_i k_\ell (m_j)_2}{(B)_2} \{1 + O(B^{-1})\}.
\end{aligned}$$

Under $Q_{i\ell}$, the two assigned document sets are independent. A union bound and $k_i, k_\ell \leq K$ therefore give

$$\begin{aligned}
Q_{i\ell}(D^c) &\leq \sum_{j \in \mathcal{S}_J} \mathbb{P}(j \in \mathcal{J}_i \mid \mathcal{S}_J) \mathbb{P}(j \in \mathcal{J}_\ell \mid \mathcal{S}_J) \\
&\leq \frac{C_\star}{B^2} \sum_{j \in \mathcal{S}_J} m_j^2.
\end{aligned}$$

Under $P_{i\ell}$, the corresponding union bound gives

$$\begin{aligned} P_{i\ell}(D^c) &\le \sum_{j \in \mathcal{S}_J} \mathbb{P}(j \in \mathcal{J}_i \cap \mathcal{J}_\ell \mid \mathcal{S}_J) \\ &\le \frac{C_\star}{B^2} \sum_{j \in \mathcal{S}_J} (m_j)_2. \end{aligned}$$

Since $m_j \le \bar{m}$ and $\sum_{j \in \mathcal{S}_J} m_j = B$, it follows that

$$Q_{i\ell}(D^c) + P_{i\ell}(D^c) \le \frac{C_\star}{B^2} \sum_{j \in \mathcal{S}_J} \{m_j^2 + (m_j)_2\} \le \frac{C_\star(2\bar{m} - 1)}{B} = O(B^{-1}).$$

Combining this bound with (C.20) yields

$$\begin{aligned} 2\|P_{i\ell} - Q_{i\ell}\|_{\mathrm{TV}} &= \sum_{(A,C) \in D} |P_{i\ell}(A,C) - Q_{i\ell}(A,C)| \\ &\quad + \sum_{(A,C) \in D^c} |P_{i\ell}(A,C) - Q_{i\ell}(A,C)| \\ &\le \frac{C_\star}{B} Q_{i\ell}(D) + P_{i\ell}(D^c) + Q_{i\ell}(D^c) = O(B^{-1}). \end{aligned} \tag{C.21}$$

The bound is uniform over $i \neq \ell$ and the selection realizations in $\mathcal{H}_J$.

Because $Q_{i\ell}$ is the product of the exact marginals of $P_{i\ell}$,

$$\mathrm{Cov}(U_i, U_\ell \mid \mathcal{S}_J) = \mathbb{E}_{P_{i\ell}}[U_i U_\ell] - \mathbb{E}_{Q_{i\ell}}[U_i U_\ell].$$

The standard inequality

$$|\mathbb{E}_P h - \mathbb{E}_Q h| \le 2\|h\|_\infty \|P - Q\|_{\mathrm{TV}},$$

together with $|U_i| \le K(K-1)M_J^{\mathrm{ov}}$, therefore gives

$$|\,\mathrm{Cov}(U_i, U_\ell \mid \mathcal{S}_J)| \le \frac{C_\star (M_J^{\mathrm{ov}})^2}{B}, \qquad \mathrm{Var}(U_i \mid \mathcal{S}_J) \le C_\star (M_J^{\mathrm{ov}})^2.$$

The sums in (C.18) run only over active workers, so $n \le \sum_i k_i = B$. It follows that, on $\mathcal{H}_J$,

$$\begin{aligned} \mathrm{Var}(R_J^{\mathrm{alloc}}(a) \mid \mathcal{S}_J) &\le C_\star \left\{ n (M_J^{\mathrm{ov}})^2 + \frac{n^2 (M_J^{\mathrm{ov}})^2}{B} \right\} \\ &\le C_\star B (M_J^{\mathrm{ov}})^2 \le \frac{C_\star}{\bar{B}}, \end{aligned} \tag{C.22}$$

where the last inequality uses (C.17) and $B \le 2\bar{B}$.

Finally, $R_J^{\mathrm{alloc}}(a)$ has conditional mean zero given $\mathcal{S}_J$. Since $\mathcal{H}_J \in \sigma(\mathcal{S}_J)$, conditional Chebyshev's inequality gives, for every $\epsilon > 0$,

$$\begin{aligned} \mathbb{P}\big(|R_J^{\mathrm{alloc}}(a)| > \epsilon\big) &\le \mathbb{P}(\mathcal{H}_J^c) + \mathbb{E}\big[\mathbf{1}_{\mathcal{H}_J} \mathbb{P}\big(|R_J^{\mathrm{alloc}}(a)| > \epsilon \,\big|\, \mathcal{S}_J\big)\big] \\ &\le \mathbb{P}(\mathcal{H}_J^c) + \frac{C_\star}{\epsilon^2 \bar{B}} \longrightarrow 0. \end{aligned}$$

□

Proof of Theorem 2. Recall from (C.13) that $\mathbb{P}(\mathcal{H}_J^c) \to 0$. Throughout the proof, $C_0 < \infty$ denotes a

constant depending only on the fixed bounds and may change from line to line.

By Lemma 3 and the definitions of λ_j and β_{jk}^{ov}, we can decompose the variance into a "Diagonal term" $D_J(a)$ and a "Worker reuse term" $C_J(a)$

$$v_{\varepsilon,J}(a) = D_J(a) + C_J(a),$$

where

$$D_J(a) := \frac{1}{J}\sum_{j=1}^{J} \frac{I_j}{\pi_j}\frac{a_j^2\nu_j}{\lambda_j}, \qquad C_J(a) := \sum_{j\neq k} \beta_{jk}^{\mathrm{ov}} I_j I_k N_{jk}.$$

The diagonal term depends only on the document sample and is unaffected by worker allocation. We establish its concentration around its mean using the usual Horvitz–Thompson variance formula under Poisson sampling. By Assumption **(D1)**, the selection indicators are independent, so

$$\mathbb{E}[D_J(a)] = \frac{1}{J}\sum_{j=1}^{J} \frac{a_j^2\nu_j}{\lambda_j} \longrightarrow v_{\mathrm{diag}}(a;\pi,m),$$

and

$$\mathrm{Var}(D_J(a)) = \frac{1}{J^2}\sum_{j=1}^{J} \frac{1-\pi_j}{\pi_j}\left(\frac{a_j^2\nu_j}{\lambda_j}\right)^2 \leq \frac{C_0}{\bar{B}},$$

where we used $1/\pi_j = Jm_j/(\lambda_j\bar{B}) \leq J\bar{m}/(c_\lambda\bar{B})$. Consequently,

$$D_J(a) = v_{\mathrm{diag}}(a;\pi,m) + o_p(1).$$

For the Worker reuse, or the off-diagonal term, add and subtract the conditional mean of N_{jk} given $\mathcal{S}_J$ to isolate fluctuation from allocation:

$$C_J(a) = R_J^{\mathrm{alloc}}(a) + \sum_{j\neq k} \beta_{jk}^{\mathrm{ov}} I_j I_k \mathbb{E}[N_{jk} \mid \mathcal{S}_J].$$

Lemma 8 gives $R_J^{\mathrm{alloc}}(a) = o_p(1)$. On $\mathcal{H}_J$, for distinct selected documents $j,k \in \mathcal{S}_J$, define

$$\delta_{jk} := \mathbb{E}[N_{jk} \mid \mathcal{S}_J] - (K-1)\frac{m_j m_k}{B}.$$

By (C.15),

$$|\delta_{jk}| \leq C_0 \frac{m_j m_k}{B^2},$$

uniformly over these document pairs and sampling realizations in $\mathcal{H}_J$.

Write $b_{jk} := a_j a_k \tau_{jk}$ and define

$$U_J(a) := \frac{1}{J^2}\sum_{j\neq k} \frac{I_j I_k}{\pi_j \pi_k} b_{jk}.$$

The array $\{b_{jk}\}$ is symmetric and uniformly bounded. Define the remainder from approximating the conditional overlaps by

$$R_J^{\mathrm{mean}}(a) := \begin{cases} \displaystyle\sum_{\substack{j,k\in\mathcal{S}_J \\ j\neq k}} \beta_{jk}^{\mathrm{ov}}\delta_{jk}, & \text{on } \mathcal{H}_J, \\ 0, & \text{on } \mathcal{H}_J^{\mathrm{c}}. \end{cases}$$

Substituting the conditional-overlap approximation into the decomposition of $C_J(a)$ gives, on $\mathcal{H}_J$,

$$C_J(a) = R_J^{\text{alloc}}(a) + (K-1)\frac{\bar{B}}{B}U_J(a) + R_J^{\text{mean}}(a).$$

To bound the remainder, recall $M_J^{\text{ov}} = \max_{j\neq k} |\beta_{jk}^{\text{ov}}|$. Since $\sum_{j\in\mathcal{S}_J} m_j = B$, on $\mathcal{H}_J$,

$$\begin{aligned} |R_J^{\text{mean}}(a)| &\leq \frac{C_0 M_J^{\text{ov}}}{B^2} \sum_{\substack{j,k\in\mathcal{S}_J \\ j\neq k}} m_j m_k \\ &\leq C_0 M_J^{\text{ov}} \leq \frac{C_0}{\bar{B}}, \end{aligned}$$

where the last inequality follows from (C.17). Because $R_J^{\text{mean}}(a) = 0$ outside $\mathcal{H}_J$, it follows that

$$R_J^{\text{mean}}(a) = O_p(\bar{B}^{-1}) = o_p(1).$$

It remains to control the document-selection fluctuation in $U_J(a)$. Define

$$T_J(a) := \mathbb{E}[U_J(a)] = \frac{1}{J^2}\sum_{j\neq k} b_{jk}, \qquad \xi_j := \frac{I_j}{\pi_j} - 1, \qquad \bar{b}_j := \frac{1}{J}\sum_{k\neq j} b_{jk}.$$

Here ξ_j is the centered selection weight of document j, and $\bar{b}_j$ summarizes its pairwise contributions across the corpus. Expanding $(1+\xi_j)(1+\xi_k)-1 = \xi_j + \xi_k + \xi_j\xi_k$, and using symmetry and ordered-pair summation, gives

$$U_J(a) - T_J(a) = \frac{2}{J}\sum_{j=1}^{J} \bar{b}_j\xi_j + \frac{1}{J^2}\sum_{j\neq k} b_{jk}\xi_j\xi_k. \tag{C.23}$$

This is the Hoeffding decomposition with respect to the independent selection indicators. As for a second-order U-statistic, the first term collects the single-document projections: $\mathbb{E}[U_J(a) \mid I_j] - T_J(a) = (2/J)\bar{b}_j\xi_j$. The second term captures the remaining pairwise interactions. It is degenerate: its conditional mean given any individual I_j is zero, because every product retains an independent centered factor.

The ξ_j are independent and centered, with $\mathbb{E}[\xi_j^2] = (1-\pi_j)/\pi_j \leq C_0 J/\bar{B}$. Since the $\bar{b}_j$ are uniformly bounded, the linear term satisfies

$$\mathbb{E}\left[\left(\frac{2}{J}\sum_{j=1}^{J}\bar{b}_j\xi_j\right)^2\right] = \frac{4}{J^2}\sum_{j=1}^{J}\bar{b}_j^2\mathbb{E}[\xi_j^2] \leq \frac{C_0}{\bar{B}}.$$

For the quadratic term, independence and centering eliminate every cross-product containing an index that appears only once. In particular, two pairs sharing just one document contribute zero; only identical or reversed pairs remain. Symmetry therefore gives

$$\mathbb{E}\left[\left(\frac{1}{J^2}\sum_{j\neq k} b_{jk}\xi_j\xi_k\right)^2\right] = \frac{2}{J^4}\sum_{j\neq k} b_{jk}^2\mathbb{E}[\xi_j^2]\mathbb{E}[\xi_k^2] \leq \frac{C_0}{\bar{B}^2}.$$

Thus the linear and quadratic terms are $O_p(\bar{B}^{-1/2})$ and $O_p(\bar{B}^{-1})$, respectively. Both vanish, so $U_J(a) - T_J(a) = o_p(1)$. The maintained array limit $T_J(a) \to \tau(a)$ then yields

$$U_J(a) = \tau(a) + o_p(1) = O_p(1).$$

Finally, Lemma 5 gives $B/\bar{B} \xrightarrow{p} 1$. On $\mathcal{H}_J$, we have $\bar{B}/B \le 2$ and $|\bar{B}/B - 1| \le 2|B/\bar{B} - 1|$. The preceding decomposition of $C_J(a)$ therefore implies, on this event,

$$|C_J(a) - (K-1)\tau(a)| \le |R_J^{\text{alloc}}(a)| + |R_J^{\text{mean}}(a)| + 2(K-1)\left\{|U_J(a) - \tau(a)| + |\tau(a)|\left|\frac{B}{\bar{B}} - 1\right|\right\}.$$

Every term on the right converges to zero in probability. Since $\mathbb{P}(\mathcal{H}_J^c) \to 0$, it follows that, for every $\epsilon > 0$,

$$\mathbb{P}(|C_J(a) - (K-1)\tau(a)| > \epsilon) \le \mathbb{P}(\mathcal{H}_J^c) + \mathbb{P}(\mathcal{H}_J \cap \{|C_J(a) - (K-1)\tau(a)| > \epsilon\}) \longrightarrow 0.$$

Together with the diagonal convergence, this proves

$$v_{\varepsilon,J}(a) \xrightarrow{p} v_{\text{diag}}(a; \pi, m) + (K-1)\tau(a).$$

Thus Condition **(RW4-M)** holds with $v_\varepsilon(a) = v_{\text{diag}}(a; \pi, m) + (K-1)\tau(a)$, completing the proof. □

D Variance estimation and feasible inference

This section proves the unbiasedness and asymptotic consistency of the variance estimators introduced in the main text.

D.1 Unbiasedness

For notational convenience, set $\widehat{r}_j^2 = 0$ when $I_j = 0$.

Lemma 9 (Unbiasedness of the variance estimators)**.** *Suppose Assumptions **(RW0)**, **(S1)**–**(S2)**, and **(D2)** hold, the allocation is simple, and $m_j \ge 2$ whenever $\pi_j > 0$. Then, for every selected document j,*

$$\mathbb{E}[\widehat{\sigma}_j^2 \mid \mathcal{G}_J] = \nu_j, \qquad \mathbb{E}[\widehat{r}_j^2 \mid \mathcal{G}_J] = r_j^2,$$

and, for every pair of distinct selected documents,

$$\mathbb{E}[\widehat{\Gamma}_{jk} \mid \mathcal{G}_J] = N_{jk}\tau_{jk}.$$

Consequently,

$$\mathbb{E}[\widehat{v}_{\varepsilon,J}(a) \mid \mathcal{G}_J] = v_{\varepsilon,J}(a). \tag{D.1}$$

*If Assumption **(D1)** also holds, then*

$$\mathbb{E}[\widehat{v}_{r,J}(a)] = v_{r,J}(a), \tag{D.2}$$

and hence

$$\mathbb{E}[\widehat{v}_J(a)] = v_{r,J}(a) + \bar{v}_{\varepsilon,J}(a) = \bar{B}\ \text{Var}\Big(\widehat{\Phi}_J(a) - \Phi_J(a)\Big). \tag{D.3}$$

Proof. Condition on $\mathcal{G}_J$, and write $U_{ij} := Y_{ij}^* - \mu_j$. By (B.4) and the maintained response model, response vectors belonging to distinct workers are conditionally independent, and, for $j \ne k$, the relevant conditional moments satisfy

$$\mathbb{E}[U_{ij} \mid \mathcal{G}_J] = 0, \qquad \mathbb{E}[U_{ij}^2 \mid \mathcal{G}_J] = \nu_j, \qquad \mathbb{E}[U_{ij}U_{ik} \mid \mathcal{G}_J] = \tau_{jk}.$$

For each selected document j, graph simplicity ensures that its $m_j \geq 2$ responses come from distinct workers. Conditional on $\mathcal{G}_J$, these responses are independent with common mean μ_j and variance ν_j. Hence the usual sample-variance identity gives

$$\mathbb{E}[\widehat{\sigma}_j^2 \mid \mathcal{G}_J] = \nu_j.$$

Moreover,

$$\mathbb{E}[(\bar{Y}_j - g_j)^2 \mid \mathcal{G}_J] = \mathrm{Var}(\bar{Y}_j \mid \mathcal{G}_J) + (\mu_j - g_j)^2 = \frac{\nu_j}{m_j} + r_j^2.$$

Since $\widehat{r}_j^2 = (\bar{Y}_j - g_j)^2 - \widehat{\sigma}_j^2/m_j$, these two identities give $\mathbb{E}[\widehat{r}_j^2 \mid \mathcal{G}_J] = r_j^2$.

For the overlap identity, fix distinct selected documents j, k, and put $N := N_{jk}$. The claim is immediate if $N = 0$. If $N \geq 1$, fix $i \in M_j \cap M_k$, and define

$$\bar{U}_{j,-i} := \frac{1}{m_j - 1} \sum_{\ell \in M_j \setminus \{i\}} U_{\ell j}, \qquad \bar{U}_{k,-i} := \frac{1}{m_k - 1} \sum_{q \in M_k \setminus \{i\}} U_{qk}.$$

Since $Y_{ij}^* - \bar{Y}_{j,-i} = U_{ij} - \bar{U}_{j,-i}$ (and analogously for document k), the two mixed terms have conditional expectation zero: worker i's responses are independent of the centered responses entering either leave-out average. The two leave-out averages can nevertheless be correlated. In their product, only terms involving the same worker can contribute to the conditional expectation. There are exactly $N - 1$ such terms, one for each remaining shared worker, each with conditional expectation τ_{jk}. Therefore

$$\begin{aligned}
&\mathbb{E}\big[(Y_{ij}^* - \bar{Y}_{j,-i})(Y_{ik}^* - \bar{Y}_{k,-i}) \,\big|\, \mathcal{G}_J\big] \\
&\quad = \mathbb{E}[U_{ij}U_{ik} \mid \mathcal{G}_J] + \mathbb{E}[\bar{U}_{j,-i}\bar{U}_{k,-i} \mid \mathcal{G}_J] \\
&\qquad - \underbrace{\mathbb{E}[U_{ij}\bar{U}_{k,-i} \mid \mathcal{G}_J]}_{=0} - \underbrace{\mathbb{E}[\bar{U}_{j,-i}U_{ik} \mid \mathcal{G}_J]}_{=0} \\
&\quad = \tau_{jk} + \frac{N-1}{(m_j - 1)(m_k - 1)}\tau_{jk} \\
&\quad = \left\{1 + \frac{N-1}{(m_j - 1)(m_k - 1)}\right\}\tau_{jk}.
\end{aligned}$$

Summing over the N shared workers and applying the definition of $\kappa_{jk}^{\mathrm{loo}}$ gives

$$\mathbb{E}[\widehat{\Gamma}_{jk} \mid \mathcal{G}_J] = \kappa_{jk}^{\mathrm{loo}} N \left\{1 + \frac{N-1}{(m_j - 1)(m_k - 1)}\right\}\tau_{jk} = N\tau_{jk}.$$

Conditional on $\mathcal{G}_J$, all coefficients and overlap counts in $\widehat{v}_{\varepsilon,J}(a)$ are fixed. Applying the local identities term by term in (4.16) therefore proves (D.1).

Now suppose Assumption **(D1)** also holds. Since I_j is $\mathcal{G}_J$-measurable, the local identity and the convention for unselected documents give

$$\mathbb{E}[I_j\widehat{r}_j^2 \mid \mathcal{G}_J] = I_j r_j^2, \qquad \mathbb{E}[I_j\widehat{r}_j^2] = \pi_j r_j^2.$$

Substitution into (4.15) proves (D.2). Taking expectations in (D.1), adding the two components, and applying Lemma 4 proves (D.3). □

D.2 Concentration

Recall that

$$\lambda_j := \frac{J\pi_j m_j}{\bar{B}}$$

compares document j's expected label expenditure with $\bar{B}/J$, the expected expenditure under an equal division of the expected annotation budget across documents.

Assumption 2 (Feasible-inference regularity)**.** *The expected budget satisfies $\bar{B} \to \infty$, and, for fixed finite constants $\bar{m}$, $\bar{a}$, and $\bar{r}$,*

$$2 \le m_j \le \bar{m}, \qquad |a_j| \le \bar{a}, \qquad |r_j| \le \bar{r}$$

uniformly over j and J. There is a deterministic sequence $c_J \in (0,1]$ such that

$$\min_{j \le J} \lambda_j \ge c_J, \qquad c_J^3 \bar{B} \longrightarrow \infty.$$

The condition permits the minimum relative expenditure to approach zero. Under the balanced-expenditure condition in Theorem 2, c_J can be fixed, so the rate reduces to $\bar{B} \to \infty$. Bounded outcomes and Assumption **(RW2)** imply the stated residual bound.

Proposition 4 (Concentration of the variance estimators)**.** *Suppose the bounded-response-scale assumption and Assumptions **(RW0)**, **(S1)**–**(S2)**, **(D1)**, and **(D2)** hold. Suppose the allocation is simple and assigns at most $K < \infty$ documents to each worker. Under Assumption 2,*

$$\widehat{v}_{r,J}(a) - v_{r,J}(a) \xrightarrow{p} 0, \qquad \widehat{v}_{\varepsilon,J}(a) - v_{\varepsilon,J}(a) \xrightarrow{p} 0.$$

Proof. Throughout the proof, $C_0 < \infty$ denotes a constant depending only on the fixed bounds and may change from line to line. Let $R_Y < \infty$ be the width of the maintained outcome interval. Then $|Y_{ij}^* - g_j| \le R_Y + \bar{r}$.

Sampling component. Introduce the infeasible estimator where we suppose we can observe r_j

$$\widehat{v}_{r,J}^{\mathrm{or}}(a) := \frac{\bar{B}}{J^2} \sum_{j=1}^{J} \frac{I_j(1-\pi_j)}{\pi_j^2} a_j^2 r_j^2.$$

We first establish that the infeasible estimator is consistent

Independent Bernoulli selection gives $\mathbb{E}[\widehat{v}_{r,J}^{\mathrm{or}}(a)] = v_{r,J}(a)$ and

$$\mathrm{Var}(\widehat{v}_{r,J}^{\mathrm{or}}(a)) = \frac{\bar{B}^2}{J^4} \sum_{j=1}^{J} \frac{(1-\pi_j)^3}{\pi_j^3} a_j^4 r_j^4 \le \frac{C_0}{c_J^3 \bar{B}} \longrightarrow 0, \tag{D.4}$$

where we used $\pi_j = \lambda_j \bar{B}/(J m_j)$.

Now, we replace r_j^2 by $\widehat{r}_j^2$, and establish our estimator converges to the infeasible estimator in probaability. Write

$$R_J^{\mathrm{res}}(a) := \widehat{v}_{r,J}(a) - \widehat{v}_{r,J}^{\mathrm{or}}(a) = \sum_{j=1}^{J} \omega_j^{\mathrm{res}} I_j (\widehat{r}_j^2 - r_j^2), \qquad \omega_j^{\mathrm{res}} := \frac{\bar{B}}{J^2} \frac{1-\pi_j}{\pi_j^2} a_j^2.$$

Lemma 9 gives

$$\mathbb{E}[R_J^{\mathrm{res}}(a) \mid \mathcal{G}_J] = 0. \tag{D.5}$$

For the replacement bound, expand the corrected residual as

$$\widehat{r}_j^2 = \frac{1}{m_j(m_j-1)} \sum_{\substack{i,\ell \in M_j \\ i \neq \ell}} (Y_{ij}^* - g_j)(Y_{\ell j}^* - g_j). \tag{D.6}$$

For an active worker i, let $\mathbf{Y}_i := (Y_{ij}^* : j \in \mathcal{J}_i)$. Conditional on $\mathcal{G}_J$, these worker response vectors are independent by (B.4).

Let $R_J^{\mathrm{res},(i)}(a)$ denote the value of $R_J^{\mathrm{res}}(a)$ obtained by replacing worker i's responses with those of a new worker drawn independently from the reference population, evaluating the same assigned documents with fresh response shocks. The graph and all other workers' responses are held fixed. Under the conditional law, this replaces $\mathbf{Y}_i$ by an independent copy. We use this construction to apply the Efron–Stein inequality (Boucheron et al., 2013, Theorem 3.1) below.

Because $|Y_{ij}^* - g_j| \leq R_Y + \bar{r}$, replacing one label on document j changes only $2(m_j - 1)$ products in (D.6), each by at most $2(R_Y + \bar{r})^2$. Hence

$$|R_J^{\mathrm{res}}(a) - R_J^{\mathrm{res},(i)}(a)| \leq C_0 \sum_{j \in \mathcal{J}_i} \frac{\omega_j^{\mathrm{res}}}{m_j}.$$

Applying the Efron–Stein inequality (Boucheron et al., 2013, Theorem 3.1) conditionally on $\mathcal{G}_J$, and then using the preceding replacement bound, gives

$$\begin{aligned}
\mathrm{Var}(R_J^{\mathrm{res}}(a) \mid \mathcal{G}_J) &\leq \frac{1}{2} \sum_i \mathbb{E}\Big[\{R_J^{\mathrm{res}}(a) - R_J^{\mathrm{res},(i)}(a)\}^2 \,\Big|\, \mathcal{G}_J \Big] \\
&\leq C_0 \sum_i \left(\sum_{j \in \mathcal{J}_i} \frac{\omega_j^{\mathrm{res}}}{m_j} \right)^2 \\
&\leq C_0 \sum_i \sum_{j \in \mathcal{J}_i} \frac{(\omega_j^{\mathrm{res}})^2}{m_j^2} \\
&= C_0 \sum_{j \in \mathcal{S}_J} \frac{(\omega_j^{\mathrm{res}})^2}{m_j}.
\end{aligned} \tag{D.7}$$

The third inequality uses Cauchy–Schwarz and $|\mathcal{J}_i| \leq K$, with the fixed factor K absorbed into C_0. The last equality counts each selected document in exactly m_j worker neighborhoods.

because each selected document appears in exactly m_j worker neighborhoods. Therefore, Taking expec-

tations in (D.7) and using the conditional centering in (D.5) gives

$$\begin{aligned}
\mathbb{E}[R_J^{\mathrm{res}}(a)^2] &= \mathbb{E}[\mathrm{Var}(R_J^{\mathrm{res}}(a) \mid \mathcal{G}_J)] \\
&\le C_0\, \mathbb{E}\left[\sum_{j=1}^{J} I_j \frac{(\omega_j^{\mathrm{res}})^2}{m_j}\right] \\
&= C_0 \sum_{j=1}^{J} \pi_j \frac{(\omega_j^{\mathrm{res}})^2}{m_j} \\
&= C_0 \frac{\bar{B}^2}{J^4} \sum_{j=1}^{J} \frac{(1-\pi_j)^2 a_j^4}{\pi_j^3 m_j} \\
&= \frac{C_0}{J\bar{B}} \sum_{j=1}^{J} \frac{(1-\pi_j)^2 a_j^4 m_j^2}{\lambda_j^3} \\
&\le \frac{C_0}{c_J^3 \bar{B}} \longrightarrow 0.
\end{aligned} \tag{D.8}$$

Here we used $\mathbb{E}[I_j] = \pi_j$, the definition of ω_j^{res}, and $\pi_j = \lambda_j \bar{B}/(Jm_j)$. The final inequality uses $\lambda_j \ge c_J$, the boundedness of a_j and m_j, and $(1-\pi_j)^2 \le 1$, with fixed constants absorbed into C_0. The limit follows from $c_J^3 \bar{B} \to \infty$.

Thus $R_J^{\mathrm{res}}(a) = o_p(1)$. Combining this with (D.4) gives

$$\widehat{v}_{r,J}(a) - v_{r,J}(a) = R_J^{\mathrm{res}}(a) + \left\{\widehat{v}_{r,J}^{\mathrm{or}}(a) - v_{r,J}(a)\right\} = o_p(1),$$

which proves the sampling claim.

Measurement component. Let $\Delta_{\varepsilon,J}(a) := \widehat{v}_{\varepsilon,J}(a) - v_{\varepsilon,J}(a)$. Lemma 9 gives

$$\mathbb{E}[\Delta_{\varepsilon,J}(a) \mid \mathcal{G}_J] = 0. \tag{D.9}$$

We first show that replacing one worker's responses changes each $\widehat{\sigma}_j^2$ or $\widehat{\Gamma}_{jk}$ by at most a constant independent of J.

Responses lie in an interval of width R_Y, and every leave-out mean lies in the same interval. The sample variance can be written in terms of squared differences between responses:

$$\widehat{\sigma}_j^2 = \frac{1}{2m_j(m_j-1)} \sum_{\substack{\ell,\ell' \in M_j \\ \ell \ne \ell'}} (Y_{\ell j}^* - Y_{\ell' j}^*)^2.$$

Therefore,

$$\begin{aligned}
0 \le \widehat{\sigma}_j^2 &\le \frac{1}{2m_j(m_j-1)} \sum_{\substack{\ell,\ell' \in M_j \\ \ell \ne \ell'}} R_Y^2 = \frac{R_Y^2}{2}, \\
|\widehat{\Gamma}_{jk}| &\le \kappa_{jk}^{\mathrm{loo}} \sum_{i \in M_j \cap M_k} |Y_{ij}^* - \bar{Y}_{j,-i}|\, |Y_{ik}^* - \bar{Y}_{k,-i}| \\
&\le \kappa_{jk}^{\mathrm{loo}} N_{jk} R_Y^2 \le N_{jk} R_Y^2 \le \bar{m} R_Y^2.
\end{aligned} \tag{D.10}$$

The first line uses the $m_j(m_j-1)$ ordered pairs of distinct workers. The last line uses $0 \le \kappa_{jk}^{\mathrm{loo}} \le 1$ and $N_{jk} \le \min(m_j, m_k) \le \bar{m}$. These bounds hold before and after replacement, so the changes in the two

estimates are bounded by $R_Y^2/2$ and $2\bar{m}R_Y^2$, respectively.

We next show that replacing one worker's responses affects only a bounded number of these estimates. For each selected document j, let

$$\mathcal{N}_j := \{k \neq j : N_{jk} > 0\}$$

denote the set of documents sharing at least one worker with j. Each of the m_j workers assigned to j labels at most $K-1$ other documents, so

$$|\mathcal{N}_j| \leq m_j(K-1) \leq \bar{m}(K-1). \tag{D.11}$$

Replacing worker i's responses can change $\widehat{\sigma}_j^2$ only for the at most K documents in $\mathcal{J}_i$. To count the covariance estimates $\widehat{\Gamma}_{jk}$ that can change, define

$$\mathcal{A}_i := \{(j,k) : j,k \in \mathcal{S}_J,\ j \neq k,\ N_{jk} > 0,\ i \in M_j \cup M_k\}.$$

Since the pairs are ordered,

$$|\mathcal{A}_i| \leq 2\sum_{j \in \mathcal{J}_i} |\mathcal{N}_j| \leq 2K\bar{m}(K-1).$$

With the same block replacement, let $\widehat{v}_{\varepsilon,J}^{(i)}(a)$ denote the estimator recomputed after replacing $\mathbf{Y}_i$ by an independent copy from its conditional law given $\mathcal{G}_J$, with the graph fixed, and write

$$\omega_{j,J}(a) := \frac{\bar{B}|a_j|}{J\pi_j m_j} = \frac{|a_j|}{\lambda_j}.$$

The coefficient identities

$$\frac{\bar{B}a_j^2}{J^2\pi_j^2 m_j} = \frac{m_j}{\bar{B}}\,\omega_{j,J}(a)^2, \qquad \frac{\bar{B}|a_j a_k|}{J^2\pi_j\pi_k m_j m_k} = \frac{1}{\bar{B}}\,\omega_{j,J}(a)\omega_{k,J}(a).$$

Thus an affected diagonal summand changes by at most $C_0\omega_{j,J}(a)^2/\bar{B}$, while an affected off-diagonal summand (j,k) changes by at most $C_0\omega_{j,J}(a)\omega_{k,J}(a)/\bar{B}$. Together with (D.11), Cauchy–Schwarz over the bounded number of affected terms gives Summing the absolute changes over the affected terms gives

$$\begin{aligned}
&|\widehat{v}_{\varepsilon,J}(a) - \widehat{v}_{\varepsilon,J}^{(i)}(a)|^2 \\
&\quad\leq \frac{C_0}{\bar{B}^2}\left\{\sum_{j\in\mathcal{J}_i}\omega_{j,J}(a)^2 + \sum_{(j,k)\in\mathcal{A}_i}\omega_{j,J}(a)\omega_{k,J}(a)\right\}^2 \\
&\quad\leq \frac{C_0\big(|\mathcal{J}_i| + |\mathcal{A}_i|\big)}{\bar{B}^2}\left\{\sum_{j\in\mathcal{J}_i}\omega_{j,J}(a)^4 + \sum_{(j,k)\in\mathcal{A}_i}\omega_{j,J}(a)^2\omega_{k,J}(a)^2\right\} \\
&\quad\leq \frac{C_0}{\bar{B}^2}\left\{\sum_{j\in\mathcal{J}_i}\omega_{j,J}(a)^4 + \sum_{(j,k)\in\mathcal{A}_i}\omega_{j,J}(a)^2\omega_{k,J}(a)^2\right\}.
\end{aligned} \tag{D.12}$$

The second inequality applies Cauchy–Schwarz to the $|\mathcal{J}_i| + |\mathcal{A}_i|$ terms. The final inequality uses

$$|\mathcal{J}_i| + |\mathcal{A}_i| \leq K + 2K\bar{m}(K-1),$$

with this fixed bound absorbed into C_0.

The first term in (D.12) satisfies

$$\sum_i \sum_{j \in \mathcal{J}_i} \omega_{j,J}(a)^4 = \sum_{j \in \mathcal{S}_J} m_j \omega_{j,J}(a)^4 \leq \bar{m} \sum_{j \in \mathcal{S}_J} \omega_{j,J}(a)^4.$$

Each ordered overlapping pair (j,k) belongs to at most $|M_j \cup M_k| \leq 2\bar{m}$ of the sets $\mathcal{A}_i$. Moreover, symmetry of the overlap relation and $2\omega_{j,J}(a)^2\omega_{k,J}(a)^2 \leq \omega_{j,J}(a)^4 + \omega_{k,J}(a)^4$ give

$$\sum_{\substack{j \neq k \\ N_{jk}>0}} \omega_{j,J}(a)^2 \omega_{k,J}(a)^2 \leq \sum_{j \in \mathcal{S}_J} |\mathcal{N}_j| \omega_{j,J}(a)^4 \leq \bar{m}(K-1) \sum_{j \in \mathcal{S}_J} \omega_{j,J}(a)^4.$$

Conditional Efron–Stein applied to (D.12) therefore yields

$$\operatorname{Var}(\widehat{v}_{\varepsilon,J}(a) \mid \mathcal{G}_J) \leq \frac{C_0}{\bar{B}^2} \sum_{j \in \mathcal{S}_J} \omega_{j,J}(a)^4. \tag{D.13}$$

Since $v_{\varepsilon,J}(a)$ is $\mathcal{G}_J$-measurable, (D.9) and (D.13) give

$$\begin{aligned} \mathbb{E}[\Delta_{\varepsilon,J}(a)^2] &= \mathbb{E}[\operatorname{Var}(\widehat{v}_{\varepsilon,J}(a) \mid \mathcal{G}_J)] \\ &\leq \frac{C_0}{\bar{B}^2} \sum_{j=1}^{J} \pi_j \frac{|a_j|^4}{\lambda_j^4} = \frac{C_0}{J\bar{B}} \sum_{j=1}^{J} \frac{|a_j|^4}{m_j \lambda_j^3} \leq \frac{C_0}{c_J^3 \bar{B}} \longrightarrow 0. \end{aligned}$$

The first inequality uses $\mathbb{E}[\sum_{j \in \mathcal{S}_J} \omega_{j,J}(a)^4] = \sum_{j=1}^{J} \pi_j \omega_{j,J}(a)^4$. Thus $\Delta_{\varepsilon,J}(a) = o_p(1)$, proving the measurement claim and the proposition. □

D.3 Joint inference for weighted corpus averages

In this section, we establish joint inference results for a fixed-dimensional vector of weighted corpus averages and its smooth transformations.

D.3.1 Moment vector and linear combinations

Let $a^{(1)}, \ldots, a^{(q)}$ be target-weight arrays, each satisfying the maintained boundedness and prespecification conditions on a, where $q < \infty$ is fixed. Define

$$\boldsymbol{\Phi}_J := \big(\Phi_J(a^{(1)}), \ldots, \Phi_J(a^{(q)})\big)^\top, \qquad \Phi_J(a^{(r)}) = \frac{1}{J} \sum_{j=1}^{J} a_j^{(r)} \mu_j.$$

For each coordinate r, let $g_j^{(r)}$ satisfy the maintained conditions on frozen predictions. Using the same document sample and worker allocation, apply (4.1) coordinatewise:

$$\widehat{\Phi}_{Jr} := \frac{1}{J} \sum_{j=1}^{J} a_j^{(r)} g_j^{(r)} + \frac{1}{J} \sum_{j \in \mathcal{S}_J} \frac{a_j^{(r)}}{\pi_j} \big(\bar{Y}_j - g_j^{(r)}\big).$$

Collect these estimates in $\widehat{\boldsymbol{\Phi}}_J := (\widehat{\Phi}_{J1}, \ldots, \widehat{\Phi}_{Jq})^\top$.

For any $t \in \mathbb{R}^q$ fixed independently of J, let

$$a_j(t) := \sum_{r=1}^{q} t_r a_j^{(r)}.$$

Linearity gives

$$t^\top \widehat{\mathbf{\Phi}}_J = \frac{1}{J}\sum_{j=1}^{J}\sum_{r=1}^{q} t_r a_j^{(r)} g_j^{(r)} + \frac{1}{J}\sum_{j\in\mathcal{S}_J}\frac{1}{\pi_j}\left\{a_j(t)\bar{Y}_j - \sum_{r=1}^{q} t_r a_j^{(r)} g_j^{(r)}\right\}. \tag{D.14}$$

This is the scalar estimator (4.1) applied to responses $a_j(t)Y_{ij}^*$, frozen predictions $\sum_{r=1}^{q} t_r a_j^{(r)} g_j^{(r)}$, and unit outer weights. Its target is

$$\frac{1}{J}\sum_{j=1}^{J} a_j(t)\mu_j = t^\top \mathbf{\Phi}_J.$$

For each fixed t, these transformed responses and predictions remain uniformly bounded and preserve the maintained response and design conditions. The preceding scalar arguments therefore apply under the corresponding sampling- and measurement-variance limit conditions.

D.3.2 Covariance estimation

The scalar variance estimator also supplies the covariances between coordinates. The identity

$$\mathrm{Cov}(X,Y) = \frac{1}{2}\left\{\mathrm{Var}(X+Y) - \mathrm{Var}(X) - \mathrm{Var}(Y)\right\}$$

shows that it is enough to estimate the variance of each coordinate and each coordinate sum.

For each fixed $t \in \mathbb{R}^q$, let $\widehat{V}_J(t)$ denote the raw scalar variance estimator in (4.17), evaluated with responses $a_j(t)Y_{ij}^*$, frozen predictions $\sum_{r=1}^{q} t_r a_j^{(r)} g_j^{(r)}$, and unit outer weights. Let e_r denote the rth standard basis vector in $\mathbb{R}^q$. Define

$$\begin{aligned}\widehat{\Sigma}_{J,rr} &:= \widehat{V}_J(e_r),\\ \widehat{\Sigma}_{J,rs} &:= \frac{1}{2}\left\{\widehat{V}_J(e_r+e_s) - \widehat{V}_J(e_r) - \widehat{V}_J(e_s)\right\}, \qquad r \neq s.\end{aligned} \tag{D.15}$$

All evaluations use the same observed annotations. Consequently, the covariance estimates account for the dependence induced by shared documents and worker reuse through the existing scalar variance routine.

D.3.3 Joint limit and covariance consistency

The preceding scalar results yield joint normality and consistency of the covariance estimator.

Corollary 2 (Joint inference for weighted means)**.** *Maintain the conditions of Theorem 2, with the limits in* (4.10) *required for each fixed linear combination constructed above. Suppose also that the sampling-variance limit* (4.6) *holds for each such combination and that $m_j \geq 2$ for every document.*

For each fixed $t \in \mathbb{R}^q$, let $V(t)$ denote the sum of the limiting sampling and measurement variances of the corresponding scalar estimator. Define

$$\Sigma_{rr} := V(e_r), \qquad \Sigma_{rs} := \frac{1}{2}\{V(e_r+e_s) - V(e_r) - V(e_s)\}, \quad r \neq s. \tag{D.16}$$

Then Σ is nonnegative semidefinite, and

$$\sqrt{\bar{B}}\,(\widehat{\mathbf{\Phi}}_J - \mathbf{\Phi}_J) \Rightarrow \mathcal{N}_q(0,\Sigma), \qquad \widehat{\Sigma}_J \xrightarrow{p} \Sigma. \tag{D.17}$$

Proof. For each fixed t, the linear combination in (D.14) has the scalar prediction-assisted form, with bounded transformed responses and frozen predictions. Theorem 2 establishes its measurement-variance limit. Corollary 1 therefore gives the scalar Gaussian limit whenever $V(t) > 0$. If $V(t) = 0$, both component variance limits vanish; Chebyshev's inequality for the sampling term and conditional Chebyshev's inequality for the measurement term give convergence of the scaled error to zero in probability.

Both variance components are quadratic in t. Their limits therefore satisfy $V(t) = t^\top \Sigma t$, with Σ defined in (D.16). Since $V(t) \geq 0$ for every t, Σ is Positive semidefinite. The Cramér–Wold device gives the joint Gaussian limit.

The scalar variance-consistency result gives

$$\widehat{V}_J(e_r) \xrightarrow{p} V(e_r), \qquad \widehat{V}_J(e_r + e_s) \xrightarrow{p} V(e_r + e_s).$$

Applying the covariance identity in (D.15) gives consistency of every entry of $\widehat{\Sigma}_J$. Because q is fixed, this establishes $\widehat{\Sigma}_J \xrightarrow{p} \Sigma$. □

D.4 Smooth transformations and implicit parameters

Throughout this subsection, suppose the conclusions of Corollary 2 hold.

Proposition 5 (Smooth transformations)**.** *Suppose $\boldsymbol{\Phi}_J \to \boldsymbol{\Phi}_0$, and let $H : \mathcal{O} \subseteq \mathbb{R}^q \to \mathbb{R}^p$ be continuously differentiable on an open neighborhood of $\boldsymbol{\Phi}_0$. Define*

$$\boldsymbol{\beta}_J := H(\boldsymbol{\Phi}_J), \qquad \widehat{\boldsymbol{\beta}}_J := H(\widehat{\boldsymbol{\Phi}}_J),$$

and write

$$G := DH(\boldsymbol{\Phi}_0), \qquad \widehat{G}_J := DH(\widehat{\boldsymbol{\Phi}}_J).$$

Then

$$\sqrt{\bar{B}}\,(\widehat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}_J) \Rightarrow \mathcal{N}_p(0, \Omega), \qquad \Omega := G\Sigma G^\top, \tag{D.18}$$

and

$$\widehat{\Omega}_J := \widehat{G}_J \widehat{\Sigma}_J \widehat{G}_J^\top \xrightarrow{p} \Omega. \tag{D.19}$$

Proof. The joint limit gives $\widehat{\boldsymbol{\Phi}}_J - \boldsymbol{\Phi}_J = O_p(\bar{B}^{-1/2})$. Continuous differentiability therefore yields

$$\sqrt{\bar{B}}\,(\widehat{\boldsymbol{\beta}}_J - \boldsymbol{\beta}_J) = G\sqrt{\bar{B}}\,(\widehat{\boldsymbol{\Phi}}_J - \boldsymbol{\Phi}_J) + o_p(1).$$

The Gaussian limit follows by Slutsky's theorem. Continuity of DH gives $\widehat{G}_J \xrightarrow{p} G$; together with $\widehat{\Sigma}_J \xrightarrow{p} \Sigma$, this proves covariance consistency. □

Implicitly defined parameters. The same argument applies to a parameter defined by

$$F_J(\boldsymbol{\beta}_J, \boldsymbol{\Phi}_J) = 0,$$

where $F_J : \mathbb{R}^p \times \mathbb{R}^q \to \mathbb{R}^p$ is known under the fixed-corpus convention. Let $\widehat{\boldsymbol{\beta}}_J$ be a consistent local solution of $F_J(\boldsymbol{b}, \widehat{\boldsymbol{\Phi}}_J) = 0$. Suppose F_J is continuously differentiable on neighborhoods of a common positive radius around $(\boldsymbol{\beta}_J, \boldsymbol{\Phi}_J)$, with first derivatives locally equicontinuous uniformly in J. A sufficient condition is that F_J is twice continuously differentiable on these neighborhoods, with second derivatives uniformly bounded over the neighborhoods and across J. Assume their values at the target converge, with a nonsingular limit for $D_\beta F_J$.

The standard first-order expansion for estimating equations (**?**, Section 3.1) then gives (D.18) with

$$G = -\lim_{J\to\infty} \{D_\beta F_J(\boldsymbol{\beta}_J, \boldsymbol{\Phi}_J)\}^{-1} D_\Phi F_J(\boldsymbol{\beta}_J, \boldsymbol{\Phi}_J).$$

A consistent covariance estimator is

$$\begin{aligned} \widehat{G}_J &:= -\left\{D_\beta F_J(\widehat{\boldsymbol{\beta}}_J, \widehat{\boldsymbol{\Phi}}_J)\right\}^{-1} D_\Phi F_J(\widehat{\boldsymbol{\beta}}_J, \widehat{\boldsymbol{\Phi}}_J), \\ \widehat{\Omega}_J &:= \widehat{G}_J \widehat{\Sigma}_J \widehat{G}_J^\top \xrightarrow{p} G\Sigma G^\top. \end{aligned} \tag{D.20}$$

For either construction, a coordinate with positive limiting variance has estimated standard error

$$\widehat{\mathrm{se}}(\widehat{\beta}_{Jr}) = \sqrt{\widehat{\Omega}_{J,rr}/B}.$$

Wald inference for fixed linear restrictions follows whenever the limiting covariance of the tested restrictions is positive definite.

E Graph implementation

E.1 An edge-pair implementation of uniform allocation

When the rejection sampler described in the main text is computationally costly, we approximate the uniform graph allocation using the standard edge-switch chain (**?**, Section 6). We apply existing mixing results to this implementation and show that vanishing graph-sampling error preserves the baseline design's asymptotic conclusions.

Implementation by degree-preserving swaps. Conditional on a realized document sample $\mathcal{S}_J$ with $B = \sum_{j\in\mathcal{S}_J} m_j > 0$ for which the prescribed document depths and capacity-filled worker loads are feasible, write

$$\mathfrak{A}_J^{\mathrm{cf}} := \mathfrak{A}_J(\boldsymbol{d}_J, \boldsymbol{k}_J^{\mathrm{cf}})$$

for the nonempty set of simple bipartite graphs with these labelled document and worker vertices and these degree sequences, and let $\mathsf{U}_J^A := \mathrm{Unif}(\mathfrak{A}_J^{\mathrm{cf}})$. A feasible starting graph may, for example, be found by a deterministic bipartite-flow construction.

If $\mathfrak{A}_J^{\mathrm{cf}}$ is a singleton, return its unique graph and set $P_J^e = I$. Otherwise, starting from any feasible graph, use the standard edge-switch chain (**?**, Section 6) with the following attempted transition:

1. With probability one half, leave the graph unchanged.
2. Otherwise, choose an unordered pair of distinct current edges uniformly from the $\binom{B}{2}$ possibilities. If the edges are vertex-disjoint and replacing them by the two cross-edges preserves simplicity, perform that swap; otherwise leave the graph unchanged.

Denote this edge-pair kernel by P_J^e. Choose the starting graph and the number of attempted transitions from the sample and margins before running the chain, and use randomization independent of worker response blocks. Every transition preserves the prescribed document depths, worker loads, and simplicity. Thus the implementation retains these features of the theoretical design at every run length; mixing controls how closely the final graph follows the required uniform law.

We now verify that the edge-pair procedure has the same guarantees from the standard swap-chain theory. Its kernel is symmetric, hence a uniformly distributed graph remains uniformly distributed after

another attempted transition. Any feasible graph can be reached from any other through a sequence of valid swaps (Erdős et al., 2018). The probability of staying put rules out a fixed periodic alternation between groups of graphs. These properties imply that the graph distribution converges to U_J^A from every feasible starting graph, by the standard convergence theorem for finite Markov chains (**?**, Theorem 4.9).

The chain studied by Erdős et al. (2018) proposes vertex pairs, whereas the implementation above proposes edge pairs. The following comparison allows us to use their mixing theorem for this proposal rule. Suppose $\mathfrak{A}_J^{\mathrm{cf}}$ is non-singleton (hence $B \geq 2$, $|\mathcal{S}_J| \geq 2$, and $n \geq 2$), and let P_J^v denote their lazy vertex-pair swap chain. For distinct graphs $G, H \in \mathfrak{A}_J^{\mathrm{cf}}$ that differ by one valid swap,

$$P_J^e(G,H) = \varrho_J P_J^v(G,H), \qquad \varrho_J := \frac{\binom{|\mathcal{S}_J|}{2}\binom{n}{2}}{\binom{B}{2}}; \tag{E.1}$$

If $G \neq H$ and H cannot be obtained from G by one valid swap, both transition probabilities are zero. Thus both procedures allow exactly the same changes of graph.

For each current graph G, the probability of staying put is one minus the sum of the probabilities of moving to other graphs. The preceding comparison therefore also determines the holding probabilities:

$$P_J^e(G,G) = 1 - \varrho_J\big[1 - P_J^v(G,G)\big].$$

Combining the probabilities of moving and staying gives

$$P_J^e = I + \varrho_J(P_J^v - I).$$

For either chain, define its spectral gap by

$$\operatorname{gap}(P) := 1 - \lambda_2(P),$$

where $\lambda_2(P)$ is the second-largest eigenvalue. For these lazy chains, the gap controls how quickly departures from the uniform distribution decay. The matrix identity above transforms each eigenvalue λ of P_J^v into $1 + \varrho_J(\lambda - 1)$ for P_J^e. Since $\varrho_J > 0$, their ordering is preserved, giving

$$\operatorname{gap}(P_J^e) = \varrho_J \operatorname{gap}(P_J^v). \tag{E.2}$$

To transfer the polynomial mixing guarantee, it is enough to bound the change in the inverse spectral gap by a polynomial in B. Since the non-singleton case has at least two document vertices and two worker vertices,

$$\varrho_J^{-1} = \frac{\binom{B}{2}}{\binom{|\mathcal{S}_J|}{2}\binom{n}{2}} \leq \binom{B}{2}.$$

The gap comparison therefore gives

$$\frac{1}{\operatorname{gap}(P_J^e)} \leq \binom{B}{2} \frac{1}{\operatorname{gap}(P_J^v)}.$$

Thus any polynomial bound on the inverse spectral gap of the vertex-pair chain also holds, with at most an additional quadratic factor, for the edge-pair chain.

The same degree bounds also ensure that the published sufficient condition for rapid mixing eventually

holds. Put

$$\Delta_J := \max\left\{\max_{j\in\mathcal{S}_J} m_j, \max_{i\le n} k_i\right\}.$$

If $K \ge 2$, then on $\mathcal{H}_J$, for all sufficiently large rows, capacity filling gives $\Delta_J \ge K \ge 2$. Bounded depths and loads, together with the growing label count on this event, give $\Delta_J \le \sqrt{B/2}$. Thus Erdős et al. (2018, Theorem 2) applies to P_J^v, and the gap comparison transfers its polynomial mixing guarantee to P_J^e. In particular, for $\eta \in (0,1)$, the mixing time

$$\tau_J^e(\eta) := \min\left\{t : \sup_{G\in\mathfrak{A}_J^{\mathrm{cf}}} \left\|\delta_G (P_J^e)^t - \mathsf{U}_J^A\right\|_{\mathrm{TV}} \le \eta\right\}$$

is bounded by a polynomial in B and $\log(\eta^{-1})$ on the regular event for all sufficiently large rows. It should be noted that, this is an asymptotic implementability guarantee. It does not supply a universal finite-sample burn-in constant or a stopping diagnostic based on the observed chain.

Consequences for the statistical theory. The mixing result concerns the assignment graph. The next proposition translates that approximation into a guarantee for the complete design-and-response experiment. It separates conclusions that require only a simple exogenous graph, which remain exact, from the asymptotic conclusions established under uniform allocation, which are preserved when the mixing error vanishes.

Proposition 6 (Finite-run graph implementation)**.** *Suppose $K \ge 2$, $\Pr(\mathcal{H}_J^c) \to 0$, and let $0 < \varepsilon_J^{\mathrm{mix}} \downarrow 0$. On the regular allocation event $\mathcal{H}_J$, choose a feasible starting graph and an integer t_J measurably from the sample and margins, with $t_J \ge \tau_J^e(\varepsilon_J^{\mathrm{mix}})$, and run P_J^e for t_J attempted transitions. Off $\mathcal{H}_J$, use any simple exogenous allocation with worker loads bounded by K, retaining the standing fallback when the capacity-filled margins are infeasible. If $\widetilde{\mathbb{P}}_J$ and $\mathbb{P}_J$ denote the laws of the complete design-and-response experiment (the sample, margins, final assignment graph, and responses) under, respectively, this implementation and exact uniform graph allocation, then*

$$\left\|\widetilde{\mathbb{P}}_J - \mathbb{P}_J\right\|_{\mathrm{TV}} \le \varepsilon_J^{\mathrm{mix}} + \Pr(\mathcal{H}_J^c) \longrightarrow 0. \tag{E.3}$$

Consequently every baseline-design probability limit, weak limit, studentization result, and coverage conclusion proved under exact uniform allocation also holds under the finite-run implementation.

Proof. On $\mathcal{H}_J$, given the sampled documents and the prescribed document and worker degrees, we run the swap procedure for $t_J \ge \tau_J^e(\varepsilon_J^{\mathrm{mix}})$ attempted transitions. By the previous result, the resulting graph distribution is therefore within $\varepsilon_J^{\mathrm{mix}}$ of the uniform distribution U_J^A in total variation. On $\mathcal{H}_J^c$ the discrepancy is at most one. the feasible implementation procedure above, and the ideal uniform draw use the same distribution of sampled documents and annotation dpeths which we conditioned upon. Weighting the error bounds by its corresponding probability then gives the right-hand side of (E.3).

Given the realized graph and preceding design variables, responses are generated in the same way under the finite-run implementation and ideal uniform allocation. The same error bound therefore remains valid after including the responses. In particular, the probabilities of any event defined by the resulting data or statistics—such as a confidence interval covering the target—differ between the two procedures by at most $\varepsilon_J^{\mathrm{mix}} + \Pr(\mathcal{H}_J^c)$. Since this bound tends to zero, the asserted asymptotic conclusions continue to hold. □

F First-order allocation for weighted means

This section gives the complete allocation characterization for the scalar weighted-mean target under the baseline Poisson design in Section 5. We maintain the design restrictions stated there.

Documents with zero target weight are removed before assigning the budget, with the remaining weights rescaled to preserve the target.[4] Henceforth, J and a_j denote the resulting corpus size and weights, so $a_j \neq 0$.

Motivated by the variance expansion in (5.1), we define the first-order oracle as a minimizer of

$$L_J(\pi, m; a) := \sum_{j=1}^{J} a_j^2 \left\{ \frac{1-\pi_j}{\pi_j} r_j^2 + \frac{\nu_j}{\pi_j m_j} \right\}, \tag{F.1}$$

where $r_j = \mu_j - g_j$.

Let $s_j := \pi_j m_j$ denote expected label expenditure on document j. We consider the continuous relaxation with $2 \leq m_{\min} \leq \bar{m} < \infty$ and

$$0 < \pi_j \leq 1, \qquad m_{\min} \leq m_j \leq \bar{m}, \qquad s_j \geq c_\lambda \frac{\bar{B}}{J}, \qquad \sum_{j=1}^{J} s_j = \bar{B}.$$

Write

$$\underline{s}_J := c_\lambda \frac{\bar{B}}{J}, \qquad \overline{s}_J := \bar{m}. \tag{F.2}$$

Assume $0 < \bar{B} \leq J\bar{m}$ and $0 < c_\lambda \leq 1$, so this continuous policy class is nonempty.

To state the result, define the coverage-first implementation

$$\pi_j^\star(s) = \min\left\{1, \frac{s}{m_{\min}}\right\}, \qquad m_j^\star(s) = \frac{s}{\pi_j^\star(s)}, \tag{F.3}$$

and its associated loss

$$\phi_j(s) := \frac{\nu_j}{s} + r_j^2 \left(\frac{m_{\min}}{s} - 1\right)_+ = \begin{cases} \dfrac{\nu_j + m_{\min} r_j^2}{s} - r_j^2, & s \leq m_{\min}, \\ \dfrac{\nu_j}{s}, & s \geq m_{\min}. \end{cases} \tag{F.4}$$

Proposition 7 (Continuous first-order allocation)**.** *An optimal policy can be implemented using* (F.3)*, with expenditures solving the convex program*

$$\min_{\{s_j\}} \sum_{j=1}^{J} a_j^2 \phi_j(s_j) \quad \textit{subject to} \quad \sum_{j=1}^{J} s_j = \bar{B}, \qquad \underline{s}_J \leq s_j \leq \overline{s}_J. \tag{F.5}$$

There is a common multiplier $\xi \geq 0$ such that each $s_j^\star$ minimizes $a_j^2 \phi_j(s) + \xi s$ over $[\underline{s}_J, \overline{s}_J]$, with $\sum_j s_j^\star = \bar{B}$.

[4] Let J_0 be the original corpus size, $\mathcal{J}_+ := \{j : a_j \neq 0\}$, and $J_+ := |\mathcal{J}_+|$. For $j \in \mathcal{J}_+$, set $a_j^+ := (J_+/J_0)a_j$. Then

$$\frac{1}{J_+} \sum_{j \in \mathcal{J}_+} a_j^+ \mu_j = \frac{1}{J_0} \sum_{j=1}^{J_0} a_j \mu_j.$$

If all weights are zero, the target is known and no allocation is needed.

Define

$$b_{j,-} := a_j^2(\nu_j + m_{\min} r_j^2), \qquad b_{j,+} := a_j^2 \nu_j.$$

For $\xi > 0$, let

$$s_j^{\mathrm{unc}}(\xi) := \begin{cases} \sqrt{b_{j,-}/\xi}, & \xi > b_{j,-}/m_{\min}^2, \\ m_{\min}, & b_{j,+}/m_{\min}^2 \le \xi \le b_{j,-}/m_{\min}^2, \\ \sqrt{b_{j,+}/\xi}, & 0 < \xi < b_{j,+}/m_{\min}^2. \end{cases} \tag{F.6}$$

Impose the spending limits by raising values below $\underline{s}_J$ to that lower bound and reducing values above $\overline{s}_J$ to that upper bound:

$$s_j^\star = \min\left\{\overline{s}_J, \max\left\{\underline{s}_J, s_j^{\mathrm{unc}}(\xi)\right\}\right\}, \qquad \sum_{j=1}^{J} s_j^\star = \bar{B}. \tag{F.7}$$

If $\xi = 0$, choose expenditures that attain each document's minimum loss over its allowed interval while meeting the total budget:

$$s_j^\star \in \arg\min_{\underline{s}_J \le s \le \overline{s}_J} a_j^2 \phi_j(s), \qquad \sum_{j=1}^{J} s_j^\star = \bar{B}.$$

Proof. At fixed expenditure $s = \pi_j m_j$, the derivative of document j's contribution with respect to coverage is

$$\frac{\partial}{\partial \pi_j} a_j^2 \left\{ \frac{\nu_j}{s} + r_j^2 \left(\frac{1}{\pi_j} - 1 \right) \right\} = -\frac{a_j^2 r_j^2}{\pi_j^2} \le 0.$$

The depth restrictions give the feasible coverage interval

$$\frac{s}{\bar{m}} \le \pi_j \le \min\left\{1, \frac{s}{m_{\min}}\right\}.$$

Thus the largest feasible coverage minimizes the loss, giving (F.3). This choice is unique when $r_j \neq 0$; when $r_j = 0$, all feasible coverage–depth combinations at the same expenditure have the same loss. Substitution gives (F.4) and reduces the problem to (F.5).

For $\xi \ge 0$, define

$$h_j(s;\xi) := a_j^2 \phi_j(s) + \xi s.$$

Away from $m_{\min}$,

$$\frac{\partial h_j(s;\xi)}{\partial s} = \begin{cases} \xi - b_{j,-}/s^2, & s < m_{\min}, \\ \xi - b_{j,+}/s^2, & s > m_{\min}. \end{cases}$$

The derivative is nondecreasing on each branch and increases by

$$\frac{b_{j,-} - b_{j,+}}{m_{\min}^2} = \frac{a_j^2 r_j^2}{m_{\min}} \ge 0$$

at their common endpoint. Since h_j is continuous there, it is convex. Taking $\xi = 0$ also shows that the weighted loss $a_j^2 \phi_j$ is nonincreasing.

Since the objective is convex and finite on $(0, \infty)^J$ and all constraints are affine, the subgradient KKT conditions are necessary and sufficient for global optimality (**?**, Corollaries 28.2.2 and 28.3.1). We verify these conditions by minimizing $h_j(s;\xi)$ over each document's spending interval and choosing the common multiplier to satisfy the budget.

Suppose $\xi > 0$ and $b_{j,-} > 0$. Setting the derivative equal to zero on each smooth branch gives the

square-root cases in (F.6). The minimum occurs at $m_{\min}$ when the left derivative is nonpositive and the right derivative is nonnegative:

$$\xi - \frac{b_{j,-}}{m_{\min}^2} \le 0 \le \xi - \frac{b_{j,+}}{m_{\min}^2},$$

which gives the middle case. Convexity then implies that an unrestricted minimum outside the allowed spending interval is replaced by the nearest endpoint, as in (F.7). If $b_{j,-} = 0$, then $h_j(s;\xi) = \xi s$, whose minimum is at $\underline{s}_J$, also as given by that formula. Write $s_j(\xi)$ for this bounded spending choice.

We next show that a multiplier can be chosen to meet the budget. Let d_j be the smallest expenditure attaining the minimum of $a_j^2\phi_j$ over its allowed interval. Continuity and monotonicity imply that the full set of minimizing expenditures is $[d_j, \overline{s}_J]$. The explicit spending rule is continuous and nonincreasing in $\xi > 0$, with

$$\lim_{\xi \downarrow 0} s_j(\xi) = d_j.$$

Moreover, $s_j(\xi) = \underline{s}_J$ for every j whenever

$$\xi > \max_{k \le J} \frac{b_{k,-}}{\underline{s}_J^2}.$$

If $\bar{B} < \sum_j d_j$, continuity and $J\underline{s}_J = c_\lambda \bar{B} \le \bar{B}$ therefore give a positive multiplier for which $\sum_j s_j(\xi) = \bar{B}$. The case $\bar{B} = J\underline{s}_J$ is included because the lower bounds are attained at a finite multiplier. If instead $\bar{B} \ge \sum_j d_j$, take $\xi = 0$ and choose $s_j^\star \in [d_j, \overline{s}_J]$ whose sum is $\bar{B}$; such a choice exists because $\bar{B} \le J\overline{s}_J$.

Finally, for any feasible allocation s, the document-level minimizing property and the common budget give

$$\sum_{j=1}^{J} a_j^2 \phi_j(s_j^\star) + \xi \bar{B} \le \sum_{j=1}^{J} a_j^2 \phi_j(s_j) + \xi \bar{B}.$$

Cancelling $\xi\bar{B}$ proves global optimality. Implementing these expenditures through (F.3) completes the proof. □

The allocation problem has the bounded separable convex form studied by **?**, whose Lemma 1 gives the corresponding optimality conditions using one-sided derivatives. For the present objective, (F.6) and (F.7) give each document's expenditure as a function of the common multiplier. In the case $\xi > 0$, the budget equation $\sum_j s_j(\xi) = \bar{B}$ can be solved by bisection because total expenditure is continuous and nonincreasing in ξ. For a discussion of multiplier-search methods, see Patriksson (2008, Section 3.1).

F.1 A default without document-specific difficulty estimates

The first-order oracle depends on the unknown document-specific quantities r_j^2 and ν_j. To describe a simple allocation when these quantities cannot be estimated from a pilot or comparable annotated data, let $\mathcal{I}_J$ denote the information available when the design is chosen, and let $\mathbb{E}_0(\cdot \mid \mathcal{I}_J)$ denote the researcher's working pre-annotation expectation. This expectation is used only to choose the design; it does not alter the fixed-corpus inferential model. The conditional optimization below is pointwise in $\mathcal{I}_J$; in particular, candidate expenditures are $\mathcal{I}_J$-measurable.

On the minimum-depth portion of the allocation problem, the relevant combined difficulty is

$$d_j^{\text{diff}} := \nu_j + m_{\min} r_j^2.$$

Suppose the working expectation does not distinguish documents by this quantity:

$$\mathbb{E}_0(d_j^{\text{diff}} \mid \mathcal{I}_J) = d_J^0 \in (0, \infty), \qquad j = 1, \ldots, J. \tag{F.8}$$

Define

$$\|a\|_{1,J} := \sum_{k=1}^{J} |a_k|, \qquad w_j := \frac{J|a_j|}{\|a\|_{1,J}}, \qquad s_j^0 := \frac{\bar{B}}{J} w_j = \bar{B} \frac{|a_j|}{\|a\|_{1,J}}.$$

Corollary 3 (Default under common expected difficulty)**.** *Suppose* (F.8) *holds and*

$$c_\lambda \le w_j, \qquad j = 1, \ldots, J, \qquad \frac{\bar{B}}{J m_{\min}} \max_{1 \le j \le J} w_j \le 1. \tag{F.9}$$

Then $s^0 = (s_1^0, \ldots, s_J^0)$ *uniquely minimizes the working expected first-order criterion*

$$\sum_{j=1}^{J} a_j^2 \mathbb{E}_0\{\phi_j(s_j) \mid \mathcal{I}_J\} \tag{F.10}$$

over the feasible set in (F.5)*. A coverage-first solution is*

$$m_j^0 = m_{\min}, \qquad \pi_j^0 = \frac{s_j^0}{m_{\min}}, \qquad j = 1, \ldots, J. \tag{F.11}$$

Proof. For every $s > 0$,

$$\phi_j(s) \ge \frac{d_j^{\text{diff}}}{s} - r_j^2,$$

with equality whenever $s \le m_{\min}$. Taking the working expectation conditional on $\mathcal{I}_J$ and using (F.8) gives

$$\mathbb{E}_0\{\phi_j(s) \mid \mathcal{I}_J\} \ge \frac{d_J^0}{s} - \mathbb{E}_0(r_j^2 \mid \mathcal{I}_J).$$

For any feasible expenditures, Cauchy–Schwarz implies

$$\sum_{j=1}^{J} \frac{a_j^2}{s_j} \ge \frac{\left(\sum_{j=1}^{J} |a_j|\right)^2}{\sum_{j=1}^{J} s_j} = \frac{\|a\|_{1,J}^2}{\bar{B}},$$

with equality only when $s_j = \bar{B}|a_j|/\|a\|_{1,J} = s_j^0$ for every j. The first part of (F.9) gives $\underline{s}_J \le s_j^0$, while the second gives $s_j^0 \le m_{\min} \le \bar{m} = \overline{s}_J$. Thus s^0 is feasible and attains equality in both preceding inequalities. It is therefore the unique minimizer of (F.10). Equation (F.3) gives (F.11). □