

# Regression Discontinuity Designs

Classic Theories and Recent Development

Ye Wang

# Outline

- ▶ We start from considering the ideal experiment behind sharp RDD.

# Outline

- ▶ We start from considering the ideal experiment behind sharp RDD.
- ▶ We then proceed to discuss the asymptotic performance of the classic RDD estimators.

# Outline

- ▶ We start from considering the ideal experiment behind sharp RDD.
- ▶ We then proceed to discuss the asymptotic performance of the classic RDD estimators.
- ▶ Next, we discuss a special case in RDD where the running variable is discrete.

# Outline

- ▶ We start from considering the ideal experiment behind sharp RDD.
- ▶ We then proceed to discuss the asymptotic performance of the classic RDD estimators.
- ▶ Next, we discuss a special case in RDD where the running variable is discrete.
- ▶ We finally introduce two modern perspectives to understand RDD: local randomization and noise-induced randomization.

# Roadmap

- ▶ Thistlethwaite and Campbell (1960) first develop this idea in psychology.

# Roadmap

- ▶ Thistlethwaite and Campbell (1960) first develop this idea in psychology.
- ▶ Hahn, Todd, and Van der Klaauw (2001) formally discuss the identification assumptions in RDD.

# Roadmap

- ▶ Thistlethwaite and Campbell (1960) first develop this idea in psychology.
- ▶ Hahn, Todd, and Van der Klaauw (2001) formally discuss the identification assumptions in RDD.
- ▶ Porter (2003) proposes the local regression estimator based on the idea in Fan and Gijbels (1996).



# Roadmap

- ▶ Thistlethwaite and Campbell (1960) first develop this idea in psychology.
- ▶ Hahn, Todd, and Van der Klaauw (2001) formally discuss the identification assumptions in RDD.
- ▶ Porter (2003) proposes the local regression estimator based on the idea in Fan and Gijbels (1996).
- ▶ G. Imbens and Kalyanaraman (2012) introduce the first data-driven bandwidth selector.

# Roadmap

- ▶ Thistlethwaite and Campbell (1960) first develop this idea in psychology.
- ▶ Hahn, Todd, and Van der Klaauw (2001) formally discuss the identification assumptions in RDD.
- ▶ Porter (2003) proposes the local regression estimator based on the idea in Fan and Gijbels (1996).
- ▶ G. Imbens and Kalyanaraman (2012) introduce the first data-driven bandwidth selector.
- ▶ Calonico, Cattaneo, and Titiunik (2014) address the insufficiency in G. Imbens and Kalyanaraman (2012) and derive the asymptotic distributions for a group of RDD estimators.

# Roadmap

- ▶ Thistlethwaite and Campbell (1960) first develop this idea in psychology.
- ▶ Hahn, Todd, and Van der Klaauw (2001) formally discuss the identification assumptions in RDD.
- ▶ Porter (2003) proposes the local regression estimator based on the idea in Fan and Gijbels (1996).
- ▶ G. Imbens and Kalyanaraman (2012) introduce the first data-driven bandwidth selector.
- ▶ Calonico, Cattaneo, and Titiunik (2014) address the insufficiency in G. Imbens and Kalyanaraman (2012) and derive the asymptotic distributions for a group of RDD estimators.
- ▶ Calonico, Cattaneo, and Farrell (2020) suggest a bandwidth selector that is optimal for inference.

# Roadmap

- ▶ Thistlethwaite and Campbell (1960) first develop this idea in psychology.
- ▶ Hahn, Todd, and Van der Klaauw (2001) formally discuss the identification assumptions in RDD.
- ▶ Porter (2003) proposes the local regression estimator based on the idea in Fan and Gijbels (1996).
- ▶ G. Imbens and Kalyanaraman (2012) introduce the first data-driven bandwidth selector.
- ▶ Calonico, Cattaneo, and Titiunik (2014) address the insufficiency in G. Imbens and Kalyanaraman (2012) and derive the asymptotic distributions for a group of RDD estimators.
- ▶ Calonico, Cattaneo, and Farrell (2020) suggest a bandwidth selector that is optimal for inference.
- ▶ McCrary (2008) develops a test for self-selection in RDD.

# Roadmap

- ▶ Thistlethwaite and Campbell (1960) first develop this idea in psychology.
- ▶ Hahn, Todd, and Van der Klaauw (2001) formally discuss the identification assumptions in RDD.
- ▶ Porter (2003) proposes the local regression estimator based on the idea in Fan and Gijbels (1996).
- ▶ G. Imbens and Kalyanaraman (2012) introduce the first data-driven bandwidth selector.
- ▶ Calonico, Cattaneo, and Titiunik (2014) address the insufficiency in G. Imbens and Kalyanaraman (2012) and derive the asymptotic distributions for a group of RDD estimators.
- ▶ Calonico, Cattaneo, and Farrell (2020) suggest a bandwidth selector that is optimal for inference.
- ▶ McCrary (2008) develops a test for self-selection in RDD.
- ▶ Lee (2008) first applies RDD to analyze congressional elections in the US.

# Roadmap

- ▶ Theoretical results in Calonico, Cattaneo, and Titiunik (2014) can be directly applied to analyze fuzzy RDD and kink

# Roadmap

- ▶ Theoretical results in Calonico, Cattaneo, and Titiunik (2014) can be directly applied to analyze fuzzy RDD and kink
- ▶ Card et al. (2015) first introduce kink and fuzzy kink designs into econometrics.

# Roadmap

- ▶ Theoretical results in Calonico, Cattaneo, and Titiunik (2014) can be directly applied to analyze fuzzy RDD and kink
- ▶ Card et al. (2015) first introduce kink and fuzzy kink designs into econometrics.
- ▶ Cattaneo et al. (2016) and Cattaneo et al. (2020) discuss RDD with multiple cutoffs and their extrapolation.



# Roadmap

- ▶ Theoretical results in Calonico, Cattaneo, and Titiunik (2014) can be directly applied to analyze fuzzy RDD and kink
- ▶ Card et al. (2015) first introduce kink and fuzzy kink designs into econometrics.
- ▶ Cattaneo et al. (2016) and Cattaneo et al. (2020) discuss RDD with multiple cutoffs and their extrapolation.
- ▶ Kolesár and Rothe (2018) solve the problem of inference in RDD with a discrete running variable.

# Roadmap

- ▶ Theoretical results in Calonico, Cattaneo, and Titiunik (2014) can be directly applied to analyze fuzzy RDD and kink
- ▶ Card et al. (2015) first introduce kink and fuzzy kink designs into econometrics.
- ▶ Cattaneo et al. (2016) and Cattaneo et al. (2020) discuss RDD with multiple cutoffs and their extrapolation.
- ▶ Kolesár and Rothe (2018) solve the problem of inference in RDD with a discrete running variable.
- ▶ Cattaneo, Frandsen, and Titiunik (2015) suggest the perspective of local randomization for RDD.

# Roadmap

- ▶ Theoretical results in Calonico, Cattaneo, and Titiunik (2014) can be directly applied to analyze fuzzy RDD and kink
- ▶ Card et al. (2015) first introduce kink and fuzzy kink designs into econometrics.
- ▶ Cattaneo et al. (2016) and Cattaneo et al. (2020) discuss RDD with multiple cutoffs and their extrapolation.
- ▶ Kolesár and Rothe (2018) solve the problem of inference in RDD with a discrete running variable.
- ▶ Cattaneo, Frandsen, and Titiunik (2015) suggest the perspective of local randomization for RDD.
- ▶ G. Imbens and Wager (2019) and Eckles et al. (2020) develop the perspective of noise-induced randomization.

## Set up

- ▶ We possess a dataset with  $N$  units.

## Set up

- ▶ We possess a dataset with  $N$  units.
- ▶ We observe  $(Y_i, D_i, Z_i)$  for each unit  $i$ ,  $i \in \{1, 2, \dots, N\}$ .
- ▶  $Z_i$  is called the running variable with density  $f(z)$ , which decides the value of the binary treatment  $D_i$ .

## Set up

- ▶ We possess a dataset with  $N$  units.
- ▶ We observe  $(Y_i, D_i, Z_i)$  for each unit  $i$ ,  $i \in \{1, 2, \dots, N\}$ .
- ▶  $Z_i$  is called the running variable with density  $f(z)$ , which decides the value of the binary treatment  $D_i$ .
- ▶ There exists a cutoff  $c$  (often set to be 0) such that  $D_i = \mathbf{1}\{Z_i \geq c\}$ .

## Set up

- ▶ We possess a dataset with  $N$  units.
- ▶ We observe  $(Y_i, D_i, Z_i)$  for each unit  $i$ ,  $i \in \{1, 2, \dots, N\}$ .
- ▶  $Z_i$  is called the running variable with density  $f(z)$ , which decides the value of the binary treatment  $D_i$ .
- ▶ There exists a cutoff  $c$  (often set to be 0) such that  $D_i = \mathbf{1}\{Z_i \geq c\}$ .
- ▶ The causal parameter of interest,  $\tau_{SRD}$ , is defined as

$$\mathbb{E}[Y_i(1) - Y_i(0)|Z_i = c].$$

## Set up

- ▶ We possess a dataset with  $N$  units.
- ▶ We observe  $(Y_i, D_i, Z_i)$  for each unit  $i$ ,  $i \in \{1, 2, \dots, N\}$ .
- ▶  $Z_i$  is called the running variable with density  $f(z)$ , which decides the value of the binary treatment  $D_i$ .
- ▶ There exists a cutoff  $c$  (often set to be 0) such that  $D_i = \mathbf{1}\{Z_i \geq c\}$ .
- ▶ The causal parameter of interest,  $\tau_{SRD}$ , is defined as

$$\mathbb{E}[Y_i(1) - Y_i(0)|Z_i = c].$$

- ▶ Notice that the causal parameter is a local one by definition.



# Identification

- Identification in the classic RDD relies the assumption of continuity (Hahn, Todd, and Van der Klaauw 2001).

# Identification

- ▶ Identification in the classic RDD relies the assumption of continuity (Hahn, Todd, and Van der Klaauw 2001).
- ▶ Define  $\mu(z) = \mathbb{E}[Y_i|Z_i = z]$ ,  $\mu_+ = \lim_{z \rightarrow c^+} \mu(z)$ , and  $\mu_- = \lim_{z \rightarrow c^-} \mu(z)$ .

# Identification

- ▶ Identification in the classic RDD relies the assumption of continuity (Hahn, Todd, and Van der Klaauw 2001).
- ▶ Define  $\mu(z) = \mathbb{E}[Y_i|Z_i = z]$ ,  $\mu_+ = \lim_{z \rightarrow c^+} \mu(z)$ , and  $\mu_- = \lim_{z \rightarrow c^-} \mu(z)$ .
- ▶ Obviously, if  $\mu_+ = \mathbb{E}[Y_i(1)|Z_i = c]$  and  $\mu_- = \mathbb{E}[Y_i(0)|Z_i = c]$ , then

$$\tau_{SRD} = \mu_+ - \mu_-.$$

# Identification

- ▶ Identification in the classic RDD relies the assumption of continuity (Hahn, Todd, and Van der Klaauw 2001).
- ▶ Define  $\mu(z) = \mathbb{E}[Y_i|Z_i = z]$ ,  $\mu_+ = \lim_{z \rightarrow c^+} \mu(z)$ , and  $\mu_- = \lim_{z \rightarrow c^-} \mu(z)$ .
- ▶ Obviously, if  $\mu_+ = \mathbb{E}[Y_i(1)|Z_i = c]$  and  $\mu_- = \mathbb{E}[Y_i(0)|Z_i = c]$ , then

$$\tau_{SRD} = \mu_+ - \mu_-.$$

- ▶ All we need to do is to estimate  $\mu_+$  and  $\mu_-$ .

## Ideal experiment behind RDD

- ▶ Intuitively, the assumption of continuity requires that all the confounders (observable or not) vary smoothly around the cutoff  $c$ .

# Ideal experiment behind RDD

- ▶ Intuitively, the assumption of continuity requires that all the confounders (observable or not) vary smoothly around the cutoff  $c$ .
- ▶ We can treat RDD as a random assignment in the block defined by  $Z_i = c$ .

## Ideal experiment behind RDD

- ▶ Intuitively, the assumption of continuity requires that all the confounders (observable or not) vary smoothly around the cutoff  $c$ .
- ▶ We can treat RDD as a random assignment in the block defined by  $Z_i = c$ .
- ▶ Suppose there are 1,000 congressional elections, in all of which both candidates win 50% of all votes.

## Ideal experiment behind RDD

- ▶ Intuitively, the assumption of continuity requires that all the confounders (observable or not) vary smoothly around the cutoff  $c$ .
- ▶ We can treat RDD as a random assignment in the block defined by  $Z_i = c$ .
- ▶ Suppose there are 1,000 congressional elections, in all of which both candidates win 50% of all votes.
- ▶ We then randomly select a winner for each election with a coin flip, which implies that  $D_i \perp \{Y_i(1), Y_i(0)\} | Z_i = c$ .



## Ideal experiment behind RDD

- ▶ Intuitively, the assumption of continuity requires that all the confounders (observable or not) vary smoothly around the cutoff  $c$ .
- ▶ We can treat RDD as a random assignment in the block defined by  $Z_i = c$ .
- ▶ Suppose there are 1,000 congressional elections, in all of which both candidates win 50% of all votes.
- ▶ We then randomly select a winner for each election with a coin flip, which implies that  $D_i \perp \{Y_i(1), Y_i(0)\} | Z_i = c$ .
- ▶ The difference in the outcome reflects the impact of the winner's attributes (e.g party affiliation).

## Ideal experiment behind RDD

- ▶ Unfortunately, we do not have that many elections with a 50-50 split in practice.

## Ideal experiment behind RDD

- ▶ Unfortunately, we do not have that many elections with a 50-50 split in practice.
- ▶ All we can do is to approximate the two limits using data at hand.

## Ideal experiment behind RDD

- ▶ Unfortunately, we do not have that many elections with a 50-50 split in practice.
- ▶ All we can do is to approximate the two limits using data at hand.
- ▶ The RDD estimate is thus biased by definition.

## Ideal experiment behind RDD

- ▶ Unfortunately, we do not have that many elections with a 50-50 split in practice.
- ▶ All we can do is to approximate the two limits using data at hand.
- ▶ The RDD estimate is thus biased by definition.
- ▶ Yet the bias diminishes to zero as we have a larger sample, which usually leads to a smaller bandwidth.

## Ideal experiment behind RDD

- ▶ Unfortunately, we do not have that many elections with a 50-50 split in practice.
- ▶ All we can do is to approximate the two limits using data at hand.
- ▶ The RDD estimate is thus biased by definition.
- ▶ Yet the bias diminishes to zero as we have a larger sample, which usually leads to a smaller bandwidth.
- ▶ In other words, we approach the ideal experiment as sample size increases.

## Ideal experiment behind RDD

- ▶ Unfortunately, we do not have that many elections with a 50-50 split in practice.
- ▶ All we can do is to approximate the two limits using data at hand.
- ▶ The RDD estimate is thus biased by definition.
- ▶ Yet the bias diminishes to zero as we have a larger sample, which usually leads to a smaller bandwidth.
- ▶ In other words, we approach the ideal experiment as sample size increases.
- ▶ Remember that we need extra structural assumptions (continuity) for the trick to work.

# Estimation

- ▶ To estimate  $\mu_+$  and  $\mu_-$ , the most common choice is local regression.



# Estimation

- ▶ To estimate  $\mu_+$  and  $\mu_-$ , the most common choice is local regression.
- ▶ Suppose we have selected a bandwidth  $h$ , we just need to fit kernelized regression within the windows  $[c - h, c]$  and  $[c, c + h]$ :

$$\begin{aligned} & (\hat{\mu}_+, \hat{\beta}_+) \\ &= \arg \min_{\alpha, \beta} \sum_{i=1}^N \mathbf{1}\{Z_i \geq c\} (Y_i - \mu - \beta(Z_i - c))^2 \mathbf{K}\left(\frac{Z_i - c}{h_N}\right) \end{aligned}$$

- ▶  $(\hat{\mu}_-, \hat{\beta}_-)$  are similarly estimated.

# Estimation

- ▶ To estimate  $\mu_+$  and  $\mu_-$ , the most common choice is local regression.
- ▶ Suppose we have selected a bandwidth  $h$ , we just need to fit kernelized regression within the windows  $[c - h, c]$  and  $[c, c + h]$ :

$$\begin{aligned} & (\hat{\mu}_+, \hat{\beta}_+) \\ &= \arg \min_{\alpha, \beta} \sum_{i=1}^N \mathbf{1}\{Z_i \geq c\} (Y_i - \mu - \beta(Z_i - c))^2 \mathbf{K}\left(\frac{Z_i - c}{h_N}\right) \end{aligned}$$

- ▶  $(\hat{\mu}_-, \hat{\beta}_-)$  are similarly estimated.
- ▶ Then,

$$\hat{\tau}_{SRD} = \hat{\mu}_+ - \hat{\mu}_-.$$

## Estimation


- ▶ The two regressions are local since we apply the kernel weights.

## Estimation

- ▶ The two regressions are local since we apply the kernel weights.
- ▶ Intuitively, observations that are closer to the cutoff should receive a larger weight.

## Estimation

- ▶ The two regressions are local since we apply the kernel weights.
- ▶ Intuitively, observations that are closer to the cutoff should receive a larger weight.
- ▶ There are multiple choices for the kernel.



## Estimation in practice

- ▶ In practice, some scholars select the bandwidth using an existent algorithm and then manually fit two regressions within the selected window.

## Estimation in practice

- ▶ In practice, some scholars select the bandwidth using an existent algorithm and then manually fit two regressions within the selected window.
- ▶ What is the problem of this approach?

## Estimation in practice

- ▶ In practice, some scholars select the bandwidth using an existent algorithm and then manually fit two regressions within the selected window.
- ▶ What is the problem of this approach?
- ▶ It is inconsistent with the bandwidth selector and often leads to larger biases.



## Estimation in practice

- ▶ In practice, some scholars select the bandwidth using an existent algorithm and then manually fit two regressions within the selected window.
- ▶ What is the problem of this approach?
- ▶ It is inconsistent with the bandwidth selector and often leads to larger biases.
- ▶ One may add higher order terms of  $(Z_i - c)$  into the regressions.

## Estimation in practice

- ▶ In practice, some scholars select the bandwidth using an existent algorithm and then manually fit two regressions within the selected window.
- ▶ What is the problem of this approach?
- ▶ It is inconsistent with the bandwidth selector and often leads to larger biases.
- ▶ One may add higher order terms of  $(Z_i - c)$  into the regressions.
- ▶ But for sharp RDD, linear regression has the optimal rate of convergence due to its nice performance on the boundary (Porter 2003).

## Estimation in practice

- ▶ Never fit global polynomials to estimate the two intercepts (Gelman and Imbens 2019).

## Estimation in practice

- ▶ Never fit global polynomials to estimate the two intercepts (Gelman and Imbens 2019).
- ▶ We are interested in local quantities rather than global fitness.

## Estimation in practice

- ▶ Never fit global polynomials to estimate the two intercepts (Gelman and Imbens 2019).
- ▶ We are interested in local quantities rather than global fitness.
- ▶ Estimates of the intercepts may be driven by points that are far away from the cutoff if you use global polynomials.

## Estimation in practice

- ▶ Never fit global polynomials to estimate the two intercepts (Gelman and Imbens 2019).
- ▶ We are interested in local quantities rather than global fitness.
- ▶ Estimates of the intercepts may be driven by points that are far away from the cutoff if you use global polynomials.
- ▶ It is always critical to draw plots in RDD.

## Estimation in practice

- ▶ Never fit global polynomials to estimate the two intercepts (Gelman and Imbens 2019).
- ▶ We are interested in local quantities rather than global fitness.
- ▶ Estimates of the intercepts may be driven by points that are far away from the cutoff if you use global polynomials.
- ▶ It is always critical to draw plots in RDD.
- ▶ The result would not be convincing if we cannot see the jump of the outcome variable from the plot.

## Test the assumption of continuity

- ▶ As the assumption of continuity is crucial for identification in RDD, we should test its validity in practice.



## Test the assumption of continuity

- ▶ As the assumption of continuity is crucial for identification in RDD, we should test its validity in practice.
- ▶ A popular approach is the placebo test: variables that are not affected by the treatment should change smoothly over the cutoff.
- ▶ For example, we can run the estimator for a covariate  $X$  to see whether the result is significant.

## Test the assumption of continuity


- ▶ As the assumption of continuity is crucial for identification in RDD, we should test its validity in practice.
- ▶ A popular approach is the placebo test: variables that are not affected by the treatment should change smoothly over the cutoff.
- ▶ For example, we can run the estimator for a covariate  $X$  to see whether the result is significant.
- ▶ See Meyersson (2014) for examples

## Test the assumption of continuity

- ▶ The assumption is violated if units in the sample self-select into one side of the cutoff.

# Test the assumption of continuity

- ▶ The assumption is violated if units in the sample self-select into one side of the cutoff.
- ▶ For example, students may cheat to meet the requirement of a scholarship.
- ▶ As a result, the density function  $f(z)$  will not change smoothly across the cutoff.



## Test the assumption of continuity

- ▶ McCrary (2008) develop the first formal test of the discontinuity in density.

## Test the assumption of continuity

- ▶ McCrary (2008) develop the first formal test of the discontinuity in density.
- ▶ Essentially, we are testing whether

$$\theta = \ln \lim_{z \rightarrow 0^+} f(z) - \ln \lim_{z \rightarrow 0^-} f(z)$$

has a significant deviation from 0.

## Test the assumption of continuity

- ▶ McCrary (2008) develop the first formal test of the discontinuity in density.
- ▶ Essentially, we are testing whether

$$\theta = \ln \lim_{z \rightarrow 0^+} f(z) - \ln \lim_{z \rightarrow 0^-} f(z)$$

has a significant deviation from 0.

- ▶ Similarly, we need to estimate the two boundary points of  $\ln f(z)$  using local regression (notice that  $Y_i$  is not needed).

## Test the assumption of continuity

- ▶ McCrary (2008) develop the first formal test of the discontinuity in density.
- ▶ Essentially, we are testing whether

$$\theta = \ln \lim_{z \rightarrow 0^+} f(z) - \ln \lim_{z \rightarrow 0^-} f(z)$$

has a significant deviation from 0.

- ▶ Similarly, we need to estimate the two boundary points of  $\ln f(z)$  using local regression (notice that  $Y_i$  is not needed).
- ▶ Cattaneo, Jansson, and Ma (2020) provide an augmented algorithm for non-parametric density estimation.



## An example

- ▶ We will work with the Meyersson (2014) paper: “Islamic Rule and the Empowerment of the Poor and the Pious”
- ▶ The paper shows a (local) result: the victory of Islamic parties in Turkey resulted in better outcomes for women.
- ▶ Running variable: the difference in vote share between the largest Islamic party and the largest secular party (not two party)
- ▶ Outcome that we'll look at: high school education

## Set up the data


```
##      Min. 1st Qu.  Median      Mean 3rd Qu.      Max.      NA's  
## -1.0000 -0.4600 -0.3102 -0.2786 -0.1061  0.9905      544
```

# Estimation

```
## Call: rdrobust
##
## Number of Obs.          2630
## BW type                mserd
## Kernel                  Triangular
## VCE method              NN
##
## Number of Obs.          2315          315
## Eff. Number of Obs.      529           266
## Order est. (p)           1             1
## Order bias (q)           2             2
## BW est. (h)              0.172         0.172
## BW bias (b)              0.286         0.286
## rho (h/b)               0.603         0.603
## Unique Obs.             2313          315
##
## =====
##      Method      Coef. Std. Err.      z    P>|z|      [ 95% C.I. ]
## =====
##   Conventional    0.030     0.014    2.116    0.034    [0.002 , 0.058]
##      Robust       -         -    1.776    0.076    [-0.003 , 0.063]
## =====
```

# Plot it

```
rdplot(d$hischshr1520f, d$iwm94, p = 4)
```




## Placebo tests

- Do placebo tests on other covariates and other outcomes.

```
## $coef
##                               Coeff
## Conventional    0.004097863
## Bias-Corrected 0.008070629
## Robust          0.008070629
##
## $se
##                               Std. Err.
## Conventional    0.01227104
## Bias-Corrected 0.01227104
## Robust          0.01408919
```

# Placebo plot



## More Placebos

```
## $coef
##                               Coeff
## Conventional    0.01285314
## Bias-Corrected 0.01263466
## Robust          0.01263466
##
```

```
## $se
##                               Std. Err.
## Conventional    0.01403235
## Bias-Corrected 0.01403235
## Robust          0.01691543
```

```
## $coef
##                               Coeff
## Conventional    0.0039695120
## Bias-Corrected 0.0008440962
## Robust          0.0008440962
##
```

```
## $se
##                               Std. Err.
## Conventional    0.01537558
## Bias-Corrected 0.01537558
```

## Sorting


- Density tests are also a good way to examine the possibility of sorting.

```
##
## Manipulation testing using local polynomial density estimation
##
## Number of obs =          2660
## Model =              unrestricted
## Kernel =             triangular
## BW method =          estimated
## VCE method =         jackknife
##
## c = 0                Left of c                Right of c
## Number of obs        2332                    328
## Eff. Number of obs   769                      314
## Order est. (p)        2                       2
## Order bias (q)        3                       3
## BW est. (h)           0.25                    0.282
##
```



# Density Plot

```
rdplotdensity(rd_density, d$iwm94)
```



```
## Call: rdrobust
##
## Number of Obs.          2630
## BW type                mserd
## Kernel                  Triangular
## VCE method              NN
##
## Number of Obs.          2315      315
## Eff. Number of Obs.     428       238
## Order est. (p)          1         1
## Order bias (q)          2         2
## BW est. (h)             0.139     0.139
## BW bias (b)             0.286     0.286
## rho (h/b)              0.486     0.486
## Unique Obs.            2313      315
##
## =====
##      Method      Coef. Std. Err.      z    P>|z|      [ 95% C.I. ]
## =====
##   Conventional    0.020    0.242    0.082    0.934    [-0.455 , 0.495]
##      Robust       -        -    0.205    0.838    [-0.689 , 0.849]
## =====
```

## Bias of RDD estimation

- ▶ Let's introduce some notations for deriving the bias of RDD estimation.
- ▶ Denote  $(Y_1, Y_2, \dots, Y_N)$  as  $\mathbf{Y}$ ,  $(Z_1, Z_2, \dots, Z_N)$  as  $\mathbf{Z}$ , and

$$\mathbf{R} = (\boldsymbol{\iota}, \mathbf{Z})$$

$$\mathbf{W}_+ = \{\mathbf{1}\{Z_i \geq c\} K\left(\frac{Z_i - c}{h_N}\right)\}_{N \times N}$$

$$\mathbf{M} = (\mu(Z_1), \mu(Z_2), \dots, \mu(Z_N))'$$

where  $\boldsymbol{\iota}$  is a vector with  $N$  1s and  $\mathbf{W}$  is a diagonal matrix of weights.

- ▶ We use  $\mu_+^{(k)}$  to denote the  $k$ -th order derivative of  $\mu_+$  (similar for  $\mu_-$ ) and  $\sigma^2(z)$  to denote  $\text{Var}[Y_i | Z_i = z]$ .

## Bias of RDD estimation

- ▶ Notice that  $\mathbf{Y} = \mathbf{M} + \mathbf{e}$ .

## Bias of RDD estimation

- ▶ Notice that  $\mathbf{Y} = \mathbf{M} + \mathbf{e}$ .
- ▶ The estimate  $\hat{\mu}_+$  equals to the first row of

$$\begin{aligned} & (\mathbf{R}'\mathbf{W}_+\mathbf{R})^{-1}(\mathbf{R}'\mathbf{W}_+\mathbf{Y}) \\ &= (\mathbf{R}'\mathbf{W}_+\mathbf{R})^{-1}(\mathbf{R}'\mathbf{W}_+\mathbf{M}) + (\mathbf{R}'\mathbf{W}_+\mathbf{R})^{-1}(\mathbf{R}'\mathbf{W}_+\mathbf{e}) \end{aligned}$$

## Bias of RDD estimation

- ▶ Notice that  $\mathbf{Y} = \mathbf{M} + \mathbf{e}$ .
- ▶ The estimate  $\hat{\mu}_+$  equals to the first row of

$$\begin{aligned} & (\mathbf{R}'\mathbf{W}_+\mathbf{R})^{-1}(\mathbf{R}'\mathbf{W}_+\mathbf{Y}) \\ &= (\mathbf{R}'\mathbf{W}_+\mathbf{R})^{-1}(\mathbf{R}'\mathbf{W}_+\mathbf{M}) + (\mathbf{R}'\mathbf{W}_+\mathbf{R})^{-1}(\mathbf{R}'\mathbf{W}_+\mathbf{e}) \end{aligned}$$

- ▶ Expectation of the second term is zero and we have the Taylor expansion for  $\mu(Z_i)$ :

$$\mu(Z_i) = \mu_+ + \mu_+^{(1)}(0)Z_i + \frac{\mu_+^{(2)}(0)}{2}Z_i^2 + \nu_i$$

## Bias of RDD estimation

- ▶ Notice that  $\mathbf{Y} = \mathbf{M} + \mathbf{e}$ .
- ▶ The estimate  $\hat{\mu}_+$  equals to the first row of

$$\begin{aligned} & (\mathbf{R}'\mathbf{W}_+\mathbf{R})^{-1}(\mathbf{R}'\mathbf{W}_+\mathbf{Y}) \\ &= (\mathbf{R}'\mathbf{W}_+\mathbf{R})^{-1}(\mathbf{R}'\mathbf{W}_+\mathbf{M}) + (\mathbf{R}'\mathbf{W}_+\mathbf{R})^{-1}(\mathbf{R}'\mathbf{W}_+\mathbf{e}) \end{aligned}$$

- ▶ Expectation of the second term is zero and we have the Taylor expansion for  $\mu(Z_i)$ :

$$\mu(Z_i) = \mu_+ + \mu_+^{(1)}(0)Z_i + \frac{\mu_+^{(2)}(0)}{2}Z_i^2 + \nu_i$$

- ▶ Hence,

$$\mathbf{M} = \mathbf{R} \begin{pmatrix} \mu_+ \\ \mu_+^{(1)}(0) \end{pmatrix} + \mathbf{S}_2 \frac{\mu_+^{(2)}(0)}{2} + \boldsymbol{\nu}$$

where  $\mathbf{S}_2 = (Z_1^2, Z_2^2, \dots, Z_N^2)$  and  $\boldsymbol{\nu} = (\nu_1, \nu_2, \dots, \nu_N)$ .

## Bias of RDD estimation

- Now, the estimation bias of  $\hat{\mu}_+$ ,  $\mathbb{E}[\hat{\mu}_+] - \mu_+$ , is the first row of

$$\mu_+^{(1)}(0) + (\mathbf{R}'\mathbf{W}_+\mathbf{R})^{-1} \left( \mathbf{R}'\mathbf{W}_+\mathbf{S}_2 \frac{\mu_+^{(2)}(0)}{2} \right) + (\mathbf{R}'\mathbf{W}_+\mathbf{R})^{-1} (\mathbf{R}'\mathbf{W}_+\nu)$$



## Bias of RDD estimation

- ▶ Now, the estimation bias of  $\hat{\mu}_+$ ,  $\mathbb{E}[\hat{\mu}_+] - \mu_+$ , is the first row of

$$\mu_+^{(1)}(0) + (\mathbf{R}'\mathbf{W}_+\mathbf{R})^{-1} \left( \mathbf{R}'\mathbf{W}_+\mathbf{S}_2 \frac{\mu_+^{(2)}(0)}{2} \right) + (\mathbf{R}'\mathbf{W}_+\mathbf{R})^{-1} (\mathbf{R}'\mathbf{W}_+\nu)$$

- ▶ The convergence rates of these two terms rely on the properties of the kernel.

## Bias of RDD estimation

- Now, the estimation bias of  $\hat{\mu}_+$ ,  $\mathbb{E}[\hat{\mu}_+] - \mu_+$ , is the first row of

$$\mu_+^{(1)}(0) + (\mathbf{R}'\mathbf{W}_+\mathbf{R})^{-1} \left( \mathbf{R}'\mathbf{W}_+\mathbf{S}_2 \frac{\mu_+^{(2)}(0)}{2} \right) + (\mathbf{R}'\mathbf{W}_+\mathbf{R})^{-1} (\mathbf{R}'\mathbf{W}_+\nu)$$

- The convergence rates of these two terms rely on the properties of the kernel.
- Via some cumbersome calculation, we can see that

$$\mathbb{E}[\hat{\mu}_+] - \mu_+ = C_1 \mu_+^{(2)}(0) h^2 + o_p(h^2)$$

## Bias of RDD estimation

- We can similarly derive the variance of  $\hat{\mu}_+$  using the properties of regression:

$$\mathbb{V}[\hat{\mu}_+] = \frac{C_2}{Nh} \frac{\sigma_+^2(0)}{f_+(0)} + o_p\left(\frac{1}{Nh}\right)$$

## Bias of RDD estimation

- ▶ We can similarly derive the variance of  $\hat{\mu}_+$  using the properties of regression:

$$\mathbb{V}[\hat{\mu}_+] = \frac{C_2}{Nh} \frac{\sigma_+^2(0)}{f_+(0)} + o_p\left(\frac{1}{Nh}\right)$$

- ▶ Obviously, the bias and the variance of  $\hat{\mu}_-$  have similar forms.

## Bias of RDD estimation

- ▶ We can similarly derive the variance of  $\hat{\mu}_+$  using the properties of regression:

$$\mathbb{V}[\hat{\mu}_+] = \frac{C_2}{Nh} \frac{\sigma_+^2(0)}{f_+(0)} + o_p\left(\frac{1}{Nh}\right)$$

- ▶ Obviously, the bias and the variance of  $\hat{\mu}_-$  have similar forms.
- ▶ More generally, we can estimate the  $k$ -th order derivative of  $\mu_+$  and  $\mu_-$  with a  $p$ -th order local regression.
- ▶ The bias will be of order  $p + 1$ .

# Bandwidth selection for optimizing MSE

- ▶ Before G. Imbens and Kalyanaraman (2012), the practice is to minimize the regression's MSE on the entire support using cross-validation.

# Bandwidth selection for optimizing MSE

- ▶ Before G. Imbens and Kalyanaraman (2012), the practice is to minimize the regression's MSE on the entire support using cross-validation.
- ▶ G. Imbens and Kalyanaraman (2012) argue that we should select a bandwidth to minimize the MSE of estimation:

$$\begin{aligned}MSE(h_N) &= \mathbb{E} [\hat{\tau}_{SRD} - \tau_{SRD} | \mathbf{Z}]^2 \\&= (\mathbb{E} [\hat{\tau}_{SRD} | \mathbf{Z}] - \tau_{SRD})^2 + \text{Var} [\hat{\tau}_{SRD} | \mathbf{Z}] \\&= \text{Bias}^2 + \text{Variance}.\end{aligned}$$

# Bandwidth selection for optimizing MSE

- ▶ G. Imbens and Kalyanaraman (2012) show that in practice we can minimize the asymptotic MSE:

$$AMSE(h_N) = C_1 h_N^4 \left( \mu_+^{(2)}(0) - \mu_-^{(2)}(0) \right)^2 + \frac{C_2}{N h_N} \frac{\sigma^2(0)}{f(0)}$$



## Bandwidth selection for optimizing MSE

- ▶ G. Imbens and Kalyanaraman (2012) show that in practice we can minimize the asymptotic MSE:

$$AMSE(h_N) = C_1 h_N^4 \left( \mu_+^{(2)}(0) - \mu_-^{(2)}(0) \right)^2 + \frac{C_2}{N h_N} \frac{\sigma^2(0)}{f(0)}$$

- ▶ From the expression we can solve the optimal bandwidth:

$$h_N^* = C \left( \frac{\frac{\sigma^2(0)}{f(0)}}{\left( \mu_+^{(2)}(0) - \mu_-^{(2)}(0) \right)^2} \right)^{\frac{1}{5}} N^{-\frac{1}{5}}.$$

## Bandwidth selection for optimizing MSE

- ▶ G. Imbens and Kalyanaraman (2012) show that in practice we can minimize the asymptotic MSE:

$$AMSE(h_N) = C_1 h_N^4 \left( \mu_+^{(2)}(0) - \mu_-^{(2)}(0) \right)^2 + \frac{C_2}{N h_N} \frac{\sigma^2(0)}{f(0)}$$

- ▶ From the expression we can solve the optimal bandwidth:

$$h_N^* = C \left( \frac{\frac{\sigma^2(0)}{f(0)}}{\left( \mu_+^{(2)}(0) - \mu_-^{(2)}(0) \right)^2} \right)^{\frac{1}{5}} N^{-\frac{1}{5}}.$$

- ▶ In practice, we can estimate  $h_N^*$  with a plug-in estimator.

## Bias correction

- ▶ G. Imbens and Kalyanaraman (2012) do not prove the asymptotic normality of the RDD estimate.

## Bias correction

- ▶ G. Imbens and Kalyanaraman (2012) do not prove the asymptotic normality of the RDD estimate.
- ▶ Calonico, Cattaneo, and Titiunik (2014) study the asymptotic distribution of the studentized RDD estimate:

$$\frac{\hat{\tau}_{SRD} - \tau_{SRD}}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}}.$$

## Bias correction

- ▶ G. Imbens and Kalyanaraman (2012) do not prove the asymptotic normality of the RDD estimate.
- ▶ Calonico, Cattaneo, and Titiunik (2014) study the asymptotic distribution of the studentized RDD estimate:

$$\frac{\hat{\tau}_{SRD} - \tau_{SRD}}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}},$$

- ▶ They show that the bandwidth selected via the previous algorithm is too wide to guarantee the asymptotic normality of the studentized estimate.

## Bias correction

- ▶ G. Imbens and Kalyanaraman (2012) do not prove the asymptotic normality of the RDD estimate.
- ▶ Calonico, Cattaneo, and Titiunik (2014) study the asymptotic distribution of the studentized RDD estimate:

$$\frac{\hat{\tau}_{SRD} - \tau_{SRD}}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}},$$

- ▶ They show that the bandwidth selected via the previous algorithm is too wide to guarantee the asymptotic normality of the studentized estimate.
- ▶ We need  $h_N = o_p(N^{-\frac{1}{5}})$  while the algorithm leads to  $h_N = O_p(N^{-\frac{1}{5}})$ .

## Bias correction

- ▶ G. Imbens and Kalyanaraman (2012) do not prove the asymptotic normality of the RDD estimate.
- ▶ Calonico, Cattaneo, and Titiunik (2014) study the asymptotic distribution of the studentized RDD estimate:

$$\frac{\hat{\tau}_{SRD} - \tau_{SRD}}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}},$$

- ▶ They show that the bandwidth selected via the previous algorithm is too wide to guarantee the asymptotic normality of the studentized estimate.
- ▶ We need  $h_N = o_p(N^{-\frac{1}{5}})$  while the algorithm leads to  $h_N = O_p(N^{-\frac{1}{5}})$ .
- ▶ Consequently, the studentized estimate will be asymptotically biased.

## Bias correction

► Intuitively,

$$\frac{\hat{\tau}_{SRD} - \tau_{SRD}}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}} = \frac{\hat{\tau}_{SRD} - \mathbb{E}[\hat{\tau}_{SRD}]}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}} + \frac{\mathbb{E}[\hat{\tau}_{SRD}] - \tau_{SRD}}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}}.$$



## Bias correction

- Intuitively,

$$\frac{\hat{\tau}_{SRD} - \tau_{SRD}}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}} = \frac{\hat{\tau}_{SRD} - \mathbb{E}[\hat{\tau}_{SRD}]}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}} + \frac{\mathbb{E}[\hat{\tau}_{SRD}] - \tau_{SRD}}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}}.$$

- The first term is a weighted average of residuals and converges to  $N(0,1)$  by CLT.

## Bias correction

- Intuitively,

$$\frac{\hat{\tau}_{SRD} - \tau_{SRD}}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}} = \frac{\hat{\tau}_{SRD} - \mathbb{E}[\hat{\tau}_{SRD}]}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}} + \frac{\mathbb{E}[\hat{\tau}_{SRD}] - \tau_{SRD}}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}}.$$

- The first term is a weighted average of residuals and converges to  $N(0, 1)$  by CLT.
- We need to guarantee that the second term is  $o_p(1)$ .

## Bias correction

- Intuitively,

$$\frac{\hat{\tau}_{SRD} - \tau_{SRD}}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}} = \frac{\hat{\tau}_{SRD} - \mathbb{E}[\hat{\tau}_{SRD}]}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}} + \frac{\mathbb{E}[\hat{\tau}_{SRD}] - \tau_{SRD}}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}}.$$

- The first term is a weighted average of residuals and converges to  $N(0, 1)$  by CLT.
- We need to guarantee that the second term is  $o_p(1)$ .
- Remember that the numerator is  $O_p(h^2)$  and the denominator is  $O_p(\frac{1}{\sqrt{Nh}})$ , thus the total bias is  $O_p(\sqrt{Nh^5})$ .

## Bias correction

- ▶ Calonico, Cattaneo, and Titiunik (2014) provide a bias-correction estimator  $\hat{\tau}_{SRD}^{bc}$ .

## Bias correction

- ▶ Calonico, Cattaneo, and Titiunik (2014) provide a bias-correction estimator  $\hat{\tau}_{SRD}^{bc}$ .
- ▶ Intuitively, we use another local regression estimator with bandwidth  $b_N$  to estimate the first-order derivative of  $\mu_+$  and  $\mu_-$  and subtract them from  $\hat{\mu}_+$  and  $\hat{\mu}_-$ .

## Bias correction

- ▶ Calonico, Cattaneo, and Titiunik (2014) provide a bias-correction estimator  $\hat{\tau}_{SRD}^{bc}$ .
- ▶ Intuitively, we use another local regression estimator with bandwidth  $b_N$  to estimate the first-order derivative of  $\mu_+$  and  $\mu_-$  and subtract them from  $\hat{\mu}_+$  and  $\hat{\mu}_-$ .
- ▶ Of course, bias correction introduces extra uncertainty (from the extra local regression) into the estimate, hence the variance has to be adjusted accordingly.

## Bias correction

- ▶ Calonico, Cattaneo, and Titiunik (2014) provide a bias-correction estimator  $\hat{\tau}_{SRD}^{bc}$ .
- ▶ Intuitively, we use another local regression estimator with bandwidth  $b_N$  to estimate the first-order derivative of  $\mu_+$  and  $\mu_-$  and subtract them from  $\hat{\mu}_+$  and  $\hat{\mu}_-$ .
- ▶ Of course, bias correction introduces extra uncertainty (from the extra local regression) into the estimate, hence the variance has to be adjusted accordingly.
- ▶ They propose two variance estimators, one based on regression analysis and the other based on the idea of nearest neighborhood matching (Abadie and Imbens 2006).

## Bias correction

- Calonico, Cattaneo, and Titiunik (2014) prove that

$$\frac{\hat{\tau}_{SRD}^{bc} - \tau_{SRD}}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}^{bc}]}} \rightarrow N(0, 1)$$

as long as  $N \min\{b_N^5, h_N^5\} \max\{b_N^2, h_N^2\} \rightarrow 0$ .



## Bias correction

- Calonico, Cattaneo, and Titiunik (2014) prove that

$$\frac{\hat{\tau}_{SRD}^{bc} - \tau_{SRD}}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}^{bc}]}} \rightarrow N(0, 1)$$

as long as  $N \min\{b_N^5, h_N^5\} \max\{b_N^2, h_N^2\} \rightarrow 0$ .

- Now the numerator is  $O_p(h^3 + h^2 b^2)$  and the denominator is  $O_p(\frac{1}{\sqrt{Nh}} + \frac{1}{\sqrt{Nh^4 b^5}})$ .

## Bias correction

- Calonico, Cattaneo, and Titiunik (2014) prove that

$$\frac{\hat{\tau}_{SRD}^{bc} - \tau_{SRD}}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}^{bc}]}} \rightarrow N(0, 1)$$

as long as  $N \min\{b_N^5, h_N^5\} \max\{b_N^2, h_N^2\} \rightarrow 0$ .

- Now the numerator is  $O_p(h^3 + h^2 b^2)$  and the denominator is  $O_p(\frac{1}{\sqrt{Nh}} + \frac{1}{\sqrt{Nh^4 b^5}})$ .
- In other words, we can still use the algorithm in G. Imbens and Kalyanaraman (2012) to select the bandwidth.

## Bias correction

- ▶ Calonico, Cattaneo, and Titiunik (2014) prove that

$$\frac{\hat{\tau}_{SRD}^{bc} - \tau_{SRD}}{\sqrt{\mathbb{V}[\hat{\tau}_{SRD}^{bc}]}} \rightarrow N(0, 1)$$

as long as  $N \min\{b_N^5, h_N^5\} \max\{b_N^2, h_N^2\} \rightarrow 0$ .

- ▶ Now the numerator is  $O_p(h^3 + h^2 b^2)$  and the denominator is  $O_p(\frac{1}{\sqrt{Nh}} + \frac{1}{\sqrt{Nh^4 b^5}})$ .
- ▶ In other words, we can still use the algorithm in G. Imbens and Kalyanaraman (2012) to select the bandwidth.
- ▶ We just need to modify the obtained estimate to ensure asymptotic normality.

## RDD with a discrete running variable

- ▶ The classic theory developed in Calonico, Cattaneo, and Titiunik (2014) runs into difficulties when the running variable is discrete.

## RDD with a discrete running variable

- ▶ The classic theory developed in Calonico, Cattaneo, and Titiunik (2014) runs into difficulties when the running variable is discrete.
- ▶ By definition, there cannot be more observations around the cutoff point as  $N$  increases.

## RDD with a discrete running variable

- ▶ The classic theory developed in Calonico, Cattaneo, and Titiunik (2014) runs into difficulties when the running variable is discrete.
- ▶ By definition, there cannot be more observations around the cutoff point as  $N$  increases.
- ▶ Kolesár and Rothe (2018) provide a finite-sample confidence interval in this case.

## RDD with a discrete running variable

- ▶ The classic theory developed in Calonico, Cattaneo, and Titiunik (2014) runs into difficulties when the running variable is discrete.
- ▶ By definition, there cannot be more observations around the cutoff point as  $N$  increases.
- ▶ Kolesár and Rothe (2018) provide a finite-sample confidence interval in this case.
- ▶ The CI does not rely on asymptotics and holds for any fixed  $N$ .

## RDD with a discrete running variable

- ▶ The classic theory developed in Calonico, Cattaneo, and Titiunik (2014) runs into difficulties when the running variable is discrete.
- ▶ By definition, there cannot be more observations around the cutoff point as  $N$  increases.
- ▶ Kolesár and Rothe (2018) provide a finite-sample confidence interval in this case.
- ▶ The CI does not rely on asymptotics and holds for any fixed  $N$ .
- ▶ The intuition is to bound the curvature of  $\mu(z)$  and consider the worst scenario.



## RDD with a discrete running variable

- ▶ First notice that the local regression estimator is simply a weighted average of the outcome:

$$\hat{\tau}_{SRD} = \sum_{Z_i \geq 0} \hat{\gamma}_+(Z_i) Y_i - \sum_{Z_i \leq 0} \hat{\gamma}_-(Z_i) Y_i$$

where  $\hat{\gamma}_{i,+}$  and  $\hat{\gamma}_{i,-}$  are weights that are only dependent on  $Z$ .

- ▶ In particular, if we use the rectangular kernel, we have

$$\hat{\gamma}_+(Z_i) = \frac{\sum_{0 \leq Z_j \leq h_N} Z_j^2 - Z_i \sum_{0 \leq Z_j \leq h_N} Z_j}{\sum_{0 \leq Z_j \leq h_N} Z_j^2 - (\sum_{0 \leq Z_j \leq h_N} Z_j)^2 / N^*}.$$

where  $N^* = \#\{j : 0 \leq Z_j \leq h_N\}$ .

## RDD with a discrete running variable

- ▶ Let's assume that  $|\mu^{(2)}(z)| \leq C$ , then the bias of  $\hat{\tau}_{SRD}$  is bounded by

$$B = C \sum_{i=1}^N |\hat{\gamma}_+(Z_i) + \hat{\gamma}_-(Z_i)| Z_i^2$$

## RDD with a discrete running variable

- ▶ Let's assume that  $|\mu^{(2)}(z)| \leq C$ , then the bias of  $\hat{\tau}_{SRD}$  is bounded by

$$B = C \sum_{i=1}^N |\hat{\gamma}_+(Z_i) + \hat{\gamma}_-(Z_i)| Z_i^2$$

- ▶ Kolesár and Rothe (2018) show that the one-sided CI of  $\hat{\tau}_{SRD}$  has the form  $[\hat{c}, \infty]$ , where

$$\hat{c} = \hat{\tau}_{SRD} - B - z_{1-\alpha} \sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}$$

## RDD with a discrete running variable

- ▶ Let's assume that  $|\mu^{(2)}(z)| \leq C$ , then the bias of  $\hat{\tau}_{SRD}$  is bounded by

$$B = C \sum_{i=1}^N |\hat{\gamma}_+(Z_i) + \hat{\gamma}_-(Z_i)| Z_i^2$$

- ▶ Kolesár and Rothe (2018) show that the one-sided CI of  $\hat{\tau}_{SRD}$  has the form  $[\hat{c}, \infty]$ , where

$$\hat{c} = \hat{\tau}_{SRD} - B - z_{1-\alpha} \sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}$$

- ▶ Comparing the result with that from Calonico, Cattaneo, and Titiunik (2014), the only difference is that we need to bound the bias using the “smooth parameter”  $C$ .

## RDD with a discrete running variable

- ▶ Let's assume that  $|\mu^{(2)}(z)| \leq C$ , then the bias of  $\hat{\tau}_{SRD}$  is bounded by

$$B = C \sum_{i=1}^N |\hat{\gamma}_+(Z_i) + \hat{\gamma}_-(Z_i)| Z_i^2$$

- ▶ Kolesár and Rothe (2018) show that the one-sided CI of  $\hat{\tau}_{SRD}$  has the form  $[\hat{c}, \infty]$ , where

$$\hat{c} = \hat{\tau}_{SRD} - B - z_{1-\alpha} \sqrt{\mathbb{V}[\hat{\tau}_{SRD}]}$$

- ▶ Comparing the result with that from Calonico, Cattaneo, and Titiunik (2014), the only difference is that we need to bound the bias using the “smooth parameter”  $C$ .
- ▶ The problem here leads to some novel perspectives to understand RDD.

## RDD as local randomization

- ▶ We have discussed that the classic RDD can be seen as a simple experiment when  $Z = 0$ .

## RDD as local randomization

- ▶ We have discussed that the classic RDD can be seen as a simple experiment when  $Z = 0$ .
- ▶ Yet the conventional analysis is not a design-based approach as we do not try to model the assignment process.

## RDD as local randomization

- ▶ We have discussed that the classic RDD can be seen as a simple experiment when  $Z = 0$ .
- ▶ Yet the conventional analysis is not a design-based approach as we do not try to model the assignment process.
- ▶ A recent perspective treats RDD as an experiment that is implemented where  $Z \in [-h, h]$ .



## RDD as local randomization

- ▶ We have discussed that the classic RDD can be seen as a simple experiment when  $Z = 0$ .
- ▶ Yet the conventional analysis is not a design-based approach as we do not try to model the assignment process.
- ▶ A recent perspective treats RDD as an experiment that is implemented where  $Z \in [-h, h]$ .
- ▶ If we know  $h$ , classic approaches (regression, weighting, matching, etc.) can be applied to estimate the treatment effect.

## RDD as local randomization

- ▶ We have discussed that the classic RDD can be seen as a simple experiment when  $Z = 0$ .
- ▶ Yet the conventional analysis is not a design-based approach as we do not try to model the assignment process.
- ▶ A recent perspective treats RDD as an experiment that is implemented where  $Z \in [-h, h]$ .
- ▶ If we know  $h$ , classic approaches (regression, weighting, matching, etc.) can be applied to estimate the treatment effect.
- ▶ Cattaneo, Frandsen, and Titiunik (2015) suggest that we should find  $h$  by balancing all the covariates.

## RDD as local randomization

- ▶ We have discussed that the classic RDD can be seen as a simple experiment when  $Z = 0$ .
- ▶ Yet the conventional analysis is not a design-based approach as we do not try to model the assignment process.
- ▶ A recent perspective treats RDD as an experiment that is implemented where  $Z \in [-h, h]$ .
- ▶ If we know  $h$ , classic approaches (regression, weighting, matching, etc.) can be applied to estimate the treatment effect.
- ▶ Cattaneo, Frandsen, and Titiunik (2015) suggest that we should find  $h$  by balancing all the covariates.
- ▶ The perspective is criticized by Eckles et al. (2020): if the treatment is randomly assigned on  $[-h, h]$ , then  $\mu(z)$  should be constant on the interval.

## RDD as noise-induced randomization

- ▶ Eckles et al. (2020) argue that  $Z$  should be seen as a noisy measure of the true confounder  $U$ .

## RDD as noise-induced randomization

- ▶ Eckles et al. (2020) argue that  $Z$  should be seen as a noisy measure of the true confounder  $U$ .
- ▶ For example,  $U_i$  is one's true capability and  $Z_i$  is her test score.

## RDD as noise-induced randomization

- ▶ Eckles et al. (2020) argue that  $Z$  should be seen as a noisy measure of the true confounder  $U$ .
- ▶ For example,  $U_i$  is one's true capability and  $Z_i$  is her test score.
- ▶ We assume the measurement error is normally distributed

$$Z_i|U_i \sim \mathcal{N}(U_i, \nu^2)$$

and conditionally independent to the potential outcomes

$$\{Y_i(1), Y_i(0)\} \perp Z_i|U_i$$

## RDD as noise-induced randomization

- ▶ Eckles et al. (2020) argue that  $Z$  should be seen as a noisy measure of the true confounder  $U$ .
- ▶ For example,  $U_i$  is one's true capability and  $Z_i$  is her test score.
- ▶ We assume the measurement error is normally distributed

$$Z_i|U_i \sim \mathcal{N}(U_i, \nu^2)$$

and conditionally independent to the potential outcomes

$$\{Y_i(1), Y_i(0)\} \perp Z_i|U_i$$

- ▶ The assumptions imply that

$$\{Y_i(1), Y_i(0)\} \perp D_i|U_i$$

and the only issue is that  $U_i$  is unobservable.

## RDD as noise-induced randomization

- ▶ Eckles et al. (2020) claim that it is possible to estimate the causal effect using the noisy measure  $Z_i$ .



## RDD as noise-induced randomization

- ▶ Eckles et al. (2020) claim that it is possible to estimate the causal effect using the noisy measure  $Z_i$ .
- ▶ Remember that when the confounders are observable, all we need to do is to balance the confounders across treatment groups.

## RDD as noise-induced randomization

- ▶ Eckles et al. (2020) claim that it is possible to estimate the causal effect using the noisy measure  $Z_i$ .
- ▶ Remember that when the confounders are observable, all we need to do is to balance the confounders across treatment groups.
- ▶ Now the confounder is unobservable, hence we should balance a functional  $f$  of its noisy measure  $Z_i$ .

## RDD as noise-induced randomization

- ▶ Eckles et al. (2020) claim that it is possible to estimate the causal effect using the noisy measure  $Z_i$ .
- ▶ Remember that when the confounders are observable, all we need to do is to balance the confounders across treatment groups.
- ▶ Now the confounder is unobservable, hence we should balance a functional  $f$  of its noisy measure  $Z_i$ .
- ▶  $Z_i$  contains information of  $U_i$  and we know their relationship.

## RDD as noise-induced randomization

- ▶ Eckles et al. (2020) claim that it is possible to estimate the causal effect using the noisy measure  $Z_i$ .
- ▶ Remember that when the confounders are observable, all we need to do is to balance the confounders across treatment groups.
- ▶ Now the confounder is unobservable, hence we should balance a functional  $f$  of its noisy measure  $Z_i$ .
- ▶  $Z_i$  contains information of  $U_i$  and we know their relationship.
- ▶ If  $f(Z_i)$  is balanced and  $f$  is properly chosen, we expect  $U_i$  to be balanced as well.

## RDD as noise-induced randomization

- ▶ Eckles et al. (2020) claim that it is possible to estimate the causal effect using the noisy measure  $Z_i$ .
- ▶ Remember that when the confounders are observable, all we need to do is to balance the confounders across treatment groups.
- ▶ Now the confounder is unobservable, hence we should balance a functional  $f$  of its noisy measure  $Z_i$ .
- ▶  $Z_i$  contains information of  $U_i$  and we know their relationship.
- ▶ If  $f(Z_i)$  is balanced and  $f$  is properly chosen, we expect  $U_i$  to be balanced as well.
- ▶ Essentially, we are learning the information of unobservables from observable variables.

## RDD as noise-induced randomization

- ▶ Eckles et al. (2020) claim that it is possible to estimate the causal effect using the noisy measure  $Z_i$ .
- ▶ Remember that when the confounders are observable, all we need to do is to balance the confounders across treatment groups.
- ▶ Now the confounder is unobservable, hence we should balance a functional  $f$  of its noisy measure  $Z_i$ .
- ▶  $Z_i$  contains information of  $U_i$  and we know their relationship.
- ▶ If  $f(Z_i)$  is balanced and  $f$  is properly chosen, we expect  $U_i$  to be balanced as well.
- ▶ Essentially, we are learning the information of unobservables from observable variables.
- ▶ This is an idea called “manifold learning.”

## RDD as noise-induced randomization

- ▶ Eckles et al. (2020) also focus on estimators with the following form:

$$\hat{\tau}_{SRD} = \sum_{Z_i \geq 0} \hat{\gamma}_+(Z_i) Y_i - \sum_{Z_i \leq 0} \hat{\gamma}_-(Z_i) Y_i$$

## RDD as noise-induced randomization

- ▶ Eckles et al. (2020) also focus on estimators with the following form:

$$\hat{\tau}_{SRD} = \sum_{Z_i \geq 0} \hat{\gamma}_+(Z_i) Y_i - \sum_{Z_i \leq 0} \hat{\gamma}_-(Z_i) Y_i$$

- ▶  $\hat{\gamma}_{i,+}$  and  $\hat{\gamma}_{i,-}$  need to balance the confounder  $U_i$ , thus we require

$$\int_{-\infty}^0 \hat{\gamma}_-(z) dF(z) = 1, \int_0^{\infty} \hat{\gamma}_+(z) dF(z) = 1$$

$$\left| \int_{-\infty}^0 \hat{\gamma}_-(z) \phi(z|u) dz - \int_0^{\infty} \hat{\gamma}_+(z) \phi(z|u) dz \right| \leq t \text{ for all } u$$



## RDD as noise-induced randomization

- ▶ Eckles et al. (2020) also focus on estimators with the following form:

$$\hat{\tau}_{SRD} = \sum_{Z_i \geq 0} \hat{\gamma}_+(Z_i) Y_i - \sum_{Z_i \leq 0} \hat{\gamma}_-(Z_i) Y_i$$

- ▶  $\hat{\gamma}_{i,+}$  and  $\hat{\gamma}_{i,-}$  need to balance the confounder  $U_i$ , thus we require

$$\int_{-\infty}^0 \hat{\gamma}_-(z) dF(z) = 1, \int_0^{\infty} \hat{\gamma}_+(z) dF(z) = 1$$
$$\left| \int_{-\infty}^0 \hat{\gamma}_-(z) \phi(z|u) dz - \int_0^{\infty} \hat{\gamma}_+(z) \phi(z|u) dz \right| \leq t \text{ for all } u$$

- ▶ Note that the classic local regression estimator is a special case of  $\hat{\tau}_{SRD}$  where the weights are estimated based on OLS.

## RDD as noise-induced randomization

- ▶ Nevertheless, the regression weights are not optimal for the MSE  $\mathbb{E}[\hat{\tau}_{SRD} - \tau_{SRD}]^2$ .

## RDD as noise-induced randomization

- ▶ Nevertheless, the regression weights are not optimal for the MSE  $\mathbb{E}[\hat{\tau}_{SRD} - \tau_{SRD}]^2$ .
- ▶ Eckles et al. (2020) suggest that researchers should directly minimize the MSE under the constraints above:

$$\begin{aligned} & \hat{\gamma}_+(\cdot), \hat{\gamma}_-(\cdot) \\ &= \arg \max_{\gamma_+, \gamma_-} \left[ \frac{\sigma^2}{N} \left( \int_{-\infty}^0 \hat{\gamma}_-^2(z) dF(z) + \int_0^{\infty} \hat{\gamma}_+^2(z) dF(z) \right) + \frac{t^2}{M^2} \right] \end{aligned}$$

where  $\sigma^2 = \text{Var}[Y_i|Z_i]$  and  $M$  is a constant.

## RDD as noise-induced randomization

- ▶ Nevertheless, the regression weights are not optimal for the MSE  $\mathbb{E}[\hat{\tau}_{SRD} - \tau_{SRD}]^2$ .
- ▶ Eckles et al. (2020) suggest that researchers should directly minimize the MSE under the constraints above:

$$\begin{aligned} & \hat{\gamma}_+(\cdot), \hat{\gamma}_-(\cdot) \\ &= \arg \max_{\gamma_+, \gamma_-} \left[ \frac{\sigma^2}{N} \left( \int_{-\infty}^0 \hat{\gamma}_-^2(z) dF(z) + \int_0^{\infty} \hat{\gamma}_+^2(z) dF(z) \right) + \frac{t^2}{M^2} \right] \end{aligned}$$

where  $\sigma^2 = \text{Var}[Y_i|Z_i]$  and  $M$  is a constant.

- ▶ It is a standard convex optimization problem and can be easily solved in statistical software.

## RDD as noise-induced randomization

- ▶ Nevertheless, the regression weights are not optimal for the MSE  $\mathbb{E}[\hat{\tau}_{SRD} - \tau_{SRD}]^2$ .
- ▶ Eckles et al. (2020) suggest that researchers should directly minimize the MSE under the constraints above:

$$\begin{aligned} & \hat{\gamma}_+(\cdot), \hat{\gamma}_-(\cdot) \\ &= \arg \max_{\gamma_+, \gamma_-} \left[ \frac{\sigma^2}{N} \left( \int_{-\infty}^0 \hat{\gamma}_-^2(z) dF(z) + \int_0^{\infty} \hat{\gamma}_+^2(z) dF(z) \right) + \frac{t^2}{M^2} \right] \end{aligned}$$

where  $\sigma^2 = \text{Var}[Y_i|Z_i]$  and  $M$  is a constant.

- ▶ It is a standard convex optimization problem and can be easily solved in statistical software.
- ▶ Eckles et al. (2020) show that the studentized  $\hat{\tau}_{SRD}$  is asymptotically normal.

## RDD as noise-induced randomization

- ▶ The approach in Eckles et al. (2020) is purely design-based and requires no assumption on the smoothness of the regression function.

## RDD as noise-induced randomization

- ▶ The approach in Eckles et al. (2020) is purely design-based and requires no assumption on the smoothness of the regression function.
- ▶ It applies to cases with either a continuous running variable or a discrete running variable.

# RDD as noise-induced randomization

- ▶ The approach in Eckles et al. (2020) is purely design-based and requires no assumption on the smoothness of the regression function.
- ▶ It applies to cases with either a continuous running variable or a discrete running variable.
- ▶ Inference is also straightforward in this method.



## RDD as noise-induced randomization

- ▶ The approach in Eckles et al. (2020) is purely design-based and requires no assumption on the smoothness of the regression function.
- ▶ It applies to cases with either a continuous running variable or a discrete running variable.
- ▶ Inference is also straightforward in this method.
- ▶ However, it requires the knowledge of the measurement error, which might be unrealistic in practice.

# RDD as noise-induced randomization

- ▶ The approach in Eckles et al. (2020) is purely design-based and requires no assumption on the smoothness of the regression function.
- ▶ It applies to cases with either a continuous running variable or a discrete running variable.
- ▶ Inference is also straightforward in this method.
- ▶ However, it requires the knowledge of the measurement error, which might be unrealistic in practice.
- ▶ The future direction is to learn the conditional distribution from data.

# References I

Abadie, Alberto, and Guido W Imbens. 2006. "Large Sample Properties of Matching Estimators for Average Treatment Effects." *Econometrica* 74 (1). Wiley Online Library: 235–67.

Calonico, Sebastian, Matias D Cattaneo, and Max H Farrell. 2020. "Optimal Bandwidth Choice for Robust Bias-Corrected Inference in Regression Discontinuity Designs." *The Econometrics Journal* 23 (2). Oxford University Press: 192–210.

Calonico, Sebastian, Matias D Cattaneo, and Rocio Titiunik. 2014. "Robust Nonparametric Confidence Intervals for Regression-Discontinuity Designs." *Econometrica* 82 (6). Wiley Online Library: 2295–2326.

Card, David, David S Lee, Zhuan Pei, and Andrea Weber. 2015. "Inference on Causal Effects in a Generalized Regression Kink

## References II

Design." *Econometrica* 83 (6). Wiley Online Library: 2453–83.

Cattaneo, Matias D, Brigham R Frandsen, and Rocio Titiunik. 2015. "Randomization Inference in the Regression Discontinuity Design: An Application to Party Advantages in the Us Senate." *Journal of Causal Inference* 3 (1). De Gruyter: 1–24.

Cattaneo, Matias D, Michael Jansson, and Xinwei Ma. 2020. "Simple Local Polynomial Density Estimators." *Journal of the American Statistical Association* 115 (531). Taylor & Francis: 1449–55.

Cattaneo, Matias D, Luke Keele, Rocío Titiunik, and Gonzalo Vazquez-Bare. 2020. "Extrapolating Treatment Effects in Multi-Cutoff Regression Discontinuity Designs." *Journal of the American Statistical Association*. Taylor & Francis, 1–12.

Cattaneo, Matias D, Rocío Titiunik, Gonzalo Vazquez-Bare, and Luke Keele. 2016. "Interpreting Regression Discontinuity Designs with Multiple Cutoffs." *The Journal of Politics* 78 (4). University of

## References III

Chicago Press Chicago, IL: 1229–48.

Eckles, Dean, Nikolaos Ignatiadis, Stefan Wager, and Han Wu. 2020. “Noise-Induced Randomization in Regression Discontinuity Designs.” *arXiv Preprint arXiv:2004.09458*.

Fan, Jianqing, and Irene Gijbels. 1996. *Local Polynomial Modelling and Its Applications: Monographs on Statistics and Applied Probability 66*. Vol. 66. CRC Press.

Gelman, Andrew, and Guido Imbens. 2019. “Why High-Order Polynomials Should Not Be Used in Regression Discontinuity Designs.” *Journal of Business & Economic Statistics* 37 (3). Taylor & Francis: 447–56.

Hahn, Jinyong, Petra Todd, and Wilbert Van der Klaauw. 2001. “Identification and Estimation of Treatment Effects with a Regression-Discontinuity Design.” *Econometrica* 69 (1). JSTOR:

## References IV

201–9.

Imbens, Guido, and Karthik Kalyanaraman. 2012. “Optimal Bandwidth Choice for the Regression Discontinuity Estimator.” *The Review of Economic Studies* 79 (3). Oxford University Press: 933–59.

Imbens, Guido, and Stefan Wager. 2019. “Optimized Regression Discontinuity Designs.” *Review of Economics and Statistics* 101 (2). MIT Press: 264–78.

Kolesár, Michal, and Christoph Rothe. 2018. “Inference in Regression Discontinuity Designs with a Discrete Running Variable.” *American Economic Review* 108 (8): 2277–2304.

Lee, David S. 2008. “Randomized Experiments from Non-Random Selection in Us House Elections.” *Journal of Econometrics* 142 (2).

## References V

Elsevier: 675–97.

McCrary, Justin. 2008. “Manipulation of the Running Variable in the Regression Discontinuity Design: A Density Test.” *Journal of Econometrics* 142 (2). Elsevier: 698–714.

Meyersson, Erik. 2014. “Islamic Rule and the Empowerment of the Poor and Pious.” *Econometrica* 82 (1). Wiley Online Library: 229–69.

Porter, Jack. 2003. “Estimation in the Regression Discontinuity Model.” *Unpublished Manuscript, Department of Economics, University of Wisconsin at Madison* 2003. Citeseer: 5–19.

Thistlethwaite, Donald L, and Donald T Campbell. 1960. “Regression-Discontinuity Analysis: An Alternative to the Ex Post Facto Experiment.” *Journal of Educational Psychology* 51 (6). American Psychological Association: 309.