

# Math 5223 Homework 5

Jiasong Zhu

October 20, 2025

## Problem 1

*Proof.* (a) Let  $r(\theta, \varphi) = (R \sin \varphi \cos \theta, R \sin \varphi \sin \theta, R \cos \varphi)$ ,  $-\pi < \theta < \pi, 0 < \varphi < \pi$ . Note that

$$r_\theta = \frac{\partial r}{\partial \theta} = (-R \sin \varphi \sin \theta, R \sin \varphi \cos \theta, 0),$$

$$r_\varphi = \frac{\partial r}{\partial \varphi} = (R \cos \varphi \cos \theta, R \cos \varphi \sin \theta, -R \sin \varphi).$$

Compute,

$$\langle \mathbf{r}_\theta, \mathbf{r}_\theta \rangle = R^2 \sin^2 \varphi (\sin^2 \theta + \cos^2 \theta) = R^2 \sin^2 \varphi,$$

$$\langle \mathbf{r}_\varphi, \mathbf{r}_\varphi \rangle = R^2 (\cos^2 \varphi + \sin^2 \varphi) = R^2,$$

$$\langle \mathbf{r}_\theta, \mathbf{r}_\varphi \rangle = 0.$$

Hence, in the coordinates  $(\theta, \varphi)$ , the round metric is

$$g_R = \langle \mathbf{r}_\varphi, \mathbf{r}_\varphi \rangle d\varphi^2 + 2\langle \mathbf{r}_\theta, \mathbf{r}_\varphi \rangle d\theta d\varphi + \langle \mathbf{r}_\theta, \mathbf{r}_\theta \rangle d\theta^2 = R^2 d\varphi^2 + R^2 \sin^2 \varphi d\theta^2,$$

as desired.

(b) In the coordinates  $(\theta, \varphi)$ , the only nonzero components of the metric are

$$g_{\varphi\varphi} = R^2, \quad g_{\theta\theta} = R^2 \sin^2 \varphi,$$

so the inverse metric satisfies

$$g^{\varphi\varphi} = \frac{1}{R^2}, \quad g^{\theta\theta} = \frac{1}{R^2 \sin^2 \varphi}.$$

Since  $g_{\theta\theta}$  depends only on  $\varphi$  and  $g_{\varphi\varphi}$  is constant,

$$\partial_\varphi g_{\theta\theta} = 2R^2 \sin \varphi \cos \varphi, \quad \partial_\theta g_{\theta\theta} = 0, \quad \partial_\varphi g_{\varphi\varphi} = 0, \quad \partial_\theta g_{\varphi\varphi} = 0$$

Hence the only nonzero Christoffel symbols are

$$\Gamma^\varphi_{\theta\theta} = \frac{1}{2} g^{\varphi\varphi} (0 + 0 - \partial_\varphi g_{\theta\theta}) = -\frac{1}{2} \cdot \frac{1}{R^2} \cdot 2R^2 \sin \varphi \cos \varphi = -\sin \varphi \cos \varphi,$$

$$\Gamma^\theta_{\theta\varphi} = \Gamma^\theta_{\varphi\theta} = \frac{1}{2} g^{\theta\theta} \partial_\varphi g_{\theta\theta} = \frac{1}{2} \cdot \frac{1}{R^2 \sin^2 \varphi} \cdot 2R^2 \sin \varphi \cos \varphi = \cot \varphi.$$

(c) With the metric  $g_R = R^2 d\varphi^2 + R^2 \sin^2 \varphi d\theta^2$  the only nonzero Christoffel symbols are

$$\Gamma^\varphi_{\theta\theta} = -\sin \varphi \cos \varphi, \quad \Gamma^\theta_{\theta\varphi} = \Gamma^\theta_{\varphi\theta} = \cot \varphi.$$

Hence the geodesic equations in coordinates  $(\theta(t), \varphi(t))$  are

$$\begin{aligned} \theta'' + 2\Gamma^\theta_{\theta\varphi} \theta' \varphi' &= \theta'' + 2 \cot \varphi \theta' \varphi' = 0, \\ \varphi'' + \Gamma^\varphi_{\theta\theta} (\theta')^2 &= \varphi'' - \sin \varphi \cos \varphi (\theta')^2 = 0. \end{aligned}$$

Consider a meridian:  $\theta(t) \equiv \theta_0$  and  $\varphi(t) = t$ . Then  $\theta' = \theta'' = 0$ , so the first equation holds trivially, and the second becomes  $\varphi'' = 0$ , which is satisfied by  $\varphi(t) = at + b$ . Therefore  $(\theta(t), \varphi(t)) = (\theta_0, t)$  is a geodesic.  $\square$

## Problem 2

*Proof.* On the unit sphere with spherical coordinates  $(\theta, \varphi)$ , the round metric is  $g = d\varphi^2 + \sin^2 \varphi d\theta^2$ . From Problem 1(b), the only nonzero Christoffel symbols are

$$\Gamma^\varphi_{\theta\theta} = -\sin \varphi \cos \varphi, \quad \Gamma^\theta_{\theta\varphi} = \Gamma^\theta_{\varphi\theta} = \cot \varphi.$$

Let  $V = \frac{\partial}{\partial \varphi}$ . Using  $\nabla_{\partial_i} \partial_j = \Gamma^k_{ij} \partial_k$  for the coordinate frame, we have

$$\nabla_{\frac{\partial}{\partial \theta}} V = \nabla_{\partial_\theta} \partial_\varphi = \Gamma^\theta_{\theta\varphi} \partial_\theta + \Gamma^\varphi_{\theta\varphi} \partial_\varphi = \cot \varphi \frac{\partial}{\partial \theta},$$

since  $\Gamma^\varphi_{\theta\varphi} = 0$ . Also,

$$\nabla_{\frac{\partial}{\partial \varphi}} V = \nabla_{\partial_\varphi} \partial_\varphi = \Gamma^\theta_{\varphi\varphi} \partial_\theta + \Gamma^\varphi_{\varphi\varphi} \partial_\varphi = 0,$$

because  $\Gamma^\theta_{\varphi\varphi} = \Gamma^\varphi_{\varphi\varphi} = 0$ .

Along the equator  $\varphi = \pi/2$ , the tangent is  $\partial_\theta$  and

$$\nabla_{\partial_\theta} V = \cot(\pi/2) \partial_\theta = 0,$$

so  $V$  is parallel along the equator. Along any meridian  $\theta = \theta_0$ , the tangent is  $\partial_\varphi$  and  $\nabla_{\partial_\varphi} V = 0$ , so  $V$  is parallel along each meridian.  $\square$

### Problem 3

*Proof.* Fix  $p \in M$ . Using the standard identification  $T_{(p,0)}(TM) \cong T_p M \oplus T_p M$  (horizontal  $\oplus$  vertical), write a tangent vector as  $(X, W)$ . Consider the map  $G : TM \rightarrow M$ ,  $G(q, V) = \exp_q V$ . Then

$$dG_{(p,0)}(X, W) = X + W.$$

Indeed, if  $V \equiv 0$  and  $q(s)$  is a curve with  $q(0) = p$ ,  $q'(0) = X$ , then  $G(q(s), 0) = q(s)$ , hence  $\frac{d}{ds}\big|_0 G = X$ . If  $q \equiv p$  and  $V(s)$  is a curve in  $T_p M$  with  $V(0) = 0$ ,  $V'(0) = W$ , then  $\frac{d}{ds}\big|_0 \exp_p V(s) = d(\exp_p)_0(W) = W$  since  $d(\exp_p)_0 = \text{id}_{T_p M}$ . By linearity the general variation gives  $X + W$ .

Thus, for  $F(q, V) = (q, G(q, V))$ ,

$$dF_{(p,0)}(X, W) = (X, dG_{(p,0)}(X, W)) = (X, X + W) \in T_p M \oplus T_p M \cong T_{(p,p)}(M \times M).$$

With respect to the decompositions  $T_{(p,0)}(TM) \cong T_p M \oplus T_p M$  and  $T_{(p,p)}(M \times M) \cong T_p M \oplus T_p M$ , this is the block matrix

$$dF_{(p,0)} = \begin{pmatrix} \text{id} & 0 \\ \text{id} & \text{id} \end{pmatrix}, \quad \text{whose inverse is } \begin{pmatrix} \text{id} & 0 \\ -\text{id} & \text{id} \end{pmatrix}.$$

Hence  $dF_{(p,0)}$  is an isomorphism.

By the inverse function theorem, there exist neighborhoods  $\mathcal{U}_p \subset TM$  of  $(p, 0)$  and  $\mathcal{V}_p \subset M \times M$  of  $(p, p)$  such that  $F : \mathcal{U}_p \rightarrow \mathcal{V}_p$  is a diffeomorphism. Since  $p$  was arbitrary, shrinking and taking the union over  $p$  shows that  $F$  is a local diffeomorphism from a neighborhood of the zero section in  $TM$  onto a neighborhood of the diagonal  $\Delta \subset M \times M$ . □

### Problem 4

*Proof.* (a) Let  $T(s, t) = \partial_t \Gamma(s, t)$ . Note that

$$\begin{aligned} \frac{d}{ds}\bigg|_{s=0} L(\Gamma_s) &= \frac{d}{ds}\bigg|_{s=0} \sum_{i=1}^{k-1} \int_{a_i}^{a_{i+1}} |T(s, t)|, dt \\ &= \sum_{i=1}^{k-1} \int_{a_i}^{a_{i+1}} \frac{\langle \nabla_{\partial_s} T, T \rangle}{|T|} \bigg|_{s=0} dt \\ &\stackrel{|T|=1 \text{ at } s=0}{=} \sum_{i=1}^{k-1} \int_{a_i}^{a_{i+1}} \langle \nabla_{\partial_s} T, \dot{\gamma} \rangle, dt. \end{aligned}$$

Since the Levi-Civita connection is torsion free, we have  $\nabla_{\partial_s} T = \nabla_{\partial_t} \partial_s \Gamma$ . Thus, with  $V(t) = \partial_s \Gamma(0, t)$ , we have

$$\left. \frac{d}{ds} \right|_{s=0} L(\Gamma_s) = \sum_{i=1}^{k-1} \int_{a_i}^{a_{i+1}} \langle D_t V, \dot{\gamma} \rangle, dt.$$

By integration by part,

$$\int_{a_i}^{a_{i+1}} \langle D_t V, \dot{\gamma} \rangle, dt = [\langle V, \dot{\gamma} \rangle]_{t=a_i^+}^{t=a_{i+1}^-} - \int_{a_i}^{a_{i+1}} \langle V, D_t \dot{\gamma} \rangle, dt.$$

Take the sum over all  $i$ , we have

$$\sum_{i=1}^{k-1} [\langle V, \dot{\gamma} \rangle]_{a_i^+}^{a_{i+1}^-} = \langle V(b), \dot{\gamma}(b) \rangle - \langle V(a), \dot{\gamma}(a) \rangle - \sum_{i=2}^{k-1} \langle V(a_i), \Delta_i \dot{\gamma} \rangle,$$

since the interior endpoint contributions telescope and leave the jumps of  $\dot{\gamma}$  at the break points  $a_i$ . Therefore,

$$\left. \frac{d}{ds} \right|_{s=0} L(\Gamma_s) = - \int_a^b \langle V, D_t \dot{\gamma} \rangle, dt + \langle V(b), \dot{\gamma}(b) \rangle - \langle V(a), \dot{\gamma}(a) \rangle - \sum_{i=1}^{k-1} \langle V(a_i), \Delta_i \dot{\gamma} \rangle.$$

(b) Fix any  $w \in T_q N$ . Take a smooth curve  $\eta : (-\delta, \delta) \rightarrow N$  with  $\eta(0) = q$  and  $\eta'(0) = w$ . For  $|s|$  small, join  $p$  to  $\eta(s)$  by a unit-speed minimizing geodesic  $\gamma_s$ , and define a smooth variation

$$\Gamma : (-\varepsilon, \varepsilon) \times [0, 1] \rightarrow M, \quad \Gamma(s, t) = \gamma_s(t).$$

Let  $V(t) := \partial_s \Gamma(0, t)$  be its variation field along  $\gamma$ . Then  $V(0) = 0$  and  $V(1) = \eta'(0) = w \in T_q N$ .

Since  $q$  realizes the distance from  $p$  to  $N$ , the length function  $L(\Gamma_s)$  has a minimum at  $s = 0$ , hence

$$\left. \frac{d}{ds} \right|_{s=0} L(\Gamma_s) = 0.$$

Applying the first variation formula from part (a),

$$\left. \frac{d}{ds} \right|_{s=0} L(\Gamma_s) = - \int_0^1 \langle V, D_t \dot{\gamma} \rangle dt + \langle V(1), \dot{\gamma}(1) \rangle - \langle V(0), \dot{\gamma}(0) \rangle.$$

Because  $\gamma$  is a geodesic,  $D_t \dot{\gamma} = 0$ , and  $V(0) = 0$ , we obtain

$$0 = \langle V(1), \dot{\gamma}(1) \rangle = \langle w, \dot{\gamma}(1) \rangle.$$

This holds for every  $w \in T_q N$ , so  $\dot{\gamma}(1) \perp T_q N$ . Therefore any minimizing geodesic from  $p$  to a nearest point  $q \in N$  meets  $N$  orthogonally.  $\square$

## Problem 5

*Proof.*

□