

Kinetic Resolution of Propargylic Ethers via [2,3]-Wittig Rearrangement to Synthesize Chiral α -Hydroxyallenes

Xi Xu, Shunxi Dong, LiLi Feng, Sijing Wang, Xiaohua Liu,* and Xiaoming Feng*



Cite This: *Org. Lett.* 2020, 22, 2692–2696



Read Online

ACCESS |




Metrics & More



Article Recommendations



Supporting Information




ABSTRACT: An efficient kinetic resolution of propargyloxy dicarbonyl compounds via asymmetric [2,3]-Wittig rearrangement was achieved by using a chiral N,N' -dioxide/ Ni^{II} complex catalyst. Various chiral α -allenyl alcohols were obtained in high enantioselectivities under mild conditions. The utility of this method was readily demonstrated in the asymmetric synthesis of the chiral 2,5-dihydrofuran derivative.

α -Hydroxyallenes widely exist in a series of natural biological products and also serve as versatile synthetic intermediates in organic transformations.¹ The construction of α -hydroxyallenes has attracted considerable attention for decades, and several catalytic enantioselective synthetic methods have been developed, such as desymmetrization reaction,² Aldol-like reaction,³ arylboration reaction,⁴ Petasis reaction,⁵ and rearrangement reaction.⁶ Among them, [2,3]-Wittig rearrangement^{7,8} of propargylic ethers⁹ provides an efficient and facile route to elaborated functionalized α -hydroxyallenes. Recently, our group described an asymmetric propargyl [2,3]-Wittig rearrangement of oxindole derivatives using a chiral N,N' -dioxide/ Ni^{II} complex, and the corresponding 3-hydroxy 3-substituted oxindoles bearing an allenyl group were obtained in good results (Scheme 1a).^{6c} Moreover, chiral dihydrofuran derivatives represent important structural motifs existing in natural products,¹⁰ such as diplobifuranyllone B,¹¹ (+)-furanomycin,¹² and cryptoresinol (Scheme 1c),¹³ and intramolecular hydroalkoxylation of chiral α -hydroxyallenes¹⁴ is usually one of the most powerful strategies to construct these scaffolds. In consequence, development of new and efficient catalytic asymmetric approaches for their rapid access is in high demand.


With this aim, we envisioned that chiral Lewis acid catalysts were potentially capable to realize kinetic resolution of racemic propargyloxymalonate derivatives via asymmetric [2,3]-Wittig rearrangement. As shown in Scheme 1b, chiral Lewis acid complexes could recognize one enantiomer and promote it to rearrange into an axial chiral α -hydroxyallene product via chirality transfer, leaving behind the enrichment of the other enantiomer. However, on the one hand, control of the regio-

Scheme 1. Asymmetric Propargylic [2,3]-Wittig Rearrangement and Representative Related Compounds


(a) Propargylic [2,3]-Wittig rearrangement to synthesize chiral 3-allenyl 3-hydroxyoxindoles^{6c}



(b) Propargylic [2,3]-Wittig rearrangement via kinetic resolution to synthesize α -hydroxyallenes (This work)



(c) Representative natural products containing a dihydrofuran skeleton



Received: February 18, 2020

Published: March 13, 2020




ACS Publications

© 2020 American Chemical Society

2692

<https://dx.doi.org/10.1021/acs.orglett.0c00649>
Org. Lett. 2020, 22, 2692–2696

Table 1. Optimization of the Reaction Conditions^a

entry	ligand	metal salts	1a/2a			3a	
			yield (%) ^b	1a:2a	ee (%) ^c	yield (%) ^b	ee (%) ^c
1	L-PiCHPh ₂	Ni(OTf) ₂	52	26/74	29/56	29	92
2	L-PimtBu	Ni(OTf) ₂	69	5/95	–/11	32	38
3	L-PiPr ₃	Ni(OTf) ₂	89	99/1	8/–	14	98
4	L-PiPr ₃	Ni(ClO ₄) ₂ ·6H ₂ O	75	98/2	26/–	24	99
5 ^d	L-PiPr ₃	Ni(ClO ₄) ₂ ·6H ₂ O	50	93/7	93/–	46	97
6	tBu-box	Ni(OTf) ₂	84	99/1	0/–	–	–

^aUnless otherwise noted, the reactions were performed with ligand/metal salts (1:1, 10 mol %), **1a** (0.1 mmol), and Et₃N (1.2 equiv) in EtOAc (1.0 mL) at 50 °C for 48 h. ^bIsolated yield, and the ratio of **1a** to **2a** in the bracket represents was determined via HPLC analysis. ^cDetermined by HPLC analysis on a chiral stationary phase. ^dThe reaction was performed in 0.3 mL of EtOAc.

and stereoselectivity was the foreseeable challenge in developing such a synergistic kinetic resolution/[2,3]-Wittig rearrangement reaction having several competitive side reactions. On the other hand, the propargylic ethers show low reactivity in that the alkyne sp centers distort the transition-state geometries. Herein, we wish to disclose the results in this endeavor. A chiral *N,N'*-dioxide-Ni^{II} complex^{15,16} was found to be efficient in promoting asymmetric propargyl [2,3]-Wittig rearrangement via kinetic resolution of α -propargyloxy dicarbonyl compounds with high to excellent enantioselectivities. This methodology was further applied in the asymmetric synthesis of a chiral 2,5-dihydrofuran derivative.


Initially, we commenced our investigation with the kinetic resolution of racemic α -propargyloxy dicarbonyl compound **1a** by [2,3]-Wittig rearrangement (Table 1). L-PiCHPh₂/Ni(OTf)₂, which is efficient in our previous work,^{6c} was tested in EtOAc (1.0 mL) at 50 °C in the presence of Et₃N. [2,3]-Wittig rearrangement of the racemic compound **1a** took place smoothly, giving the corresponding chiral α -hydroxyallene **3a** in 29% yield with 92% ee (entry 1). Meanwhile, the unreacted α -propargyloxy ether **1a** along with dihydrofuran byproduct **2a** was isolated together (the ratio of **1a** to **2a**, ca. 26:74, 29% ee for **1a**, 56% ee for **2a**).¹⁷ Due to a similar *R_f* value, column separation of **1a** and **2a** was unsuccessful. Next, other metal salts were investigated by coordinating with L-PiCHPh₂, but no better results were given (see SI, p S4 for further details). Then, chiral *N,N'*-dioxide ligands were examined (entries 2 and 3), and it was found that both the amino acid skeleton and steric hindrance of amide moiety of the ligands had a significant influence on the regio- and enantioselectivity. When L-PimtBu was used, the separated major compound was **2a** (**1a**:**2a** = 5:95, 69% yield in total, 11% ee for **2a**), and allene **3a** was isolated in 32% yield with 38% ee (entry 2). In sharp contrast, the use of L-PiPr₃ prevented the formation of dihydrofuran **2a**, albeit the allene **3a** was afforded in 14% yield with 98% ee (entry 3). Switching Ni(OTf)₂ to Ni(ClO₄)₂·6H₂O resulted in the improved ee value for recovered **1a** and a

slightly better isolated yield for allene (entry 4). To our delight, when the reaction was performed with a higher concentration, the α -propargyloxy ether **1a** was recovered in 50% yield (**1a**:**2a**, 93/7) with 93% ee, and allene **3a** was separated in 46% yield with 97% ee (entry 5; see SI for further details). In comparison, the Ni^{II}/bisoxazoline catalyst was not able to promote the titled rearrangement reaction (entry 6).

Having identified the optimized reaction conditions in hand, we investigated the substrate scope of the reaction. Due to the separation problem, only the allene products **3** were isolated and fully characterized. As depicted in Scheme 2, varying the diester groups affected the reactivity. Substrate **1a** with methyl ester and substrate **1b** bearing ethyl ester had higher activity than substrate **1c** with sterically congested *tert*-butyl ester, but their products **3a–3c** were isolated in high enantioselectivities (97–99% ee). Aryl substituents at the terminal position α -propargyloxy ethers were investigated, and the position of the substituents had a limited impact on the yield and selectivity (**3e–3g**, 43–45% yields, and 97–99% ee). Substrates either with an electron-donating substituent (methyl or methoxyl) or with an electron-withdrawing substituent (chloro, ester, and trifluoromethyl) underwent kinetic resolution/asymmetric [2,3]-Wittig rearrangement smoothly and afforded the corresponding α -allenyl alcohols (**3d**, **3h–j**) in good yields (42–47%) with high enantioselectivities (91–98% ee). Moreover, 1-naphthyl-substituted product **3k** and 2-thiophenyl-containing product **3l** were formed with good ee values (90% ee and 95% ee). Regardless of the steric hindrance of R² groups at the stereogenic center (isopropyl, *n*-butyl, and benzyl), all these propargylic ethers could give excellent enantioselectivities (**3m–3o**, 96–98% ee). The absolute configuration of **3o** was determined to be (aR) by X-ray crystallographic analysis (CCDC 1951577).


To further demonstrate the potential of this protocol, we first carried out the gram-scale successive kinetic resolution/[2,3]-Wittig rearrangement of racemic 4-phenylbut-3-yn-2-ol derived **1a** for the synthesis of both enantiomers of chiral α -allenyl alcohol **3a** (Scheme 3a). By treatment of 8.0 mmol of

Scheme 2. Kinetic Resolution and Asymmetric [2,3]-Wittig Rearrangement of Racemic Propargyl Ethers^a



^aUnless otherwise noted, the reactions were performed with $\text{L-PiPr}_3/\text{Ni}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ (10 mol %, 1:1), **1** (0.2 mmol), and Et_3N (1.2 equiv) in EtOAc (0.6 mL) at 50 °C. ^b**1** (0.1 mmol) in EtOAc (0.3 mL).

Scheme 3. Gram-Scale Reaction and Derivatization



(±)-**1a** in the presence of 10 mol % $\text{L-PiPr}_3/\text{Ni}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ complex for 89 h, the isolated (aR)-**3a** was obtained in 37% yield with 97% ee. The recovered crude materials were subsequently subjected to the catalyst system of *ent*- $\text{L-PiPr}_3/\text{Ni}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ complex for 84 h to give (aS)-**3a** with 93% ee. Moreover, the asymmetric synthesis of the basic skeleton of natural product cryptoresinol was accomplished from the isolated allenyl alcohol product. As shown in Scheme 3b, the product **3o** could be easily transformed into dihydrofuran **2o** in 98% yield and 97% ee by treating with AgNO_3 and CaCO_3 . Then, the de-esterification of dihydrofuran **2o** followed by reduction with DIBAL provided the desired product **4** in 45% total yield and high enantioselectivity (92%/97% ee). Although moderate diastereoselectivity (39/61 dr) was afforded, it

should be noted that the diastereomers could be isolated by flash chromatography.

The absolute configuration of the recovered **1a** was assigned to be (S) by comparison of the optical rotations with (R)-3-butyn-2-ol (98% ee) derived **1a**.¹⁸ It implied that the (R)-**1a** preferred to undergo [2,3]-Wittig rearrangement in the presence of $\text{L-PiPr}_3/\text{Ni}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ catalyst, which delivered (aR)-**3a** as the product. Based on this information and our previous studies,^{6c,19} a possible catalytic model was proposed. As depicted in Figure 1, the substrate **1o** undergoes




Figure 1. Proposed asymmetric induction models for **3o**.

deprotonation assisted by Et_3N to form the enolization intermediate, which coordinates tightly to the Lewis acid catalyst through the ether oxygen atom and the enolate ion from the auxiliary ester group. Due to the steric hindrance between the Bn group and amino acid skeleton unit, (S)-**1o** was unlikely to coordinate with the $\text{L-PiPr}_3/\text{Ni}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ catalyst (Figure 1, left); in contrast, the bonded (R)-**1o** was ready to undergo rearrangement (right), delivering (aR)-**3o** via center-chirality transfer to axial chirality.

In conclusion, we have successfully developed an asymmetric [2,3]-Wittig rearrangement via kinetic resolution of racemic propargyloxy dicarbonyl compounds. The current protocol provides an efficient and facile access to chiral α -hydroxyallenes with various substituents in high enantioselectivities. These functionalized α -hydroxyallenes could be easily converted to related 2,5-dihydrofuran derivatives. Further studies on other type of asymmetric [2,3]-Wittig rearrangement and its application are underway.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.0c00649>.

Experimental details; analytical data (NMR, HPLC, ESI-MS, IR, CD) (PDF)

Accession Codes

CCDC 1951577 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Authors

Xiaohua Liu – Key Laboratory of Green Chemistry & Technology, Ministry of Education, College of Chemistry,

Sichuan University, Chengdu 610064, China; orcid.org/0000-0001-9555-0555; Email: liuxh@scu.edu.cn

Xiaoming Feng – Key Laboratory of Green Chemistry & Technology, Ministry of Education, College of Chemistry, Sichuan University, Chengdu 610064, China; orcid.org/0000-0003-4507-0478; Email: xmfeng@scu.edu.cn

Authors

Xi Xu – Key Laboratory of Green Chemistry & Technology, Ministry of Education, College of Chemistry, Sichuan University, Chengdu 610064, China

Shunxi Dong – Key Laboratory of Green Chemistry & Technology, Ministry of Education, College of Chemistry, Sichuan University, Chengdu 610064, China; orcid.org/0000-0002-3018-3085

LiLi Feng – Key Laboratory of Green Chemistry & Technology, Ministry of Education, College of Chemistry, Sichuan University, Chengdu 610064, China

Sijing Wang – Key Laboratory of Green Chemistry & Technology, Ministry of Education, College of Chemistry, Sichuan University, Chengdu 610064, China

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acs.orglett.0c00649>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We appreciate the National Natural Science Foundation of China (U19A2014 and 21801175) for financial support.

REFERENCES

- (1) For selected reviews, see: (a) Hoffmann-Röder, A.; Krause, N. Synthesis and Properties of Allenic Natural Products and Pharmaceuticals. *Angew. Chem., Int. Ed.* **2004**, *43*, 1196–1216. (b) Ma, S. M. Some Typical Advances in the Synthetic Applications of Allenes. *Chem. Rev.* **2005**, *105*, 2829–2871. (c) Yu, S. H.; Ma, S. M. How easy are the syntheses of allenenes? *Chem. Commun.* **2011**, *47*, 5384–5418. (d) Neff, R. K.; Frantz, D. E. Recent Advances in the Catalytic Syntheses of Allenes: A Critical Assessment. *ACS Catal.* **2014**, *4*, 519–528. (e) Ye, J. T.; Ma, S. M. Conquering three-carbon axial chirality of allenenes. *Org. Chem. Front.* **2014**, *1*, 1210–1224. (f) Ye, J. T.; Ma, S. M. Palladium-Catalyzed Cyclization Reactions of Allenes in the Presence of Unsaturated Carbon–Carbon Bonds. *Acc. Chem. Res.* **2014**, *47*, 989–1000. (g) Chu, W.-D.; Zhang, Y.; Wang, J. B. Recent advances in catalytic asymmetric synthesis of allenenes. *Catal. Sci. Technol.* **2017**, *7*, 4570–4579.
- (2) Sapu, C. M.; Bäckvall, J.; Deska, J. Enantioselective Enzymatic Desymmetrization of Prochiral Allenic Diols. *Angew. Chem., Int. Ed.* **2011**, *50*, 9731–9734.
- (3) (a) Tap, A.; Blond, A.; Wakchaure, V. N.; List, B. Chiral Allenes via Alkynyllogous Mukaiyama Aldol Reaction. *Angew. Chem., Int. Ed.* **2016**, *55*, 8962–8965. (b) Wang, G.; Liu, X. H.; Chen, Y. S.; Yang, J.; Li, J.; Lin, L. L.; Feng, X. M. Diastereoselective and Enantioselective Alleno-aldol Reaction of Allenates with Isatins to Synthesis of Carbinol Allenates Catalyzed by Gold. *ACS Catal.* **2016**, *6*, 2482–2486. (c) Tang, Y.; Xu, J.; Yang, J.; Lin, L. L.; Feng, X. M.; Liu, X. H. Asymmetric Three-Component Reaction for the Synthesis of Tetrasubstituted Allenates via Allenate-Copper Intermediates. *Chem.* **2018**, *4*, 1658–1672. (d) Li, S.; Tang, Z. W.; Wang, Y.; Wang, D.; Wang, Z. L.; Yu, C. X.; Li, T. J.; Wei, D. H.; Yao, C. S. NHC-Catalyzed Aldol-Like Reactions of Allenates with Isatins: Regiospecific Syntheses of γ -Functionalized Allenates. *Org. Lett.* **2019**, *21*, 1306–1310.
- (4) Liao, Y.; Yin, X. M.; Wang, X. H.; Yu, W. Z.; Fang, D. M.; Hu, L. R.; Wang, M.; Liao, J. Enantioselective Synthesis of Multisubstituted Allenes by Cooperative Cu/Pd-Catalyzed 1,4-Arylboration of 1,3-Enynes. *Angew. Chem., Int. Ed.* **2020**, *59*, 1176–1180.
- (5) Jiang, Y.; Diagne, A. B.; Thomson, R. J.; Schaus, S. E. Enantioselective Synthesis of Allenes by Catalytic Traceless Petasis Reactions. *J. Am. Chem. Soc.* **2017**, *139*, 1998–2005.
- (6) (a) Moniz, G. A.; Wood, J. L. Catalyst-Based Control of [2,3]- and [3,3]-Rearrangement in α -Diazoketone-Derived Propargyloxy Enols. *J. Am. Chem. Soc.* **2001**, *123*, 5095–5097. (b) Li, Z. J.; Boyarskikh, V.; Hansen, J. H.; Autschbach, J.; Musaev, D. G.; Davies, H. M. L. Scope and Mechanistic Analysis of the Enantioselective Synthesis of Allenes by Rhodium-Catalyzed Tandem Ylide Formation/[2,3]-Sigmatropic Rearrangement between Donor/Acceptor Carbenoids and Propargylic Alcohols. *J. Am. Chem. Soc.* **2012**, *134*, 15497–15504. (c) Xu, X.; Zhang, J. L.; Dong, S. X.; Lin, L. L.; Lin, X. B.; Liu, X. H.; Feng, X. M. Nickel(II)-catalyzed asymmetric propargyl [2,3]-Wittig rearrangement of oxindole derivatives: A chiral amplification effect. *Angew. Chem., Int. Ed.* **2018**, *57*, 8734–8738.
- (7) For recent reviews on [2,3]-sigmatropic rearrangement, see: (a) Nakai, T.; Mikami, K. [2,3]-Wittig sigmatropic rearrangements in organic synthesis. *Chem. Rev.* **1986**, *86*, 885–902. (b) Doyle, M. P. Catalytic Methods for Metal Carbene Transformations. *Chem. Rev.* **1986**, *86*, 919–939. (c) Sweeney, J. B. Sigmatropic rearrangements of ‘onium’ ylides. *Chem. Soc. Rev.* **2009**, *38*, 1027–1038. (d) Moyano, A.; El-Hamdouni, N.; Atlamsani, A. Asymmetric Organocatalytic Rearrangement Reactions. *Chem. - Eur. J.* **2010**, *16*, 5260–5273. (e) Jones, A. C.; May, J. A.; Sarpong, R.; Stoltz, B. M. Toward a Symphony of Reactivity: Cascades Involving Catalysis and Sigmatropic Rearrangements. *Angew. Chem., Int. Ed.* **2014**, *53*, 2556–2591. (f) West, T. H.; Spoehrle, S. S. M.; Kasten, K.; Taylor, J. E.; Smith, A. D. Catalytic Stereoselective [2,3]-Rearrangement Reactions. *ACS Catal.* **2015**, *5*, 7446–7479.
- (8) For some examples of catalytic asymmetric [2,3]-Wittig rearrangement, see: (a) McNally, A.; Evans, B.; Gaunt, M. J. Organocatalytic Sigmatropic Reactions: Development of a [2,3]-Wittig Rearrangement through Secondary Amine Catalysis. *Angew. Chem., Int. Ed.* **2006**, *45*, 2116–2119. (b) Kitamura, M.; Hirokawa, Y.; Maezaki, N. Asymmetric [2,3]-Wittig Rearrangement of Oxygenated Allyl Benzyl Ethers in the Presence of a Chiral di-*t*Bu-bis(oxazoline) Ligand: A Novel Synthetic Approach to THF Lignans. *Chem. - Eur. J.* **2009**, *15*, 9911–9917. (c) Denmark, S. E.; Cullen, L. R. Development of a Phase-Transfer-Catalyzed, [2,3]-Wittig Rearrangement. *J. Org. Chem.* **2015**, *80*, 11818–11848. (d) Ošeka, M.; Kimm, M.; Kaabel, S.; Järving, I.; Rissanen, K.; Kanger, T. Asymmetric Organocatalytic Wittig [2,3]-Rearrangement of Oxindoles. *Org. Lett.* **2016**, *18*, 1358–1361. (e) Kennedy, C. R.; Guidera, J. A.; Jacobsen, E. N. Synergistic Ion-Binding Catalysis Demonstrated via an Enantioselective, Catalytic [2,3]-Wittig Rearrangement. *ACS Cent. Sci.* **2016**, *2*, 416–423. (f) Ošeka, M.; Kimm, M.; Järving, I.; Lippur, K.; Kanger, T. Two Catalytic Methods of an Asymmetric Wittig [2,3]-Rearrangement. *J. Org. Chem.* **2017**, *82*, 2889–2897. (g) Peňáška, T.; Mojzes, M. M.; Filo, J.; Jurdáková, H.; Mečiarová, M.; Šebesta, R. Organocatalysts Effect on the Stereoselectivity of [2,3]-Wittig Rearrangement. *Eur. J. Org. Chem.* **2019**, *2019*, 605–610. (h) Kimm, M.; Ošeka, M.; Kaabel, S.; Metsala, A.; Järving, I.; Kanger, T. [2,3]-Wittig Rearrangement as a Formal Asymmetric Alkylation of α -Branched Ketones. *Org. Lett.* **2019**, *21*, 4976–4980.
- (9) (a) Ma, M.; Peng, L. L.; Li, C. K.; Zhang, X.; Wang, J. B. Highly Stereoselective [2,3]-Sigmatropic Rearrangement of Sulfur Ylide Generated through Cu(I) Carbene and Sulfides. *J. Am. Chem. Soc.* **2005**, *127*, 15016–15017. (b) Zhang, Z. K.; Sheng, Z.; Yu, W. Z.; Wu, G. J.; Zhang, R.; Chu, W. D.; Zhang, Y.; Wang, J. B. Catalytic asymmetric trifluoromethylthiolation via enantioselective [2,3]-sigmatropic rearrangement of sulfonium ylides. *Nat. Chem.* **2017**, *9*, 970–976. (c) Rodríguez, R. I.; Ramírez, E.; Fernández-Salas, J. A.; Sánchez-Obregón, R.; Yuste, F.; Alemán, J. Asymmetric [2,3]-Wittig Rearrangement: Synthesis of Homoallylic, Allenylic, and Enynyl α -Benzyl Alcohols. *Org. Lett.* **2018**, *20*, 8047–8051.

- (10) (a) Dominguez, J.; Dunitz, J. D.; Gerlach, H.; Prelog, V. Stoffwechselprodukte von Actinomyceten. 32. Mitteilung. Über die Konstitution von Nonactin. *Helv. Chim. Acta* **1962**, *45*, 129–138. (b) Kobayashi, J.; Ishibashi, M.; Murayama, T.; Takamatsu, M.; Iwamura, M.; Ohizumi, Y.; Sasaki, T. Amphidinolide E, a Novel Antileukemic 19-Membered Macrolide from the Cultured Symbiotic Dinoflagellate Amphidinium sp. *J. Org. Chem.* **1990**, *55*, 3421–3423. (c) Bauer, I.; Maranda, L.; Young, K. A.; Shimizu, Y.; Fairchild, C.; Cornell, L.; MacBeth, J.; Huang, S. Isolation and Structure of Caribenolide I, a Highly Potent Antitumor Macrolide from a Cultured Free-Swimming Caribbean Dinoflagellate, Amphidinium sp. S1–36–5. *J. Org. Chem.* **1995**, *60*, 1084–1086. (d) Malmström, J.; Christophersen, C.; Barrero, A. F.; Oltra, J. E.; Justicia, J.; Rosales, A. Bioactive Metabolites from a Marine-Derived Strain of the Fungus *Emericella Variecolor*. *J. Nat. Prod.* **2002**, *65*, 364–367. (e) Jacques, R.; Pal, R.; Parker, N. A.; Sear, C. E.; Smith, P. W.; Ribaucourt, A.; Hodgson, D. M. Recent applications in natural product synthesis of dihydrofuran and -pyran formation by ring-closing alkene metathesis. *Org. Biomol. Chem.* **2016**, *14*, 5875–5893. (f) Shi, T. D.; Wijeratne, E. M. K.; Solano, C.; Ambrose, A. J.; Ross, A. B.; Norwood, C.; Orido, C. K.; Grigoryan, T.; Tillotson, J.; Kang, M.; Luo, G.; Keegan, B. M.; Hu, W. H.; Blagg, B. S. J.; Zhang, D. D.; Gunatilaka, A. A. L.; Chapman, E. An Isoform-Selective PTP1B Inhibitor Derived from Nitrogen-Atom Augmentation of Radicicol. *Biochemistry* **2019**, *58*, 3225–3231. (g) Shi, T. D.; Teng, S. H.; Reddy, A. G. K.; Guo, X.; Zhang, Y. T.; Moore, K. T.; Buckley, T.; Mason, D. J.; Wang, W.; Chapman, E.; Hu, W. H. Catalytic asymmetric synthesis of 2,5-dihydrofurans using synergistic bifunctional Ag catalysis. *Org. Biomol. Chem.* **2019**, *17*, 8737–8744.
- (11) (a) Evidente, A.; Andolfi, A.; Fiore, M.; Spanu, E.; Maddau, L.; Franceschini, A.; Marras, F.; Motta, A. Diplobifuranylones A and B, 5'-Monosubstituted Tetrahydro-2H-bifuranyl-5-ones Produced by *Diplodia corticola*, a Fungus Pathogen of Cork Oak. *J. Nat. Prod.* **2006**, *69*, 671–674. (b) Cheng, X. P.; Quintanilla, C. D.; Zhang, L. M. Total Synthesis and Structure Revision of Diplobifuranylon B. *J. Org. Chem.* **2019**, *84*, 11054–11060.
- (12) (a) Katagiri, K.; Tori, K.; Kimura, Y.; Yoshida, T.; Nagasaki, T.; Minato, H. A New Antibiotic. Furanomycin, an Isoleucine Antagonist. *J. Med. Chem.* **1967**, *10*, 1149–1154. (b) Joullie, M. M.; Wang, P. C.; Semple, J. E. Total Synthesis and Revised Structural Assignment of (+)-furanomycin. *J. Am. Chem. Soc.* **1980**, *102*, 887–889.
- (13) (a) Takahashi, K.; Yasue, M.; Ogiyama, K. A Norlignan, Cryptoresinol, from the Heartwood of *Cryptomeria Japonica*. *Phytochemistry* **1988**, *27*, 1550–1552. (b) Ohashi, H.; Kawai, S.; Sakurai, Y.; Yasue, M. Norlignan from the Knot Resin of *Araucaria Angustifolia*. *Phytochemistry* **1992**, *31*, 1371–1373. (c) Yoshida, K.; Nishiguchi, M.; Hishiyama, S.; Kato, A.; Takahashi, K. Generation and alteration of norlignans in a transition zone during the drying of a *Cryptomeria japonica* log. *J. Wood Sci.* **2006**, *52*, 372–375.
- (14) For selected examples of the metal-catalyzed cyclization of α -allenyl alcohols, see: (a) Olsson, L.-I.; Claesson, A. Synthesis of 2,5-Dihydrofurans and 5,6-Dihydro-2H-pyrans by Silver(I)-Catalyzed Cyclization of Allenic Alcohols. *Synthesis* **1979**, 1979, 743–745. (b) Hoffmann-Röder, A.; Krause, N. The golden gate to catalysis. *Org. Biomol. Chem.* **2005**, *3*, 387–391. (c) Alcaide, B.; Almendros, P.; Luna, A.; Soriano, E. An Alternative to Precious Metals: $\text{Hg}(\text{ClO}_4)_2 \cdot 3\text{H}_2\text{O}$ as a Cheap and Water-Tolerant Catalyst for the Cycloisomerization of Allenols. *J. Org. Chem.* **2015**, *80*, 7050–7057. (d) El-Sepelgy, O.; Brzozowska, A.; Sklyaruk, J.; Jang, Y. K.; Zubar, V.; Rueping, M. Cooperative Metal–Ligand Catalyzed Intramolecular Hydroamination and Hydroalkoxylation of Allenes Using a Stable Iron Catalyst. *Org. Lett.* **2018**, *20*, 696–699.
- (15) For reviews of N,N' -dioxide ligands, see: (a) Liu, X. H.; Lin, L. L.; Feng, X. M. Chiral N,N' -Dioxides: New Ligands and Organocatalysts for Catalytic Asymmetric Reactions. *Acc. Chem. Res.* **2011**, *44*, 574–587. (b) Liu, X. H.; Lin, L. L.; Feng, X. M. Chiral N,N' -dioxide ligands: synthesis, coordination chemistry and asymmetric catalysis. *Org. Chem. Front.* **2014**, *1*, 298–302. (c) Liu, X. H.; Zheng, H. F.; Xia, Y.; Lin, L. L.; Feng, X. M. Asymmetric Cycloaddition and Cyclization Reactions Catalyzed by Chiral N,N' -Dioxide-Metal Complexes. *Acc. Chem. Res.* **2017**, *50*, 2621–2631.
- (16) Lin, X. B.; Tang, Y.; Yang, W.; Tan, F.; Lin, L. L.; Liu, X. H.; Feng, X. M. Chiral Nickel(II) Complex Catalyzed Enantioselective Doyle–Kirmse Reaction of α -Diazo Pyrazoleamides. *J. Am. Chem. Soc.* **2018**, *140*, 3299–3305.
- (17) Preliminary studies indicated that **2a** was obtained from starting material **1a** rather than rearrangement product **3a**. Moreover, with **L-PiCHPh₂** as the ligand, (**S**)-**1a** preferably transformed into **2a**; in contrast, (**R**)-**1a** preferably rearranged into (**aR**)-**3a** (see SI pp S8 and S17 for details). Currently, the exact pathway for the formation of **2a** is not clear; two possible routes were proposed, including conia-ene and carbanion nucleophilic attack to alkyne. For selected examples of carbanion nucleophilic attack to alkyne, see: (a) Gao, K.; Wu, J. Sc(OTf)₃-Catalyzed or *t*-BuOK Promoted Tandem Reaction of 2-(2-(Alkynyl)benzylidene)malonate with Indole. *Org. Lett.* **2008**, *10*, 2251–2254. (b) Chaudhari, T. Y.; Ginotra, S. K.; Tandon, V. Facile access to functionalized indenenes and fused quinolines by regioselective 5-enolexo-dig Michael addition and cyclization reactions. *Org. Biomol. Chem.* **2017**, *15*, 9319–9330. For selected examples of conia-ene reaction, see: (c) Matsuzawa, A.; Mashiko, T.; Kumagai, N.; Shibasaki, M. La/Ag Heterobimetallic Cooperative Catalysis: A Catalytic Asymmetric Conia-Ene Reaction. *Angew. Chem., Int. Ed.* **2011**, *50*, 7616–7619. (d) Cao, M.; Yesilcimen, A.; Wasa, M. Enantioselective Conia-Ene-Type Cyclizations of Alkynyl Ketones through Cooperative Action of $\text{B}(\text{C}_6\text{F}_5)_3$, *N*-Alkylamine and a Zn-Based Catalyst. *J. Am. Chem. Soc.* **2019**, *141*, 4199–4203.
- (18) The procedure for the synthesis of (+)-3-butyn-2-ol (98% ee) derived **1a**; see SI for details.
- (19) Liu, Y. B.; Liu, X. H.; Hu, H. P.; Guo, J.; Xia, Y.; Lin, L. L.; Feng, X. M. Synergistic Kinetic Resolution and Asymmetric Propargyl Claisen Rearrangement for the Synthesis of Chiral Allenes. *Angew. Chem., Int. Ed.* **2016**, *55*, 4054–4058.