

Survival Analysis Homework5

Zhuodiao Kuang

2023-10-20

Problem1

Construct 95% CI for the hazard ratio from a PH model shown in the follow table for the risk reduction between two treatment groups.

Analysis of Maximum Likelihood Estimates - discrete								
Parameter		DF	Parameter Estimate	Standard Error	Chi-Square	Pr > ChiSq	Hazard Ratio	Label
trt	6-MP	1	-1.62822	0.43313	14.1316	0.0002	0.196	trt 6-MP

According to the table, the parameter estimate and the standard error are -1.628 and 0.433, respectively.

We've already learned that the 95% CI confidence interval for the coefficient estimates is as follows:

$$[\hat{\beta} - z_{0.975}se(\hat{\beta}), \hat{\beta} + z_{0.975}se(\hat{\beta})]$$

Then, from $\frac{h(t|Z=1)}{h(t|Z=0)} = e^{\beta}$, we can obtain the hazard ratio as 0.196, with its 95% CI:

$$[e^{\hat{\beta} - z_{0.975}se(\hat{\beta})}, e^{\hat{\beta} + z_{0.975}se(\hat{\beta})}]$$

```
est <- -1.628
se <- 0.433
z <- qnorm(0.975)
lq <- exp(est-z*se)
uq <- exp(est+z*se)
lq;uq
```

```
## [1] 0.08402303
```

```
## [1] 0.4587107
```

So, the 95% CI for the hazard ratio is [0.084, 0.459].

Problem2

The observed survival data (T_i, Δ_i, Z_i) $i = 1, 2, 3, 4, 5, 6$ are $(16, 1, 1)$, $(20, 0, 1)$, $(12, 1, 0)$, $(14, 0, 0)$, $(11, 1, 0)$, $(9, 1, 1)$. Please construct the partial likelihood.

$$L_p(\beta) = \prod_{i=1}^n \left\{ \frac{e^{\beta z_i}}{\sum_{l \in R(t_i)} e^{\beta' z_l}} \right\}^{\Delta_i}$$
$$\therefore \Delta_1 = \Delta_3 = \Delta_5 = \Delta_6 = 1$$
$$\therefore \Delta_2 = \Delta_4 = 0, n = 6$$

$$\therefore L_p(\beta) = \frac{e^{\beta}}{e^{\beta} + e^{\beta}} \frac{1}{1 + 1 + e^{\beta} + e^{\beta}} \frac{1}{1 + 2 + 2e^{\beta}} \frac{e^{\beta}}{e^{\beta} + 3 + 2e^{\beta}}$$
$$= \frac{1}{2} \frac{1}{2 + 2e^{\beta}} \frac{1}{3 + 2e^{\beta}} \frac{e^{\beta}}{3 + 3e^{\beta}}$$

Problem3

Show that PH model score test is the same as the log-rank test for an indicator covariate when there is no ties.