

## 泰勒公式

局部泰勒公式:  $x \rightarrow x_0$  时  $f(x) = f(x_0) + f'(x_0)(x-x_0) + \frac{f''(x_0)(x-x_0)^2}{2} + \dots + \frac{f^{(n)}(x_0)(x-x_0)^n}{n!} + o((x-x_0)^n)$  (佩亚诺余项)

证明: 令  $T_n(x) = f(x_0) + f'(x_0)(x-x_0) + \dots + \frac{f^{(n)}(x_0)(x-x_0)^n}{n!}$

$\lim_{x \rightarrow x_0} \frac{f(x) - T_n(x)}{(x-x_0)^n} = \lim_{x \rightarrow x_0} \frac{f^{(n)}(x) - f^{(n)}(x_0)}{n!} = 0$  证等式等同于证差为0或无穷小量 借分母低阶无穷小量凑出必然的结构

证毕

麦克劳林公式:  $x_0 = 0$  时  $f(x) = f(0) + f'(0)x + \dots + \frac{f^{(n)}(0)x^n}{n!} + o(x^n)$

常用麦克劳林公式:  $e^x = 1 + x + \dots + \frac{1}{n!}x^n + o(x^n)$

$\ln(1+x) = x - \frac{1}{2}x^2 + \dots + (-1)^{n-1} \frac{x^n}{n} + o(x^n)$

$(1+x)^\alpha = 1 + \alpha x + \frac{\alpha(\alpha-1)}{2}x^2 + \dots + \frac{\alpha!}{(\alpha-n)!n!}x^n + o(x^n)$

$\sin x = x - \frac{1}{6}x^3 + \dots + (-1)^{n-1} \frac{x^{2n-1}}{(2n-1)!} + \begin{cases} o(x^{2n-1}) \\ 0 + o(x^{2n}) \end{cases}$

$\cos x = 1 - \frac{1}{2}x^2 + \dots + (-1)^n \frac{x^{2n}}{(2n)!} + \begin{cases} o(x^{2n}) \\ 0 + o(x^{2n+1}) \end{cases}$

局部泰勒展开的唯一性定理: 若  $f(x) = A_0 + A_1(x-x_0) + A_2(x-x_0)^2 + \dots + A_n(x-x_0)^n + o((x-x_0)^n)$ ,  $x \rightarrow x_0$

则有  $A_k = \frac{f^{(k)}(x_0)}{k!}$

也就是说用多项式去逼近某个函数, 局部的泰勒展开公式只有一种

证明: 显然  $A_0 = f(x_0)$  即证  $\frac{f(x) - f(x_0)}{x - x_0} = A_1 + A_2(x-x_0) + \dots + A_n(x-x_0)^{n-1} + o((x-x_0)^{n-1})$ ,  $x \rightarrow x_0$

$\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} = \lim_{x \rightarrow x_0} f'(x)$

$= f'(x_0) = A_1$

$= \lim_{x \rightarrow x_0} (\text{右式})$

同理  $f^{(k)}(x_0) = k! A_k$

证毕

求  $e^x$  的麦克劳林展开式

令  $t = -x^2$   $e^t = \dots$

$= \dots$

求  $e^{\cos x}$  的麦克劳林展开式

保证  $x \rightarrow 0$  时,  $t \rightarrow 0$ , 即最后一项为无穷小量

带拉格朗日余项的泰勒展开式:  $f(x) = f(x_0) + \dots + \frac{f^{(n)}(x_0)(x-x_0)^n}{n!} + \frac{f^{(n+1)}(\xi)(x-x_0)^{n+1}}{(n+1)!}$

证明: 令  $F(t) = f(t) - f(x_0) - \dots - \frac{f^{(n)}(x_0)(t-x_0)^n}{n!}$

$G(t) = (t-x_0)^{n+1}$

显然  $F(x_0) = 0$ ,  $G(x_0) = 0$

$\frac{F(x) - F(x_0)}{G(x) - G(x_0)} = \frac{F'(x_1) - F'(x_0)}{G'(x_1) - G'(x_0)} = \frac{F^{(n+1)}(\xi)}{G^{(n+1)}(\xi)}$

即  $F(x) = \frac{f^{(n+1)}(\xi)}{(n+1)!} (x-x_0)^{n+1}$

证毕

常见函数的拉格朗日余项:  $x \rightarrow 0$

$$e^x \sim \frac{e^{\xi}}{(n+1)!} x^{n+1}$$

$$\ln(1+x) \sim \frac{1}{(n+1)} \cdot \frac{1}{(1+\xi)^{n+1}} x^{n+1}$$

$$(1+x)^\alpha \sim \frac{\alpha!}{(\alpha-n-1)! (n+1)!} x^{n+1}$$

$$\sin x \sim (-1)^n \frac{\cos \xi}{(2n+1)!} x^{2n+1}$$

$$\cos x \sim (-1)^{n-1} \frac{\sin \xi}{(2n+2)!} x^{2n+2}$$

常用求极限方法

1° 选择合适换元方法

2° 提前提出确定极限

3° 无穷小量代换

4° 泰勒展开至适当级数

5° 洛必达法则

It's all one ghetto, man, giant gutter in outer space.

It's all one ghetto, man, giant gutter in outer space.

It's all one ghetto, man, giant gutter in outer space.