

EXERCISE CLASS : Linear regression

Exercice 1 (MCO et re-centrage).

- 1) Quels sont les vecteurs $\mathbf{y} \in \mathbb{R}^n$ tels que $\text{var}_n(\mathbf{y}) = 0$ (var_n est la variance empirique) ?
- 2) Pour une matrice $X \in \mathbb{R}^{n \times p}$, que vaut $\text{Ker}(X^\top X)$?
- 3) Pour une matrice $X \in \mathbb{R}^{n \times (p+1)}$, $n > 1$ et $p \geq 1$, qui possède comme première colonne une colonne de 1, notons $(1, \tilde{x}_i^\top) \in \mathbb{R}^{p+1}$, les lignes de X . Montrer que $X^\top X$ non-inversible est équivalent à

$$\text{cov}_n(\tilde{X}) = n^{-1} \sum_{i=1}^n (\tilde{x}_i - \hat{\mu}_n)(\tilde{x}_i - \hat{\mu}_n)^\top \quad \text{non-inversible,}$$

$$\text{où } \hat{\mu}_n = n^{-1} \sum_{i=1}^n \tilde{x}_i.$$

- 4) Supposons que $\text{cov}_n(\tilde{X})$ soit inversible. Exprimer l'estimateur des moindres carrés calculé sur des variables centrées.

Exercice 2 (MCO et invariance).

On suppose que X est de rang plein et on note $\hat{\theta}_n$ l'estimateur OLS. On note $\tilde{X} = (\tilde{X}_1, \dots, \tilde{X}_p)$. On change l'échelle d'une des variables : \tilde{X}_k est remplacé par $\tilde{X}_k b$, où $b > 0$.

- 1) Soit $X_b = (1, \tilde{X}_1, \dots, \tilde{X}_k b, \dots, \tilde{X}_p)$. Montrer que $X_b = XD$ où D est une matrice diagonale que l'on précisera.
- 2) Soit $\hat{\theta}_{b,n}$ l'estimateur OLS associé à X_b . Exprimer $\hat{\theta}_{b,n}$ en fonction de $\hat{\theta}_n$ et D .
- 3) Donner la variance de $\hat{\theta}_{b,n}$.
- 4) On a vu que l'estimateur $\hat{\theta}_n$ était affecté par un changement d'échelle. Qu'en est-il de la valeur prédite par le modèle ?
- 5) Peut-on dire la même chose de l'estimateur Ridge ?

Exercice 3 (prediction intervals). For $i = 1, \dots, n$, we consider $y_i \in \mathbb{R}$ and $x_i = (x_{i,0}, \dots, x_{i,p})^\top \in \mathbb{R}^{p+1}$ with $x_{i,0} = 1$. The OLS estimator is any coefficient vector $\hat{\theta}_n = (\hat{\theta}_{n,0}, \dots, \hat{\theta}_{n,p})^\top \in \mathbb{R}^{p+1}$ such that

$$\hat{\theta}_n \in \arg \min_{\theta \in \mathbb{R}^{p+1}} \sum_{i=1}^n (y_i - x_i^\top \theta)^2.$$

With the notations

$$X = \begin{pmatrix} x_1^\top \\ \vdots \\ x_n^\top \end{pmatrix} = \begin{pmatrix} x_{1,0} & \dots & x_{1,p} \\ \vdots & & \vdots \\ x_{n,0} & \dots & x_{n,p} \end{pmatrix} \in \mathbb{R}^{n \times (p+1)}, \quad Y = \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix}.$$

We have

$$\hat{\theta}_n \in \arg \min_{\theta \in \mathbb{R}^{p+1}} \|Y - X\theta\|.$$

We assume the following Gaussian model, for all $i = 1, \dots, n$, $y_i = x_i^\top \theta^* + \epsilon_i$ with $(\epsilon_i) \sim_{iid} \mathcal{N}(0, \sigma^2)$ such that $\ker(X) = \{0\}$.

- 1) Let $x = (1, \tilde{x}^\top)^\top$ with $\tilde{x} \in \mathbb{R}^p$. Give $\hat{p}(x)$ the predicted value at x by the OLS.
- 2) Give the distribution of $\hat{p}(x)$. The mean $p(x)$ and variance $u(x)$ should be made explicit.

3) Define

$$\hat{\sigma}_n^2 = \frac{1}{n - (p + 1)} \sum_{i=1}^n (y_i - x_i^T \hat{\theta}_n)^2,$$

and recall that $\hat{\sigma}_n^2(n - (p + 1))$ is independent from $\hat{\theta}_n$ and follows a chi-squared distribution with $n - (p + 1)$ degrees of freedom. Show that

$$\frac{(\hat{p}(x) - p(x))}{\hat{\sigma}_n \sqrt{x^T (X^T X)^{-1} x}} \sim t(n - (p + 1)),$$

where $t(k)$ is the Student distribution with k degrees of freedom (hint $\mathcal{N}(0, 1)/\sqrt{\chi_k^2/k} \simeq t(k)$ if $\mathcal{N}(0, 1) \perp \chi_k^2$).

4) Let y be the output associated to the predictor x . The value y is supposed to be independent from the sample (y_i) . Show that

$$\frac{y - \hat{p}(x)}{\hat{\sigma}_n \sqrt{1 + x^T (X^T X)^{-1} x}} \sim t(n - (p + 1)).$$

5) Build confidence intervals for $p(x)$ and Y . The last one is often called prediction interval.

Exercise 4 (ridge). Under the same setting as the previous exercise

- 1) Show that the ridge is unique and that $\hat{\theta}_n^{(rdg)} = (X^T X + \lambda I_p)^{-1} X^T Y$.
- 2) Give the bias and the variance of the Ridge.
- 3) In the Gaussian regression model, show that $\hat{\theta}_n^{(rdg)}$ is distributed according to a normal distribution of which the mean and variance shall be specified.
- 4) Let $k \in \{1, \dots, p\}$ and $\alpha \in (0, 1/2)$. Assuming that the variance of the noise is $\sigma^2 = 1$, give a confidence interval for $\hat{\theta}_{n,k}^{(rdg)}$ with level α .