Hordeolum, Chalazia, and Cysts

Lorne Yudcovitch, OD, MS, FAAO

1

Hordeolum

- External hordeolum
  - Infection of a Zeiss gland
- Internal hordeolum
  - Infection of a meibomian gland
- *Staphylococcus aureus* is the infectious agent in 90-95% of cases of hordeolum
- Abscess ("Bag of pus")

2

Hordeolum

Subjective

- Patients usually complain of a localized painful swelling on one eyelid
- In some cases, the complaint may start as a generalized edema and erythema of the lid that later becomes localized.
- A history of similar problems is common.
- Constitutional signs and symptoms are inconsistent with a hordeolum diagnosis

3

Hordeolum

Epidemiology:


Found more frequently in persons who have

- **Diabetes**
- **Other debilitating illness**
- **Chronic blepharitis**
- **Seborrhea**
- **High serum lipids (High lipid levels increase the blockage rate of sebaceous glands, but lowering of serum lipid levels in these patients has not decreased frequency of recurrence)**

4

External Hordeolum

- **Localized, raised, red bump near lid margin**
- **Tender to touch**
- **"Points" to skin side of lid**



5


External Hordeolum



6

External Hordeolum

Hordeolum associated with staphylococcal blepharitis




7

External Hordeolum

- Localized infection





8


External Hordeolum

- No treatment; usually are self-limited
  - Most eventually point and drain by themselves
  - Discharge anteriorly, close to the lashes
- Warm soaks (qid for 15 min) are the mainstays of treatment
- Topical antibiotics may be used for recurrent lesions and for those that are actively draining
- Epilation of eyelash associated with infected follicle
  - May enhance drainage of pus

9

Internal Hordeolum

- Localized, red, raised bump
- Tender to the touch
- Points to the conjunctival side of the lid



10


Internal Hordeolum




11

Internal Hordeolum











12

Internal Hordeolum








13

Internal Hordeolum

- Large internal hordeolum discharging anteriorly through skin



- Infection can spread from the gland to the lid connective tissue → preseptal cellulitis



14

Internal Hordeolum

- Usually are self-limited
- Most eventually point and drain by themselves.
- Warm soaks (qid for 15 min) are the mainstays of treatment.
- Topical antibiotics may be used for recurrent lesions and for those that are actively draining
- Surgical drainage of pointed lesions can speed the healing process
- Systemic antibiotics are indicated only when inflammation has spread beyond the immediate area of the hordeolum

15

Internal Hordeolum


Drainage of a hordeolum

- Stab incision at the site of pointing using an 18-gauge needle or a #11 blade
- External incisions lead to scarring, so making external eyelid incisions or punctures is inadvisable, unless the hordeolum already is pointing externally
- A large abscess may have multiple pockets and require multiple stabs.
- Internal incisions should be made vertically to minimize the area of cornea swept by a scar during blinking
- Leave the incision open with a clean margin

16


Chalazion

- AKA meibomian cyst
- Lipogranulomatous infiltration of meibomian gland
- May follow an internal hordeolum
- Signs
  - Firm nodule within tarsal plate
  - Non-tender, non-inflamed
- Some refer to a internal hordeolum as an acute chalazion



17

Chalazion




18

Chalazia




19

Chalazion



20


Chalazion





21

Chalazion



22

Chalazion



23

Chalazion

- Warm compresses
  - 10-15 minutes
  - At least qid but q2h better
  - Work best if lesion is recent
- Steroid injection into lesion
  - 0.1-0.2 ml triamcinolone injected through conjunctiva
- Incision and curettage
  - Local anesthetic
  - Chalazion clamp
  - Incision from conjunctival side
  - Curettage - scrape out material
  - Topical antibiotic ointment - X 2 days





24


Incision of Chalazion



25

Cyst of Moll


- AKA suderiferous cyst or apocrine hidrocystoma
- Arises from a blocked sweat-secreting gland associated with the cilia (gland of Moll)



26

Cyst of Moll

- Dome shaped
- Non-tender
- No inflammation
- Translucent
- Fluid-filled
- Anterior lid margin



27

Cyst of Moll





Image courtesy of Shaina Hewitt, PUCO Class of 2019

28

Cyst of Moll




29

Cyst of Moll

Treatment

- **Primarily a cosmetic concern**
- **Can aspirate with a fine needle but recurrence is common**
- **Excision with removal of the sac to avoid recurrence**




30

Cyst of Zeis

- Arises from a blocked oil-secreting gland associated with the cilia (gland of Zeis)
- Contains oily secretions and is less translucent than a cyst of Moll

31


Cyst of Zeis



32


Sebaceous Cyst

- Forms when an oil gland in the skin is blocked
- Can be found on any part of the body
- Yellowish-white lumps on the skin
  - Have a central punctum
- If squeezed, exude a stringy, cheesy white material




33


Sebaceous Cyst



34

Sebaceous Cyst








35

Differentiate sebaceous cyst from xanthelasma (aka xanthoma)…



Sebaceous cyst



Xanthelasma

36

Sebaceous Cyst

- Primarily a cosmetic problem
- The total sac of the cyst must be removed
- If the cheesy material is just squeezed out, the cyst will reform
- In the case of an infected cyst, an oral antibiotic should be prescribed

37

Sebaceous Cyst

- Good hygiene may prevent the formation of sebaceous cysts
  - Some individuals are predisposed to development of these cysts, so even careful hygiene does not prevent them

38

THANK YOU!

39