Allergic Lid Disease

Lorne Yudcovitch, O.D., M.S., F.A.A.O.


1

ALLERGIC DISEASE BACKGROUND

2

Types of allergic reactions


- Type I - immediate type reaction:
- Reaction resulting from the release of pharmacologically active substances such as histamine, or eosinophillic chemotactic factor (ecf) from IgE-sensitized basophils and mast cells after contact with specific antigens
- Occurs within minutes to hours



3

Types of allergic reactions

- Type IV - delayed type reaction:
- Reactions caused by sensitized lymphocytes (T-cells) after contact with antigens which result from direct cytotoxicity or from the release of lymphokines
- Cell-mediated reaction
- Occurs over hours to days



4

A key allergic clinical finding…

- **Dennie–Morgan fold (aka Dennie-Morgan line)**
- A fold or line in the skin below the lower eyelid caused by edema in allergic dermatitis





5

CONTACT DERMATITIS

6

Contact Dermatitis








7

CONTACT DERMATITIS



- Very common immunological disease
- Results from direct contact with environmental allergen
- Dermatitis = any inflammation of the epidermis
- Contact dermatitis = any skin inflammation that occurs when the skin surface comes in contact with a substance originating outside the body
- Episodes can be triggered by specific events

8

Irritant Contact Dermatitis




- More common
- Contact with substance that directly damages skin
- Nonspecific response of skin to chemical damage that releases mediators of inflammation from epidermal cells
- Occurs within 1 – 24 hours; can occur within minutes

9


Irritant Contact Dermatitis


- Hx is very important!
- Red rash with pain, burning, stinging, discomfort; maybe no itching
- Often affects hands
- Occupations: cleaning, hospital care, food prep, hairdressing
- Women > men
- Common irritants
- Soap and water, solvents, microtrauma, dry air, cleaners



10

Irritant Contact Dermatitis



11

Irritant Contact Dermatitis

- Erythema
- Edema
- Glazed, parched, scalded appearance
- Skin usually dry with scaling
- No diagnostic test
  - Patch test negative
- Dx based on exclusion of other cutaneous diseases and clinical appearance of site exposed to a known irritant



12

Irritant Contact Dermatitis





13

Irritant Contact Dermatitis

TX/Management





- Avoid irritant via barriers
- Non-sensitizing moisturizers
  - Plain petroleum jelly
- Mild skin cleansers
  - Aquanil, Cetaphil cleanser, Neutrogena cleanser
- Soaps with low irritant effect
  - Dove, Cetaphil
- Alcohol based hand cleansers with moisturizer
- Avoid inappropriate solvents or abrasives to clean hands
- No topical steroids!!!

14

ICD Management










15

Allergic Contact Dermatitis:

- Occurs in sensitized individuals
- Involves a cell mediated immunity response
- Requires a latency period
- The re-exposure precipitates the dermatitis
  - Response is usually in 48 hrs

Subjective

- Hx is very important!
- Red rash with ITCHING



16

Allergic Contact Dermatitis:

List of possible allergens is endless

- Plants (poison oak*, ivy)
- Metals (nickel*, gold)
- Hair products
- Skin care products
- Rubber/latex
- Dyes
- Chromates for leather tanning
- Fragrances
- Preservatives
- Acrylates
- Cosmetics







17

Allergic Contact Dermatitis to Nickel





18

Allergic Contact Dermatitis to Hair Spray



19

Allergic Contact Dermatitis to Cosmetics



20

Examples of medications that may *cause* Allergic Contact Dermatitis

Topical ocular drugs

- Antazoline
- Neomycin
- Phenylephrine
- Atropine
- Pilocarpine
- Anti-glaucoma drugs





21

Allergic Contact Dermatitis due to Topical Ocular Drug



22

ACD due to
Topical Neomycin



23

ACD to Topical Ocular Atropine



24

ACD to Topical Ocular Mydriatic



25

Allergic Contact Dermatitis

Objective

- Pruritis (= itch!)
- Erythema
- Edema
- Chemosis of conjunctiva
- Vesicles
- Weeping rash → crusting, scaling → darkened, cracked (fissured), leathery appearance
- Keratitis may develop



26

Allergic Contact Dermatitis to Contact Lens Solution




27

Conjunctival Chemosis



28

Conjunctival Chemosis




29

Allergic Contact Dermatitis

DX


- Elicited from history
- Patch skin tests
  - Application of suspected materials to skin on patches
  - Left on 48 hours, then read again between 72 hrs – 1 wk
  - Positive: development of erythema, papules, edema, vesicles
  - Correlation with history, location of eruption, clinical course to determine if true etiology
- DDX from many similar appearing rashes



Skin Allergy Test


30

Conjunctival Chemosis



31

ACD





32

Allergic Contact Dermatitis

TX/Management

- Remove allergen
- Cold compresses 4 – 6X/day
- Topical steroid ung or cream BID – TID for 5 – 10 days
  - < 1% hydrocortisone
- Oral antihistamine
  - Diphenhydramine (Benadryl)
- Ocular lubricants 4 – 6X/day for comfort or keratitis
- Topical immunomodulator BID
  - Pimecromlimus (Elidel cream)
  - Tacrolimus (Protopic)





33

ACD Management









34

ATOPIC DERMATITIS

- Eczematous skin eruption AKA eczema
- Onset in infancy: most common skin disease in children
- Inherited condition
  - Usually in conjunction with other atopic conditions
- INFLAMED PATCHES OF VERY ITCHY SKIN



35

Atopic Dermatitis




36

Atopic Dermatitis

Clinical manifestations

- Lesions vary with age



Early infancy:

- Rash that is exudative, vesicular to maculopapular, see fine scales
- Location often on face (cheeks common), scalp, knees in infants
- ITCHING

37


Atopic Dermatitis



38

Atopic Dermatitis





39

Atopic Dermatitis

Later stages (as pt. ages):



- Dermatitis
- Skin becomes dry and lichenified
- Excoriated areas (abraded)
- Pruritus→ ITCHING
- Healing - hypo or hyper-pigmentation
- Location most often seen on elbows, neck, back of knees, ankles in adults

40

Atopic Dermatitis



41

Atopic Dermatitis
Eczematous flakes



42


Atopic Dermatitis



43

Atopic Dermatitis








44

Atopic Dermatitis

- Patient is more prone to secondary infection due to Staphylococcus or Streptococcus
- Unusual susceptibility to vaccinia, herpes simplex and molluscum contagiosum viruses
- Possible ocular manifestations of atopic dermatitis
  - Primarily lids
- Clinical Course unpredictable, but usually subsides by age 3 or 4 with exacerbations and remissions frequently recurring throughout childhood, adolescence and adulthood

45

Atopic Dermatitis



- ITCHY rash
- Erythema
- Edema
- Exudative lesions
- Xerosis
- Lichenfication
- Excoriation and crusting
- DX - tests not needed if condition presents with clear clinical markers
  - Patch test

46

Patch Test





47

Atopic Dermatitis TX/Management





- Cool damp compresses x 15-30 min or short, cool shower/bath
- White petrolatum moisturizer
- Topical steroid
  - 1% hydrocortisone ung BID – TID
- Oral antihistamine
  - Diphenhydramine (Benadryl)
- Topical antibiotic if lesions secondarily infected
- Topical immunomodulator
  - Pimecromlimus (Elidel cream)
  - Tacrolimus (Protopic)
- Nonmedical considerations:
  - Clothing
  - Cool temperature
  - Humidifier



48

AD Management



49

URTICARIA AND ANGIOEDEMA (Hives and Giant Urticaria)

50

Urticaria


- Urticaria = ITCHY WELTS = HIVES:
- Cutaneous eruption
- Multiple pathogenic mechanisms
- May or may not be immunologic in origin
- Only 70% have a specific identifiable cause
- Acute urticaria -- immunologic
- Chronic urticaria -- unknown



Urticaria

51

Urticaria



52

Urticaria

- 10 to 25% of the population
- Intense pruritus
- Raised wheals surrounded by erythematous flare
  - Size 1 to 5 mm to greater than 20 mm
  - Dilation and increased permeability of blood vessels in superficial dermis with edema in surrounding epidermis
  - May have a ring shape
- May occur on any part of the body
- May be transient in location

53


Urticaria





54

Dermatographic Urticaria
aka Dermatographism aka “skin writing”





55


ANGIOEDEMA: WHEN THE INFLAMMATION INVOLVES DEEPER LAYERS OF SKIN

- Skin appears normal
- Dilation and increased permeability of blood vessels in the deep dermis with resulting edema of dermis
- More extensive edematous areas
- Little pruritus - sometimes described as painful or burning
- Most often involves face, lips, eyelid, tongue, throat, hands, feet
- May produce respiratory distress due to upper airway edema



56

Angioedema



57

Angioedema



58

URTICARIA AND ANGIOEDEMA CAUSES

- Drug reactions
- Foods, food additives
- Inhalant, ingested or contact antigens
- Infections
- Physical
  - Cold
  - Heat
  - Exercise – induced
  - Solar
  - Water
- Insect bites
- Psychogenic
- Systemic disease





59

Angioedema



60

URTICARIA AND ANGIOEDEMA

DDX:

- Need to be concerned that the edema is not caused by renal or cardiac disease
- Contact dermatitis



TX:

- Try to find the cause and avoid
- Acute phase:
  - Cold compresses
  - Antihistamine
  - Systemic steroids - prednisone unpredictable
  - Epinephrine sc for acute pharyngeal or laryngeal angioedema

61

THANK YOU!

62