

AROUND SEGAL CONJECTURE IN p -ADIC GEOMETRY

ZHOUEHANG MAO

ABSTRACT. This article records multiple results coming from interplay between de-completed topological periodic cyclic homology, Segal conjecture, and F -smoothness.

We establish completeness of motivic filtration on de-completed topological periodic cyclic homology of commutative rings with weakly finitely generated absolute cotangent complex. When the ring in question is in addition F -smooth, we show that Segal conjecture holds for its topological Hochschild homology. We also identify our de-completed topological periodic cyclic homology with Manam’s Frobenius untwisted topological periodic cyclic homology for quasiregular semiperfectoid rings.

We find a crystalline degeneration of Segal conjecture which corresponds to such a statement for F -smoothness. On the other hand, inspired by constructions for topological Hochschild homology, the theory of cyclotomic synthetic spectra allows us to produce a relative conjugate filtration on Hodge–Tate cohomology and its variants, and in the same time, a relative conjugate filtration on topological Hochschild homology and its variants. As a consequence, we deduce transitivity of weak and strong F -smoothness.

1. INTRODUCTION

This article stems from studies of de-completed topological periodic cyclic homology in [Mao24], which turns out to be closely related to Segal conjecture. The interplay between de-completed topological periodic cyclic homology, Segal conjecture, and F -smoothness leads us to multiple results on both the commutative and noncommutative world, which is the main theme of this article.

Roughly speaking, in [BMS18], the authors produce a motivic filtration on topological periodic cyclic homology, whose associated graded pieces are given by Nygaard-completed absolute prismatic cohomology. In this sense, we say that topological periodic cyclic homology *corresponds to* Nygaard-completed absolute prismatic cohomology. Our work gave a construction of a localizing invariant corresponding to non-Hodge-completed de Rham cohomology. Via the same techniques, we extended Devalapurkar–Hahn–Raksit–Yuan’s de-completed topological Hochschild homology to a localizing invariant as well. In particular, it gives rise to de-completed topological periodic cyclic homology $\mathrm{TP}_{/\mathbb{Z}}^{\mathrm{poly}}$ (in this article, we take tacit (p, v_1) -completion for this de-completion). There are two natural questions out of it:

- (1) When is the motivic filtration on de-completed topological periodic cyclic homology complete?
- (2) When is de-completed topological periodic cyclic homology already “Nygaard-complete”?

It turns out that the “Nygaard-completeness” of de-completed topological periodic cyclic homology is closely related to Segal conjecture relative to $\mathrm{THH}(\mathbb{Z})$ (Remark 2.15). Our first result addresses the first question, which says that the motivic filtration on de-completed topological periodic cyclic homology is complete if the ring in question satisfies a very weak finite generation condition:

Theorem 1.1 (Lemma 3.16 and Theorem 3.19). *Let R be an (animated) commutative ring such that the cotangent complex $L_{R/\mathbb{Z}}$ is, as an R -module spectrum, p -completely (weakly) finitely generated. Then the motivic filtration on $\mathrm{TP}_{/\mathbb{Z}}^{\mathrm{poly}}(R)$ is complete.*

Received by the editors September 21, 2026.

Note: This document has been written using GNU $\mathrm{T}_{\mathrm{E}}\mathrm{X}_{\mathrm{MACS}}_{\mathrm{I}}$ [H+98].

This includes all (animated) commutative rings R whose derived modulo p is (weakly) F -finite (Example 3.25). The concept of F -smoothness was inspired by Segal conjecture. We prove a sufficient condition for Segal conjecture for THH of an F -smooth ring:

Theorem 1.2 (Lemma 3.16 and Theorem 3.20). *Let R be an F -smooth p -quasisyntomic ring such that the cotangent complex $L_{R/\mathbb{Z}}$ is, as an R -module spectrum, p -completely (weakly) finitely generated. Then the cyclotomic spectrum $\mathrm{THH}(R)$ satisfies p -complete Segal conjecture.*

This incorporates qualitative versions of known results such as smooth algebras over a perfectoid ring, or regular rings whose reduction mod p is F -finite (Example 3.26). This also includes a DVR which is not F -finite (Example 3.28).

On the other hand, in [BM23], it is established that, a p -torsion-free p -quasisyntomic ring R is F -smooth if its reduction R/p is F -smooth. We show that, the argument in [HW22] implies an analogue for Segal conjecture.

Theorem 1.3 (Proposition 4.2). *Let M be a right $\mathrm{THH}(\mathrm{BP}\langle n \rangle)$ -module in cyclotomic spectra such that $M/(p, v_1, \dots, v_n)$ is bounded below. Suppose that the cyclotomic spectrum*

$$M \otimes_{\mathrm{THH}(\mathrm{BP}\langle n \rangle)}^{\mathbb{L}} \mathrm{THH}(\mathbb{F}_p)$$

satisfies Segal conjecture, then the cyclotomic spectrum $M/(p, v_1, \dots, v_n)$ satisfies Segal conjecture.

Actually, in this case, its further reduction modulo v_{n+1} satisfies Lichtenbaum–Quillen conjecture as well. For a cyclotomic spectrum, Lichtenbaum–Quillen conjecture implies Segal conjecture, and the reverse implication requires (weak) canonical vanishing in general. However, if the cyclotomic spectrum in question admits a ring structure and a Bökstedt element, in which case we call it a *Bökstedt ring*, it was shown in [BHLS23] that canonical vanishing is automatic. An observation is that, their argument also adapts to modules over a Bökstedt ring (Lemma 2.17). In particular, the cyclotomic rings $\mathrm{THH}(\mathbb{F}_p)$ and $\mathrm{THH}(\mathbb{Z})$ are Bökstedt rings after reduction modulo (p, v_1) , and thus for a DG-category $\mathcal{C}$, we may deduce boundedness of $\mathrm{TR}(\mathcal{C})/(p, v_1)$ when $\mathrm{THH}(\mathcal{C})$ satisfies Segal conjecture (Corollary 2.18).

In addition to automatic canonical vanishing of (Segal-conjecture-satisfying) modules over a Bökstedt ring, their de-completed $(-)^{t\mathbb{T}}$ has a simple formula out of $(-)^{h\mathbb{T}}$ (Proposition 2.21). In the special case $\mathrm{TP}_{/\mathbb{Z}}^{\mathrm{poly}}$, we have

Theorem 1.4 (Corollaries 2.22 to 2.24). *Let $\mathcal{C}$ be a $\mathbb{Z}$ -linear DG-category. Then we have*

$$\mathrm{TP}_{/\mathbb{Z}}^{\mathrm{poly}}(\mathcal{C}) \simeq (\mathrm{TC}^-(\mathcal{C}) \otimes_{\mathrm{TC}^-(\mathbb{Z}), \varphi}^{\mathbb{L}} \mathrm{TP}(\mathbb{Z}))_{(p, v_1)}^{\wedge},$$

and the same if we replace $\mathbb{Z}$ by a perfectoid ring, and in this case, we may also replace v_1 by $\mathrm{Ker}(\theta)$ of Fontaine’s map on the perfectoid ring in question. Moreover, when R is a quasiregular semiperfectoid ring, we can identify $\mathrm{TP}_{/\mathbb{Z}}^{\mathrm{poly}}(R)$ with $\mathrm{TP}^{(-1)}(R)$ in [Man24] as expected.

Moreover, combining with ideas in [AKN23] on continuity of syntomic cohomology, we give a conceptual explanation of continuity of TC for modules over a Bökstedt ring (Proposition 2.26 and Corollary 2.27).

Finally, the theory of cyclotomic syntomic spectra in [AR24] allows us to systematically consider variants of prismatic cohomology and topological Hochschild homology on the same foot. In particular, we produce relative conjugate filtration on both Hodge–Tate cohomology and $\mathrm{THH}_{\Delta}^{\Phi_{C_p}}$ (a de-completion of THH^{tC_p}): the former is a mixed characteristic analogue of the one in [Bha12], while the later is a mixed characteristic and relative generalization of Kaledin’s conjugate filtration.

Theorem 1.5 (Examples 5.4 and 5.5 and Proposition 5.8). *Let k be an (animated) commutative ring.*

- (1) Let R be an (animated) commutative k -algebra. Then there exists a complete exhaustive filtration on $\overline{\Delta}_R\{i\}$ with the j -th associated graded piece given by

$$\left(\bigwedge_R^{i-j} L_{R/k}\right) \otimes_k^{\mathbb{L}} \overline{\Delta}_k\{j\}[j-i].$$

- (2) Let $\mathcal{C}$ be a dualizable presentable stable k -linear ∞ -category. Then there exists an exhaustive filtration on $\mathrm{THH}_{\Delta}(\mathcal{C})^{\Phi_{C_p}}$ with the j -th associated graded piece given by

$$\mathrm{HH}(\mathcal{C}/k) \otimes_k^{\mathbb{L}} \overline{\Delta}_k\{j\}[2j].$$

Moreover, if the k -module spectrum $L_{k/\mathbb{Z}}$ is weakly finitely generated, then this filtration is complete.

A similar construction leads to a relative filtration on $\mathrm{gr}_N^* \Delta$ and THH :

Theorem 1.6 (Examples 6.2 and 6.3 and Proposition 6.4). *Let k be an (animated) commutative ring.*

- (1) Let R be an (animated) commutative k -algebra. Then there exists a complete exhaustive filtration on $\overline{\Delta}_R\{i\}$ with the j -th associated graded piece given by

$$\left(\bigwedge_R^{i-j} L_{R/k}\right) \otimes_k^{\mathbb{L}} \mathrm{gr}_N^j \Delta_k\{j\}[j-i].$$

- (2) Let $\mathcal{C}$ be a dualizable presentable stable k -linear ∞ -category. Then there exists an exhaustive filtration on $\mathrm{THH}(\mathcal{C})$ with the j -th associated graded piece given by

$$\mathrm{HH}(\mathcal{C}/k) \otimes_k^{\mathbb{L}} \mathrm{gr}_N^j \Delta_k\{j\}[2j].$$

Moreover, if $\mathrm{THH}(\mathcal{C})$ is t -bounded below, then this filtration is complete.

As a consequence, we get transitivity of weak F -smoothness and F -smoothness, which was our original motivation to produce relative conjugate filtrations, inspired by Tess BOUIS's proof of transitivity of p -Cartier smoothness in [Bou23, Lem 2.10]. Before that, we first establish an animated version of transitivity of p -Cartier smoothness.

Theorem 1.7 (Theorem 6.30). *Let k be an (animated) ring, and $R \rightarrow A$ a map of (animated) k -algebras, such that $R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ is t -bounded, and $A \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ has bounded Tor-amplitude over $R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$. Suppose that*

- the R -module spectrum $L_{R/k}$ is p -completely flat; and that
- the A -module spectrum $L_{A/R}$ is p -completely flat.

If the p -completed derived de Rham cohomology $\mathrm{dR}_{A/R}$ is p -completely Hodge-complete, then the p -completed derived de Rham cohomology $\mathrm{dR}_{A/k}$ is p -completely “Hodge-complete relative to” $\mathrm{dR}_{R/k}$. In particular, if the p -completed derived de Rham cohomology $\mathrm{dR}_{R/k}$ is p -completely Hodge-complete, then so is $\mathrm{dR}_{A/k}$.

Now we phrase our transitivity of weak F -smoothness and F -smoothness.

Theorem 1.8 (Theorem 6.12). *Let $k \rightarrow R$ be a map of (animated) rings. Suppose that*

- the (animated) ring k is weakly F -smooth; and that
- the R -module spectrum $L_{R/k}$ is p -completely flat.

Then the (animated) ring R is weakly F -smooth as well.

Inspired by relative Segal conjecture, we introduce relative Nygaard-completeness (Definition 6.13), and prove that it is implied by p -Cartier smoothness with weak F -smoothness of the base.

Theorem 1.9 (Theorem 6.15 and Corollary 6.16). *Let $k \rightarrow R$ be a map of (animated) rings, such that $k \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ is t -bounded, and $R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ has bounded Tor-amplitude over $k \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$. Suppose that*

- the (animated) ring k is weakly F -smooth; and that
- the R -module spectrum $L_{R/k}$ is p -completely flat.

If the p -completed derived de Rham cohomology $dR_{R/k}$ is p -completely Hodge-complete, then the absolute prismatic cohomology $\Delta_R\{*\}$ is Nygaard-complete relative to $\Delta_k\{*\}$. In particular, if k is in addition F -smooth, then so is R .

We have formulated Theorem 1.9 to be clearly an absolute version of Theorem 1.7, which generalizes [BM23, Prop 4.9]. There are variants of Theorems 1.7 and 1.9 without bounds on Tor-amplitude of ring maps. For this, we introduce the concept of universal convergence of Hodge and Nygaard filtrations (Definitions 6.20 and 6.39), which generalizes “having a universal Cartier isomorphism” in [BLM18, Def 9.5.9]. These notions are to explore, but as an exhibition of its utility, we have

Theorem 1.10 (Corollary 6.44). *Let $k \rightarrow R$ be a map of rings such that the induced map $k \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \rightarrow R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ of animated $\mathbb{F}_p$ -algebras is relatively perfect. If the ring k is weakly F -smooth, the absolute prismatic cohomology Δ_k is Nygaard-complete, and the Nygaard filtration on $\widehat{\Delta_k}\{*\}$ converges universally, then the same is true for R .*

This is a version of [BM23, Prop 4.7] without flatness, giving an answer to the paragraph right before *loc. cit.*

Acknowledgments. We thank Benjamin ANTIEAU, Tess BOUIS, Daniel FINK, Rok GREGORIC, Hyungseop KIM and Georg TAMME for discussions. The author acknowledges support from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through the Collaborative Research Centre TRR 326 *Geometry and Arithmetic of Uniformized Structures*, project number 444845124.

2. DE-COMPLETED TP REVISITED

Devalapurkar–Hahn–Raksit–Yuan defines de-completed topological periodic cyclic homology of animated rings in terms of genuine equivariant homotopy theory, and [Mao24, §5] extended it to DG-categories. In this section, we first reformulate it via an equivalent description (Definition 2.1) without genuine equivariant homotopy theory, explaining its relation to Segal conjecture (Remark 2.15 and Proposition 2.16). Then in Corollary 2.24, we identify it with $TP^{(-1)}$ introduced in [Man24] (of which [Gre25, §1] gives an excellent exposition) in the general context of modules over a Bökstedt ring, a concept introduced in [BHLS23]. Finally, we give a conceptual explanation (Corollary 2.27) of continuity of TC, which applies to filtered systems which are not uniformly t -bounded below.

Definition 2.1. Let $\mathcal{C}$ be a dualizable presentable stable $\mathbb{Z}$ -linear ∞ -category. Then its $((p, v_1)$ -completed) *topological polynomial periodic cyclic homology* $TP_{/\mathbb{Z}}^{\text{poly}}(\mathcal{C})$ is defined to be the (p, v_1) -completion of the spectrum

$$\mathrm{THH}_{\Delta}(\mathcal{C})^{h\mathbb{T}} := (\mathrm{THH}(\mathcal{C}) \otimes_{\mathrm{THH}(\mathbb{Z}), \varphi_p}^{\mathbb{L}} \mathrm{THH}(\mathbb{Z})^{tC_p})^{h\mathbb{T}},$$

where THH_{Δ} is de-completed THH à la Devalapurkar–Hahn–Raksit–Yuan, extended to DG-categories in [Mao24, §5]. In this paper, we only use

$$\mathrm{THH}_{\Delta}(\mathcal{C})^{\Phi C_p} = \mathrm{THH}(\mathcal{C}) \otimes_{\mathrm{THH}(\mathbb{Z})}^{\mathbb{L}} \mathrm{THH}(\mathbb{Z})^{tC_p}$$

to simplify the cumbersome expression on the right hand side.

Remark 2.2. Let $\mathcal{C}$ be a dualizable presentable stable $\mathbb{Z}$ -linear ∞ -category. Then the *assembly map* out of $TP_{/\mathbb{Z}}^{\text{poly}}(\mathcal{C})$ is given by the (p, v_1) -completion of the map

$$(\mathrm{THH}_{\Delta}(\mathcal{C})^{\Phi C_p})^{h\mathbb{T}} \longrightarrow (\mathrm{THH}(\mathcal{C})^{tC_p})^{h\mathbb{T}}$$

induced by the cyclotomic Frobenius map $\mathrm{THH}(\mathcal{C}) \rightarrow \mathrm{THH}(\mathcal{C})^{tC_p}$, where the target can be (p, v_1) -completely identified with $\mathrm{TP}(\mathcal{C})$ by the Tate orbit lemma¹.

Remark 2.3. In Definition 2.1 and Remark 2.2, we may replace the base $\mathrm{THH}(\mathbb{Z})$ by $\mathrm{THH}(\mathrm{BP}\langle n \rangle)$, and replace (p, v_1) -completion by $(p, v_1, \dots, v_{n+1})$ -completion.

Construction 2.4. *Let R be an animated ring. Then by [AR24] (on polynomial rings, from which one can left Kan extend to general animated rings), the filtered cyclotomic spectrum $\mathrm{Fil}_M \mathrm{THH}(R)$ acquires an $\mathbb{E}_\infty$ -cyclotomic synthetic spectrum structure. This gives rise to a synthetic spectrum structure on*

$$\mathrm{Fil}_M \mathrm{THH}_\Delta(R)^{\Phi C_p} := \mathrm{Fil}_M \mathrm{THH}(R) \otimes_{\mathrm{Fil}_{\mathrm{ev}} \mathrm{THH}(\mathbb{Z}), \varphi_p}^{\mathbb{L}} \mathrm{Fil}_{\mathrm{ev}} \mathrm{THH}(\mathbb{Z})^{tC_p}$$

with a $\mathbb{T}_{\mathrm{ev}}$ -action, where $\mathrm{Fil}_M \mathrm{THH}(\mathbb{Z})^{tC_p} \simeq (\mathrm{Fil}_M \mathrm{THH}(\mathbb{Z}))^{tC_{p, \mathrm{ev}}}$. In particular, it gives rise to filtrations on $\mathrm{THH}_\Delta(R)^{\Phi C_p}$ and on $\mathrm{TP}_{/\mathbb{Z}}^{\mathrm{poly}}(R)$ as well, called motivic filtrations. By construction, the first filtration is exhaustive. To see that the second filtration is also exhaustive, we need some preparations for $\mathrm{Fil}_M \mathrm{THH}(\mathbb{Z})$.

Remark 2.5. Let R be an animated ring. Recall that

$$\pi_*(\mathrm{THH}(\mathbb{Z}) \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p) = \mathbb{F}_p[\mu_1] \otimes \Lambda(\lambda_1)$$

with $\deg \mu_1 = 2p$ and $\deg \lambda_1 = 2p - 1$, and the Frobenius map

$$\mathrm{THH}(\mathbb{Z}) \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \rightarrow \mathrm{THH}(\mathbb{Z})^{tC_p} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$$

is inverting the class μ_1 . As in [HRW22, §6], these classes lift to synthetic classes (non-equivariantly), still denoted by λ_1 and μ_1 respectively, and the Frobenius map is inverting μ_1 as well. This realizes $\mathrm{Fil}_M(\mathrm{THH}_\Delta(R)^{\Phi C_p} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ as inverting the class μ_1 in $\mathrm{Fil}_M(\mathrm{THH}(R) \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$, and in particular, realizes $\mathrm{THH}_\Delta(R)^{\Phi C_p} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ as inverting the class μ_1 in $\mathrm{THH}(R) \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$.

Lemma 2.6. *Let M be a $\mathrm{Fil}_M \mathrm{THH}(\mathbb{Z})$ -module in $\mathrm{SynSp}_{\mathbb{T}_{\mathrm{ev}}}$, and*

$$N := M \otimes_{\mathrm{Fil}_M \mathrm{THH}(\mathbb{Z})}^{\mathbb{L}} \mathrm{Fil}_M \mathrm{THH}(\mathbb{Z})^{tC_p}.$$

Then the norm map

$$\Sigma_{\mathrm{S}_{\mathrm{ev}}}(1) \otimes_{\mathrm{S}_{\mathrm{ev}}} N_{h\mathbb{T}_{\mathrm{ev}}} \longrightarrow N^{h\mathbb{T}_{\mathrm{ev}}}$$

is an equivalence of synthetic spectra after $(v_0, \tilde{v}_1)$ -completion. Consequently, the functor

$$\begin{aligned} \mathrm{Mod}_{\mathrm{Fil}_M \mathrm{THH}(\mathbb{Z})}(\mathrm{SynSp}_{\mathbb{T}_{\mathrm{ev}}}) &\longrightarrow \mathrm{SynSp}_{(v_0, \tilde{v}_1)}^\wedge \\ M &\longmapsto (N^{h\mathbb{T}_{\mathrm{ev}}})_{(v_0, \tilde{v}_1)}^\wedge \end{aligned}$$

preserves small colimits, where v_0 and $\tilde{v}_1$ are the synthetic analogues of the homotopy classes p and v_1 in $\mathbb{S}$ and $\mathbb{S}/p$.

Proof. By [AKN24, Cons 2.58], we have a fiber sequence

$$\Sigma_{\mathrm{S}_{\mathrm{ev}}}(1) \otimes_{\mathrm{S}_{\mathrm{ev}}} N_{h\mathbb{T}_{\mathrm{ev}}} \longrightarrow N^{h\mathbb{T}_{\mathrm{ev}}} \longrightarrow N^{t\mathbb{T}_{\mathrm{ev}}}.$$

The result follows, as the synthetic spectrum $N^{t\mathbb{T}_{\mathrm{ev}}}$ carries a $(\mathrm{Fil}_M \mathrm{THH}(\mathbb{Z})^{tC_p})^{t\mathbb{T}_{\mathrm{ev}}}$ -module structure, and the $\mathbb{E}_\infty$ -algebra $((\mathrm{Fil}_M \mathrm{THH}(\mathbb{Z})^{tC_p})^{t\mathbb{T}_{\mathrm{ev}}})_{(v_0, \tilde{v}_1)}^\wedge$ in synthetic spectra is zero.

A way to see the later: the $\mathbb{T}_{\mathrm{ev}}$ -equivariant lift $\tilde{\mu}_1$ of the class μ_1 in $\mathrm{Fil}_M \mathrm{THH}(\mathbb{Z})/(v_0, \tilde{v}_1)$ is sent to an invertible $\mathbb{T}_{\mathrm{ev}}$ -equivariant class in $\mathrm{Fil}_M \mathrm{THH}(\mathbb{Z})^{tC_p}/(v_0, \tilde{v}_1)$, thus to an invertible class in

$$(\mathrm{Fil}_M \mathrm{THH}(\mathbb{Z})^{tC_p})^{t\mathbb{T}_{\mathrm{ev}}}/(v_0, \tilde{v}_1).$$

¹In full generality, we need the t -boundedness of $\mathrm{TR}(\mathbb{Z})/(p, v_1)$, a special case of Lichtenbaum–Quillen for $\mathrm{TR}(\mathrm{BP}\langle n \rangle)/(p, v_1, \dots, v_{n+1})$ established in [HW22].

On the other hand, it becomes a zero class in $(\mathrm{Fil}_M \mathrm{THH}(\mathbb{Z}))^{t\mathbb{T}^{\mathrm{ev}}}$, which sends to the zero class along the map

$$(\mathrm{Fil}_M \mathrm{THH}(\mathbb{Z}))^{t\mathbb{T}^{\mathrm{ev}}}/(v_0, \tilde{v}_1) \rightarrow (\mathrm{Fil}_M \mathrm{THH}(\mathbb{Z})^{tC_p})^{t\mathbb{T}^{\mathrm{ev}}}/(v_0, \tilde{v}_1).$$

□

Corollary 2.7. *Let R be an animated ring. Then the motivic filtration on $\mathrm{TP}_{/\mathbb{Z}}^{\mathrm{poly}}(R)$ is exhaustive.*

Remark 2.8. Let R be a polynomial ring. Then the map

$$\mathrm{THH}_{\Delta}(R)^{\Phi_{C_p}} \longrightarrow \mathrm{THH}(R)^{tC_p}$$

induced by the cyclotomic Frobenius $\mathrm{THH}(R) \rightarrow \mathrm{THH}(R)^{tC_p}$ is an equivalence after (p, v_1) -completion by results in [Mao24], and passing to associated graded pieces of motivic filtrations, it is an equivalence as well by, say, [BM23, Prop 2.6] and F -smoothness of R . It follows that the motivic filtrations considered in Construction 2.4 coincides with the one considered in [Mao24].

We may replace the (p, v_1) -completed base $\mathrm{THH}(\mathbb{Z})$ (or more precisely, its reduction modulo (p, v_1)) by a Bökstedt ring, a notion that appears in [BHLS23].

Definition 2.9. A *Bökstedt ring of (height $n \in \mathbb{N}$)* is given by the datum (R, μ) where

- (1) R is a p -nilpotent t -bounded-below $h\mathbb{A}_2$ -ring in cyclotomic spectra, which satisfies Segal conjecture;
- (2) μ is an element of $\pi_{2p^n}(R)$ which is a Bökstedt class [BHLS23, Def 2.31].

Remark 2.10. In practice, Bökstedt rings come from modulo out a sequence of homotopy classes from an $\mathbb{E}_2$ -algebra in cyclotomic spectra, such as $\mathrm{THH}(\mathrm{BP}\langle n \rangle)/(p, v_1, \dots, v_{n+1})$. Thanks to [Bur22], we may raise powers of homotopy classes to improve associativity and commutativity. In particular, one might freely replace $h\mathbb{A}_2$ by $\mathbb{E}_1$ or even $\mathbb{E}_2$ in Definition 2.9.

Recall that every Bökstedt ring (R, μ) has the following properties [BHLS23, Lem 2.33 & 2.36 & Prop 2.30]:

- The spectrum R/μ is t -bounded.
- The map $R[\mu^{-1}] \rightarrow R^{tC_p}$ induced by the cyclotomic Frobenius $\varphi_p : R \rightarrow R^{tC_p}$ is an equivalence.
- The cyclotomic spectrum R is cyclotomically t -bounded, i.e. the spectrum $\mathrm{TR}(R)$ is t -bounded.

And conversely, every p -nilpotent $h\mathbb{A}_2$ -ring in cyclotomic spectra, which is cyclotomically t -bounded, admits a Bökstedt ring structure.

Example 2.11. $(\mathrm{THH}(\mathbb{F}_p)/p, (\mu_0 \in \mathrm{THH}_2(\mathbb{F}_p)))$ is a Bökstedt ring.

Example 2.12. $(\mathrm{THH}(\mathbb{Z})/(p, v_1), (\mu_1 \in \mathrm{THH}_{2p}(\mathbb{Z}; \mathbb{F}_p)))$ is a Bökstedt ring, or more generally, $(\mathrm{THH}(\mathrm{BP}\langle n \rangle)/(p, v_1, \dots, v_{n+1}), (\mu_1 \in \mathrm{THH}_{2p^{n+1}}(\mathrm{BP}\langle n \rangle; \mathbb{F}_p)))$. This is a consequence of [HW22, Thm G].

Example 2.13 ([Mao24, §5]). Let S be a perfectoid ring. Then $(\mathrm{THH}(S)/(p, v_1), (\mu_0 \in \pi_2(\mathrm{THH}(S)_p^\wedge)))$ is a Bökstedt ring, and the map $\mathrm{THH}(\mathbb{Z})/p \rightarrow \mathrm{THH}(S)/p$ carries the class μ_1 to μ_0^p up to a unit.

The “Nygaard-completeness” of $\mathrm{TP}^{\mathrm{poly}}$ is closely related to the following relative version of Segal conjecture (Remark 2.15).

Definition 2.14. Let (R, μ) be a Bökstedt ring. We say that a right R -module M in cyclotomic spectra satisfies *Segal conjecture relative to R* if the map

$$M[\mu^{-1}] = M \otimes_R R^{tC_p} \longrightarrow M^{tC_p}$$

induced by the cyclotomic Frobenius $\varphi_p : M \rightarrow M^{tC_p}$ is an equivalence. Equivalently, it is saying that $\mathrm{fib}(\varphi_p)[\mu^{-1}] = 0$.

Remark 2.15. Let $\mathcal{C}$ be a dualizable presentable stable $\mathbb{Z}$ -linear ∞ -category. Then by definition, the assembly map $\mathrm{TP}^{\mathrm{poly}}(\mathcal{C}) \rightarrow \mathrm{TP}(\mathcal{C})_{(p, v_1)}^\wedge$ is an equivalence if the cyclotomic spectrum $\mathrm{THH}(\mathcal{C})/(p, v_1)$ satisfies Segal conjecture relative to $\mathrm{THH}(\mathbb{Z})/(p, v_1)$.

The following statement compares absolute and relative Segal conjecture, which follows from adapting the proof of [BHLS23, Lem 2.33].

Proposition 2.16. *Let (R, μ) be a Bökstedt ring. Then a right R -module M in cyclotomic spectrum satisfies Segal conjecture if and only the following two conditions hold:*

- (1) *The spectrum M/μ is t -bounded above.*
- (2) *The R -module M in cyclotomic spectrum satisfies Segal conjecture relative to R .*

Proof. Let $N := \mathrm{fib}(\varphi_p)$. The canonical map $N \rightarrow M$ is an equivalence after modulo μ , as μ acts invertibly on M^{tC_p} . Suppose that M satisfies Segal conjecture, then N is t -bounded above, and in this case, so is N/μ . Thus it suffices to establish the equivalence of Segal conjecture and relative Segal conjecture under the t -bounded-aboveness of N/μ . Indeed, for every $m \in \mathbb{Z}$, the sequential tower

$$\pi_m(N) \xrightarrow{\mu} \pi_{m+2p^n}(N) \xrightarrow{\mu} \pi_{m+4p^n}(N) \xrightarrow{\mu} \cdots,$$

which computes $\pi_m(N[\mu^{-1}])$, stabilizes (uniformly for $m \in \mathbb{N}$, say). This implies that, the spectrum N is t -bounded above (i.e. Segal conjecture) if and only if $N[\mu^{-1}] = 0$ (i.e. relative Segal conjecture). $\square$

We note that, for modules over a Bökstedt ring, it satisfies (absolute) Segal conjecture if and only if it satisfies it has bounded TR.

Lemma 2.17 ([BHLS23, Lem 2.36]). *Let (R, μ) be a Bökstedt ring, and M a right R -module in $\mathrm{Sp}^{B\mathbb{T}}$. Suppose that the spectrum M/μ is b -truncated. Then M satisfies strong canonical vanishing with parameter b .*

Proof. The proof of [BHLS23, Lem 2.36] works verbatim. We briefly summarize as follows. First pick a lift $\hat{\mu} \in \pi_{2p^n}(\Sigma R_{h\mathbb{T}})$, and there exists $d \gg 0$ such that $a_\lambda^d \hat{\mu} = 0$. Then the composite $\mathbb{T}$ -equivariant map

$$(1) \quad \tau_{>b} M \xrightarrow{c} M \xrightarrow{a_\lambda^d} \Sigma^{d\lambda} M$$

is null homotopic, as the first map c factors as

$$\tau_{>b} M \longrightarrow \Sigma^{2p^n} M \xrightarrow{\mathrm{Nm}(\hat{\mu})} M$$

since the composite $\mathbb{T}$ -equivariant map $\tau_{>b} M \xrightarrow{c} M \rightarrow M/\mathrm{Nm}(\hat{\mu})$ is null-homotopic, by b -truncatedness of M/μ , where Nm is the norm map $R_{h\mathbb{T}}[1] \rightarrow R^{h\mathbb{T}}$, and thus $\mathrm{Nm}(\hat{\mu})$ is an equivariant lift of μ . But now, the composite $\mathbb{T}$ -equivariant map

$$\Sigma^{2p^n} M \longrightarrow M \xrightarrow{a_\lambda^d} \Sigma^{d\lambda} M$$

is null-homotopic. This gives rise to a null homotopy of the composite $\mathbb{T}$ -equivariant map (1). $\square$

Corollary 2.18. *Let M be a right $\mathrm{THH}(\mathrm{BP}\langle n \rangle)$ -module in cyclotomic spectra such that $M/(p, v_1, \dots, v_n)$ satisfies Segal conjecture. Then the spectrum $\mathrm{TR}(M)/(p, v_1, \dots, v_{n+1})$ is t -bounded above.*

Proof. Apply Proposition 2.16 and Lemma 2.17 to the Bökstedt ring $\mathrm{THH}(\mathrm{BP}\langle n \rangle)/(p, v_1, \dots, v_{n+1})$ as in Example 2.12. $\square$

Example 2.19. Let $\mathcal{C}$ be a bounded smooth DG-category (over $\mathbb{Z}$). Then by Corollary 2.18 and one of main results in [Mao24], the spectrum $\mathrm{TR}(\mathcal{C})/(p, v_1)$ is t -bounded.

Example 2.20. Let S be a perfectoid ring, and $\mathcal{C}$ a bounded smooth S -linear DG-category. Then by Corollary 2.18 and one of main results in [Mao24], the spectrum $\mathrm{TR}(\mathcal{C})/(p, v_1)$ is t -bounded. In particular, when S is a perfect $\mathbb{F}_p$ -algebra, $\mathrm{TR}(\mathcal{C})$ is p -completely t -bounded above: since $\mathrm{TR}(\mathbb{F}_p) = \mathbb{Z}_p$, we see that $v_1 = 0$ on $\mathrm{TR}(\mathcal{C})$.

For modules over a Bökstedt ring, we can simply define the de-completed $\mathbb{T}$ -Tate construction by taking homotopy $\mathbb{T}$ -fixed points. This is justified by the following, an analogue of Lemma 2.6.

Proposition 2.21. *Let (R, μ) be a Bökstedt ring. Then*

(1) *the functor*

$$\begin{aligned} \mathrm{RMod}_R(\mathrm{Sp}^{B\mathbb{T}}) &\longrightarrow \mathrm{Sp} \\ M &\longmapsto (M \otimes_R^{\mathbb{L}} R^{tC_p})^{h\mathbb{T}} \end{aligned}$$

preserves sifted colimits; and

(2) *letting M be an R -module in $\mathrm{Sp}^{B\mathbb{T}}$, the assembly map*

$$M^{h\mathbb{T}} \otimes_{R^{h\mathbb{T}}, \varphi := \varphi_p^{h\mathbb{T}}}^{\mathbb{L}} R^{t\mathbb{T}} \longrightarrow (M \otimes_R^{\mathbb{L}} R^{tC_p})^{h\mathbb{T}}$$

is an equivalence.

Proof. We prove the both statements in the same time by examining the vertical morphism

$$\begin{array}{ccccc} (\Sigma M_{h\mathbb{T}}) \otimes_{R^{h\mathbb{T}}, \varphi}^{\mathbb{L}} R^{t\mathbb{T}} & \longrightarrow & M^{h\mathbb{T}} \otimes_{R^{h\mathbb{T}}, \varphi} R^{t\mathbb{T}} & \longrightarrow & M^{t\mathbb{T}} \otimes_{R^{h\mathbb{T}}, \varphi} R^{t\mathbb{T}} \\ \downarrow & & \downarrow & & \downarrow \\ \Sigma(M \otimes_R^{\mathbb{L}} R^{tC_p})_{h\mathbb{T}} & \longrightarrow & (M \otimes_R^{\mathbb{L}} R^{tC_p})^{h\mathbb{T}} & \longrightarrow & (M \otimes_R^{\mathbb{L}} R^{tC_p})^{t\mathbb{T}} \end{array}$$

of horizontal fiber sequences, where the right $R^{h\mathbb{T}}$ -module structure on $M^{t\mathbb{T}}$ is induced by the canonical map $\mathrm{can} : R^{h\mathbb{T}} \rightarrow R^{t\mathbb{T}}$, while the base change on the first row is along the Frobenius map $\varphi := \varphi_p^{h\mathbb{T}} : R^{h\mathbb{T}} \rightarrow R^{t\mathbb{T}}$.

Let $\tilde{\mu} \in \pi_{2p^n}(R^{h\mathbb{T}})$ be an equivariant lift of μ , which further admits a lift to $\pi_{2p^n}(\Sigma R_{h\mathbb{T}})$. Then by assumption, the map $\varphi : R^{h\mathbb{T}} \rightarrow R^{t\mathbb{T}}$ is precisely localization at the class $\tilde{\mu}$. However, since $\tilde{\mu}$ lifts along $\mathrm{Nm} : \Sigma R_{h\mathbb{T}} \rightarrow R^{h\mathbb{T}}$, we have $\mathrm{can}(\tilde{\mu}) = 0$. This implies that

$$M^{t\mathbb{T}} \otimes_{R^{h\mathbb{T}}, \varphi}^{\mathbb{L}} R^{t\mathbb{T}} = 0$$

(compare with Lemma 2.6). Moreover, by Tate orbit and fixed point lemma, we have $(R^{tC_p})^{t\mathbb{T}} \simeq 0$, and thus

$$(M \otimes_R^{\mathbb{L}} R^{tC_p})^{t\mathbb{T}} = 0.$$

It follows that the two left horizontal maps are equivalences. The construction $\Sigma(-)_{h\mathbb{T}}$ preserves sifted colimits, and in particular, inverting $\tilde{\mu}$, thus the leftmost vertical map is an equivalence. Thus all arrows on the left square are equivalences, and the result follows. $\square$

Corollary 2.22. *Let $\mathcal{C}$ be a dualizable presentable stable $\mathbb{Z}$ -linear ∞ -category. Then the canonical map*

$$\mathrm{TC}^-(\mathcal{C}) \otimes_{\mathrm{TC}^-(\mathbb{Z}), \varphi}^{\mathbb{L}} \mathrm{TP}(\mathbb{Z}) \longrightarrow \mathrm{TP}_{/\mathbb{Z}}^{\mathrm{poly}}(\mathcal{C})$$

is an equivalence after (p, v_1) -completion².

Proof. It follows from applying Proposition 2.21 to the Bökstedt ring $\mathrm{THH}(\mathbb{Z})/(p, v_1)$ as in Example 2.12. $\square$

²We do not complete it with respect to any “secondary filtration” as hinted in [Man24, Rem 1.6].

Corollary 2.23. *Let S be a perfectoid ring, and $\mathcal{C}$ a dualizable presentable stable S -linear ∞ -category. Then the canonical map*

$$\mathrm{TC}^-(\mathcal{C}) \otimes_{\mathrm{TC}^-(S), \varphi}^{\mathbb{L}} \mathrm{TP}(S) \longrightarrow \mathrm{TP}_{/\mathbb{Z}}^{\mathrm{poly}}(\mathcal{C})$$

is an equivalence after (p, I) -completion, where $I := \mathrm{fib}(\Delta_S \rightarrow \overline{\Delta}_S = S)$.

Proof. Applying Proposition 2.21 to the Bökstedt ring $\mathrm{THH}(S)/(p, v_1)$ as in Example 2.13, we get the result up to (p, v_1) -completion. However, the commutative ring $\pi_*(\mathrm{TP}(S)_p^\wedge)$, we have $v_1 \doteq I^p$ up to a unit (cf. [Mao24, §5]), thus (p, v_1) -completion is the same as (p, I) -completion for $\mathrm{TP}(S)_p^\wedge$ -modules. $\square$

Corollary 2.24. *Let R be a quasiregular semiperfectoid ring. Then the canonical map*

$$\mathrm{TC}^-(R) \longrightarrow \mathrm{TP}^{\mathrm{poly}}(R)$$

is a localization after (p, I) -completion, where $I := \mathrm{fib}(\Delta_R \rightarrow \overline{\Delta}_R)$, which can be identified with $\mathrm{TP}^{(-1)}(R)$ in [Man24].

Proof. Pick a surjection $S \twoheadrightarrow R$ for a perfectoid ring S . The result follows from Corollary 2.23 as the Frobenius map $\varphi : \mathrm{TC}^-(S)_p^\wedge \rightarrow \mathrm{TP}(S)_p^\wedge$ is already a localization (at a class in $\pi_2(\mathrm{TC}^-(S)_p^\wedge)$). The identification of this localization with $\mathrm{TP}^{(-1)}(R)$ is established in [Gre25, Rem 1.3]. $\square$

Finally, we give a conceptual explanation³ of continuity of TC established in [CMM21] via de-completed TC^- and TP, which is much more akin to [AKN23, Prop 7.12]. Actually, the argument is parallel, as a consequence of the following simple category-theoretic observation:

Lemma 2.25. *Let $\mathcal{C}$ be an ∞ -category with finite limits. Let*

$$\begin{array}{ccc} x_{00} & \longrightarrow & x_{01} \\ \downarrow & \nearrow & \downarrow \\ x_{10} & \longrightarrow & x_{11} \end{array}$$

*be a diagram in $\mathcal{C}$, where the outer square is a Cartesian diagram, while the two inner triangles are **not** commutative.*

Then the map from the limit of the upper triangle (viewed as a diagram $\partial\Delta^2 \rightarrow \mathcal{C}$) to the limit of the lower triangle (viewed as a diagram $\partial\Delta^2 \rightarrow \mathcal{C}$ as well) is an equivalence.

Proof. It follows directly from “decomposing the diagram” (e.g. we decompose the lower triangle via $\partial\Delta^2 = \Lambda_2^2 \amalg_{\partial\Delta^1} \Delta^1$, where the composite functor $\Lambda_2^2 \rightarrow \partial\Delta^2 \rightarrow \mathcal{C}$ is the lower cospan $(x_{10} \rightarrow x_{11} \leftarrow x_{01})$, and the composite $\Delta^1 \rightarrow \partial\Delta^2 \rightarrow \mathcal{C}$ is the “diagonal” arrow $(x_{10} \rightarrow x_{01})$, and x_{00} is the limit of the first composite functor $\Lambda_2^2 \rightarrow \mathcal{C}$). Alternatively, the proof of [AKN23, Prop 7.12] works in this case as well. $\square$

Proposition 2.26. *Let (R, μ) be a Bökstedt ring. Then the composite functor*

$$\mathrm{RMod}_R(\mathrm{CycSp}_p) \longrightarrow \mathrm{CycSp}_p \xrightarrow{\mathrm{TC}} \mathrm{Sp}$$

preserves sifted colimits.

Proof. Let M be a right R -module in cyclotomic spectra. Then the Frobenius map $\varphi : M^{h\mathbb{T}} \rightarrow M^{t\mathbb{T}}$ factors as $M^{h\mathbb{T}} \xrightarrow{\hat{\varphi}} (M \otimes_R^{\mathbb{L}} R^{tC_p})^{h\mathbb{T}} \rightarrow M^{t\mathbb{T}}$. Now we formulate the triangle $\partial\Delta^2 \rightarrow \mathrm{Sp}$ given by

³We thank Georg TAMME for asking us such a question.

$$\begin{array}{ccc}
& & (M \otimes_R^{\mathbb{L}} R^{tC_p})^{h\mathbb{T}} \\
& \nearrow & \downarrow \\
M^{h\mathbb{T}} & \xrightarrow{\text{can}} & M^{t\mathbb{T}}
\end{array}$$

where the “diagonal” map is $\tilde{\varphi}$. Note that $\text{TC}(M)$ is precisely the limit of this triangle. Let $P(M)$ denote the pullback

$$M^{h\mathbb{T}} \times_{\text{can}, M^{t\mathbb{T}}} (M \otimes_R^{\mathbb{L}} R^{tC_p})^{h\mathbb{T}},$$

which gives rise to a functor $P : \text{RMod}_R(\text{CycSp}_p) \rightarrow \text{Sp}$, understood as “de-completed homotopy fixed points”. By Lemma 2.25, the spectrum $\text{TC}(M)$ can also be computed by the limit of the upper triangle

$$\begin{array}{ccc}
P(M) & \longrightarrow & (M \otimes_R^{\mathbb{L}} R^{tC_p})^{h\mathbb{T}} \\
\downarrow & \nearrow & \\
M^{h\mathbb{T}} & &
\end{array}$$

Since we have a fiber sequence $\Sigma M_{h\mathbb{T}} \rightarrow P(M) \rightarrow (M \otimes_R^{\mathbb{L}} R^{tC_p})^{h\mathbb{T}}$, it follows from Proposition 2.21 that P preserves sifted colimits, and so does TC on $\text{RMod}_R(\text{CycSp}_p)$, as $\text{TC}(M)$ is an equalizer of $P(M) \rightrightarrows (M \otimes_R^{\mathbb{L}} R^{tC_p})^{h\mathbb{T}}$, where we invoke Proposition 2.21 again. $\square$

Corollary 2.27. *Let $n \in \mathbb{Z}_{\geq -1}$. Then the exact functor*

$$\text{RMod}_{\text{THH}(\text{BP}\langle n \rangle)}(\text{CycSp}_p) \rightarrow \text{CycSp}_{(p, v_1, \dots, v_{n+1})}^{\wedge} \xrightarrow{\text{TC}(-)^{\wedge}} \text{Sp}_{(p, v_1, \dots, v_{n+1})}^{\wedge}$$

preserves small colimits.

Proof. It follows from applying Proposition 2.26 to the Bökstedt ring $\text{THH}(\text{BP}\langle n \rangle)/(p, v_1, \dots, v_{n+1})$ as in Example 2.12. $\square$

Example 2.28. Let (M_{α}) be a uniformly t -bounded below filtered diagram of $\text{THH}(\text{BP}\langle n \rangle)$ -modules in CycSp . Then $(\text{colim}_{\alpha} \text{TC}(M_{\alpha}))_p^{\wedge}$ and $\text{TC}(\text{colim}_{\alpha} M_{\alpha})_p^{\wedge}$ are t -bounded below. It follows that they are automatically $(p, v_1, \dots, v_{n+1})$ -complete, and hence the canonical map

$$\text{colim}_{\alpha} \text{TC}(M_{\alpha}) \longrightarrow \text{TC}\left(\text{colim}_{\alpha} M_{\alpha}\right)$$

is an equivalence after p -completion. This reproduces [CMM21, Thm 2.7] for $\text{THH}(\text{BP}\langle n \rangle)$ -modules in CycSp . When the filtered system is not uniformly t -bounded below, Corollary 2.27 is potentially more informative.

3. COMPLETENESS OF MOTIVIC FILTRATION

In this section, we will give a sufficient condition (Theorem 3.19) for completeness of motivic filtration on de-completed topological periodic cyclic homology. We also establishes a Segal conjecture (Theorem 3.20) when the ring in question is additionally F -smooth. For this purpose, we first introduce a weak finite generation condition (Definition 3.13) for modules, based on Shannon’s notion of rank of flat modules. Then we deduce a connectivity (Corollary 3.18) for associated graded pieces of absolute prismatic cohomology with respect to Nygaard filtration under this finite generation of cotangent complex. The main completeness result then follows.

We start with a higher algebra version of Shannon’s rank of flat modules:

Definition 3.1 ([Sha70]). Let R be a connective $\mathbb{E}_1$ -ring. We say that a flat right R -module P is of *finite rank* (abbrev. $\mathrm{rk}_R(P) < \infty$) if there exists a natural number $r \in \mathbb{N}$ such that P is a filtered colimit of finite free right R -modules of rank $\leq r$, and in this case, we say that P is of rank $\leq r$ (abbrev. $\mathrm{rk}_R(P) \leq r$).

Example 3.2 (Ind-Zariski localizations). The flat $\mathbb{Z}$ -module $\mathbb{Q}$ is of rank ≤ 1 , as we can write $\mathbb{Q}$ as the sequential colimit of $\mathbb{Z} \hookrightarrow (2!)^{-1}\mathbb{Z} \hookrightarrow (3!)^{-1}\mathbb{Z} \hookrightarrow \dots$. More generally, let R be an $\mathbb{E}_1$ -ring, M a finite free right R -module, and $S \subseteq \pi_0(R)$ a multiplicative subset satisfying right Ore condition in $\pi_*(R)$. Then the localization $S^{-1}M$, as a flat right R -module spectrum, is of rank ≤ 1 .

Example 3.3 (Standard étale maps). Let R be a commutative ring, $g \in R[t]$ an R -polynomial, and $f \in R[t]$ a monic R -polynomial. Let $S := R[t]_f/(g)$. Then $\mathrm{rk}_R(S) \leq \deg(g)$. Recall that, when the derivative f' is invertible in S , such a map $R \rightarrow S$ is called a *standard étale map* [Sta21, Tag 00UB].

Remark 3.4. It follows from definition that

- Let $R \rightarrow S$ be a map of connective $\mathbb{E}_1$ -rings, and M a flat right R -module. Then $\mathrm{rk}_S(M \otimes_R^{\mathbb{L}} S) \leq \mathrm{rk}_R(M)$.
- Let R be a connective $\mathbb{E}_1$ -ring, and M and N two flat right R -modules. Then $\mathrm{rk}_R(M \oplus N) \leq \mathrm{rk}_R(M) + \mathrm{rk}_R(N)$.
- Let R be a connective $\mathbb{E}_\infty$ -ring, and M and N two flat R -modules. Then $\mathrm{rk}_R(M \otimes_R^{\mathbb{L}} N) \leq \mathrm{rk}_R(M) \mathrm{rk}_R(N)$.
- Let R and S be two connective $\mathbb{E}_1$ -rings, M a flat right R -module, and N a flat right S -module. Then $M \times N$ is a flat $R \times S$ -module, with $\mathrm{rk}_{R \times S}(M \times N) \leq \max\{\mathrm{rk}_R(M), \mathrm{rk}_S(N)\}$.

We give a criterion for flat modules being of finite rank, as a higher algebra generalization of [Sha70, Thm 2].

Notation 3.5. Let R be a connective $\mathbb{E}_1$ -ring, and $n \in \mathbb{N}$. We will denote by $R^{\oplus n}$, or R^n when there is no ambiguity, the finite free (right) R -module of rank n . More generally, let M be a right R -module spectrum. We will denote by $M^{\oplus n}$ the direct sum of n copies of M .

Proposition 3.6. Let R be a connective $\mathbb{E}_1$ -ring, $r \in \mathbb{N}$, and P a flat right R -module. Then P is of rank $\leq r$ if and only if, for every finite free right R -module L , every map $L \rightarrow P$ factors through some R^s for $s \leq r$.

Proof. “**Only if**” part: We write P as a filtered colimit $\mathrm{colim}_\alpha L_\alpha$ with every L_α being finite free of rank $\leq r$. By compactness of L in RMod_R , we have

$$\mathrm{Hom}_{\mathrm{RMod}_R}(L, P) = \mathrm{colim}_\alpha \mathrm{Hom}_{\mathrm{RMod}_R}(L, L_\alpha)$$

and the result follows.

“**If**” part: It suffices to check that the slice category $(\mathrm{RMod}_R^{\mathrm{free}, \mathrm{rk} \leq r})_{/P}$ is filtered. By Lazard’s theorem [Lur17, Thm 7.2.2.15], the category $(\mathrm{RMod}_R^{\mathrm{free}, \mathrm{rk} < \infty})_{/P}$ is filtered. In particular, any diagram $q : K \rightarrow (\mathrm{RMod}_R^{\mathrm{free}, \mathrm{rk} \leq r})_{/P}$ extends to a diagram $q^\triangleright : K^\triangleright \rightarrow (\mathrm{RMod}_R^{\mathrm{free}, \mathrm{rk} < \infty})_{/P}$. By assumption, the map $q^\triangleright(\infty) \rightarrow P$ factors through some R^s for $s \leq r$, where $\infty \in K^\triangleright$ is the cocone point. The result then follows. $\square$

It follows that

Corollary 3.7. Let R be a connective $\mathbb{E}_1$ -ring, $r \in \mathbb{N}$, and P a flat right R -module. Then P is of rank $\leq r$ if and only if so is the flat right $\pi_0(R)$ -module $\pi_0(P)$.

As mentioned in the last paragraph of [Sha70] that, the rank of flat modules coincides with the usual rank over principal ideal domains. Actually, the same argument works over Bézout domains, and in particular, over valuation rings, which we record as follows.

Lemma 3.8. *Let R be a Bézout domain (e.g. a valuation ring), $K := \text{Frac}(R)$, and M a flat right R -module. Then $\text{rk}_R(M) = \dim_K(M \otimes_R K)$.*

Proof. First, suppose that $\text{rk}_R(M) \leq r$ for some $r \in \mathbb{N}$. Then M is a filtered colimit of finite free R -modules of rank $\leq r$, which implies that $M \otimes_R K$ is a filtered colimit of K -vector spaces of dimension $\leq r$, thus $\dim_K(M \otimes_R K) \leq r$.

On the other hand, suppose that $\dim_K(M \otimes_R K) \leq r$. Since M is flat, we know that the canonical map $M \rightarrow M \otimes_R K$ is injective. Given any $(r+1)$ -tuple $(x_1, \dots, x_{r+1}) \in M^{\oplus(r+1)}$, we see that the tuple $(x_1 \otimes 1, \dots, x_{r+1} \otimes 1) \in (M \otimes_R K)^{\oplus(r+1)}$ is K -linear dependent. This implies that, there exists a tuple $(a_1, \dots, a_{r+1}) \in R^{r+1}$ such that $a_1 x_1 + \dots + a_{r+1} x_{r+1} = 0$. Since R is a Bézout domain, we may divide a_i by $\gcd(a_1, \dots, a_{r+1})$, which reduces to the case that $\{a_1, \dots, a_{r+1}\} \subseteq R$ generates the unit ideal.

We now show that the map $f : R^{r+1} \rightarrow M, (b_1, \dots, b_{r+1}) \mapsto \sum x_i b_i$ factors through R^r . Indeed, we note that $(a_1, \dots, a_{r+1}) \in \ker(f)$, which implies that f factors through $Q := R^{r+1}/(a_1, \dots, a_{r+1})R$. However, since $\{a_1, \dots, a_{r+1}\} \subseteq R$ generates the unit ideal, the map $R \rightarrow R^{n+1}, r \mapsto (a_1 r, \dots, a_{r+1} r)$ of R -modules admits a retract. It follows that Q is a proper direct summand of R^{r+1} . Since R is a Bézout domain, we deduce that Q is a finite free R -module of rank r . By [Sha70, Thm 2] (or Proposition 3.6), we see that $\text{rk}_R(M) \leq r$. $\square$

We deduce a descent property of this finite generation condition.

Proposition 3.9. *Let $R \rightarrow S$ be a faithfully flat map of connective $\mathbb{E}_\infty$ -rings with $\text{rk}_R(S) < \infty$, and M a flat R -module. Suppose that the flat S -module $M \otimes_R^{\mathbb{L}} S$ is of finite rank, then so is the flat R -module M , with inequality*

$$\text{rk}_R(M) \leq \text{rk}_R(S) \text{rk}_S(M \otimes_R^{\mathbb{L}} S).$$

Proof. Suppose that $\text{rk}_S(M \otimes_R^{\mathbb{L}} S) \leq r$ and $\text{rk}_R(S) \leq \mu$. By Corollary 3.7, we may assume that R is concentrated in degree 0. Let $(x_1, \dots, x_n) \in M^{\oplus n}$ corresponding to a map $R^n \rightarrow M$ of R -modules. By [Sha70, Thm 2] (or Proposition 3.6), there exists $(y_1, \dots, y_s) \in (M \otimes_R S)^{\oplus s}$ with $s \leq r$, such that $x_i \otimes 1$ lies in $\text{Span}\{y_1, \dots, y_s\}$. Since S is a filtered colimit of finite free R -modules of rank $\leq \mu$, we may choose some map $R^t \rightarrow S$ with $t \leq \mu$, and each $y_i \in M \otimes_R S$ admits a lift $\tilde{y}_i \in M \otimes_R R^{\oplus t} \cong M^{\oplus t}$, and we write $\tilde{y}_i = (z_{i1}, \dots, z_{it})$.

We claim that each x_j belongs to $\text{Span}\{z_{11}, z_{12}, \dots, z_{st}\}$. Indeed, by construction, we have $\tilde{y}_i$ lies in $\text{Span}\{z_{i1} \otimes 1, \dots, z_{it} \otimes 1\} \subseteq M \otimes_R R^{\oplus t}$, and thus $y_i \in \text{Span}\{z_{i1} \otimes 1, \dots, z_{it} \otimes 1\} \subseteq M \otimes_R S$. Consequently, each $x_j \otimes 1 \in M \otimes_R S$ belongs to $\text{Span}\{z_{11} \otimes 1, z_{12} \otimes 1, \dots, z_{st} \otimes 1\}$. The result then follows from the faithful flatness of $R \rightarrow S$. $\square$

Example 3.10 (Zariski descent). Let R be a connective $\mathbb{E}_\infty$ -ring, $d \in \mathbb{N}$, and $(f_1, \dots, f_d) \in \pi_0(R)^d$ a tuple of elements generating the unit ideal of $\pi_0(R)$. Then the map $R \rightarrow \prod_{i=1}^d R_{f_i} =: S$ is a Zariski cover. By Example 3.2 and Remark 3.4, the flat R -module S is of rank $\leq d$.

Let M be a flat R -module such that each localization M_{f_i} is a flat R_{f_i} -module of finite rank. Then by Remark 3.4, the flat S -module $M \otimes_R^{\mathbb{L}} S$ is of finite rank as well, and it follows from Proposition 3.9 that the flat R -module M is of finite rank.

Example 3.11 (Étale descent). Let $R \rightarrow S$ be a faithfully flat étale map of connective $\mathbb{E}_\infty$ -rings. By [Sta21, Tag 00UE], we may choose a Zariski cover $S \rightarrow \prod_{i=1}^d S_{f_i} =: T$ such that each composite $R \rightarrow S \rightarrow S_{f_i}$ is a

standard étale map on π_0 . By Example 3.3, Remark 3.4, and Corollary 3.7 that the flat R -module T is of finite rank.

Let M be a flat R -module such that the flat S -module $M \otimes_R^{\mathbb{L}} S$ is of finite rank. Then by Remark 3.4, the flat T -module $M \otimes_R^{\mathbb{L}} T$ is of finite rank, and it follows from Proposition 3.9 that the flat R -module M is of finite rank as well.

Remark 3.12. There is a variant of Definition 3.1 when R is a connective $\mathbb{E}_\infty$ -rings, where we replace finite free right R -modules of rank $\leq r$ by finite *projective* R -modules of rank $\leq r$. For that version, previously statements adapt, and one can generalize Bézout domains to Prüfer domains in Lemma 3.8.

Now we come back to our weakening of finite generation condition.

Definition 3.13. Let R be a connective $\mathbb{E}_1$ -ring, and $r \in \mathbb{N}$. We say that a connective right R -module M is *weakly finitely generated*⁴ (of rank $\leq r$) if, there exists a flat right R -module P of finite rank (of rank $\leq r$), along with a surjective⁵ map $P \rightarrow M$ of R -modules.

Example 3.14. Let R be a connective $\mathbb{E}_1$ -ring, and M a connective right R -module which is finitely generated (i.e. $\pi_0(M)$ is a finitely generated right $\pi_0(R)$ -module). Then M is weakly finitely generated. This justifies the terminology. Slightly more generally, weak finite generation is a condition on π_0 under countable generation hypothesis. We collect relevant results in Section A.

It follows from Proposition 3.6 that

Corollary 3.15. *Let R be a connective $\mathbb{E}_1$ -ring, and M a flat right R -module. Then M is weakly finitely generated if and only if it is of finite rank.*

A merit of this finite generation condition is an asymptotic connectivity of exterior powers.

Lemma 3.16. *Let R be an animated ring, and M a connective R -module weakly finitely generated. Then the R -module spectra $\left\{ \left(\bigwedge_R^j M \right) [-j] \mid j \in \mathbb{N} \right\}$ are uniformly t -bounded below, and the bound only depends on the rank of M .*

Proof. By definition, there exists a natural number $r \in \mathbb{N}$, a flat R -module P which is a filtered colimit of finite free R -modules of rank $\leq r$, and a surjective map $P \rightarrow M$ of connective R -modules. Let $K := \text{fib}(P \rightarrow M)$, which is connective as well, and we get a fiber sequence $P \rightarrow M \rightarrow K[1]$. It follows that, for every $j \in \mathbb{N}$, the R -module spectrum $\left(\bigwedge_R^j M \right) [-j]$ admits a finitary filtration (cf. [Ill71, §V.4]), whose associated graded pieces are given by

$$\left(\bigwedge_R^i P \right) \otimes_R^{\mathbb{L}} \left(\bigwedge_R^{j-i} K[1] \right) [-j] \simeq \left(\bigwedge_R^i P \right) \otimes_R^{\mathbb{L}} (\Gamma_R^{j-i} K) [-i],$$

where we used Quillen and Illusie's décalage [Ill71, I.4.3.2.1], see also [Lur18, Prop 25.2.4.2]. Note that $\bigwedge_R^i P = 0$ for $i > r$ by Lazard's theorem [Lur17, Thm 7.2.2.15], as $\bigwedge_R^i(-)$ preserves filtered colimits of R -module spectra, and it vanishes on finite free R -modules of rank $\leq r$. It follows that $\left(\bigwedge_R^j M \right) [-j]$ is $(-r)$ -connective. $\square$

⁴As pointed out by ChatGPT-6 Astra, our concept is different from the one in [BPŠ22, §8.2] under the same name. In the special case of R being a commutative chain ring and M being an R -module concentrated in degree 0, this coincides with *Fleischer rank* in [Cou13, Rem 3.1].

⁵A map of connective R -modules is *surjective* if it is surjective on π_0 .

Let R be an animated ring. [BL22, Rem 5.5.9] showed that $\mathrm{gr}_N^m \Delta_R\{n\}$ is bounded below. Combining with Lemma 3.16, we adapt the same argument to show that, under such a weak finite generation, they are uniformly bounded below. Recall that the $(p$ -completed) diffracted Hodge cohomology $\Omega_R^\mathcal{D}$ admits an increasing filtration $\mathrm{Fil}_*^{\mathrm{conj}} \Omega_R^\mathcal{D}$, called the *conjugate filtration*, [BL22, Cons 4.7.1]. Connectivity of exterior powers gives rise to that of several cohomology theories:

Lemma 3.17. *Let R be an animated ring such that the spectra*

$$\left\{ \left(\bigwedge_R^n L_{R/\mathbb{Z}} \right) [-n] \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \mid n \in \mathbb{Z} \right\}$$

is uniformly t -bounded below. Then the R -module spectra

$$\left\{ \mathrm{Fil}_s^{\mathrm{conj}} \Omega_R^\mathcal{D} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \mid s \in \mathbb{N} \right\}$$

are uniformly t -bounded below, and the bound is the same as the previous bound.

Proof. [BL22, Rem 4.5.3] identifies the associated graded pieces $\mathrm{gr}_*^{\mathrm{conj}} \Omega_R^\mathcal{D}$ with $(p$ -completed) shifted exterior powers $(\bigwedge_R^* L_{R/\mathbb{Z}})[-*]$, and the result then follows. $\square$

Corollary 3.18. *Let R be an animated ring such that the spectra*

$$\left\{ \left(\bigwedge_R^n L_{R/\mathbb{Z}} \right) [-n] \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \mid n \in \mathbb{Z} \right\}$$

is uniformly t -bounded below. Then the R -module spectra

$$\left\{ \mathrm{gr}_N^m \Delta_R\{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \mid (m, n) \in \mathbb{Z} \times \mathbb{Z} \right\}$$

and

$$\left\{ \mathrm{Fil}_m^{\mathrm{conj}} \overline{\Delta}_R\{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \mid (m, n) \in \mathbb{Z} \times \mathbb{Z} \right\}$$

are uniformly t -bounded below, and the bound only depends on the previous uniform t -bound. In particular, the spectra $\{\widehat{\Delta}_R\{n\} \mid n \in \mathbb{Z}\}$ and $\{\Delta_R\{n\} \mid n \in \mathbb{Z}\}$ are p -completely⁶ uniformly t -bounded below.

Proof. It follows from Lemma 3.17 and the fiber sequences

$$\mathrm{gr}_N^m \Delta_R\{n\} \longrightarrow \mathrm{Fil}_m^{\mathrm{conj}} \Omega_R^\mathcal{D} \xrightarrow{\Theta+m} \mathrm{Fil}_{m-1}^{\mathrm{conj}} \Omega_R^\mathcal{D}$$

and

$$\mathrm{Fil}_m^{\mathrm{conj}} \overline{\Delta}_R\{n\} \longrightarrow \mathrm{Fil}_m^{\mathrm{conj}} \Omega_R^\mathcal{D} \xrightarrow{\Theta+n} \mathrm{Fil}_m^{\mathrm{conj}} \Omega_R^\mathcal{D}$$

as established in [BL22, Rem 4.7.5 & 5.5.8]. $\square$

These connectivity estimates are sufficient for establishing the completeness of motivic filtration on $\mathrm{TP}_{/\mathbb{Z}}^{\mathrm{poly}}$. Recall that it is exhaustive (Corollary 2.7).

Theorem 3.19. *Let R be an animated ring such that the spectra*

$$\left\{ \left(\bigwedge_R^n L_{R/\mathbb{Z}} \right) [-n] \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \mid n \in \mathbb{Z} \right\}$$

is uniformly t -bounded below. Then $\mathrm{Fil}_M^n (\mathrm{THH}_{\Delta}(R)^{\Phi C_p})_p^{\wedge}$ is linearly increasingly connective as $n \rightarrow \infty$, and the motivic filtration on $\mathrm{TP}_{/\mathbb{Z}}^{\mathrm{poly}}(R)$ is complete.

⁶That is to say, they are uniformly t -bounded below after $(-)\otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$.

Proof. Recall that the motivic filtration on $\mathrm{THH}(R)$ is complete [BL22, Thm 6.2.4 & Prop 6.2.12]. By Remark 2.5, we are reduced to show that, after inverting μ_1 in $\mathrm{Fil}_M(\mathrm{THH}(R) \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$, the motivic filtration remains complete. We prove this by connectivity. Indeed, the r -th filtration piece of $\mathrm{Fil}_M(\mathrm{THH}(R) \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)[\mu_1^{-1}]$ is the sequential colimit of

$$\mathrm{Fil}_M^r(\mathrm{THH}(R) \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p) \xrightarrow{\mu_1} \mathrm{Fil}_M^{r+p}(\mathrm{THH}(R) \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)[-2p] \xrightarrow{\mu_1} \dots$$

We note that $\{\mathrm{Fil}_M^n \mathrm{THH}(R)[-2n] | n \in \mathbb{Z}\}$ is p -completely t -bounded below by motivic completeness on $\mathrm{THH}(R)$ and Corollary 3.18. It follows that

$$\{\mathrm{Fil}_M^{r+n} \mathrm{THH}(R)[-2n]\}_{n \in \mathbb{Z}} = \{\Sigma^{2r} \mathrm{Fil}_M^n \mathrm{THH}(R)[-2n]\}_{n \in \mathbb{N}}$$

is p -completely uniformly t -bounded below, and the given bound tends to ∞ as $r \rightarrow \infty$ with linear growth. In particular, the motivic filtration on $(\mathrm{THH}_{\Delta}(R)^{\Phi_{C_p}})_p^{\wedge}$ is complete, and it remains complete after taking $(-)^{h\mathrm{Te}_v}$. $\square$

This also implies that, when R is additionally F -smooth à la Bhatt–Mathew [BM23], the cyclotomic spectrum $\mathrm{THH}(R)/(p, v_1)$ satisfies Segal conjecture relative to $\mathrm{THH}(\mathbb{Z})/(p, v_1)$, by passing to associated graded pieces of motivic filtrations. Actually, the same argument implies that absolute Segal conjecture holds, which is slightly stronger by Proposition 2.16.

Theorem 3.20. *Let R be an F -smooth p -quasisyntomic commutative ring. Suppose such that the spectra*

$$\left\{ \left(\bigwedge_R^n L_{R/\mathbb{Z}} \right) [-n] \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p | n \in \mathbb{Z} \right\}$$

is uniformly t -bounded below. Then the fiber $\mathrm{fib}(\mathrm{THH}(R) \rightarrow \mathrm{THH}(R)^{tC_p})$ of the cyclotomic Frobenius map is p -completely t -bounded above.

Proof. [BM23, Rem 4.11] showed⁷ that the filtered Frobenius map

$$\mathrm{Fil}_N^* \Delta_R \longrightarrow \Delta_R^{[*]}$$

induces an equivalence

$$\mathrm{gr}_N^i \Delta_R \longrightarrow \tau_{\geq -i} \mathrm{gr}^i \Delta_R^{[*]} \simeq \tau_{\geq -i} \overline{\Delta}_R \{i\}.$$

By Corollary 3.18, the Hodge–Tate cohomology $\overline{\Delta}_R \{i\}$ for all i ’s are uniformly t -bounded below. It follows that, the map

$$\mathrm{gr}_M^* \mathrm{THH}(R) \longrightarrow \mathrm{gr}_M^* \mathrm{THH}(R)^{tC_p}$$

is an equivalence after p -completion as $* \gg 0$. The result then follows from the completeness of motivic filtrations on THH and THH^{tC_p} , [BL22, Thm 6.2.4 & Prop 6.2.12] along with the trick of tensoring with $\mathbb{T}/C_p$ as in [Rig25, Lem 3.8]. $\square$

We give some special cases of Theorems 3.19 and 3.20. We learned the following result from Daniel FINK.

Lemma 3.21. *Let $k \rightarrow R$ be a map of animated $\mathbb{F}_p$ -algebra. Then the R -module spectrum $L_{R/k}$ can be realized, functorially in maps $(k \rightarrow R) \in \mathrm{Fun}(\Delta^1, \mathrm{CAlg}_{\mathbb{F}_p}^{\mathrm{an}})$ of animated $\mathbb{F}_p$ -algebras, as a retract of the cotangent complex $L_{R/\varphi_k^* R}$ of the relative Frobenius map $\varphi_{R/k} : \varphi_k^* R \rightarrow R$, viewed as a module over the target.*

⁷We thank Hyungseop KIM for pointing this out to us.

Proof. This is implicit in the proof of [BF26, Lem 2.19]⁸. We briefly review their argument. We have the transitivity sequence

$$L_{\varphi_k^* R/k} \otimes_{\varphi_k^* R, \varphi_{R/k}}^{\mathbb{L}} R \longrightarrow L_{R/k} \longrightarrow L_{R/\varphi_k^* R}$$

associated to the composite map $k \rightarrow \varphi_k^* R \xrightarrow{\varphi_{R/k}} R$ of animated rings. It remains to specify a null-homotopy of the first map. Indeed, the first map is zero on standard maps $k \rightarrow R$ of the form $\mathbb{F}_p[x_1, \dots, x_m] \rightarrow \mathbb{F}_p[x_1, \dots, x_m, y_1, \dots, y_n]$ for $(m, n) \in \mathbb{N}^2$, which gives rise to a null-homotopy of the first map functorially in such standard maps $k \rightarrow R$. We then left Kan extend this null homotopy to all maps $k \rightarrow R$ of animated $\mathbb{F}_p$ -algebras. $\square$

We introduce the following weakened version of F -finiteness.

Definition 3.22. We say that an animated $\mathbb{F}_p$ -algebra R is *weakly F -finite* if the Frobenius map $\varphi : R \rightarrow R$ realizes the target as a weakly finitely generated module over the source.

This weakened F -finiteness gives rise to a weakened finite generation, just as usual F -finiteness:

Lemma 3.23. *Let $R \rightarrow S$ be a map of animated rings. Suppose that, as an R -modules, S is weakly finitely generated. Then the cotangent complex $L_{S/R}$ is weakly finitely generated.*

Proof. Pick a surjection $P \rightarrow S$ where P is a flat R -module of rank $\leq r$. We write P as a filtered colimit $\operatorname{colim}_{i \in \mathcal{I}} F_i$ of finite free R -modules of rank $\leq r$. Then we get a surjection

$$\operatorname{colim}_{i \in \mathcal{I}} \operatorname{LSym}_R(F_i) \longrightarrow S$$

of animated R -algebras, which gives rise to a surjection

$$\operatorname{colim}_{i \in \mathcal{I}} L_{\operatorname{LSym}_R(F_i)/R} \otimes_{\operatorname{LSym}_R(F_i)}^{\mathbb{L}} S \longrightarrow L_{S/R}$$

of S -module spectra, where the source is a filtered colimit of finite free S -modules of rank $\leq r$. The result then follows. $\square$

Corollary 3.24. *Let A be a weakly F -finite animated $\mathbb{F}_p$ -algebra. Denote by $\varphi : A \rightarrow A$ its Frobenius map. Then the cotangent complex $L_{A/\mathbb{F}_p}$ is weakly finitely generated.*

Proof. It follows from Lemmas 3.21 and 3.23. $\square$

Example 3.25. Let R be an animated ring such that the animated $\mathbb{F}_p$ -algebra $R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ is weakly F -finite. It then follows from Corollary 3.24, Lemma 3.16, and Theorem 3.19 that the motivic filtration on $\operatorname{TP}_{/\mathbb{Z}}^{\operatorname{poly}}(R)$ is complete.

Example 3.26. Let R be an F -smooth p -quasisyntomic ring such that the animated ring $\mathbb{F}_p$ -algebra $R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ is weakly F -finite. By Corollary 3.24, Lemma 3.16, and Theorem 3.20, we see that $\operatorname{THH}(R)_p^{\wedge}$ satisfies Segal conjecture. This includes qualitative versions for both smooth algebras over a perfectoid ring [Mat21, Prop 5.10] and regular rings whose reduction mod p is F -finite ([Mat21, Cor 5.3] in the excellent case, and [BM23, Cor 4.20] in general).

Example 3.27. We give an example of a Cartier smooth $\mathbb{F}_p$ -algebra which is weakly F -finite but not F -finite. Let A be a smooth $\mathbb{F}_p$ -algebra, and $S \subseteq A$ a multiplicative subset. Then the subring $R := A + t(S^{-1}A)[t] \subseteq (S^{-1}A)[t]$ is Cartier smooth, as it is the filtered colimit $\operatorname{colim}_{s \in S} A[s^{-1}t]$. The R -module $L_{R/\mathbb{F}_p}$ is not finitely generated in general (say, $A = \mathbb{F}_p[x]$ and $S = \{x^n | n \in \mathbb{N}\}$), but R is weakly F -finite.

⁸This turns out to be implicit in [GR03, Lem 6.5.13(i)] as well.

Example 3.28 (non F -finite DVR). Let k be a perfect field of char p , and let $f = \sum_{n=1}^{\infty} a_n t^n \in k[[t]]$ be a formal power series with zero constant coefficient which is transcendental over $k[t]$. Let $f_j := \sum_{n=1}^j a_n t^n$, and

$$\begin{aligned} u_j &:= \frac{(f - f_j)^p}{t^j} \\ &= \sum_{n=j+1}^{\infty} a_n^p t^{np-j} \end{aligned}$$

for $j \in \mathbb{N}$. Then the localization $k[t, u_0, \dots, u_j, \dots]_{(t, u_0, \dots, u_j, \dots)} \subseteq k[[t]]$ is a DVR, but it is not excellent, as established in [Val73]. However, it is weakly F -finite, being a filtered union of the subrings $k[t, u_0]_{(t, u_0)} \subseteq k[t, u_1]_{(t, u_1)} \subseteq k[t, u_2]_{(t, u_2)} \subseteq \dots$ (note that we have $tu_{j+1} - u_j \in tk[[t]]$).

Example 3.29 (non weakly F -finite DVR). We point out that there exists a DVR of char p with perfect residue field, which is not weakly F -finite over $\mathbb{F}_p$. This is the sort of DVR constructed in [DS18, §4.1] (cf. [Li26, §2]). Indeed, let $k := \overline{\mathbb{F}_p}$ and $K := k(x_1, \dots, x_n, \dots)$. There an embedding $\iota : K \rightarrow k((t))$ of k -algebras obtained by assigning the image of x_n recursively via a cardinality argument in [DS18, Rem 4.3]. Let $V_p := k[[t]] \cap \iota(K)$ with $\mathfrak{m}_{V_p} := tk[[t]] \cap \iota(K)$. Then $(V_p, \mathfrak{m}_{V_p})$ is a DVR over k with residue field being $V_p/\mathfrak{m}_{V_p} = k$ and fractional field $\text{Frac}(V_p)$ being $\iota(K) \cong K$, as explained in [DS18, §4.1]. It follows that

$$L_{V_p/\mathbb{F}_p} \otimes_{V_p}^{\mathbb{L}} K = L_{K/\mathbb{F}_p} = \bigoplus_{n=1}^{\infty} K dx_i,$$

and thus by Corollary 3.15 and Lemma 3.8 and the flatness of the V_p -module $L_{V_p/\mathbb{F}_p}$ established in [BLM18, Prop 9.5.1], the V_p -module $L_{V_p/\mathbb{F}_p}$ is not weakly finitely generated, and by Corollary 3.24, the $\mathbb{F}_p$ -algebra V_p is not weakly F -finite over $\mathbb{F}_p$.

4. CRYSTALLINE DEGENERATION OF SEGAL CONJECTURE

In [HW22], the authors proved Segal conjecture for $\text{THH}(\text{BP}\langle n \rangle)$. In this section, we show that their argument implies a general statement (Proposition 4.2) which allows us to deduce Segal conjecture of a “formal $\text{BP}\langle n \rangle$ -scheme” from Segal conjecture for its special fiber over $\mathbb{F}_p$. We refer to this argument as *crystalline degeneration* by virtue of its resemblance to [AKN24, §2.2].

First, the proof of [HRW22, Prop C.5.4] actually shows that

Lemma 4.1. *Let A be a connective $\mathbb{E}_1$ -ring. Then for every $n \in \mathbb{Z}$, the filtered object*

$$\pi_n \left(\text{THH} \left(\text{desc}_{\mathbb{F}_p}^{\geq *}(A) \right) / v_0 \right)$$

is eventually constant, converging to $\text{THH}(A)_p^{\wedge}$.

Proposition 4.2. *Let M be a right $\text{THH}(\text{BP}\langle n \rangle)$ -module in genuine C_p -spectra, such that $M/(p, v_1, \dots, v_n)$ is t -bounded below. Suppose that the canonical map*

$$M^{\Phi C_p} \otimes_{\text{THH}(\text{BP}\langle n \rangle)}^{\mathbb{L}} \text{THH}(\mathbb{F}_p) \longrightarrow (M \otimes_{\text{THH}(\text{BP}\langle n \rangle)}^{\mathbb{L}} \text{THH}(\mathbb{F}_p))^{tC_p}$$

is an equivalence on π_ for $* \gg 0$. Then so is the map*

$$M^{\Phi C_p} \otimes_{\text{BP}\langle n \rangle}^{\mathbb{L}} \mathbb{F}_p \longrightarrow M^{tC_p} \otimes_{\text{BP}\langle n \rangle}^{\mathbb{L}} \mathbb{F}_p$$

induced by the canonical map $M^{\Phi C_p} \rightarrow M^{tC_p}$.

Proof. We endow $\mathrm{BP}\langle n \rangle$ the $\mathbb{N}$ -graded decreasing filtration $\mathrm{desc}_{\mathbb{F}_p}^{\geq *}(\mathrm{BP}\langle n \rangle)$ coming from the Adams spectral sequence as in [HW22, Rec 4.3.2], with

$$\pi_* \left(\mathrm{gr} \left(\mathrm{desc}_{\mathbb{F}_p}^{\geq *}(\mathrm{BP}\langle n \rangle) \right) \right) = \mathbb{F}_p[v_0, \dots, v_n],$$

where $\mathrm{wt}(v_i) = 2p^i - 1$. Then the cyclotomic spectrum $\mathrm{THH}(\mathrm{BP}\langle n \rangle)$ is equipped with an $\mathbb{N}$ -graded decreasing filtration $\mathrm{Fil}_{\mathrm{cris}}^*$ as well, concretely given by

$$\mathrm{Fil}_{\mathrm{cris}}^* \mathrm{THH}(\mathrm{BP}\langle n \rangle) = \mathrm{THH} \left(\mathrm{desc}_{\mathbb{F}_p}^{\geq *}(\mathrm{BP}\langle n \rangle) \right).$$

This induces an $\mathbb{N}$ -graded decreasing filtration $\mathrm{Fil}_{\mathrm{cris}}^*$ on the canonical map $M^{\Phi C_p} \rightarrow M^{tC_p}$, and it suffices to see that the filtered map (for convenience, we omit the grading shear L_p on the source)

$$(2) \quad \mathrm{Fil}_{\mathrm{cris}}^* M^{\Phi C_p} \otimes_{\mathrm{desc}_{\mathbb{F}_p}^{\geq *}(\mathrm{BP}\langle n \rangle)}^{\mathbb{L}} \mathbb{F}_p \longrightarrow \mathrm{Fil}_{\mathrm{cris}}^* M^{tC_p} \otimes_{\mathrm{desc}_{\mathbb{F}_p}^{\geq *}(\mathrm{BP}\langle n \rangle)}^{\mathbb{L}} \mathbb{F}_p$$

is an equivalence in high enough degrees.

We first show this after passing to associated graded pieces. For this, we do not need the bounded-below-ness of the genuine C_p -spectrum M . Indeed, let $M_{\mathbb{F}_p} := M \otimes_{\mathrm{THH}(\mathrm{BP}\langle n \rangle)}^{\mathbb{L}} \mathrm{THH}(\mathbb{F}_p)$. The associated graded pieces of the map (2) is the composite graded map

$$\begin{aligned} & M_{\mathbb{F}_p}^{\Phi C_p} \otimes_{\mathrm{THH}(\mathbb{F}_p)}^{\mathbb{L}} \mathrm{THH}(\mathbb{F}_p[v_0, \dots, v_n]) / (v_0, \dots, v_n) \\ \longrightarrow & M_{\mathbb{F}_p}^{tC_p} \otimes_{\mathrm{THH}(\mathbb{F}_p)}^{\mathbb{L}} \mathrm{THH}(\mathbb{F}_p[v_0, \dots, v_n]) / (v_0, \dots, v_n) \\ \longrightarrow & (M_{\mathbb{F}_p} \otimes_{\mathrm{THH}(\mathbb{F}_p)}^{\mathbb{L}} \mathrm{THH}(\mathbb{F}_p[v_0, \dots, v_n]))^{tC_p} / (v_0, \dots, v_n). \end{aligned}$$

The first map is an equivalence in high enough degrees, as it is tensoring the canonical map $M_{\mathbb{F}_p}^{\Phi C_p} \rightarrow M_{\mathbb{F}_p}^{tC_p}$, which is an equivalence in high enough degrees by assumption, with $\mathrm{THH}(\mathbb{S}[v_0, \dots, v_n]) / (v_0, \dots, v_n)$. The second map is an equivalence by [HW22, Lem 4.1.3] (see the last paragraph of the proof of [HW22, Prop 4.2.2]).

We now show the completeness of filtrations on the source and the target of the filtered map (2). By Lemma 4.1 and that the genuine C_p -spectrum M is bounded below, the filtration $\mathrm{Fil}_{\mathrm{cris}}^*$ is $(p, v_1, \dots, v_n)$ -completely eventually constant on $M^{\Phi C_p}$ and M , thus the filtration $\mathrm{Fil}_{\mathrm{cris}}^*$ is complete on M^{tC_p} by [HW22, Cor C.4.3]. The result then follows from the equivalence

$$\mathrm{desc}_{\mathbb{F}_p}^{\geq *}(\mathrm{BP}\langle n \rangle) / (v_0, \tilde{v}_1, \dots, \tilde{v}_n) \simeq \mathbb{F}_p$$

as in the proof of [HW22, Thm 4.3.1]. $\square$

Example 4.3. Let R be a (-1) -connective $\mathbb{E}_1$ - $\mathbb{Z}$ -algebra. Suppose that the cyclotomic spectrum $\mathrm{THH}(R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ satisfies Segal conjecture. Then Proposition 4.2 implies that, the cyclotomic spectrum $\mathrm{THH}(R)$ satisfies Segal conjecture p -completely. This can be viewed as a shadow of the fact that, a p -quasisyntomic p -torsion-free commutative ring R is F -smooth if the commutative $\mathbb{F}_p$ -algebra R/p is Cartier smooth, [BM23, Cor 4.17].

5. RELATIVE CONJUGATE FILTRATION

In this section, we will present a general procedure (Construction 5.1), which allows us to produce a relative conjugate filtration (Example 5.4) on Hodge–Tate cohomology $\overline{\Delta}_R$ for any map $k \rightarrow R$ of animated rings, a mixed characteristic analogue of [Bha12, Prop 3.22]. In the same time, it also allows us to produce a relative conjugate filtration (Example 5.5) on both $\mathrm{THH}_{\Delta}(\mathcal{C})^{\Phi C_p}$ and $\mathrm{THH}(\mathcal{C}) \otimes_{\mathrm{THH}(k)}^{\mathbb{L}} \mathrm{THH}(k)^{tC_p}$ after p -completion, for every animated ring k and every k -linear DG-category $\mathcal{C}$, mixed characteristic and relative versions of Kaledin’s conjugate filtration. We also discuss the completeness of these filtrations (Propositions 5.7 and 5.8).

Construction 5.1. To simplify notation, everything is p -completed. Let k be an animated ring. The theory of synthetic THH as in [AR24] gives rise to an $\mathbb{E}_\infty$ -algebra

$$\mathrm{Fil}_M^{\geq 0}(\mathrm{THH}(k) \otimes_{\mathrm{THH}(\mathbb{Z})}^{\mathbb{L}} \mathrm{THH}(\mathbb{Z})^{tC_p}) = \mathrm{Fil}_M^{\geq 0} \mathrm{THH}_\Delta(k)^{\Phi C_p}$$

in SynSp . This gives rise to a $\mathbb{N}$ -filtered $\mathbb{E}_\infty$ -algebra⁹ $\mathrm{Fil}_{\mathrm{KO}}^* \mathrm{Fil}_M^{\geq 0} \mathrm{THH}_\Delta(k)^{\Phi C_p}$ in SynSp , concretely given by

$$\mathrm{Fil}_{\mathrm{KO}}^j \mathrm{Fil}_M^{\geq 0} \mathrm{THH}_\Delta(k)^{\Phi C_p} := \mathrm{Fil}_M^{\geq j} \mathrm{THH}_\Delta(k)^{\Phi C_p} \in \mathrm{SynSp}$$

for $j \in \mathbb{N}$, where given a filtered object $\mathrm{Fil} W$, the filtered object of $\mathrm{Fil}^{\geq j} W$ is

$$\cdots \xleftarrow{\simeq} \mathrm{Fil}_M^j W \xleftarrow{\simeq} \mathrm{Fil}_M^j W \leftarrow \mathrm{Fil}_M^{j+1} W \leftarrow \cdots,$$

that is, first restricting to $\mathbb{N}_{\geq j}$ then left Kan extending along $(\mathbb{N}_{\geq j}, \geq) \rightarrow (\mathbb{N}, \geq)$.

Let N be a $\mathrm{Fil}_M^{\geq 0} \mathrm{THH}_\Delta(k)^{\Phi C_p}$ -module in SynSp . The $\mathbb{N}$ -graded filtration on $\mathrm{Fil}_M^{\geq 0} \mathrm{THH}_\Delta(k)^{\Phi C_p}$ gives rise to an $\mathbb{Z}$ -graded filtration on

$$N' := N \otimes_{\mathrm{Fil}_M^{\geq 0} \mathrm{THH}_\Delta(k)^{\Phi C_p}}^{\mathbb{L}} \mathrm{Fil}_M \mathrm{THH}_\Delta(k)^{\Phi C_p},$$

given by

$$\mathrm{Fil}_{\mathrm{KO}}^j N' := N \otimes_{\mathrm{Fil}_M^{\geq 0} \mathrm{THH}_\Delta(k)^{\Phi C_p}}^{\mathbb{L}} \mathrm{Fil}_M^{\geq j} \mathrm{THH}_\Delta(k)^{\Phi C_p} \in \mathrm{SynSp}$$

for $j \in \mathbb{Z}$. We see that

$$\begin{aligned} \mathrm{gr}_{\mathrm{KO}}^j N' &= N \otimes_{\mathrm{Fil}_M^{\geq 0} \mathrm{THH}_\Delta(k)^{\Phi C_p}}^{\mathbb{L}} \mathrm{ins}^j \mathrm{gr}_M^j \mathrm{THH}(k)^{tC_p} \\ &= N \otimes_{\mathrm{Fil}_M^{\geq 0} \mathrm{THH}_\Delta(k)^{\Phi C_p}}^{\mathbb{L}} \mathrm{ins}^j \widehat{\Delta}_k\{j\}[2j] \in \mathrm{SynSp}, \end{aligned}$$

where ins^j is the left Kan extension along $\{j\} \hookrightarrow (\mathbb{N}, \geq)$, see [Rak26, Not 3.1.1]. Moreover, unwinding the construction, we see that the $\mathrm{Fil}_M^{\geq 0} \mathrm{THH}_\Delta(k)^{\Phi C_p}$ -module structure on $\mathrm{ins}^j \widehat{\Delta}_k\{j\}[2j]$ is induced by the $\mathrm{gr}_M^0 \mathrm{THH}_\Delta(k)^{\Phi C_p} = \widehat{\Delta}_k$ -module structure on $\widehat{\Delta}_k\{j\}[2j]$.

We may replace $\mathrm{THH}_\Delta(k)^{\Phi C_p}$ by $\mathrm{THH}(k)^{tC_p}$ throughout, and for every $\mathrm{Fil}_M^{\geq 0} \mathrm{THH}(k)^{tC_p}$ -module N in SynSp , we get an $\mathbb{Z}$ -graded filtration on

$$N' := N \otimes_{\mathrm{Fil}_M^{\geq 0} \mathrm{THH}(k)^{tC_p}}^{\mathbb{L}} \mathrm{Fil}_M \mathrm{THH}(k)^{tC_p},$$

with associated graded pieces

$$\mathrm{gr}_{\mathrm{KO}}^j N' = N \otimes_{\mathrm{Fil}_M^{\geq 0} \mathrm{THH}(k)^{tC_p}}^{\mathbb{L}} \mathrm{ins}^j \widehat{\Delta}_k\{j\}[2j] \in \mathrm{SynSp}$$

for $j \in \mathbb{Z}$.

Example 5.2. To simplify notation, everything is p -completed. Let k be an animated ring, and M a $\mathrm{Fil}_M \mathrm{THH}(k)$ -module in SynSp . Apply Construction 5.1 to the $\mathrm{Fil}_M \mathrm{THH}_\Delta(k)^{\Phi C_p}$ -module

$$M \otimes_{\mathrm{Fil}_M \mathrm{THH}(k)}^{\mathbb{L}} \mathrm{Fil}_M^{\geq 0} \mathrm{THH}_\Delta(k)^{\Phi C_p}$$

in SynSp , we get a filtered synthetic spectrum

$$\mathrm{Fil}_{\mathrm{KO}}^* \left(M \otimes_{\mathrm{Fil}_M \mathrm{THH}(k)}^{\mathbb{L}} \mathrm{Fil}_M \mathrm{THH}_\Delta(k)^{\Phi C_p} \right)$$

⁹Along with the “internal” filtration of synthetic spectra, this can be seen as a bi-filtered object, but we prefer to view it as a filtered synthetic spectrum. In addition, the subscript KO is an abbreviation for Katz–Oda, as the procedure here is the same as the procedure of producing Katz–Oda filtration on derived de Rham cohomology.

with j -th associated graded pieces given by

$$\begin{aligned}
& M \otimes_{\mathrm{Fil}_M \mathrm{THH}(k)}^{\mathbb{L}} \mathrm{ins}^j \overline{\Delta}_k \{j\} [2j] \\
&= (M \otimes_{\mathrm{Fil}_M \mathrm{THH}(k)}^{\mathbb{L}} \mathrm{ins}^0 \mathrm{gr}_M^0 \mathrm{THH}(k)) \otimes_{\mathrm{ins}^0 \mathrm{gr}_M^0 \mathrm{THH}(k)}^{\mathbb{L}} \mathrm{ins}^j \overline{\Delta}_k \{j\} [2j] \\
&= (M \otimes_{\mathrm{Fil}_M \mathrm{THH}(k)}^{\mathbb{L}} \mathrm{ins}^0 k) \otimes_{\mathrm{ins}^0 k}^{\mathbb{L}} \mathrm{ins}^j \overline{\Delta}_k \{j\} [2j],
\end{aligned}$$

for $j \in \mathbb{Z}$, where in the second equality, we take advantage of the commutative diagram

$$\begin{array}{ccc}
\mathrm{Fil}_M \mathrm{THH}(k) & \longrightarrow & \mathrm{Fil}_M \mathrm{THH}_{\Delta}(k)^{\Phi_{C_p}} \\
\downarrow & & \downarrow \\
\mathrm{ins}^0 \mathrm{gr}_M^0 \mathrm{THH}(k) & \longrightarrow & \mathrm{ins}^0 \mathrm{gr}_M^0 \mathrm{THH}_{\Delta}(k)^{\Phi_{C_p}}
\end{array}$$

of $\mathbb{E}_{\infty}$ -algebras of synthetic spectra. Unwinding the construction, we see that the $\mathrm{ins}^0 k$ -module structure on $\mathrm{ins}^j \overline{\Delta}_k \{j\} [2j]$ is given by the canonical map $\varphi_0 : k \rightarrow \overline{\Delta}_k$. Similarly, we get a filtered synthetic spectrum

$$\mathrm{Fil}_{\mathrm{KO}}^*(M \otimes_{\mathrm{Fil}_M \mathrm{THH}(k)}^{\mathbb{L}} \mathrm{Fil}_M \mathrm{THH}(k)^{t_{C_p}})$$

with j -th associated graded piece given by

$$(M \otimes_{\mathrm{Fil}_M \mathrm{THH}(k)}^{\mathbb{L}} \mathrm{ins}^0 k) \otimes_{\mathrm{ins}^0 k}^{\mathbb{L}} \mathrm{ins}^j \widehat{\overline{\Delta}}_k \{j\} [2j]$$

for $j \in \mathbb{Z}$.

Notation 5.3. Let $k \rightarrow R$ be a map of animated ring, and $\mathcal{C}$ a dualizable presentable stable k -linear ∞ -category. We will denote by $\mathrm{THH}_{\Delta/k}(\mathcal{C})^{\Phi_{C_p}}$ the $\mathbb{T}$ -equivariant spectrum

$$\mathrm{THH}(\mathcal{C}) \otimes_{\mathrm{THH}(k)}^{\mathbb{L}} \mathrm{THH}(k)^{t_{C_p}},$$

and by $\mathrm{Fil}_M \mathrm{THH}_{\Delta/k}(R)^{\Phi_{C_p}}$ the $\mathbb{T}_{\mathrm{ev}}$ -equivariant synthetic spectrum

$$\mathrm{Fil}_M \mathrm{THH}(R) \otimes_{\mathrm{Fil}_M \mathrm{THH}(k)}^{\mathbb{L}} \mathrm{Fil}_M \mathrm{THH}(k)^{t_{C_p}}.$$

Example 5.4. To simplify notation, everything is p -completed. Let $k \rightarrow R$ be a map of animated rings. Take $M = \mathrm{Fil}_M \mathrm{THH}(R)$ in Example 5.2, we get a filtered synthetic spectrum¹⁰ $\mathrm{Fil}_{\mathrm{KO}/k} \mathrm{Fil}_M \mathrm{THH}_{\Delta}(R)^{\Phi_{C_p}}$, with

$$\begin{aligned}
\mathrm{gr}_{\mathrm{KO}/k}^j \mathrm{Fil}_M \mathrm{THH}_{\Delta}(R)^{\Phi_{C_p}} &= (\mathrm{Fil}_M \mathrm{THH}(R) \otimes_{\mathrm{Fil}_M \mathrm{THH}(k)}^{\mathbb{L}} \mathrm{ins}^0 k) \otimes_{\mathrm{ins}^0 k}^{\mathbb{L}} \mathrm{ins}^j \overline{\Delta}_k \{j\} [2j] \\
&= \mathrm{Fil}_M \mathrm{HH}(R/k) \otimes_{\mathrm{ins}^0 k}^{\mathbb{L}} \mathrm{ins}^j \overline{\Delta}_k \{j\} [2j] \in \mathrm{SynSp},
\end{aligned}$$

and a filtered synthetic spectrum $\mathrm{Fil}_{\mathrm{KO}/k} \mathrm{Fil}_M \mathrm{THH}_{\Delta/k}(R)^{\Phi_{C_p}}$, with

$$\mathrm{gr}_{\mathrm{KO}/k}^j \mathrm{Fil}_M \mathrm{THH}_{\Delta/k}(R)^{\Phi_{C_p}} = \mathrm{Fil}_M \mathrm{HH}(R/k) \otimes_{\mathrm{ins}^0 k}^{\mathbb{L}} \mathrm{ins}^j \widehat{\overline{\Delta}}_k \{j\} [2j] \in \mathrm{SynSp}.$$

Passing to the i th associated graded piece of the synthetic spectra, we get filtered objects $\mathrm{Fil}_{\mathrm{KO}/k} \overline{\Delta}_R \{i\} [2i]$, with

$$\begin{aligned}
\mathrm{gr}_{\mathrm{KO}/k}^j \overline{\Delta}_R \{i\} [2i] &= \left(\bigwedge_R^{i-j} L_{R/k} \right) [i-j] \otimes_k^{\mathbb{L}} \overline{\Delta}_k \{j\} [2j] \\
&= \left(\bigwedge_R^{i-j} L_{R/k} \right) \otimes_k^{\mathbb{L}} \overline{\Delta}_k \{j\} [i+j], \\
\mathrm{gr}_{\mathrm{KO}/k}^j \overline{\Delta}_R \{i\} &= \left(\bigwedge_R^{i-j} L_{R/k} \right) \otimes_k^{\mathbb{L}} \overline{\Delta}_k \{j\} [j-i].
\end{aligned}$$

¹⁰We replace the subscript KO by KO/k to stress that the base is k .

We note that this filtration is concentrated in degrees $j \leq i$, and in the usual convention, one might invert the sign of the weight to make it an increasing filtration. One also sees that this (inverted) increasing filtration starts with zero, which is a special case of Proposition 5.7. When k is an animated $\mathbb{F}_p$ -algebra, this recovers [Bha12, Prop 3.22] for the composite map $\mathbb{F}_p \rightarrow k \rightarrow R$ of animated rings, thus we view our result as a mixed characteristic analogue of that.

Example 5.5. To simplify notation, everything is p -completed. Let k be an animated ring, and $\mathcal{C}$ a dualizable presentable k -linear stable ∞ -category. Take M to be $\mathrm{THH}(\mathcal{C})$, with induced synthetic spectrum structure from $\mathrm{Fil}_M \mathrm{THH}(k)$, in Example 5.2, we get a filtered synthetic spectrum, whose underlying filtered spectrum under forgetful functor $\mathrm{SynSp} \rightarrow \mathrm{Sp}$ is denoted by $\mathrm{Fil}_{\mathrm{KO}/k} \mathrm{THH}_{\Delta}(\mathcal{C})^{\Phi C_p}$, with

$$\mathrm{gr}_{\mathrm{KO}/k}^j \mathrm{THH}_{\Delta}(\mathcal{C})^{\Phi C_p} = \mathrm{HH}(\mathcal{C}/k) \otimes_k^{\mathbb{L}} \widehat{\Delta}_k\{j\}[2j],$$

and another filtered spectrum $\mathrm{Fil}_{\mathrm{KO}/k} \mathrm{THH}_{\Delta/k}(\mathcal{C})^{\Phi C_p}$, with

$$\mathrm{gr}_{\mathrm{KO}/k}^j \mathrm{THH}_{\Delta/k}(\mathcal{C})^{\Phi C_p} = \mathrm{HH}(\mathcal{C}/k) \otimes_k^{\mathbb{L}} \widehat{\Delta}_k\{j\}[2j].$$

We make some final remarks on completeness of Katz–Oda filtration, which is, by construction, exhaustive. The case is direct if we consider τ -complete synthetic spectra instead.

Notation 5.6. Let $\widehat{\mathrm{SynSp}} \subseteq \mathrm{SynSp}$ denote the full subcategory of τ -complete synthetic spectra, which admits a left adjoint $(-)^{\widehat{}} : \mathrm{SynSp} \rightarrow \widehat{\mathrm{SynSp}}$.

Proposition 5.7. Let k be an animated ring, and N a $\mathrm{Fil}_M^{\geq 0} \mathrm{THH}_{\Delta}(k)^{\Phi C_p}$ -module in SynSp which is weight-bounded below. Then the filtered synthetic spectra $\mathrm{Fil}_{\mathrm{KO}} N'$ in both cases of Construction 5.1 is complete after τ -completion to $\widehat{\mathrm{SynSp}}$.

Proof. Since the functor $\mathrm{gr} : \widehat{\mathrm{SynSp}} \rightarrow \mathrm{Mod}_{\mathrm{gr}(\mathbb{S}_{\mathrm{ev}})}(\mathrm{Gr}(\mathrm{Sp}))$ preserves small colimits and is conservative, it suffice to in $\mathrm{Mod}_{\mathrm{gr}(\mathbb{S}_{\mathrm{ev}})}(\mathrm{Gr}(\mathrm{Sp}))$. In this case, since N is weight-bounded below, one can see that the limit tower $(\mathrm{Fil}_{\mathrm{KO}}^n N')_n$, after passing to $\mathrm{Gr}(\mathrm{Sp})$, is degreewise eventually zero. $\square$

Now we address the motivic completeness on the nose, which is equivalent to the motivic completeness of the underlying spectrum if the motivic completeness holds after τ -completion.

Proposition 5.8. Let k be an animated ring, and M a $\mathrm{Fil}_M \mathrm{THH}(k)$ -module in SynSp which is weight-bounded below. Suppose that M is uniformly t -bounded below.

- (1) If the spectra $\{\mathrm{gr}_N^m \Delta_k\{n\} | (m, n) \in \mathbb{Z} \times \mathbb{Z}\}$ is uniformly t -bounded below (this is the case when the k -module $L_{k/\mathbb{Z}}$ is p -completely weakly finitely generated by Corollary 3.18), then the Katz–Oda filtration on

$$M \otimes_{\mathrm{Fil}_M \mathrm{THH}(k)}^{\mathbb{L}} \mathrm{Fil}_M \mathrm{THH}_{\Delta}(k)^{\Phi C_p}$$

in Example 5.2 is complete;

- (2) If the Nygaard-completed Hodge–Tate $\widehat{\Delta}\{n\}$ is t -bounded below uniformly in $n \in \mathbb{Z}$ (this is the case when the k -module $L_{k/\mathbb{Z}}$ is p -completely weakly finitely generated by Corollary 3.18), then the Katz–Oda filtration on

$$M \otimes_{\mathrm{Fil}_M \mathrm{THH}(k)}^{\mathbb{L}} \mathrm{Fil}_M \mathrm{THH}(k)^{t C_p}$$

in Example 5.2 is complete.

Proof. Both cases follows from connectivity estimates of $\mathrm{Fil}_M \mathrm{THH}_{\Delta}(k)^{\Phi C_p}$ and $\mathrm{Fil}_M \mathrm{THH}(k)^{t C_p}$. Indeed, in the first case, connectivity estimates are established in the proof of Theorem 3.19, while the second connectivity estimates follow from the motivic completeness of $\mathrm{THH}(k)^{t C_p}$ for animated rings k , cf. [BL22, Cor 6.2.15] along with the trick of tensoring with $\mathbb{T}/C_p$ as in [Rig25, Lem 3.8]. $\square$

6. TRANSITIVITY OF F -SMOOTHNESS

The goal of this section is to establish transitivity of F -smoothness. For this purpose, we first introduce a variant of conjugate filtration in Section 5 on associate graded pieces of Nygaard-filtered absolute prismatic cohomology, which leads to a transitivity of weak F -smoothness (Theorem 6.12). Then we discuss a relative Nygaard-completeness (Definition 6.13). We give an animated version of transitivity of p -Cartier smoothness (the first case of Theorem 6.30), and a transitivity of F -smoothness (Theorem 6.15 and Corollary 6.16). The strategy of two proofs are parallel:

- (1) Establish a flatness of the map between cohomology theories via conjugate filtrations (Lemmas 6.28 and 6.37).
- (2) Using this flatness to deduce completeness of Katz–Oda filtrations (Corollaries 6.29 and 6.38).
- (3) Analyze associated graded pieces of Katz–Oda filtrations.

Moreover, we define *universal convergence* of Hodge and Nygaard filtration (Definitions 6.20 and 6.39) generalizing “having a universal Cartier isomorphism” [BLM18, Def 9.5.9]. This notion gives rise to variants of transitivity of p -Cartier smoothness and F -smoothness without bounded Tor-amplitude (the first case of Theorem 6.30 and Theorem 6.42). We also establish transitivity of universal convergence (Theorems 6.31 and 6.43). Over a perfectoid base, two universal convergences could also be compared (Theorem 6.45 and Proposition 6.48).

Construction 6.1. *To simplify notation, everything is p -completed. Let k be an animated ring, and M a $\mathrm{Fil}_M \mathrm{THH}(k)$ -module in SynSp . As in Construction 5.1 and Example 5.2, the filtration given by $\mathrm{Fil}_{\mathrm{KO}}^j \mathrm{Fil}_M \mathrm{THH}(k) := \mathrm{Fil}_M^{\geq j} \mathrm{THH}(k)$ on $\mathrm{Fil}_M \mathrm{THH}(k)$ gives rise to a filtration $\mathrm{Fil}_{\mathrm{KO}} M$ on M , with*

$$\begin{aligned} \mathrm{gr}_{\mathrm{KO}}^j M &= M \otimes_{\mathrm{Fil}_M \mathrm{THH}(k)}^{\mathbb{L}} \mathrm{ins}^j \mathrm{gr}_M^j \mathrm{THH}(k) \\ &= (M \otimes_{\mathrm{Fil}_M \mathrm{THH}(k)}^{\mathbb{L}} \mathrm{ins}^0 k) \otimes_{\mathrm{ins}^0 k}^{\mathbb{L}} \mathrm{ins}^j \mathrm{gr}_N^j \Delta_k\{j\}[2j] \in \mathrm{Mod}_{\mathrm{Fil}_M \mathrm{THH}(k)}(\mathrm{SynSp}). \end{aligned}$$

Example 6.2. To simplify notation, everything is p -completed. Let k be an animated ring, and $\mathcal{C}$ a dualizable presentable k -linear stable ∞ -category. Take M to be $\mathrm{THH}(\mathcal{C})$, with induced synthetic spectrum structure from $\mathrm{Fil}_M \mathrm{THH}(k)$, in Example 5.2, we get a filtered synthetic spectrum, whose underlying filtered spectrum under forgetful functor $\mathrm{SynSp} \rightarrow \mathrm{Sp}$ is denoted by $\mathrm{Fil}_{\mathrm{KO}/k} \mathrm{THH}(\mathcal{C})$, with

$$\mathrm{gr}_{\mathrm{KO}/k}^j \mathrm{THH}(\mathcal{C}) = \mathrm{HH}(\mathcal{C}/k) \otimes_k^{\mathbb{L}} \mathrm{gr}_N^j \Delta_k\{j\}[2j].$$

Example 6.3. To simplify notation, everything is p -completed. Let $k \rightarrow R$ be a map of animated rings. Then as in Example 5.4, applying Construction 6.1 to $M = \mathrm{Fil}_M \mathrm{THH}(R)$, we get a filtered synthetic spectrum $\mathrm{Fil}_{\mathrm{KO}/k} \mathrm{Fil}_M \mathrm{THH}(R)$ which admits a commutative ring structure, with

$$\begin{aligned} \mathrm{gr}_{\mathrm{KO}/k}^j \mathrm{Fil}_M \mathrm{THH}(R) &= (\mathrm{Fil}_M \mathrm{THH}(R) \otimes_{\mathrm{Fil}_M \mathrm{THH}(k)}^{\mathbb{L}} \mathrm{ins}^0 k) \otimes_{\mathrm{ins}^0 k}^{\mathbb{L}} \mathrm{ins}^j \mathrm{gr}_N^j \Delta_k\{j\}[2j] \\ &= \mathrm{Fil}_M \mathrm{HH}(R/k) \otimes_{\mathrm{ins}^0 k}^{\mathbb{L}} \mathrm{ins}^j \mathrm{gr}_N^j \Delta_k\{j\}[2j]. \end{aligned}$$

Passing to the i th associated graded pieces of the synthetic spectra, we get filtered objects $\mathrm{Fil}_{\mathrm{KO}/k} \mathrm{gr}_N^i \Delta_k\{i\}[2i]$, with

$$\begin{aligned} \mathrm{gr}_{\mathrm{KO}/k}^j \mathrm{gr}_N^i \Delta_k\{i\}[2i] &= \left(\bigwedge_R^{i-j} L_{R/k} \right) [i-j] \otimes_k^{\mathbb{L}} \mathrm{gr}_N^j \Delta_k\{j\}[2j] \\ &= \left(\bigwedge_R^{i-j} L_{R/k} \right) \otimes_k^{\mathbb{L}} \mathrm{gr}_N^j \Delta_k\{j\}[i+j] \in D(R), \\ \mathrm{gr}_{\mathrm{KO}/k}^j \mathrm{gr}_N^i \Delta_k\{i\} &= \left(\bigwedge_R^{i-j} L_{R/k} \right) \otimes_k^{\mathbb{L}} \mathrm{gr}_N^j \Delta_k\{j\}[j-i] \in D(R), \end{aligned}$$

which is concentrated in weights $j \leq i$, where the R -module structure is induced by that on the first tensor factor, thus the filtration is complete and exhaustive, a special case of Proposition 6.4.

Again, by construction, Katz–Oda filtration on THH is always exhaustive. Completeness of Katz–Oda filtration on THH is much better than that on $\mathrm{THH}_{\Delta}^{\Phi C_p}$ and $\mathrm{THH}_{\Delta/k}^{\Phi C_p}$:

Proposition 6.4. *Let k be an animated ring, and M a $\mathrm{Fil}_M \mathrm{THH}(k)$ -module in SynSp which is weight-bounded below, and τ -complete. Then the filtered synthetic spectra $\mathrm{Fil}_{\mathrm{KO}} M$ in Construction 6.1 is complete.*

Proof. First, each synthetic spectrum $\mathrm{Fil}_{\mathrm{KO}}^j \mathrm{Fil}_M \mathrm{THH}(k)$ is complete, as the maps $\mathrm{Fil}_{\mathrm{KO}}^{j+1} M \rightarrow \mathrm{Fil}_{\mathrm{KO}}^j M$ induces an equivalence on the limit of the tower of multiplication by τ , thus the limits of these towers are the same, being zero. Thus it suffices to check the completeness of $\mathrm{Fil}_{\mathrm{KO}} M$ after τ -completion to $\widehat{\mathrm{SynSp}}$, and the proof of Proposition 5.7 adapts. $\square$

Example 6.5. Let k be an animated ring. Take R to be the monogenic polynomial k -algebra $k[t]$. Then Example 6.3 gives rise to a 2-step increasing filtration, which corresponds to a fiber sequence

$$L_{k[t]/k}^1 \otimes_k^{\mathbb{L}} \mathrm{gr}_N^{i-1} \Delta_k[-1] \longrightarrow \mathrm{gr}_N^i \Delta_{k[t]} \longrightarrow k[t] \otimes_k^{\mathbb{L}} \mathrm{gr}_N^i \Delta_k,$$

which is the fiber sequence in [BM23, Prop 4.8], used to establish the weak F -smoothness of $k[t]$ from that of k .

Example 6.5 hints a possibility of applying Example 6.3 to understand weak F -smoothness. This is indeed the case (Theorem 6.12). We first review weak F -smoothness in [BM23, Def 4.2]. We give a slightly different definition, which is equivalent to theirs by Remark 6.11.

Definition 6.6. We say that an animated ring A is *weakly F -smooth* if it is p -quasi-lci over $\mathbb{Z}$, and for each $i \in \mathbb{Z}$, the $(A \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ -module spectrum

$$\mathrm{fib} \left(\mu_1 : \mathrm{gr}_N^i \Delta_A \{j\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \longrightarrow \mathrm{gr}_N^{i+p} \Delta_A \{j+p\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \right)$$

has Tor-amplitude in degrees $\leq -(i+2)$ for every¹¹ $j \in \mathbb{Z}$, where μ_1 is induced by the class μ_1 in $\mathrm{Fil}_M(\mathrm{THH}(\mathbb{Z}) \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$, which gives rise to a map

$$\mathrm{Fil}_M(\mathrm{THH}(A) \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p) \longrightarrow \mathrm{Fil}_M(\mathrm{THH}(A) \otimes_A^{\mathbb{L}} \mathbb{F}_p)(-p)[-2p]$$

of synthetic spectra.

Remark 6.7. Let A be an animated ring which is p -quasi-lci over $\mathbb{Z}$. Then for every pair $(i, j) \in \mathbb{Z}^2$ with $i < p$, by [BL22, Cor 5.5.18], we have $\mathrm{gr}_N^i \Delta_A \{j\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \simeq \left(\bigwedge_A^i L_{A/\mathbb{Z}} \right) \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p[-i]$, which is zero when $i < 0$. Since A is p -quasi-lci over $\mathbb{Z}$, the $A \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ -module $\mathrm{gr}_N^i \Delta_A \{j\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ has Tor-amplitude ≤ 0 . It follows that the condition on $\mathrm{fib}(\mu_1)$ in Definition 6.6 is automatic when $i < 0$, and moreover, the $A \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ -module $\mathrm{fib}(\mu_1)$ has Tor-amplitude < 0 for every pair $(i, j) \in \mathbb{Z}^2$.

In [BM23, Def 4.2], the authors considered the p -complete Tor-amplitude of $\mathrm{fib}(\mu_1)$ as an A -module spectrum. To see that it is equivalent¹² to ours, we need a simple lemma for Tor-amplitude of A/I -modules as A -modules for an invertible ideal $I \subseteq A$.

¹¹This is equivalent to requiring the same for a single $j \in \mathbb{Z}$, as Breuil–Kisin twists trivialize after passing to associated graded pieces of the Nygaard filtration by [BL22, Rem 5.5.15].

¹²In an earlier version of this article, we mistakenly deduced this equivalence from the converse direction of Lemma 6.8 without assuming that the A/I -module M is extended from an A -module M_0 , cf. Example 6.9.

Lemma 6.8. *Let A be an animated ring, and $I \rightarrow A$ a generalized Cartier divisor, and M an A/I -module. Let $n \in \mathbb{Z}$ be an integer. If the A/I -module M Tor-amplitude $\leq n$, then the A -module M has Tor-amplitude $\leq n + 1$. The converse is true if there exists an A -module M_0 such that $M = M_0 \otimes_A^{\mathbb{L}} (A/I)$.*

Proof. Suppose that the A/I -module M has Tor-amplitude $\leq n$. For every coconnective A -module N , as the spectrum $(A/I) \otimes_A^{\mathbb{L}} N$ is 1-truncated, the spectrum

$$M \otimes_A^{\mathbb{L}} N \simeq M \otimes_{A/I}^{\mathbb{L}} ((A/I) \otimes_A^{\mathbb{L}} N)$$

is $(n + 1)$ -truncated.

On the other hand, suppose that $M = M_0 \otimes_A^{\mathbb{L}} (A/I)$ and the A -module M has Tor-amplitude $\leq n + 1$. Then the multiplication map $m : M \otimes_A^{\mathbb{L}} (A/I) \rightarrow M$ admits an A/I -linear retract. This implies that the A/I -module $\text{fib}(m) = M \otimes_{A/I}^{\mathbb{L}} (I/I^2)[1]$ has Tor-amplitude $\leq n + 1$. As the A/I -module I/I^2 is connectively invertible, the result follows. $\square$

Example 6.9. The $\mathbb{Z}$ -module $\mathbb{F}_p$ has Tor-amplitude ≤ 1 , while as a $\mathbb{Z}/p^2$ -module, it has unbounded Tor-amplitude, since $\mathbb{F}_p \otimes_{\mathbb{Z}/p^2}^{\mathbb{L}} \mathbb{F}_p \simeq \bigoplus_{n \in \mathbb{N}} \mathbb{F}_p[n]$. This shows that the converse of Lemma 6.8 is false in general.

Corollary 6.10. *Let A be an animated ring, and $I \rightarrow A$ a generalized Cartier divisor, and M an A/I -module. Let $n \in \mathbb{Z}$ be an integer. Then the A -module M has I -complete Tor-amplitude $\leq n$ if and only if the A -module M has Tor-amplitude $\leq n$.*

Proof. “if” part follows from definitions. We suppose that the A -module M has I -complete Tor-amplitude $\leq n$. By definition, the A/I -module $M \otimes_A^{\mathbb{L}} (A/I)$ has Tor-amplitude $\leq n$. It follows from Lemma 6.8 that the A -module $M \otimes_A^{\mathbb{L}} (A/I)$ has Tor-amplitude $\leq n + 1$. As in the proof of the converse direction of Lemma 6.8, the multiplication map $m : M \otimes_A^{\mathbb{L}} (A/I) \rightarrow M$ admits an A -linear retract $M \rightarrow M \otimes_A^{\mathbb{L}} (A/I)$ induced by $A \rightarrow A/I$, thus the A -module $\text{fib}(m) = M \otimes_A I[1]$ has Tor-amplitude $\leq n + 1$. As the A -module I is connectively invertible, the result follows. $\square$

Remark 6.11. Let A be an animated ring which is p -quasi-lci over $\mathbb{Z}$. Then the proof of [BM23, Prop 4.3] shows that A is weak F -smooth in our sense if and only if, for every $i \in \mathbb{Z}$, the $A \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ -module spectrum

$$\text{fib} \left(\varphi_i : \text{gr}_N^i \Delta_A \rightarrow \overline{\Delta}_A \{i\} \right) \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$$

has Tor-amplitude $\leq -(i + 2)$, which is equivalent to that the A -module spectrum $\text{fib}(\varphi_i) \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ has Tor-amplitude $\leq -(i + 1)$ by Lemma 6.8, which is equivalent to that the A -module spectrum $\text{fib}(\varphi_i) \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ has p -completed Tor-amplitude $\leq -(i + 1)$ by Corollary 6.10. This is equivalent to weak F -smoothness in [BM23, Def 4.2].

Theorem 6.12. *Let $k \rightarrow R$ be a map of animated rings. Suppose that*

- *the animated ring k is weakly F -smooth; and that*
- *the R -module spectrum $L_{R/k}$ is p -completely flat.*

Then the animated ring R is weakly F -smooth as well.

Proof. First, as the animated ring k is p -quasi-lci, and the R -module $L_{R/k}$ is p -completely flat, the animated ring R is p -quasi-lci as well, by transitivity of cotangent complex. For animated rings A , let $F^i(A)$ denote the fiber

$$\text{fib} \left(\mu_1 : \text{gr}_N^i \Delta_A \{i\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \longrightarrow \text{gr}_N^{i+p} \Delta_A \{i + p\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \right)$$

being an A -module. We want to invoke Example 6.3, but we have to be a bit more careful, as the map μ_1 involves a shift of Katz–Oda weight, internal filtration weight, and homotopy degree. Indeed, μ_1 comes

from a class in $\mathrm{Fil}_M(\mathrm{THH}(\mathbb{Z}) \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$, which has filtration weight p and homotopy degree $2p$. This gives rise to a class in $\mathrm{Fil}_{\mathrm{KO}} \mathrm{Fil}_M(\mathrm{THH}(k) \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$, which has Katz–Oda weight p , internal filtration weight p , and homotopy degree $2p$. In the end, we indeed get a Katz–Oda filtration $\mathrm{Fil}_{\mathrm{KO}} F^i(R)$ on the R -module $F^i(R)$, with

$$\mathrm{gr}_{\mathrm{KO}}^j F^i(R) = \left(\bigwedge_R^{i-j} L_{R/k} \right) \otimes_k^{\mathbb{L}} F^j(k)[j-i] \in D(R),$$

which corresponds to a finite-step exhaustive increasing filtration. Now for every j , the R -module spectrum

$$\left(\bigwedge_R^{i-j} L_{R/k} \right) \otimes_k^{\mathbb{L}} F^j(k)[j-i]$$

has p -complete Tor-amplitude $\leq -(j+2) + (j-i) = -(i+2)$, as for every $(R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ -module N which is concentrated in degree 0, the k -module $N \otimes_R^{\mathbb{L}} \left(\bigwedge_R^{i-j} L_{R/k} \right)$ is concentrated in degree 0 by p -complete flatness of the R -module $L_{R/k}$. The result then follows. $\square$

Now we prove transitivity of F -smoothness. For this, it seems reasonable to introduce a relative version of Nygaard-completeness, which is precisely relative Segal conjecture up to completions with respect to motivic filtrations.

Definition 6.13. Let $k \rightarrow R$ be a map of animated rings. We say that the absolute prismatic cohomology $\Delta_R\{*\}$ is *Nygaard-complete relative to $\Delta_k\{*\}$* if the canonical map

$$\mathrm{gr}_M \mathrm{THH}(R) \otimes_{\mathrm{gr}_M \mathrm{THH}(k)}^{\mathbb{L}} \mathrm{gr}_M \mathrm{THH}(k)^{tC_p} \longrightarrow \mathrm{gr}_M \mathrm{THH}(R)^{tC_p}$$

of graded $\mathbb{E}_{\infty}$ -rings is an equivalence after p -completion, or equivalently¹³, the canonical map

$$\left(\bigoplus_{n \in \mathbb{Z}} \overline{\Delta}_R\{n\} \right) \otimes_{\bigoplus_{n \in \mathbb{Z}} \overline{\Delta}_k\{n\}}^{\mathbb{L}} \left(\bigoplus_{n \in \mathbb{Z}} \widehat{\Delta}_k\{n\} \right) \longrightarrow \bigoplus_{n \in \mathbb{Z}} \widehat{\Delta}_R\{n\}$$

of graded $\mathbb{E}_{\infty}$ -rings is an equivalence after p -completion.

Remark 6.14. Let $k \rightarrow R$ be a map of animated rings. Then by definition, the absolute prismatic cohomology $\Delta_R\{*\}$ is Nygaard-complete relative to $\Delta_k\{*\}$ if and only if the filtered graded $\mathbb{E}_{\infty}$ -ring

$$\left(\bigoplus_{n \in \mathbb{Z}} \mathrm{Fil}_N \overline{\Delta}_R\{n\} \right) \otimes_{\bigoplus_{n \in \mathbb{Z}} \mathrm{Fil}_N \overline{\Delta}_k\{n\}}^{\mathbb{L}} \left(\bigoplus_{n \in \mathbb{Z}} \mathrm{Fil}_N \widehat{\Delta}_k\{n\} \right)$$

is complete up to p -completion.

The rest of this section is devoted to the following theorem.

Theorem 6.15. *Let $k \rightarrow R$ be a map of animated rings such that k is p -completely t -bounded¹⁴ and R has bounded p -complete Tor-amplitude over k . Suppose that*

- *the animated ring k is weakly F -smooth; and that*
- *the R -module spectrum $L_{R/k}$ is p -completely flat.*

If the p -completed derived de Rham cohomology $\mathrm{dR}_{R/k}$ is p -completely Hodge-complete, then the absolute prismatic cohomology $\Delta_R\{\}$ is Nygaard-complete relative to $\Delta_k\{*\}$.*

Before its proof, we mention that it implies transitivity of F -smoothness:

¹³We use the symmetric monoidal structure on the shearing, which is explained in [Rak26, Prop 3.3.4].

¹⁴We say that a spectrum $X \in \mathrm{Sp}$ is p -completely t -bounded if the spectrum X/p is t -bounded.

Corollary 6.16. *Let k be an F -smooth p -quasisyntomic ring, and $k \rightarrow R$ a p -Cartier smooth map [Bou23] such that R has bounded p^∞ -torsion¹⁵, and that R has bounded p -complete Tor-amplitude over k . Then R is F -smooth p -quasisyntomic as well.*

Proof. We examine the transitivity sequence

$$L_{k/\mathbb{Z}_p} \otimes_k^{\mathbb{L}} R \longrightarrow L_{R/\mathbb{Z}_p} \longrightarrow L_{R/k}.$$

Since k is p -quasisyntomic, the k -module $L_{k/\mathbb{Z}_p}$ has p -complete Tor-amplitude in $[0, 1]$. By assumption, the R -module $L_{R/k}$ is p -completely flat¹⁶, thus $L_{R/\mathbb{Z}_p}$ has p -complete Tor-amplitude in $[0, 1]$ as well, thus R is p -quasisyntomic. Since the ring k is F -smooth, it is weakly F -smooth [BM23, Prop 4.3]. By Theorem 6.12, the ring R is weakly F -smooth as well. Now the result follows from Theorem 6.15 and the Nygaard-completeness of the absolute prismatic cohomology $\widehat{\Delta}_k$. $\square$

Now we come to the proof of Theorem 6.15. Let $k \rightarrow R$ be a map of animated rings. The strategy is similar to Section 5, namely introducing a filtration on the graded $\mathbb{E}_\infty$ -ring $\bigoplus_{n \in \mathbb{Z}} \widehat{\Delta}_k\{*\}$, showing that it induces a complete “Katz–Oda” filtration on the map

$$\left(\bigoplus_{n \in \mathbb{Z}} \overline{\Delta}_R\{n\} \right) \otimes_{\bigoplus_{n \in \mathbb{Z}} \overline{\Delta}_k\{n\}}^{\mathbb{L}} \left(\bigoplus_{n \in \mathbb{Z}} \widehat{\Delta}_k\{n\} \right) \longrightarrow \bigoplus_{n \in \mathbb{Z}} \widehat{\Delta}_R\{n\}$$

and then show that, on each “Katz–Oda” associated graded pieces, the filtered graded spectrum is complete. A naive guess is to take the Nygaard filtration on $\bigoplus_{n \in \mathbb{Z}} \widehat{\Delta}_k\{n\}$ as in Construction 5.1 etc. Tensor products do not preserve completeness, and such a “Katz–Oda” filtration is a priori not necessarily complete. Fortunately, our assumptions allow us to control the convergence.

We start with a closer analysis of convergence of the Hodge filtration on de Rham cohomology, and to see how to deduce transitivity of p -Cartier smoothness via the usual Katz–Oda filtration. As a toy example, for a map $k \rightarrow R$ of commutative rings with the R -module $L_{R/k}$ being flat, the r -th filtered piece of the Hodge-completed derived de Rham cohomology $\widehat{\mathrm{dR}}_{R/k}$ is computed by the cochain complex

$$\cdots \longrightarrow 0 \longrightarrow \Omega_{R/k}^r \xrightarrow{d} \Omega_{R/k}^{r+1} \xrightarrow{d} \cdots,$$

which is $(-r)$ -truncated. That is to say, the convergence of the Hodge-completed derived de Rham cohomology $\widehat{\mathrm{dR}}_{R/k}$ has a linearly increasing coconnectivity estimate. This extends to the animated setting:

Notation 6.17. *Let k be an animated $\mathbb{F}_p$ -algebra. We denote by φ_k the Frobenius map $k \rightarrow k$, and by φ_k^* the (derived) base change $(-) \otimes_{k, \varphi_k}^{\mathbb{L}} k : D(k) \rightarrow D(k)$.*

Lemma 6.18. *Let $k \rightarrow R$ be a map of animated rings with R being n -truncated for some $n \in \mathbb{N}$. Suppose that the R -module $L_{R/k}$ is flat. Then for every $r \in \mathbb{N}$, the r -th Hodge-filtered piece $\mathrm{Fil}_H^r \widehat{\mathrm{dR}}_{R/k}$ is $(n-r)$ -truncated.*

Proof. Note that, for every $s \in \mathbb{N}_{>r}$, the spectrum $\mathrm{gr}_H^{[r,s]} \widehat{\mathrm{dR}}_{R/k}$ is a finite extension of

$$\mathrm{gr}_H^i \widehat{\mathrm{dR}}_{R/k} = \left(\bigwedge_R^i L_{R/k} \right) [-i]$$

for $i = r, r+1, \dots, s-1$, and for such i ’s, since the R -module $L_{R/k}$ is flat, so is the exterior power $\bigwedge_R^i L_{R/k}$, and thus $\mathrm{gr}_H^i \widehat{\mathrm{dR}}_{R/k}$ is $(n-i)$ -truncated, thus the spectrum $\mathrm{gr}_H^{[r,s]} \widehat{\mathrm{dR}}_{R/k}$ is $(n-r)$ -truncated. Taking $s \rightarrow \infty$, we see that the spectrum $\mathrm{Fil}_H^r \widehat{\mathrm{dR}}_{R/k}$ is $(n-r)$ -truncated as well. $\square$

¹⁵We do not know whether this assumption is redundant.

¹⁶The p -discreteness of $k \rightarrow R$ implies that the canonical map $(R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p) \otimes_{k \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p}^{\mathbb{L}} \pi_0(k \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p) \rightarrow \pi_0(R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ is an equivalence.

As a consequence of Lemma 6.18 and definition of Tor-amplitude, we have

Corollary 6.19. *Let $k \rightarrow R$ be a map of animated $\mathbb{F}_p$ -algebras with R being n -truncated for some $n \in \mathbb{N}$, and $m \in \mathbb{Z}$ an integer. Suppose that the R -module $L_{R/k}$ is flat. Then for every $\varphi_k^* R$ -module M of Tor-amplitude $\leq m$, and every $r \in \mathbb{N}$, the r -th Hodge-filtered piece*

$$M \otimes_{\varphi_k^* R}^{\mathbb{L}} \mathrm{Fil}_H^r \widehat{\mathrm{dR}}_{R/k}$$

is $(m + n - r)$ -truncated. In particular, the filtered spectrum $M \otimes_{\varphi_k^ R}^{\mathbb{L}} \mathrm{Fil}_H \widehat{\mathrm{dR}}_{R/k}$ is complete.*

Sometimes, we have an efficient bound of the Tor-amplitude of $\mathrm{Fil}_H^r \widehat{\mathrm{dR}}_{R/k}$, in which case a variant of Corollary 6.19 holds. This notion is partially inspired by universal Cartier isomorphisms in [BLM18, Def 9.5.14] (cf. Lemma 6.23).

Definition 6.20. Let $k \rightarrow R$ be a map of animated $\mathbb{F}_p$ -algebras. We say that a filtered $\mathrm{Fil}_H \mathrm{dR}_{R/k}$ -module $\mathrm{Fil} M$ *converges universally (of index $\alpha \in \mathbb{Z}$)* if, for every $r \in \mathbb{N}$, the $\varphi_k^* R$ -module $\mathrm{Fil}^r M$ has Tor-amplitude $\leq \alpha - r$, and it converges universally if such an integer α exists. In particular, we say that the Hodge filtration on $\widehat{\mathrm{dR}}_{R/k}$ *converges universally (resp. of index $\alpha \in \mathbb{Z}$)* if the filtered $\mathrm{Fil}_H \mathrm{dR}_{R/k}$ -module $\mathrm{Fil}_H \widehat{\mathrm{dR}}_{R/k}$ converges universally (resp. of index $\alpha \in \mathbb{Z}$).

It follows directly from definitions that

Corollary 6.21. *Let $k \rightarrow R$ be a map of animated $\mathbb{F}_p$ -algebras, and N a filtered $\mathrm{Fil}_H \mathrm{dR}_{R/k}$ -module which converges universally of index $\alpha \in \mathbb{Z}$. Then for $\varphi_k^* R$ -module M , and every $r \in \mathbb{N}$,*

- *if M is m -truncated for some $m \in \mathbb{Z}$, then the r -th Hodge-filtered piece*

$$M \otimes_{\varphi_k^* R}^{\mathbb{L}} \mathrm{Fil}^r N$$

is $(\alpha + m - r)$ -truncated. In particular, the filtered spectrum $M \otimes_{\varphi_k^ R}^{\mathbb{L}} \mathrm{Fil} N$ is complete; and*

- *if the $\varphi_k^* R$ -module M has Tor-amplitude $\leq m$ for some $m \in \mathbb{Z}$, then the r -th Hodge-filtered piece*

$$M \otimes_{\varphi_k^* R}^{\mathbb{L}} \mathrm{Fil}^r N$$

has Tor-amplitude $\leq \alpha + m - r$.

Universal convergence of Hodge filtration implies flatness of Frobenius.

Lemma 6.22. *Let $k \rightarrow R$ be a map of animated $\mathbb{F}_p$ -algebras, and $\mathrm{Fil} M$ a filtered $\mathrm{Fil}_H \mathrm{dR}_{R/k}$ -module equipped with a map $\mathrm{Fil}_H \mathrm{dR}_{R/k} \rightarrow \mathrm{Fil} M$ which induces an equivalence on associated graded pieces. Suppose that the $\mathrm{Fil}_H \mathrm{dR}_{R/k}$ -module $\mathrm{Fil} M$ converges universally of index $\alpha \in \mathbb{Z}$. Then the Frobenius map $\varphi_{R/k} : \varphi_k^* R \rightarrow R$ has Tor-amplitude $\leq \alpha$, and for every $r \in \mathbb{N}$, the $\varphi_k^* R$ -module $\bigwedge_R^r L_{R/k}$ (via forgetting along $\varphi_{R/k}$) has Tor-amplitude $\leq \alpha$.*

Proof. By assumption, for every $r \in \mathbb{N}$, the $\varphi_k^* R$ -modules $\mathrm{Fil}^r M[r]$ and $\mathrm{Fil}^{r+1} M[r+1]$ have Tor-amplitude $\leq \alpha$. Now the result follows from the fiber sequence

$$\mathrm{Fil}^r M[r] \longrightarrow \mathrm{gr}^r M[r] \longrightarrow \mathrm{Fil}^{r+1} M[r+1]$$

of $\varphi_k^* R$ -modules, and the equivalence $\bigwedge_R^r L_{R/k} = \mathrm{gr}_H^r \mathrm{dR}_{R/k}[r] \xrightarrow{\cong} \mathrm{gr}^r M[r]$ of R -modules (thus as $\varphi_k^* R$ -modules). $\square$

The Hodge-filtration on the Hodge-completed derived de Rham cohomology converges universally for Cartier smooth $\mathbb{F}_p$ -algebras with universal Cartier isomorphism [BLM18, Def 9.5.14].

Lemma 6.23. *Let R be a Cartier smooth k -algebra where $k := \mathbb{F}_p$. Suppose that, for every $r \in \mathbb{N}$, the $\varphi_k^* R$ -module $B\Omega_{R/k}^r := \text{Im} \left(\Omega_{R/k}^{r-1} \xrightarrow{d} \Omega_{R/k}^r \right)$ is flat. Then the Hodge filtration on $\widehat{dR}_{R/k}$ converges universally of index 0.*

Proof. Indeed, for every $r \in \mathbb{N}$, the $\varphi_k^* R$ -module $\text{Fil}_H^r \widehat{dR}_{R/k}$ is represented by the cochain complex

$$\cdots \longrightarrow 0 \longrightarrow \Omega_{R/k}^r \xrightarrow{d} \Omega_{R/k}^{r+1} \xrightarrow{d} \cdots$$

of $\varphi_k^* R$ -modules. We get a short exact sequence

$$\begin{array}{ccccccc} \cdots & \longrightarrow & 0 & \longrightarrow & B\Omega_{R/k}^r & \longrightarrow & 0 \longrightarrow \cdots \\ & & \parallel & & \downarrow & & \downarrow \\ \cdots & \longrightarrow & 0 & \longrightarrow & \Omega_{R/k}^r & \xrightarrow{d} & \Omega_{R/k}^{r+1} \xrightarrow{d} \cdots \\ & & \parallel & & \downarrow & & \parallel \\ \cdots & \longrightarrow & 0 & \longrightarrow & \Omega_{R/k}^r / B\Omega_{R/k}^r & \xrightarrow{d} & \Omega_{R/k}^{r+1} \xrightarrow{d} \cdots \end{array}$$

of cochain complexes of $\varphi_k^* R$ -modules. As the R -module $L_{R/k}$ is flat, by Cartier isomorphism, we see that $\varphi_k^* R$ -module represented by the bottom cochain complex has Tor-amplitude $\leq -r$. It follows from flatness of the $\varphi_k^* R$ -module $B\Omega_{R/k}^r$ that the $\varphi_k^* R$ -module $\text{Fil}_H^r \widehat{dR}_{R/k}$ has Tor-amplitude $\leq -r$ as well. $\square$

Example 6.24. Let R be a regular Noetherian k -algebra where $k := \mathbb{F}_p$. By [BLM18, Thm 9.5.1 & 9.5.18], the k -algebra R is Cartier smooth, and has a universal Cartier isomorphism. It follows from Lemma 6.23 that the Hodge filtration on $\widehat{dR}_{R/k}$ converges universally of index 0.

Example 6.25. Let R be a perfect k -algebra where $k := \mathbb{F}_p$. By [BLM18, Prop 9.5.11], the k -algebra R is Cartier smooth, and has a universal Cartier isomorphism. It follows from Lemma 6.23 that the Hodge filtration on $\widehat{dR}_{R/k}$ converges universally of index 0.

This actually admits an animated generalization:

Lemma 6.26. *Let $k \rightarrow R$ be a relative perfect map of animated $\mathbb{F}_p$ -algebras. Then the Hodge-filtration on $dR_{R/k}$ is complete, and it converges universally of index 0.*

Proof. First, by Lemma 3.21, the R -module $L_{R/k}$ is a retract of $L_{R/\varphi_k^* R}$, which is zero as the relative Frobenius map $\varphi_{R/k} : \varphi_k^* R \rightarrow R$ is an equivalence¹⁸. It follows that the Hodge filtration on $\widehat{dR}_{R/k}$ is concentrated in weight 0, and so is the conjugate filtration on $dR_{R/k}$, and thus the canonical map $dR_{R/k} \rightarrow \widehat{dR}_{R/k}$ is identified with the relative Frobenius map $\varphi_{R/k}$, and the Hodge-completeness follows. The universal convergence follows from definitions as well. $\square$

Example 6.27. Let R be a valuation k -algebra where $k := \mathbb{F}_p$. Gabber showed that the k -algebra R is Cartier smooth, [KST21, Cor A.4]. In particular, for every $r \in \mathbb{N}$, the R -module $\Omega_{R/k}^r$ is flat thus torsion-free, thus it is also torsion-free viewed as a $\varphi_k^* R$ -module. It follows that the sub- $\varphi_k^* R$ -module $B\Omega_{R/k}^r \subseteq \Omega_{R/k}^r$ is torsion-free as well, thus flat, as $\varphi_k^* R \cong R$ is a valuation ring. It then follows from Lemma 6.23 that the Hodge filtration on $\widehat{dR}_{R/k}$ converges universally of index 0.

¹⁷This is just R . It is written as $\varphi_k^* R$ to stress that the $\varphi_k^* R$ -module structures on $\Omega_{R/k}^*$ is Frobenius twisted, so that the differentials $d : \Omega_{R/k}^* \rightarrow \Omega_{R/k}^{*+1}$ are $\varphi_k^* R$ -linear.

¹⁸This argument is actually implicit in [GR03, Lem 6.5.13(i)] as well.

We now explain how to deduce an animated version of transitivity of p -Cartier smoothness established in [Bou23, Lem 2.10] from Lemma 6.18, and the proof of Theorem 6.15 shares the same strategy. The key is to bound the Tor-amplitude of the map $\mathrm{dR}_{R/k} \rightarrow \mathrm{dR}_{A/k}$ for quasi-lci maps $R \rightarrow A$:

Lemma 6.28. *Let k be an animated $\mathbb{F}_p$ -algebra, and $R \rightarrow A$ be a quasi-lci map of animated k -algebras. Then the de Rham cohomology $\mathrm{dR}_{A/k}$, being a $(\varphi_k^* A \otimes_{\varphi_k^* R}^{\mathbb{L}} \mathrm{dR}_{R/k})$ -module, admits an exhaustive increasing filtration, with associated graded pieces of the form*

$$\varphi_k^* N \otimes_{\varphi_k^* R}^{\mathbb{L}} \mathrm{dR}_{R/k}$$

for some A -module N of Tor-amplitude ≤ 0 .

Proof. We simply take the relative conjugate filtration on $\mathrm{dR}_{A/k}$ constructed in [Bha12, Prop 3.22]. The associated graded pieces are given by

$$\varphi_k^* \left(\bigwedge_A^s L_{A/R} \right) [-s] \otimes_{\varphi_k^* R}^{\mathbb{L}} \mathrm{dR}_{R/k}.$$

Now as $R \rightarrow A$ is quasi-lci, for every $s \in \mathbb{N}$, the A -module $(\bigwedge_A^s L_{A/R}) [-s]$ has Tor-amplitude ≤ 0 (cf. [BMS19, Lem 4.34]). The result then follows. $\square$

This bound of Tor-amplitude leads to completeness of the Katz–Oda filtration, under either a boundedness of Tor-amplitude, or a universal convergence of Hodge filtration.

Corollary 6.29. *Let k be an animated ring, and $R \rightarrow A$ a p -quasi-lci map of animated k -algebras, and $\mathrm{Fil} M$ a filtered $\mathrm{Fil}_H \mathrm{dR}_{R/k}$ -module. Suppose that **either***

- *the R -module A has p -complete Tor-amplitude $\leq m$ for some $m \in \mathbb{Z}$, that the R -module $L_{R/k}$ is p -completely flat, and that R is p -completely n -truncated for some $n \in \mathbb{N}$, with $\mathrm{Fil} M = \mathrm{Fil}_H \widehat{\mathrm{dR}}_{R/k}$;*
- or**
- *the $\mathrm{Fil}_H \mathrm{dR}_{R \otimes_{\mathbb{Z}} \mathbb{F}_p / k \otimes_{\mathbb{Z}} \mathbb{F}_p}$ -module $\mathrm{Fil} M \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ converges universally, and that the spectrum $\varphi_{k \otimes_{\mathbb{Z}} \mathbb{F}_p}^* (A \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ is m -truncated for some $m \in \mathbb{N}$.*

Then the Katz–Oda filtration

$$\mathrm{Fil}_{\mathrm{KO}/R}(\mathrm{dR}_{A/k} \otimes_{\mathrm{dR}_{R/k}}^{\mathbb{L}} M) := \mathrm{dR}_{A/k} \otimes_{\mathrm{dR}_{R/k}}^{\mathbb{L}} \mathrm{Fil} M$$

is p -completely complete.

Proof. First, we may replace k by $k \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$, so without loss of generality, we may equip k an animated $\mathbb{F}_p$ -algebra structure.

We fix $r \in \mathbb{N}$. Then by Lemma 6.28, the r -th Katz–Oda filtered piece

$$\mathrm{dR}_{A/k} \otimes_{\mathrm{dR}_{R/k}}^{\mathbb{L}} \mathrm{Fil}^r M$$

admits an exhaustive increasing filtration, with associated graded pieces of the form

$$(3) \quad \varphi_k^* N \otimes_{\varphi_k^* R}^{\mathbb{L}} \mathrm{Fil}^r M$$

for some A -module N of Tor-amplitude ≤ 0 . We consider the two cases separately:

- Suppose that the R -module A has Tor-amplitude $\leq m$, and that R is n -truncated, with $\mathrm{Fil} M = \mathrm{Fil}_H \widehat{\mathrm{dR}}_{R/k}$. Then every $\varphi_k^* R$ -module $\varphi_k^* N$ in (3) has Tor-amplitude $\leq m$. It follows from Corollary 6.19 and the flatness of $L_{R/k}$ that the spectrum $\varphi_k^* N \otimes_{\varphi_k^* R}^{\mathbb{L}} \mathrm{Fil}^r M$ is $(m + n - r)$ -truncated, thus so is the r -th filtered piece $\mathrm{dR}_{A/k} \otimes_{\mathrm{dR}_{R/k}}^{\mathbb{L}} \mathrm{Fil}^r M$. The result then follows.

- Suppose that $\mathrm{Fil}_H \mathrm{dR}_{R/k}$ -module $\mathrm{Fil} M$ converges universally of index $\alpha \in \mathbb{Z}$, and that the spectrum $\varphi_k^* A$ is m -truncated. Then every spectrum $\varphi_k^* N$ in (3) is m -truncated. It follows from Corollary 6.21 that the spectrum $\varphi_k^* N \otimes_{\varphi_k^* R}^{\mathbb{L}} \mathrm{Fil}^r M$ is $(\alpha + m - r)$ -truncated, and so is the r -th filtered piece $\mathrm{dR}_{A/k} \otimes_{\mathrm{dR}_{R/k}}^{\mathbb{L}} \mathrm{Fil}^r M$. The result then follows. $\square$

Now we deduce the following animated version of transitivity of p -Cartier smoothness, under either a boundedness of Tor-amplitude, or a universal convergence of Hodge filtration.

Theorem 6.30. *Let k be an animated ring, and $R \rightarrow A$ a map of animated k -algebras with the A -module $L_{A/R}$ being p -completely flat, such that **either***

- *the R -module A has bounded p -complete Tor-amplitude, the R -module $L_{R/k}$ is p -completely flat, and R is p -completely t -bounded; **or***
- *the Hodge-filtration on $\widehat{\mathrm{dR}}_{R \otimes_{\mathbb{Z}} \mathbb{F}_p / k \otimes_{\mathbb{Z}} \mathbb{F}_p}$ converges universally, the spectrum A is p -completely t -bounded, the R -module $L_{R/k}$ has p -complete Tor-amplitude ≤ 1 , and the spectrum $\varphi_{k \otimes_{\mathbb{Z}} \mathbb{F}_p}^*(A \otimes_{\mathbb{Z}} \mathbb{F}_p)$ is t -bounded¹⁹.*

If the p -completed derived de Rham cohomology $\mathrm{dR}_{A/R}$ is p -completely Hodge-complete, then the filtered spectrum

$$\mathrm{Fil}_H \mathrm{dR}_{A/k} \otimes_{\mathrm{Fil}_H \mathrm{dR}_{R/k}}^{\mathbb{L}} \mathrm{Fil}_H \widehat{\mathrm{dR}}_{R/k}$$

is p -completely complete. In particular, if the p -completed derived de Rham cohomology $\mathrm{dR}_{R/k}$ is additionally p -completely Hodge complete, then so is $\mathrm{dR}_{A/k}$.

Proof. We could replace k by $k \otimes_{\mathbb{Z}} \mathbb{F}_p$, and without loss of generality, we may endow k an animated $\mathbb{F}_p$ -algebra structure. By Corollary 6.29, it suffices to show that, for every $i \in \mathbb{N}$, the Hodge-filtration on

$$\begin{aligned} \mathrm{gr}_{\mathrm{KO}/R}^i(\mathrm{dR}_{A/k} \otimes_{\mathrm{dR}_{R/k}}^{\mathbb{L}} \widehat{\mathrm{dR}}_{R/k}) &= \mathrm{dR}_{A/k} \otimes_{\mathrm{dR}_{R/k}}^{\mathbb{L}} \mathrm{gr}_H^i \mathrm{dR}_{R/k} \\ &= \mathrm{dR}_{A/R} \otimes_R^{\mathbb{L}} \left(\bigwedge_R^i L_{R/k} \right) [-i] \\ &= \widehat{\mathrm{dR}}_{A/R} \otimes_R^{\mathbb{L}} \left(\bigwedge_R^i L_{R/k} \right) [-i] \end{aligned}$$

is complete, where the last equivalence is Hodge-completeness of $\mathrm{dR}_{A/R}$.

- In the first case, as R is t -bounded above, and A has bounded Tor-amplitude over R , the spectrum A is t -bounded above as well. The result follows from Lemma 6.18 and the flatness of the A -module $L_{A/R}$.
- In the second case, the R -module $\left(\bigwedge_R^i L_{R/k} \right) [-i]$ has Tor-amplitude ≤ 0 (cf. [BMS19, Lem 4.34]), and the result follows from Lemma 6.18 and the flatness of the A -module $L_{A/R}$ as well. $\square$

We can also establish a version of transitivity of universal convergence of Hodge filtrations via the Katz–Oda filtration:

Theorem 6.31. *Let k be an animated $\mathbb{F}_p$ -algebra, and $R \rightarrow A$ be a quasi-lci map of animated k -algebras, such that*

¹⁹This is satisfied if the Frobenius map $\varphi_{k \otimes_{\mathbb{Z}} \mathbb{F}_p}$ has bounded p -complete Tor-amplitude, as we have already assumed that A is p -completely t -bounded.

- the Hodge-filtration on $\widehat{dR}_{R/k}$ converges universally of index $\alpha \in \mathbb{N}$, with R -module $L_{R/k}$ being flat; and
- the Hodge-filtration on $\widehat{dR}_{A/R}$ converges universally of index $\beta \in \mathbb{N}$, and the Hodge-filtration on $dR_{A/R}$ is complete.

Then the Hodge filtration on

$$dR_{A/k} \otimes_{dR_{R/k}}^{\mathbb{L}} \widehat{dR}_{R/k},$$

viewed as a filtered $dR_{A/k}$ -module, converges universally of index $\alpha + \beta$.

Proof. Let $r \in \mathbb{N}$. By definition, we have to control the Tor-amplitude of the $\varphi_k^* A$ -module

$$E_r := \mathrm{Fil}_H^r(dR_{A/k} \otimes_{dR_{R/k}}^{\mathbb{L}} \widehat{dR}_{R/k}).$$

We filter this $\varphi_k^* A$ -module by the Katz–Oda filtration, and the key observation is that, the s -th Katz–Oda filtration piece of $\mathrm{Fil}_H^r(dR_{A/k} \otimes_{dR_{R/k}}^{\mathbb{L}} \widehat{dR}_{R/k})$ is the same as

$$dR_{A/k} \otimes_{dR_{R/k}}^{\mathbb{L}} \mathrm{Fil}_H^s \widehat{dR}_{R/k}$$

for $s \geq r$ (which is in general false for $s < r$)²⁰. Let $T_r := dR_{A/k} \otimes_{dR_{R/k}}^{\mathbb{L}} \mathrm{Fil}_H^r \widehat{dR}_{R/k}$. By universal convergence of the Hodge filtration on $\widehat{dR}_{R/k}$, it follows from Lemma 6.28 that the $\varphi_k^* A$ -module T_r has Tor-amplitude $\leq \alpha - r \leq \alpha + \beta - r$.

Now it remains to see that, the s -th associated graded pieces of Katz–Oda filtration on

$$\mathrm{Fil}_H^r(dR_{A/k} \otimes_{dR_{R/k}}^{\mathbb{L}} \widehat{dR}_{R/k})$$

has Tor-amplitude $\leq -r$ over $\varphi_k^* A$ for $s < r$. This is precisely given by

$$(4) \quad \mathrm{Fil}_H^{r-s} \widehat{dR}_{A/R} \otimes_R^{\mathbb{L}} \left(\bigwedge_R^s L_{R/k} \right) [-s]$$

as in the proof of Theorem 6.30, which has Tor-amplitude $\leq \beta + (s - r) - s = \beta - r$ over

$$\varphi_R^* A = \varphi_k^*(A) \otimes_{\varphi_k^* R, \varphi_{R/k}}^{\mathbb{L}} R.$$

By universal convergence of the Hodge-filtration on $\widehat{dR}_{R/k}$ and Lemma 6.22, the relative Frobenius map $\varphi_{R/k} : \varphi_k^* R \rightarrow R$ has Tor-amplitude $\leq \alpha$. It follows that (4) has Tor-amplitude $\leq \alpha + \beta - r$ over $\varphi_k^* A$. To conclude, the $\varphi_* A$ -module E_r has Tor-amplitude $\leq \alpha + \beta - r$. $\square$

Remark 6.32. We may prove Theorem 6.30 in the same line as in the proof of Theorem 6.31, which does not invoke Corollary 6.29 explicitly, but a more “quantitative” version of that, i.e. the t -boundedness of “Katz–Oda tails” T_r .

The Nygaard filtration on $\widehat{\Delta}_R$ might be more complicated. However, if we take its reduction mod p , we can take the *twisted Nygaard filtration* instead.

²⁰More generally, let $\mathrm{Fil} R$ be an $(\mathbb{N}, \geq)$ -filtered $\mathbb{E}_1$ -ring, and $\mathrm{Fil} M$ an $(\mathbb{N}, \geq)$ -filtered right $\mathrm{Fil} R$ -module, and $\mathrm{Fil} N$ a $\mathbb{Z}$ -filtered left $\mathrm{Fil} R$ -module concentrated in weights $\geq s$. Then the filtered map

$$\mathrm{Fil} M \otimes_{\mathrm{Fil} R}^{\mathbb{L}} \mathrm{Fil} N \rightarrow \mathrm{ins}^0 \mathrm{Fil}^0 M \otimes_{\mathrm{ins}^0 \mathrm{Fil}^0 R}^{\mathbb{L}} \mathrm{Fil} N$$

is an equivalence. Indeed, the description of the relative tensor product in terms of bar construction allows us to reduce to the special case that $\mathrm{Fil} R$ is the unit. Then it follows directly from cofinality.

Remark 6.33 ([BM23, Cons 3.3, Rem 3.4 & Rem 4.5]). Let R be an animated ring. Then there exists a functorial $\mathbb{N}$ -graded filtration $\text{Fil}_{\tilde{N}}(\bigoplus_{n \in \mathbb{Z}} \bar{\Delta}_R\{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$, called the *twisted Nygaard filtration*, on the graded $\mathbb{E}_{\infty}$ -ring $\bigoplus_{n \in \mathbb{Z}} \bar{\Delta}_R\{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$, with associated graded pieces given by

$$\text{cofib} \left(\mu_1 : \bigoplus_{i \in \mathbb{N}} \bigoplus_{n \in \mathbb{Z}} \text{gr}_N^{i-p} \Delta_R\{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \longrightarrow \bigoplus_{i \in \mathbb{N}} \bigoplus_{n \in \mathbb{Z}} \text{gr}_N^i \Delta_R\{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \right).$$

Then

- (1) the $\text{Fil}_N(\bigoplus_{n \in \mathbb{Z}} \bar{\Delta}_R\{n\})$ -module $\text{Fil}_{\tilde{N}}(\bigoplus_{n \in \mathbb{Z}} \bar{\Delta}_R\{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ is perfect, as it is the cofiber of a twisted endomorphism given by a class in [BM23, Ex 3.2];
- (2) the mod p -reduction $\bar{\Delta}_R\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ of Nygaard-completed absolute Hodge–Tate cohomology coincides with the completion of the twisted Nygaard completion of $\bar{\Delta}_R\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$; and
- (3) if the animated ring R is weakly F -smooth and p -completely t -bounded above, then the convergence $\text{Fil}_{\tilde{N}}(\bigoplus_{n \in \mathbb{N}} \widehat{\bar{\Delta}}_R\{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ has a linearly increasing coconnectivity estimate (the neutral t -structure on graded spectra implies uniformity in $n \in \mathbb{Z}$).

Warning 6.34. Let R be an animated ring. The $R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ -module structure on $\text{gr}_N^* \bar{\Delta}_R$ inherited from that on $\text{gr}_N^* \Delta_R$ is incompatible with the one inherited from $\bar{\Delta}_R$. They differ up to a Frobenius, as explained in [BM23, Cons 3.1]. The same for the twisted Nygaard filtration. To fix the terminology, we mean the $R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ -module structure on $\text{gr}_N^* \bar{\Delta}_R$ inherited from that on $\text{gr}_N^* \Delta_R$ by default, unless otherwise explicitly said.

Remark 6.35. By Remark 6.33, we may replace $\bar{\Delta}$ by $\bar{\Delta} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$, and Fil_N by $\text{Fil}_{\tilde{N}}$ throughout in, say, Remark 6.14.

Example 6.36 (cf. [BM23, Ex 3.2]). Let R be a p -torsion-free perfectoid ring, and let (A, I) denote its associated perfect (transversal) prism. Then the prismatic cohomology Δ_R is A , with Nygaard filtration given by the $\varphi^{-1}(I)$ -adic filtration. As the Cartier divisor $I \subseteq A$ is distinguished, we know that $\varphi^{-1}(I)^p$ becomes (p) in $\bar{\Delta}_R = R$, and the filtered ring $\text{Fil}_{\tilde{N}}(\bigoplus_{n \in \mathbb{N}} \widehat{\bar{\Delta}}_R\{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ is concretely given by

$$R\{*\}/p \longleftarrow \varphi^{-1}(I)R\{*\}/p \longleftarrow \cdots \longleftarrow \varphi^{-1}(I)^{p-1}R\{*\}/p \longleftarrow 0 \longleftarrow \cdots$$

which is concentrated in weights $[0, p-1]$.

The following lemma controls “Tor-amplitude” of the map $\bar{\Delta}_k \rightarrow \bar{\Delta}_R$ for quasi-lci maps $k \rightarrow R$ of bounded Tor-amplitude, an analogue of Lemma 6.28.

Lemma 6.37. *Let $k \rightarrow R$ be a p -quasi-lci map of animated rings. Then the graded $R \otimes_k^{\mathbb{L}} (\bigoplus_{n \in \mathbb{Z}} \bar{\Delta}_k\{n\})$ -module spectrum $\bigoplus_{n \in \mathbb{Z}} \bar{\Delta}_R\{n\}$ admits an exhaustive increasing filtration, with associated graded pieces being retracts of graded $R \otimes_k^{\mathbb{L}} (\bigoplus_{n \in \mathbb{Z}} \bar{\Delta}_k\{n\})$ -modules of the form*

$$N \otimes_k^{\mathbb{L}} \left(\bigoplus_{n \in \mathbb{Z}} \bar{\Delta}_k\{n\} \right)$$

with N being a R -module spectrum with p -complete Tor-amplitude ≤ 0 .

Proof. We simply take the Katz–Oda filtration in Example 5.4, and the associated graded pieces are direct summands (thus retracts) of the graded graded $R \otimes_k^{\mathbb{L}} (\bigoplus_{n \in \mathbb{Z}} \bar{\Delta}_k\{n\})$ -module

$$\left(\bigoplus_{s \in \mathbb{N}} \left(\bigwedge_R^s L_{R/k} \right) [-s] \right) \otimes_k^{\mathbb{L}} \left(\bigoplus_{n \in \mathbb{Z}} \bar{\Delta}_k\{n\} \right).$$

Now as $k \rightarrow R$ is p -quasi-lci, for every $s \in \mathbb{N}$, the R -module $(\bigwedge_R^s L_{R/k})[-s]$ has p -complete Tor-amplitude ≤ 0 (cf. [BMS19, Lem 4.34]), thus so does the R -module

$$N := \bigoplus_{s \in \mathbb{N}} \left(\bigwedge_R^s L_{R/k} \right) [-s].$$

□

With this Tor-amplitude of absolute prismatic cohomology, we deduce the completeness of “Katz–Oda filtration” out of twisted Nygaard filtration, just as the first case of Corollary 6.29:

Corollary 6.38. *Let $k \rightarrow R$ be a p -quasi-lci map of animated rings such that k is p -completely t -bounded and R has bounded p -complete Tor-amplitude over k . Suppose that the animated ring k is weakly F -smooth. Then the filtered graded spectrum*

$$\left(\bigoplus_{n \in \mathbb{Z}} \overline{\Delta}_R \{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \right) \otimes_{\bigoplus_{n \in \mathbb{Z}} \overline{\Delta}_k \{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p}^{\mathbb{L}} \left(\bigoplus_{n \in \mathbb{Z}} \text{Fil}_{\tilde{N}} \left(\widehat{\overline{\Delta}}_k \{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \right) \right)$$

is complete, and its convergence has a linearly increasing p -complete coconnectivity estimate.

Proof. The result follows from Remark 6.33 and Lemma 6.37 (cf. the proof of Corollary 6.29). □

Now we follow the same line as in the proof of Theorem 6.30.

Proof of Theorem 6.15 By Remarks 6.14 and 6.35 and Corollary 6.38, it suffices to check that, for every $j \in \mathbb{N}$, the filtered graded spectrum

$$F_j := \left(\bigoplus_{n \in \mathbb{Z}} \text{Fil}_{\tilde{N}} \left(\overline{\Delta}_R \{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \right) \right) \otimes_{\bigoplus_{n \in \mathbb{Z}} \text{Fil}_N \left(\overline{\Delta}_k \{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \right)}^{\mathbb{L}} \left(\bigoplus_{n \in \mathbb{Z}} \text{ins}^j \text{gr}_N^j \overline{\Delta}_k \{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \right)$$

is complete. We start with $j = 0$. By point 1 of Remark 6.33, it suffices to check the completeness of the filtered graded spectrum

$$\left(\bigoplus_{n \in \mathbb{Z}} \text{Fil}_N \overline{\Delta}_R \{n\} \right) \otimes_{\bigoplus_{n \in \mathbb{Z}} \text{Fil}_N \overline{\Delta}_k \{n\}}^{\mathbb{L}} \left(\bigoplus_{n \in \mathbb{Z}} \text{ins}^0 \text{gr}_N^0 \overline{\Delta}_k \{n\} \right)$$

after p -completion. By de Rham comparison, this filtered spectrum can be identified with the filtered graded spectrum $\bigoplus_{n \in \mathbb{Z}} \text{Fil}_H dR_{R/k} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$, which is complete.

Now we fix $j \in \mathbb{N}$. Note that we have

$$F_j = F_0 \otimes_{\bigoplus_{n \in \mathbb{Z}} \text{ins}^0 \text{gr}_N^0 \left(\overline{\Delta}_k \{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \right)}^{\mathbb{L}} \bigoplus_{n \in \mathbb{Z}} \text{ins}^j \text{gr}_N^j \left(\overline{\Delta}_k \{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \right).$$

By weak F -smoothness of k and Remark 6.33 that the filtered $\left(\bigoplus_{n \in \mathbb{Z}} \text{ins}^0 \text{gr}_N^0 \left(\overline{\Delta}_k \{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \right) \right)$ -module

$$\bigoplus_{n \in \mathbb{Z}} \text{ins}^j \text{gr}_N^j \left(\overline{\Delta}_k \{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \right)$$

has bounded Tor-amplitude, and the completeness of F_j then follows from the completeness of F_0 and the convergence estimate by Lemma 6.18. □

Finally, inspired by Definition 6.20, we introduce universal convergence for Nygaard filtration:

Definition 6.39. Let A be an animated ring. We say that a filtered graded $\mathrm{Fil}_{\tilde{N}}(\overline{\Delta}_A\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ -module $\mathrm{Fil} M$ *converges universally of index* $\alpha \in \mathbb{Z}$ if, for every $r \in \mathbb{N}$, the graded A -module $\mathrm{Fil}^r M$, inherited from the A -module structure on $\overline{\Delta}_A$ (cf. Warning 6.34) has p -complete Tor-amplitude²¹ $\leq \alpha - r$, and it *converges universally* if such an $\alpha \in \mathbb{Z}$ exists. In particular, we say that the Nygaard filtration²² on $\widehat{\Delta}_A\{*\}$ *converges universally (resp. of index* $\alpha \in \mathbb{Z})$ if the filtered $\mathrm{Fil}_{\tilde{N}}(\overline{\Delta}_A\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ -module $\mathrm{Fil}_{\tilde{N}}(\widehat{\Delta}_A\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ converges universally (resp. of index $\alpha \in \mathbb{Z}$).

Remark 6.40. Let A be an animated ring. Unlike in Warning 6.34, when we talk about universal convergence, we mean the A -module structure on $\mathrm{Fil}_{\tilde{N}}(\overline{\Delta}_A\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ inherited from the A -module structure on $\overline{\Delta}_A$, while when we talk about associated graded pieces $\mathrm{gr}_{\tilde{N}}(\overline{\Delta}_A\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$, we refer to that inherited from $\mathrm{gr}_N \Delta_A$.

Similar to Lemma 6.22, the universal convergence of Nygaard filtration gives rise to a bound of Tor-amplitude of Frobenius.

Lemma 6.41. *Let R be an animated ring, and $\mathrm{Fil} M$ a filtered graded $\mathrm{Fil}_{\tilde{N}}(\overline{\Delta}_R\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ -module equipped with a map $\mathrm{Fil}_{\tilde{N}}(\overline{\Delta}_R\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p) \rightarrow \mathrm{Fil} M$ which induces an equivalence on associated graded pieces. Suppose that the $\mathrm{Fil}_{\tilde{N}}(\overline{\Delta}_R\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ -module $\mathrm{Fil} M$ converges universally of index $\alpha \in \mathbb{Z}$. Then the Frobenius map $\psi_R : R \rightarrow R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ has p -complete Tor-amplitude $\leq \alpha$.*

Proof. We examine the fiber sequence

$$\begin{array}{ccccc} & & \mathrm{gr}_{\tilde{N}}^0(\overline{\Delta}_R\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p) & & \\ & & \downarrow \simeq & & \\ \mathrm{Fil}^0 M & \longrightarrow & \mathrm{gr}^0 M & \longrightarrow & \mathrm{Fil}^1 M[1] \end{array}$$

of R -modules, where unlike in Remark 6.40, we view the top $\mathrm{gr}_{\tilde{N}}^0(\overline{\Delta}_R\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ as an R -module inherited from the map $R \rightarrow \overline{\Delta}_R$. By assumption, we know that the map $\psi_R : R \rightarrow R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p = \mathrm{gr}_{\tilde{N}}^0(\overline{\Delta}_R\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ has Tor-amplitude $\leq \alpha$. $\square$

Similarly to the proof of the second case of Theorem 6.30, it follows from the proof of Theorem 6.15 that the following variant of transitivity without boundedness of Tor-amplitude when the Nygaard filtration on the base converges universally.

Theorem 6.42. *Let $k \rightarrow R$ be a quasi-lci map of animated rings such that R is p -completely t -bounded. Suppose that*

- *the animated ring k is weakly F -smooth and the Nygaard filtration on $\widehat{\Delta}_k\{*\}$ converges universally; and that*
- *the R -module spectrum $L_{R/k}$ is p -completely flat.*

If the p -completed derived de Rham cohomology $\mathrm{dR}_{R/k}$ is p -completely Hodge-complete, then the absolute prismatic cohomology $\Delta_R\{\}$ is Nygaard-complete relative to $\Delta_k\{*\}$.*

Adapting the proof of Theorem 6.15 in the line the proof of Theorem 6.31 (cf. Remark 6.32), we get a transitivity of universal convergence of Nygaard filtration.

Theorem 6.43. *Let $k \rightarrow R$ be a quasi-lci map of animated rings. Suppose that*

²¹By Corollary 6.10, it is equivalent to non- p -complete Tor-amplitude.

²²Here we are actually considering the twisted Nygaard filtration is actually on $\widehat{\Delta}_A\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$, and thus this is an abuse of language.

- the animated ring k is weakly F -smooth and the Nygaard filtration on $\widehat{\Delta}_k\{*\}$ converges universally of index $\alpha \in \mathbb{N}$; and that
- the Hodge filtration on $dR_{R/k}$ is p -completely complete, and the Hodge filtration on $\widehat{dR}_{R \otimes_{\mathbb{Z}} \mathbb{F}_p / k \otimes_{\mathbb{Z}} \mathbb{F}_p}$ converges universally of index $\beta \in \mathbb{N}$.

Then the filtered graded $\mathrm{Fil}_{\widehat{N}}(\overline{\Delta}_R\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ -module

$$\mathrm{Fil}_{\widehat{N}}\left(\overline{\Delta}_R\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p\right) \otimes_{\mathrm{Fil}_{\widehat{N}}\left(\overline{\Delta}_k\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p\right)}^{\mathbb{L}} \mathrm{Fil}_{\widehat{N}}\left(\widehat{\Delta}_k\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p\right)$$

converges universally.

Proof. Let E_r denote the r -th filtered piece of the filtered graded $\mathrm{Fil}_{\widehat{N}}(\overline{\Delta}_R\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ -module above. By definition, we have to control its Tor-amplitude as a graded R -module, inherited from the graded R -module structure on $\overline{\Delta}_R\{*\}$. As in the proof of Theorem 6.31, we endow E_r with Katz–Oda filtration as in the proof of Theorem 6.15, induced by the twisted Nygaard filtration on $\widehat{\Delta}_k\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$, and we have to control the tail $\mathrm{Fil}_{\mathrm{KO}}^r E_r$ and $\mathrm{gr}_{\mathrm{KO}}^s E_r$ for $0 \leq s < r$.

As in the proof of Theorem 6.31, the tail $\mathrm{Fil}_{\mathrm{KO}}^r E_r$ is simply given by

$$\left(\bigoplus_{n \in \mathbb{Z}} \overline{\Delta}_R\{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p\right) \otimes_{\bigoplus_{n \in \mathbb{Z}} \overline{\Delta}_k\{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p}^{\mathbb{L}} \left(\bigoplus_{n \in \mathbb{Z}} \mathrm{Fil}_{\widehat{N}}^r\left(\widehat{\Delta}_k\{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p\right)\right).$$

Now the base change along $\overline{\Delta}_k\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \rightarrow \overline{\Delta}_R\{*\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ above can be rewritten as the (p -completed) base change along $\overline{\Delta}_k\{*\} \rightarrow \overline{\Delta}_R\{*\}$, and it follows from Lemma 6.37 that the graded R -module $\mathrm{Fil}_{\mathrm{KO}}^r E_r$ has p -complete Tor-amplitude $\leq \alpha - r$.

Now we control $\mathrm{gr}_{\mathrm{KO}}^s E_r$ for $0 \leq s < r$. As in the proof of Theorem 6.15, this is given by

$$\left(\bigoplus_{n \in \mathbb{Z}} \mathrm{Fil}_H^{r-s} dR_{R/k} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p\right) \otimes_{\bigoplus_{n \in \mathbb{Z}} \mathrm{gr}_N^0(\overline{\Delta}_k\{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)}^{\mathbb{L}} \bigoplus_{n \in \mathbb{Z}} \mathrm{gr}_N^s(\overline{\Delta}_k\{n\} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p).$$

As a graded $\varphi_{k \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p}^*(R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ -module, it has Tor-amplitude $\leq (\beta - (r - s)) + (p - 1 - s) = p + \beta - r - 1$. Note that

$$\varphi_{k \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p}^*(R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p) = R \otimes_{k, \psi_k}^{\mathbb{L}} (k \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p),$$

and it follows from Lemma 6.41 that the R -module $\mathrm{gr}_{\mathrm{KO}}^s E_r$ has p -complete Tor-amplitude $\leq p + \alpha + \beta - 1 - r$. $\square$

As a consequence, we deduce a version of [BM23, Prop 4.7] without boundedness of Tor-amplitude (or in particular, flatness) for maps $k \rightarrow R$ whose modulo p reduction is relatively perfect.

Corollary 6.44. *Let $k \rightarrow R$ be a map of animated rings such that the induced map $k \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p \rightarrow R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p$ of animated $\mathbb{F}_p$ -algebras is relatively perfect, and that R is p -completely t -bounded. If*

- the animated ring k is weakly F -smooth, the absolute prismatic cohomology Δ_k is Nygaard-complete, and the Nygaard filtration on $\widehat{\Delta}_k\{*\}$ converges universally,

then the same is true for R . That is to say,

- the animated ring R is weakly F -smooth, the absolute prismatic cohomology Δ_R is Nygaard-complete, and the Nygaard filtration on $\widehat{\Delta}_R\{*\}$ converges universally.

Proof. It follows from Theorems 6.12, 6.42 and 6.43 and Lemma 6.26. $\square$

Finally, we compare the universal convergence of Hodge and Nygaard filtration over a perfectoid base, an analogue of [BM23, Prop 4.12] or [Bou23, Thm 2.16 & 2.18].

Theorem 6.45. *Let S be a perfectoid ring, and R an animated S -algebra. If*

- *the Hodge filtration on the Hodge-completed derived de Rham cohomology $\widehat{\mathrm{dR}}_{(R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)/(S \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)}$ converges universally,*

then

- *the Nygaard filtration on the Nygaard-completed absolute prismatic cohomology $\widehat{\Delta}_R\{*\}$ converges universally.*

Actually, this admits a partial inverse, as we will see in Proposition 6.48. As we fix a perfectoid base, we may suppress all Breuil–Kisin twists $\{*\}$. We start with some recollections on absolute prismatic cohomology of rings over a perfectoid base.

Remark 6.46. Let (A, I) be a prism with $\overline{A} := A/I$, and R an animated $\overline{A}$ -algebra. Then we have the following statements.

- (1) When the prism (A, I) is perfect, the Nygaard-filtered absolute prismatic cohomology $\mathrm{Fil}_N \Delta_R$ can be identified with the Nygaard-filtered relative prismatic cohomology $\mathrm{Fil}_N \varphi_A^* \Delta_{R/A}$, by [BL22, Thm 5.6.2].
- (2) The Hodge-filtered derived de Rham cohomology $\mathrm{dR}_{R/S}$ can be identified with

$$\mathrm{Fil}_N (\varphi_A^* \Delta_{R/A}) \otimes_{(I^{\geq 0} A)} \mathrm{ins}^0 \overline{A}$$

by [BL22, Prop 5.2.3 & Cor 5.2.8], where $(I^{\geq 0} A)$ is the ring A with I -adic filtration, plus the observation that the inclusion $d : I \rightarrow A$ induces a map $I \otimes_A^{\mathbb{L}} (I^{\geq 0} A)(1) \rightarrow I^{\geq 0} A$ of filtered $(I^{\geq 0} A)$ -modules (cf. [BL22, Rem 5.1.9]) whose cofiber is $\mathrm{ins}^0 \overline{A}$. The same comparison for the Nygaard and Hodge-completed versions.

- (3) The modulo p reduction (or equivalently, modulo $\varphi(I)$ -reduction) of de Rham comparison

$$\varphi_A^* \Delta_{R/A} \otimes_A^{\mathbb{L}} \overline{A} \simeq \mathrm{dR}_{R/\overline{A}}$$

can be rewritten as

$$\varphi_A^* \overline{\Delta}_{R/A} \otimes_{A/\varphi(I)}^{\mathbb{L}} (\overline{A} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p) \simeq \mathrm{dR}_{(R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)/(\overline{A} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)}$$

which can be upgraded to an (increasingly) filtered equivalence, where both sides are equipped with conjugate filtration. In particular, this equivalence is an equivalence of derived $\varphi_{\overline{A} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p}^*(R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$ -algebras, where the derived algebra structure on the left hand side is given by the equivalence

$$\varphi_A^*(R) \otimes_{A/\varphi(I)}^{\mathbb{L}} (\overline{A} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p) \simeq \varphi_{\overline{A} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p}^*(R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$$

of animated rings. When (A, I) is a transversal prism and R is a polynomial $\overline{A}$ -algebra, both conjugate filtrations coincide with Whitehead towers, and the general case follows from left Kan extension²³.

The key to Theorem 6.45 is the following observation.

²³More precisely, we can pick a transversal prism (A_0, I_0) equipped with a map $(A_0, I_0) \rightarrow (A, I)$ of prisms as in [BL22, Prop .2.41], and then we can left Kan extend from the case $(A, A/I \rightarrow R)$, where A is a free δ - A_0 -algebra and R is a polynomial A/I -algebra, to general animated δ - A_0 -algebras A with an animated A/I -algebra R . Strictly speaking, this construction depends on the choice of $(A_0, I_0) \rightarrow (A, I)$, but in our case, we fix a base prism (A, I) , and we will not consider the functoriality with respect to the base prism (A, I) .

Lemma 6.47. *Let (A, I) be a perfect prism, and R an animated A/I -algebra. We start with filtered object*

$$\mathrm{Fil}_N(\varphi_A^* \widehat{\Delta}_{R/A}) \otimes_{(I^{\geq 0} A)} ((I^{\geq 0} A)/d^p)$$

where d^p is the map $I^p \otimes_A^{\mathbb{L}} (I^{\geq 0} A)(p) \rightarrow I^{\geq 0} A$, the p -th composite of the map d in Remark 6.46. Then the derived base change this filtered object along $A \rightarrow A/\varphi(I)$ can be identified with $\varphi_A^* \mathrm{Fil}_{\widehat{N}}(\overline{\Delta}_R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$. Concretely, we have a fiber sequence

$$I^p \otimes_A^{\mathbb{L}} \mathrm{Fil}_N^{r-p}(\varphi_A^* \overline{\Delta}_{R/A}) \xrightarrow{d^p} \mathrm{Fil}_N^r(\varphi_A^* \overline{\Delta}_{R/A}) \rightarrow \varphi_A^* \mathrm{Fil}_{\widehat{N}}^r(\overline{\Delta}_R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p)$$

of R -modules. The same for Nygaard-completed versions.

Proof. This follows essentially from the computation in [BM23, Ex 3.2], cf. Example 6.36. $\square$

Theorem 6.45 then follows from Lemma 6.47, Remark 6.46, and Proposition 6.48. Actually, Proposition 6.48 gives rise to a partial inverse of Theorem 6.45 as well.

Proposition 6.48. *Let (A, I) be a prism with $\overline{A} := A/I$, and R an animated $\overline{A}$ -algebra. Let $\alpha \in \mathbb{N}$ be a natural number. Then the following statements are equivalent:*

- (1) *For every $r \in \mathbb{N}$, the $\varphi_A^* R$ -module $\mathrm{Fil}_H^r \widehat{\mathrm{dR}}_{R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p / \overline{A} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p}$ has p -complete Tor-amplitude $\leq \alpha - r$.*
- (2) *For every $r \in \mathbb{N}$, the r -th filtered piece F_r of*

$$\mathrm{Fil}_N(\varphi_A^* \widehat{\Delta}_{R/A}) \otimes_{(I^{\geq 0} A)} ((I^{\geq 0} A)/d^p),$$

as a $\varphi_A^(R)$ -module, has p -complete Tor-amplitude $\leq p - 1 + \alpha - r$.*

Proof. We start with supposing the first statement. By the second point of Remark 6.46, we see that the $\varphi_A^* R$ -module F_r is a finite extension of $(\mathrm{Fil}_H^i \widehat{\mathrm{dR}}_{R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p / \overline{A} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p})_{i=r-p+1}^r$, thus it has p -complete Tor-amplitude $\leq \alpha - (r - p + 1) = p - 1 + \alpha - r$.

Now we start with the second statement. Morally, we argue by realizing $\mathrm{Fil}_N(\varphi_A^* \widehat{\Delta}_{R/A})/d$, up to a shift, as a retract of $(\mathrm{Fil}_N(\varphi_A^* \widehat{\Delta}_{R/A})/d^p)/d^{24}$. Concretely, we have commutative diagram

$$(5) \quad \begin{array}{ccccc} I^{p+1} \otimes_A^{\mathbb{L}} \mathrm{Fil}_N^{r-p-1}(\varphi_A^* \widehat{\Delta}_{R/A}) & \longrightarrow & I \otimes_A^{\mathbb{L}} \mathrm{Fil}_N^{r-1}(\varphi_A^* \widehat{\Delta}_{R/A}) & \longrightarrow & I \otimes_A^{\mathbb{L}} F_{r-1} \\ \downarrow & & \downarrow & & \downarrow \\ I^p \otimes_A^{\mathbb{L}} \mathrm{Fil}_N^{r-p}(\varphi_A^* \widehat{\Delta}_{R/A}) & \longrightarrow & \mathrm{Fil}_N^r(\varphi_A^* \widehat{\Delta}_{R/A}) & \longrightarrow & F_r \\ \downarrow & & \downarrow & & \\ I^p \otimes_A^{\mathbb{L}} \mathrm{Fil}_H^{r-p} \widehat{\mathrm{dR}}_{R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p / \overline{A} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p} & \longrightarrow & \mathrm{Fil}_H^r \widehat{\mathrm{dR}}_{R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p / \overline{A} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p} & & \end{array}$$

of $\varphi_A^* R$ -modules, where rows and columns are fiber sequences. This gives rise to a fiber sequence

$$G_r := \mathrm{fib}(I \otimes_A^{\mathbb{L}} F_{r-1} \rightarrow F_r) \longrightarrow I^p \otimes_A^{\mathbb{L}} \mathrm{Fil}_H^{r-p} \widehat{\mathrm{dR}}_{R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p / \overline{A} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p} \longrightarrow \mathrm{Fil}_H^r \widehat{\mathrm{dR}}_{R \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p / \overline{A} \otimes_{\mathbb{Z}}^{\mathbb{L}} \mathbb{F}_p}$$

of $\varphi_A^* R$ -module. Now the key observation is that the second map is null-homotopic, which is a consequence of Construction 6.49 and the fact that there exists a bottom-left to top-right map

$$d^{p-1} : I^p \otimes_A^{\mathbb{L}} \mathrm{Fil}_N^{r-p}(\varphi_A^* \widehat{\Delta}_{R/A}) \longrightarrow I \otimes_A^{\mathbb{L}} \mathrm{Fil}_N^{r-1}(\varphi_A^* \widehat{\Delta}_{R/A})$$

²⁴This is not as obvious as it seems: the Moore spectrum $\mathbb{S}/2$ is not a retract of the spectrum $(\mathbb{S}/2)/2$.

in the top left square of (5), which makes the diagram commute. It follows that

$$I^p \otimes_A^{\mathbb{L}} \mathrm{Fil}_H^{r-p} \widehat{\mathrm{dR}}_{R \otimes_{\mathbb{Z}} \mathbb{F}_p / \overline{A} \otimes_{\mathbb{Z}} \mathbb{F}_p}$$

is a retract of G_r . The fiber sequence

$$F_r[-1] \longrightarrow G_r \longrightarrow I \otimes_A^{\mathbb{L}} F_{r-1}$$

of $\varphi_A^* R$ -modules implies that the $\varphi_A^* R$ -module G_r has p -complete Tor-amplitude

$$\leq \max\{(p-1+\alpha-r)-1, p-1+\alpha-(r-1)\} = p+\alpha-r,$$

and thus the $\varphi_A^* R$ -module $\mathrm{Fil}_H^{r-p} \widehat{\mathrm{dR}}_{R \otimes_{\mathbb{Z}} \mathbb{F}_p / \overline{A} \otimes_{\mathbb{Z}} \mathbb{F}_p}$ has Tor-amplitude $\leq \alpha-(r-p)$ for every $r \in \mathbb{Z}$. The result then follows. $\square$

Construction 6.49. *Let $\mathcal{C}$ be a stable ∞ -category, and*

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ C & \longrightarrow & D \end{array}$$

a commutative diagram in $\mathcal{C}$. Then a lift $C \rightarrow B$, namely a map $C \rightarrow B$ along with homotopies as below,

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & \nearrow & \downarrow \\ C & \longrightarrow & D \end{array}$$

making two triangles $(A \rightarrow C \rightarrow B)$ and $(C \rightarrow B \rightarrow D)$ commute, gives rise to a null-homotopy of the canonical map

$$\mathrm{can} : \mathrm{cofib}(A \rightarrow C) \rightarrow \mathrm{cofib}(B \rightarrow D).$$

Indeed, such a lift splits the square into two triangles, which gives rise to two “octahedral” fiber sequences

$$\mathrm{cofib}(A \rightarrow C) \longrightarrow \mathrm{cofib}(A \rightarrow B) \longrightarrow \mathrm{cofib}(C \rightarrow B)$$

and

$$\mathrm{cofib}(C \rightarrow B) \longrightarrow \mathrm{cofib}(C \rightarrow D) \longrightarrow \mathrm{cofib}(B \rightarrow D).$$

The map can is the composite of these two fiber sequences, which is null-homotopic.

APPENDIX A. WEAK FINITE GENERATION UNDER COUNTABLE GENERATION HYPOTHESIS

In this appendix, we show that, weak finite generation is a condition on π_0 for countably generated connective modules (Proposition A.1), as well as an easy characterization of weak finite generation for countably generated modules (Proposition A.2), which is comparable to relevant results in [FS88, §6]. We also give counterexamples (Examples A.4 and A.11) without countable generation, which were discovered by ChatGPT-6 Astra.

Proposition A.1. *Let R be a connective $\mathbb{E}_1$ -ring, and M a connective right R -module.*

- (1) *Suppose that the right R -module M is weakly finitely generated (of rank $\leq r$). Then the right $\pi_0(R)$ -module $\pi_0(M)$ is weakly finitely generated.*
- (2) *Conversely, suppose that the right R -module M is countably generated (i.e. the right $\pi_0(R)$ -module $\pi_0(M)$ is countably generated). If the right $\pi_0(R)$ -module $\pi_0(M)$ is weakly finitely generated, then so is the right R -module M .*

Proof. Suppose that the right R -module M is weakly finitely generated of rank $\leq r$. Then by definition, there exists a surjective map $P \rightarrow M$ of connective right R -modules with $\mathrm{rk}_R(P) \leq r$. Taking π_0 and by Corollary 3.7, we see that the right $\pi_0(R)$ -module $\pi_0(M)$ is weakly finitely generated of rank $\leq r$.

Conversely, suppose the right $\pi_0(R)$ -module $\pi_0(M)$ is generated by a countable sequence $(x_i) \in \pi_0(M)^\mathbb{N}$, and is weakly finitely generated of rank r . By definition, there exists a filtered system $(F_\alpha)_{\alpha \in J}$ of free right $\pi_0(R)$ -modules of rank $\leq r$, along with a surjection $f : \mathrm{colim}_\alpha F_\alpha \rightarrow \pi_0(M)$ of right $\pi_0(R)$ -modules.

Then by dependent choice, we can pick a chain $(\alpha_0 \rightarrow \alpha_1 \rightarrow \cdots)$ in J , such that the image of the composite map $F_{\alpha_j} \rightarrow \mathrm{colim}_\alpha F_\alpha \xrightarrow{f} \pi_0(M)$ contains $\{x_0, \dots, x_j\}$ for every $j \in \mathbb{N}$. By choosing a base for each F_{α_j} , we get a chain $(\tilde{F}_{\alpha_0} \rightarrow \tilde{F}_{\alpha_1} \rightarrow \cdots)$ of free right R -modules of rank $\leq r$, along with an identification of the base-changed map $(\tilde{F}_{\alpha_j} \rightarrow \tilde{F}_{\alpha_{j+1}}) \otimes_R^\mathbb{L} \pi_0(R)$ with the map $F_{\alpha_j} \rightarrow F_{\alpha_{j+1}}$ for every $j \in \mathbb{N}$, and compatible homotopies. This gives rise to a surjective map

$$\mathrm{colim}_{j \in \mathbb{N}} \tilde{F}_{\alpha_j} \longrightarrow M$$

of connective right R -modules, whose source is a flat right R -module of rank $\leq r$. $\square$

Countable generation hypothesis also allows us to deduce a criterion of weak finite generation similar to Proposition 3.6.

Proposition A.2. *Let R be a connective $\mathbb{E}_1$ -ring, and M a connective right R -module.*

- (1) *Suppose that the right R -module M is weakly finitely generated of rank $\leq r$. Then for every finite free right R -module F , every map $F \rightarrow M$ factors through some R^s for $s \leq r$.*
- (2) *Conversely, suppose that the right R -module M is countably generated, and for every finite free right R -module F , every map $F \rightarrow M$ factors through some R^s for $s \leq r$. Then the right R -module M is weakly finitely generated of rank $\leq r$.*

Proof. The first statement follows from Proposition 3.6. Now suppose that the right R -module M is countably generated. We pick a sequence $(x_n)_{n \in \mathbb{N}_{>0}}$ of elements of $\pi_0(M)$ generating the $\pi_0(R)$ -module $\pi_0(M)$. By dependent choice in $n \in \mathbb{N}$, we may pick sequences $(f_n : R^r \rightarrow M)_{n \in \mathbb{N}}$, $(\theta_n : R \rightarrow R^r)_{n \in \mathbb{N}_{>0}}$ and $(g_n : R^r \rightarrow R^r)_{n \in \mathbb{N}_{>0}}$ of maps of right R -modules, such that $f_0 = 0$, and for $n \in \mathbb{N}_{>0}$, the maps f_n , θ_n , and g_n of right R -modules are obtained by factoring $x_n \oplus f_{n-1} : R \oplus R^r \rightarrow M$ as

$$R \oplus R^r \xrightarrow{\theta_n + g_n} R^r \xrightarrow{f_n} M$$

by assumption. We take $P := \mathrm{colim} \left(R^r \xrightarrow{g_1} R^r \xrightarrow{g_2} R^r \xrightarrow{g_3} \cdots \right)$, and the previous data give rise to a map $f : P \rightarrow M$ of right R -modules. As the map $x_n : R \rightarrow M$ of right R -modules factors as $R \xrightarrow{\theta_n} R^r \xrightarrow{f_n} M$, we see that the map f is surjective (on π_0), and the result follows. $\square$

The countable generation in Propositions A.1 and A.2 is essential, and ChatGPT-6 Astra found counterexamples when the countable generation is missing. We start with counterexample to the second statement of Proposition A.2 without countable generation hypothesis.

Recall [ES93] that, letting R be an integral domain with $K := \mathrm{Frac}(R)$, we say that an R -module is *uniserial* if its submodules are totally ordered by inclusion, and such a module is *standard* if it is isomorphic to a subquotient of the R -module K .

Lemma A.3. *Let V be a valuation ring with $K := \mathrm{Frac}(V)$, and M an V -module. Then it is standard uniserial if and only if it is weakly finitely generated of rank ≤ 1 .*

Proof. Every sub- V -module N of K is torsion-free thus flat, and $\mathrm{rk}_V(N) \leq \mathrm{rk}_V(K) = 1$ by Lemma 3.8, every standard uniserial V -module is weakly finitely generated of rank ≤ 1 .

Conversely, suppose that M is weakly finitely generated of rank ≤ 1 . We pick a flat V -module P of rank ≤ 1 with a surjection $f : P \rightarrow M$. By Lemma 3.8, we have $\dim_K(P \otimes_V K) = \text{rk}_V(P) \leq 1$, and thus $P \hookrightarrow P \otimes_V K$ can be realized as a sub- V -module of K , and thus P is standard uniserial, and so is its quotient M . $\square$

Example A.4. By [ES93, Prop 8(i)], there exists a valuation ring V with a non-standard uniserial V -module M . This, along with Lemma A.3, shows the necessity of countable generation in Proposition A.2. Actually, the example M there is also divisible, which even exclude the weak finite generation of M by Proposition A.5.

Proposition A.5. *Let V be a valuation ring with $K := \text{Frac}(V)$, and M a uniserial V -module which is divisible. Suppose that the V -module M is weakly finitely generated, then it is of rank ≤ 1 .*

Proof. We separate the proposition into two cases.

- (1) Suppose that the V -module M contains an element $x \in M$ which is torsion-free. Then for every $y \in M$, since M is uniserial, either $Vx \subseteq Vy$ or $Vy \subseteq Vx$. In both cases, we see that, if y is torsion, then $y = 0$. Consequently, the V -module M is torsion-free, and $\dim_K(M \otimes_V K) = 1$, and thus M can be embedded into the V -module K , thus it is standard uniserial, and we invoke Lemma A.3 to deduce.
- (2) Now we suppose that V -module M is torsion. Let P be a flat V -module of rank $r > 1$ along with a surjection $f : P \rightarrow M$ of V -modules. We show that there exists a flat V -module along with a surjection $Q \rightarrow M$ such that $\text{rk}_V(Q) < r$. This implies what we want.

Pick a non-zero element $x \in P$, and let $L := P \cap Kx$ which is contained in the K -vector space $P \otimes_V K$. Then L is torsion-free thus flat, and it is of rank 1 by Lemma 3.8. Moreover, the kernel of the map $P \rightarrow P \otimes_V K \rightarrow (P \otimes_V K)/(Kx)$ is precisely L , which gives rise to an injection $P/L \hookrightarrow (P \otimes_V K)/(Kx)$ into a K -vector space. It follows that P/L is torsion-free thus flat, and it is of rank $< r$. We again separate into two cases.

- (a) Suppose that the restriction $f|_L : L \rightarrow M$ is surjective. Then we may take $Q := L$.
- (b) There exists $y \in M$ which is not contained in $f(L)$. Then by uniseriality of M , we have $f(L) \subseteq Vy$. As the V -module M is torsion, there exists a non-zero element $\alpha \in V$ such that $\alpha y = 0$, and thus $\alpha f|_L = 0$. It follows that the map $\alpha f : P \rightarrow M$ of V -modules induces a map $g : P/L \rightarrow M$ of V -modules. Since the V -module M is divisible, the map $\alpha f : P \rightarrow M$ is surjective, thus so is the induced map g . Thus we may take $Q := P/L$.

$\square$

Now we describe the counterexample to the second statement of Proposition A.1. Recall that, letting R be an integral domain with $K := \text{Frac}(R)$, an R -module M is *Matlis cotorsion* if $\text{Ext}_R^1(K, M) = 0$, it is *h-reduced* if $\text{Hom}_R(K, M) = 0$ (this is the case if M is reduced, as explained in [BP20]), and a torsion-free R -module M is *R-complete* if the canonical map

$$M \longrightarrow \lim_{s \in R \setminus 0} (M/sM)$$

is an isomorphism on π_0 . The following statement is mentioned in [FG07, §1], and we include a proof for convenience of readers.

Lemma A.6. *Let R be an integral domain with $K := \text{Frac}(R)$. Then a torsion-free R -module M is R -complete if and only if the following two conditions are satisfied.*

- (1) *The R -module M is h-reduced.*
- (2) *The R -module M is Matlis cotorsion.*

Proof. We rewrite the R -module K as the filtered colimit $\operatorname{colim}_{s \in R \setminus 0} Rs^{-1}$. Thus

$$\begin{aligned} \operatorname{RHom}_R(K, M) &= \lim_{s \in R \setminus 0} \operatorname{RHom}_R(Rs^{-1}, M) \\ &= \lim_{s \in R \setminus 0} sM \end{aligned}$$

where the limit is derived, and consequently, we get a fiber sequence

$$\operatorname{RHom}_R(K, M) \longrightarrow M \longrightarrow \lim_{s \in R \setminus 0} (M/sM)$$

in $D(R)$, which gives rise to a long exact sequence

$$0 \longrightarrow \operatorname{Hom}_R(K, M) \longrightarrow M \longrightarrow \pi_0\left(\lim_{s \in R \setminus 0} (M/sM)\right) \longrightarrow \operatorname{Ext}_R^1(K, M) \longrightarrow 0.$$

The result then follows. $\square$

Lemma A.7. *Let R be a Bézout domain, and P a flat R -module of rank $\leq r$. Then there exists a finite filtration on P whose associated graded pieces are flat R -modules of rank ≤ 1 .*

Proof. Let $K := \operatorname{Frac}(R)$, and $P_K := P \otimes_R K$. As P is a filtered colimit of finite free R -modules of rank $\leq r$, the P_K is a filtered colimit of K -vector spaces of dimension $\leq r$, thus $\dim_K(P_K) \leq r$. It follows that P_K admits a finite filtration

$$0 = F_0 \subseteq F_1 \subseteq \cdots \subseteq F_r = P_K$$

of K -vector spaces with $\dim_K(F_j/F_{j-1}) \leq 1$ for $j = 1, \dots, r$. We take $H_j := P \cap F_j$. Then the induced map $H_j/H_{j-1} \rightarrow F_j/F_{j-1}$ is injective, and thus the R -module H_j/H_{j-1} is torsion-free, therefore flat as R is a Prüfer domain.

Now we show that $\operatorname{rk}_R(H_j/H_{j-1}) \leq 1$. If $F_j = F_{j-1}$, then $H_j = H_{j-1}$ and the result is true. Otherwise, $\dim_K(F_j/F_{j-1}) = 1$, and we get an injection $H_j/H_{j-1} \hookrightarrow K$ of R -modules. Now for every finite collection $\{x_1/y_1, x_2/y_2, \dots, x_m/y_m\} \subseteq H_j/H_{j-1}$, as R is Bézout, the sub- R -module generated by them is principal, generated by

$$(y_1 \cdots y_m)^{-1} \gcd(x_1 y_2 \cdots y_m, y_1 x_2 \cdots y_m, \dots, y_1 y_2 \cdots x_m).$$

It follows that $\operatorname{rk}_R(H_j/H_{j-1}) \leq 1$. $\square$

Remark A.8. Lemma A.7 gives rise to another proof of Lemma 3.8.

Corollary A.9. *Let V be a valuation ring which is complete with respect to the valuation, and $K := \operatorname{Frac}(V)$. Then for every flat V -module P of finite rank, we have $\operatorname{Ext}_V^1(K, P) = 0$.*

Proof. The statement is true for $P = 0$. Suppose that $P \neq 0$ and $\operatorname{rk}_V(P) = 1$. Then the V -module P embeds into $P \otimes_V K$, which is a K -vector space of dimension 1 as in the proof of Lemma A.7, thus we may view P as a sub- V -module of K . We separate two cases:

- (1) If $P = K$, then as the V -module K is injective, we have $\operatorname{Ext}_V^1(K, P) = 0$.
- (2) Suppose that $P \neq K$. We may rescale the module P so that there exists $d \in V \setminus 0$ such that $Vd \subseteq P \subseteq V$. This implies that the “ V -topology” on P is the same as the subspace topology of P in V , and that P is clopen in V , thus it is complete with respect to the topology as well, and therefore V -complete. It follows that $\operatorname{Ext}_V^1(K, P) = 0$ by Lemma A.6.

The general case then follows from Lemma A.7. $\square$

Corollary A.10. *Let V be a valuation ring which is complete with respect to the valuation, and $K := \operatorname{Frac}(V)$. Let M be a weakly finitely generated connective V -module along with a map $\varphi : M \rightarrow K$ in $D(V)$ which is surjective on π_0 . Then the map $M \rightarrow K$ admits a section in $D(V)$.*

Proof. Let P be a flat V -module of finite rank along with a surjective map $P \rightarrow M$ of connective V -modules. Then the composite map $f : P \rightarrow M \rightarrow K$ is a surjective map of V -modules concentrated in degree 0. Since the V -module $\text{Ker}(f) \subseteq P$ is a torsion-free thus flat, and $\dim_K(\text{Ker}(f) \otimes_V K) \leq \dim_K(P \otimes_V K) < \infty$, by Lemma 3.8, the V -module $\text{Ker}(f)$ is also flat of finite rank, thus by Corollary A.9, we have $\text{Ext}_V^1(K, \text{Ker}(f)) = 0$, thus the surjection f admits a section, and so does the surjective map φ . $\square$

Example A.11 (ChatGPT-6 Astra). Let $\Gamma := \bigoplus_{\alpha < \omega_1} \mathbb{Z}e_\alpha$ be the totally ordered abelian group, where $\sum a_\alpha e_\alpha < \sum b_\alpha e_\alpha$ if there exists $\beta < \omega_1$ such that, for every $\beta > \alpha$, we have $a_\alpha = b_\alpha$, and $a_\beta < b_\beta$. Let k be a field. We consider the Hahn's valued field $K := k((t^\Gamma))$, which is complete with respect to its valuation v , with $V := \{x \in K \mid v(x) \geq 0\}$. Then by [Oso67, Thm A & the last paragraph of §2], and that K is not $\aleph_0$ -generated (as ω_1 is $\aleph_0$ -filtered) but $\aleph_1$ -generated, the projective dimension $\text{pd}_V(K)$ is 2.

Now we pick a connective V -module M with $\pi_0(M) = K$ with projective dimension $\text{pd}_V(M) \leq 1$ (e.g. let $F_1 \rightarrow F_0 \rightarrow K$ be a free presentation of K , and take $M := \text{cofib}(F_1 \rightarrow F_0)$). If M is weakly finitely generated, then by Corollary A.10, the Postnikov truncation map $M \rightarrow K$ admits a section in $D(V)$, which is impossible as $\text{pd}_V(M) < \text{pd}_V(K)$. Thus the connective V -module M is not weakly finitely generated, while the V -module $\pi_0(M) = K$ is weakly finitely generated of rank ≤ 1 .

REFERENCES

- [AKN23] Benjamin Antieau, Achim Krause, and Thomas Nikolaus. Prismatic cohomology relative to δ -rings. *ArXiv e-prints*, oct 2023.
- [AKN24] Benjamin Antieau, Achim Krause, and Thomas Nikolaus. On the K -theory of $\mathbb{Z}/p^n$. *ArXiv e-prints*, may 2024.
- [AR24] Benjamin Antieau and Noah Riggenbach. Cyclotomic synthetic spectra. *ArXiv e-prints*, nov 2024.
- [BF26] Manuel Blickle and Daniel Fink. L -smooth factorization for Noetherian F -finite rings. *Adv. Math.*, 500, 2026.
- [Bha12] B. Bhatt. p -adic derived de Rham cohomology. *ArXiv e-prints*, apr 2012.
- [BHLS23] Robert Burklund, Jeremy Hahn, Ishan Levy, and Tomer M. Schlank. K -theoretic counterexamples to Ravenel's telescope conjecture. *ArXiv e-prints*, oct 2023.
- [BL22] Bhargav Bhatt and Jacob Lurie. Absolute prismatic cohomology. *ArXiv e-prints*, jan 2022.
- [BLM18] Bhargav Bhatt, Jacob Lurie, and Akhil Mathew. Revisiting the de Rham-Witt complex. *ArXiv e-prints*, may 2018.
- [BM23] Bhargav Bhatt and Akhil Mathew. Syntomic complexes and p -adic étale Tate twists. *Forum Math. Pi*, 11:0, 2023.
- [BMS18] Bhargav Bhatt, Matthew Morrow, and Peter Scholze. Integral p -adic Hodge theory. *Publications Mathématiques. Institut de Hautes Études Scientifiques*, 128:219–397, 2018.
- [BMS19] Bhargav Bhatt, Matthew Morrow, and Peter Scholze. Topological Hochschild homology and integral p -adic Hodge theory. *Publications Mathématiques. Institut de Hautes Études Scientifiques*, 129:199–310, 2019.
- [Bou23] Tess Vincent Bouis. Cartier smoothness in prismatic cohomology. *J. Reine Angew. Math.*, 805:241–282, 2023.
- [BP20] Silvana Bazzoni and Leonid Positselski. Matlis category equivalences for a ring epimorphism. *J. Pure Appl. Algebra*, 224(10):106398–25, 2020.
- [BPŠ22] Silvana Bazzoni, Leonid Positselski, and Jan Štoviček. Projective covers of flat contramodules. *Int. Math. Res. Not. IMRN*, 2022(24):19527–19564, 2022.
- [Bur22] Robert Burklund. Multiplicative structures on Moore spectra. *ArXiv e-prints*, mar 2022.
- [CMM21] Dustin Clausen, Akhil Mathew, and Matthew Morrow. K -theory and topological cyclic homology of henselian pairs. *J. Amer. Math. Soc.*, 34(2):411–473, 2021.
- [Cou13] François Couchot. Indecomposable injective modules of finite Malcev rank over local commutative rings. *J. Commut. Algebra*, 5(4):481–505, 2013.
- [DS18] Rankeya Datta and Karen E. Smith. Excellence in prime characteristic. In *Local and global methods in algebraic geometry*, volume 712 of *Contemp. Math.*, pages 105–116. Amer. Math. Soc., [Providence], RI, 2018.
- [ES93] P. C. Eklof and Saharon Shelah. Explicitly non-standard uniserial modules. *J. Pure Appl. Algebra*, 86(1):35–50, 1993.
- [FG07] László Fuchs and Rüdiger Göbel. Testing for cotorsionness over domains. *Rend. Semin. Mat. Univ. Padova*, 118:85–99, 2007.

- [FS88] L. Fuchs and L. Salce. Polyserial modules over valuation domains. *Rend. Semin. Mat. Univ. Padova*, 80:243–264, 1988.
- [GR03] Ofer Gabber and Lorenzo Ramero. *Almost ring theory*, volume 1800 of *Lect. Notes Math.* Berlin: Springer, 2003.
- [Gre25] Rok Gregoric. Prismaticization via spherical loop spaces. *ArXiv e-prints*, nov 2025.
- [H+98] J. van der Hoeven et al. GNU TeXmacs. <https://www.texmacs.org>, 1998.
- [HRW22] Jeremy Hahn, Arpon Raksit, and Dylan Wilson. A motivic filtration on the topological cyclic homology of commutative ring spectra. *ArXiv e-prints*, jun 2022.
- [HW22] Jeremy Hahn and Dylan Wilson. Redshift and multiplication for truncated Brown-Peterson spectra. *Ann. of Math. (2)*, 196(3):1277–1351, 2022.
- [Ill71] Luc Illusie. *Complexe cotangent et déformations. I*. Lecture Notes in Mathematics, Vol. 239. Springer-Verlag, Berlin-New York, 1971.
- [KST21] Moritz Kerz, Florian Strunk, and Georg Tamme. Towards Vorst’s conjecture in positive characteristic. *Compos. Math.*, 157(6):1143–1171, 2021.
- [Li26] Guo Li. A Counterexample to Bhatt-Lurie’s Cohomological Dimension Conjecture. *ArXiv e-prints*, jun 2026.
- [Lur17] Jacob Lurie. Higher algebra. <https://www.math.ias.edu/~lurie/papers/HA.pdf>, sep 2017.
- [Lur18] Jacob Lurie. Spectral algebraic geometry. <https://www.math.ias.edu/~lurie/papers/SAG-rootfile.pdf>, feb 2018.
- [Man24] Deven Manam. On the Drinfeld formal group. *ArXiv e-prints*, mar 2024.
- [Mao24] Zhouhang Mao. Equivariant aspects of de-completing cyclic homology. *ArXiv e-prints*, oct 2024.
- [Mat21] Akhil Mathew. On $K(1)$ -local TR. *Compos. Math.*, 157(5):1079–1119, 2021.
- [Oso67] B. L. Osofsky. Global dimension of valuation rings. *Trans. Am. Math. Soc.*, 127:136–149, 1967.
- [Rak26] Arpon Raksit. Hochschild homology and the derived de Rham complex revisited. *Selecta Math. (N.S.)*, 32(3):0, 2026.
- [Rig25] Noah Riggenbach. K -theory of truncated polynomials. *Math. Z.*, 310(3):0, 2025.
- [Sha70] Richard T. Shannon. The rank of a flat module. *Proc. Amer. Math. Soc.*, 24:452–456, 1970.
- [Sta21] StacksProjectAuthors. The Stacks project. <https://stacks.math.columbia.edu>, 2021.
- [Val73] Paolo Valabrega. Regular local rings and excellent rings. i, ii. *J. Algebra*, 26:440–445, 1973.