

Ahmad Yazdankhah

ahmad.yazdankhah@sjsu.edu
www.cs.sjsu.edu/~yazdankhah

Mathematical Preliminaries

(Part 2)

Lecture 03
Day 03/31

CS 154
Formal Languages and Computability
Spring 2018

Agenda of Day 03

- Waiting List Enrollment ...
- Summary of Lecture 02
- Lecture 03: Teaching ...
 - Mathematical Preliminaries (Part 2)

Summary of Lecture 02: We learned ...

Sets

- A **set** is ...
 - ... a collection of **distinct** elements.
- A **list** is ...
 - ... a collection of **ordered** elements.
- A **set is known** when its **boundary** is clearly defined.
- **Universal set** of a set is ...
 - ... the set containing all possible elements under consideration.
- **Three methods** to represent sets ...
 - Roster method
 - Venn diagram
 - **Set builder**

- The **power set** of the set S is ...
 - ... the set of all subsets of S .
 - It is denoted by 2^S .
 - $|2^S| = 2^{|S|}$
- A set is **finite** if ...
 - ... its size is a natural number.
- A set is **infinite** if ...
 - ... we cannot express its size by a natural number.

Any question?

Mathematical Preliminaries

Recap from Math 42

Cartesian Products

Motivation

- Recall that in sets, "order of elements" does NOT matter.
- But in practice, we do need "ordered collections".
- As we said before, in computer science we use "Lists" for ordered collections.
- The question is how we can mathematically model lists (ordered collections)?

Introduction

- Mathematicians defined a new mathematical structure called "n-tuple".
- An n-tuple is denoted by (a_1, a_2, \dots, a_n) .
 - A special case of n-tuple is 2-tuple aka "ordered pair" (a_1, a_2) .
- We use a mathematical operation called "Cartesian product" to create n-tuples.
- This operation is named after the great French philosopher, mathematician, and physicist René Descartes (1596-1650).



Cartesian Products Definition

Definition

- Let A and B be two sets.
- The Cartesian product of A and B is the set of all ordered pairs (a, b) , where $a \in A \wedge b \in B$.
- Cartesian product of A and B is denoted by $A \times B$.

- ⓘ ▪ Definition of Cartesian product by using set builder:

$$A \times B = \{(a, b) : a \in A \wedge b \in B\}$$



Cartesian Products **Examples**

Example 24

- Let $A = \{0, 1\}$, $B = \{3, 6, 9\}$; $A \times B = ?$
- $\{0, 1\} \times \{3, 6, 9\} = \{(0, 3), (0, 6), (0, 9), (1, 3), (1, 6), (1, 9)\}$

Example 25

- Let $Q = \{q_0, q_1, q_2\}$, $\Sigma = \{a, b\}$; $Q \times \Sigma = ?$

Example 26

- Let $Q = \{q_0, q_1\}$, $\Sigma = \{a, b\}$; $Q \times (\Sigma \cup \{\lambda\}) = ?$
 - " λ " is pronounced "**lambda**".



Cartesian Products Notes

- What is the result of the following Cartesian product?

$$A = \{1, 2\}, B = \phi; A \times B = ?$$

$$A \times B = \phi$$

– In fact, the result of Cartesian product would be ϕ if one of the sets is ϕ .



- Prove it!

- Is this a true statement: $A \times B = B \times A$

Does Cartesian product have commutative property?

In general, No!

It means: $A \times B \neq B \times A$

- But in the following special conditions, they can be equal:

$$A \times B = B \times A \text{ iff } (A = B) \vee (A = \phi) \vee (B = \phi)$$

Cartesian Products **Notes**

- We can **extend** the Cartesian product to **3 sets**:

$$A \times B \times C = \{(a, b, c) : a \in A, b \in B, c \in C\}$$

- But if you **associate** the sets, we'll get:

$$(A \times B) \times C = \{((a, b), c) : a \in A, b \in B, c \in C\}$$

$$A \times (B \times C) = \{(a, (b, c)) : a \in A, b \in B, c \in C\}$$

- Therefore, Cartesian product does **NOT** honor **association rule**.

$$(A \times B) \times C \neq A \times (B \times C)$$

- In this course, we don't associate the sets.

Cartesian Products **Extension**

- We can **extend the idea** to **n sets** and define **n-tuple** as:



$$S_1 \times S_2 \times \dots \times S_n = \{(x_1, x_2, \dots, x_n) : x_1 \in S_1, \dots, x_n \in S_n\}$$

Homework

- Let $Q = \{q_0, q_1\}$, $\Sigma = \{a, b\}$, $\Gamma = \{x, y\}$
- $Q \times (\Sigma \cup \{\lambda\}) \times \Gamma = ?$




Mathematical Preliminaries

Recap from Math 42

Functions

Introduction

- In many situations in real life, there is a **relationship between two sets**.
- For example, we assign a **letter grade** to each **student** of a class.



- This **relationship** is an example of the concept of "**function**".




Functions Definition

- Let D and R be two **sets**.
- A function from D to R is a **rule** that **assigns** (or **maps**) to **some** (could be **all**) elements of D a "**unique element**" of R .
 - The set D is called the "**domain**" of the function f .
 - The set R is called the "**range**" of the function f .

- In the **previous example**:
- **Domain** is the set of students
- **Range** is the set of letter grades

D(omain)

R(ange)



Functions **Naming** and **Notation**

- We usually **name a function** by **lower-case** letters such as f , h , δ (pronounced "**delta**"), etc.
- For **example**, the function δ is denoted by: $\delta : D \rightarrow R$


Example 27

$S = \{\text{John, Tom, Marry, Ted, Adam, Eve}\}$, $G = \{A, B, C, D, F\}$

$f : S \rightarrow G$


- What is **the rule of the function** shown by the figure?

$$\begin{cases} f(\text{John}) = A \\ f(\text{Tom}) = C \\ f(\text{Marry}) = B \\ f(\text{Adam}) = A \\ f(\text{Eve}) = F \end{cases}$$



Functions Notes

- $f(\text{Ted}) = ?$
- Undefined




- So, it is possible to have some elements in the domain that is **NOT mapped** to any value of the range. (e.g. **Ted** in the domain)
- Also, it is possible to have some elements in the range that is **NOT assigned** by any value of the domain. (e.g. **D** in the range)

Functions Notes

- Is it possible for Marry to have two grades at the same time?
- Definitely, **NO**.

In this universe, it cannot happen.




- That's why, in the definition of function, we said some elements of the domain are **uniquely mapped** to an element of the range.
- In other words, if there is a mapping, it should be unique.

Functions

Example 28: Mixing Cartesian Product and Function

- Let $Q = \{q_0, q_1, q_2\}$, $\Sigma = \{a, b\}$, $\delta : Q \times \Sigma \rightarrow Q$
- What is the domain and range of δ ?
- Domain:** $Q \times \Sigma = \{q_0, q_1, q_2\} \times \{a, b\} = \{(q_0, a), (q_0, b), (q_1, a), (q_1, b), (q_2, a), (q_2, b)\}$
- Range:** $\{q_0, q_1, q_2\}$
- The **rule** of δ is shown in the following figure. **Write the rule using algebraic notation.**
- Rule of function δ :**

$$\begin{cases} \delta(q_0, a) = q_0 \\ \delta(q_0, b) = q_1 \\ \delta(q_1, b) = q_2 \\ \delta(q_2, b) = q_2 \end{cases}$$





Homework

- Let $Q = \{q_0, q_1\}$, $\Sigma = \{a\}$, $\Gamma = \{x\}$, $\delta : Q \times \{\Sigma \cup \{\lambda\}\} \times \Gamma \rightarrow Q$
- What is the domain and range of δ ?


Total Function



- A function is called "total function" if all of its domain elements are defined.

Example 29

- The following function is "total function" because all domain elements are defined.




Partial Function

- ♥ If there exist at least one element in the domain of a function that is undefined, then the function is called "partial function".

Example 30

- The following function is "partial function" because at least one element of the domain is undefined:
- $f(8) = \text{Undefined}$



Mathematical Preliminaries

Recap from Math 42


Graphs

Graph Definition

- ❗ A graph is a "mathematical construct" consisting of two sets:
 - A non-empty and finite set of vertices $V = \{v_1, v_2, \dots, v_n\}$
 - A finite set of edges $E = \{e_1, e_2, \dots, e_m\}$
 - Each edge connects two vertices.

Example 31

- $V = \{v_1, v_2, v_3\}$
- $E = \{e_{13}, e_{23}\}$




Directed Graph (aka Digraph)

- If the **direction** of the edges **matters**, then we call the graph "directed graph".
- The edges are shown by "**ordered pairs**" (start , end).
 - In **this course**, we only use **directed graphs**.

Example 32

- Draw a **digraph** with the following specifications:
 - $V = \{v_1, v_2, v_3\}$, $E = \{(v_1, v_3), (v_3, v_1), (v_3, v_2), (v_3, v_3)\}$



Graphs Terminologies




Walk

- A sequence of edges like (v_i, v_j) , (v_j, v_k) , \dots , (v_m, v_n) , is called a **walk** from v_i to v_n .
 - Note that the **end vertex** of e_i is the **start vertex** of e_{i+1} .
 - In other words, in a walk we **cannot jump**!

Example 33

- Each of the following sequences are a **walk** from v_1 to v_3 :
 - Walk 1: (v_1, v_3)
 - Walk 2: (v_1, v_3) , (v_3, v_3)
 - Walk 3: (v_1, v_3) , (v_3, v_1) , (v_1, v_3)
 - ...






Length of Walks

- The "length of a walk" is the total number of edges traversed.


Example 33 (cont'd)

- Walk 1: (v_1, v_3) ; length = 1
- Walk 2: $(v_1, v_3), (v_3, v_3)$; length = 2
- Walk 3: $(v_1, v_3), (v_3, v_1), (v_1, v_3)$; length = 3



Path

- A walk that **no edge is repeated**.




Example 34

- Which one is a **path**?
- Walk from v_1 to v_3 :
 - ✓ – Walk 1: (v_1, v_3)
 - ✓ – Walk 2: $(v_1, v_3), (v_3, v_3)$
 - ✗ – Walk 3: $(v_1, v_3), (v_3, v_1), (v_1, v_3)$

Simple Path

- A path that no vertex is repeated.
 - In other words, no vertex should be visited more than once.




Example 35

- Which one is a simple path?
- Walk from v_1 to v_3 :
 - ✓ – Walk 1: (v_1, v_3)
 - ✗ – Walk 2: $(v_1, v_3), (v_3, v_3)$
 - ✗ – Walk 3: $(v_1, v_3), (v_3, v_1), (v_1, v_3)$

Loop

- An edge from a vertex to itself is called loop.



Example 36


- Which one is a **loop**?
- Walk from v_3 to v_3 :
 - ✓ – Walk 1: (v_3, v_3)
- Is there any other loop in this graph?

Cycle

- A walk from a vertex (called **base**) to itself with no repeated edges.
- Remember that: Walk + No repeated edges = **path**
- **Rewording**: A cycle is a **path** from a vertex (called **base**) to itself.

Example 37


- Which one is a **cycle**?
- Walk from v_1 to v_1 :
 - ✗ – Walk 1: $(v_1, v_3), (v_3, v_1), (v_1, v_3), (v_3, v_1)$
 - ✓ – Walk 2: $(v_1, v_3), (v_3, v_1)$
 - ✓ – Walk 3: $(v_1, v_3), (v_3, v_3), (v_3, v_1)$





Simple Cycle

- A cycle that **no vertex other than the base is repeated**.
 - Note that the walk starts from the base and ends to the base.
 - During the walk, the base should not be repeated.



Example 38

- Which one is a **simple cycle**?
- Walk from v_1 to v_1 :


✗ – Walk 1: $(v_1, v_3), (v_3, v_1), (v_1, v_3), (v_3, v_1)$

✓ – Walk 2: $(v_1, v_3), (v_3, v_1)$

✗ – Walk 3: $(v_1, v_3), (v_3, v_3), (v_3, v_1)$


One-Dimensional Projection of a Walk

- "One-dimensional projection" (or just projection) is another way of representing a walk.



Example 39

- Represent the following walk as a one-dimensional projection.
- Walk from v_1 to v_2 : $(v_1, v_3), (v_3, v_1), (v_1, v_3), (v_3, v_2)$



- The length of this walk (= the number of edges) is clearly shown.

References

1. Linz, Peter, "An Introduction to Formal Languages and Automata, 5th ed.," Jones & Bartlett Learning, LLC, Canada, 2012
2. Kenneth H. Rosen, "Discrete Mathematics and Its Applications, 7th ed.," McGraw Hill, New York, United States, 2012
3. Sipser, Michael, "Introduction to the Theory of Computation, 3rd ed.," CENGAGE Learning, United States, 2013
ISBN-13: 978-1133187790