Mars: The Pear and the Ring

How a fast, expanded Mars set Deimos and Phobos free from a volcano on its pear

Zohar Duran

Independent Researcher, Ramla, Israel · zduran9@gmail.com

ORCID 0009-0005-3194-9387

26 October 2026

Abstract

Phobos and Deimos circle Mars on near-circular orbits of small inclination above its equatorial plane; Phobos lies inside the corotation radius and Deimos outside it. The surface of Deimos is smooth and rounded, while Phobos is scored by systems of long, parallel grooves. Both bodies are light and porous. Although Mars lost most of its water, a large reservoir of liquid water lies today 11.5–20 km beneath its crust.

In this paper we propose a single explanation: Mars, like every planet with a moon, raised a pear, and both moons left from a volcano standing high on it. The young Mars was 1.5 times its present radius, its mean density was 1,166 $\mathrm{kg\,m^{-3}}$, and it turned once in seven to nine hours; a homogeneous body of that density cannot turn faster than once in 5.27 hours. A pear rose from its equator. Its tip lagged behind the spin, and when the pear later settled it lay down westward. On the topography it runs from a base in Valles Marineris to a tip beside Pavonis Mons, 2,996 km on today's surface, and the Tharsis rise along it is two to three times steeper on the west than on the east. The gravity field points to the same place: the long axis of the Martian equator lies at 255°E, on the pear, and the figure of Mars is still offset toward it — a mild pear today. As the pear returned, its interior drained back east along its axis, through Noctis Labyrinthus and Valles Marineris, over a spill point and north through the chaos to Chryse.

The measured length is the hypotenuse of a right triangle, stretched along the raised equator of the young Mars. On the standing pear the vent of Pavonis stood 4,450 km above the equator, where gravity had fallen to about a quarter of its surface value and the outward push of the spin cancelled it: a body there weighed nothing, and moved at the speed of an orbit. Under these conditions an ordinary volcano — a shallow reservoir about 5 km down and an overpressure of 10–50 MPa, the values of Vulcanian explosions on the Earth — set two bodies free. Deimos left first, when the day was 7.3 hours, on a long orbit that crossed the ring. Phobos left second, when Mars had slowed to 8.3 hours and the corotation line had reached the vent, directly onto a circle at the line.

A round ring of condensed steam, which Mars had built from its own volatiles, stood beyond the line in the equatorial plane. It rounded both orbits and flattened their tilt from about 10°, the tilt of the vent relative to the centre of the pear's base. It held Deimos long, ground it to its present size and carried it outward, and the tide of the young Mars, 8–20 times stronger than today's, carried it beyond the final place of the line. The ring could give Phobos nothing, for Phobos already

moved at its speed; its gaseous inner edge, as thin as the Earth's air at about 115 km, exerted on Phobos a weak headwind that balanced the push of the tide, held it at the line and led it outward with the line as Mars slowed. When the ring dried, from the outside in, Phobos stood near 19,000 km and the day was about 22 hours; the line passed it, and it has been falling since — slowly for three billion years, fast in the last few hundred million.

Mars lost about 87% of its spin. It was slowed not by its moons, which carry a millionth of its spin, but by water and gas that left through the corotation line: 6–11% of its mass, a fifth to a third of the water of a young Mars rich in water. The water that never left stayed sealed beneath the crust.

Part I tells this sequence, each stage with the questions a reader will ask and their answers. Part II sets out the evidence: the pit of Pavonis, the two faces of the moons, their orbits, and the water. Part III gives detailed, falsifiable predictions — among them the composition of the samples of Phobos that the Martian Moons eXploration mission (MMX) is to return in 2031.

Keywords: Mars; Phobos; Deimos; Tharsis; Pavonis Mons; rotational fission; circumplanetary ring; corotation radius; tidal evolution; MMX

WHAT THIS PAPER PROPOSES

One pear, one volcano, one ring. A fast, expanded, water-rich Mars raised a pear that now lies westward as the Tharsis rise. From Pavonis, high on the standing pear, where a body weighs nothing, an ordinary volcano set Deimos free on a long orbit and, when Mars had slowed, Phobos on a circle at the corotation line. A round ring of condensed steam rounded both orbits; it ground Deimos and carried it out, and its gaseous inner edge led Phobos outward with the line until it dried. Since then Phobos has been falling.

What the account requires — stated once here, set out in detail in Part O:

- **A fast, expanded young Mars:** hot, rich in volatiles and not yet sorted into layers, 1.5 times today's radius, turning in 7.3–8.3 hours; and the later loss of about 87% of its spin, carried out of the system by 6–11% of its mass in water and gas through the corotation line (§B, E.2, O).
- **A pear raised by the spin of the whole planet**, which stood until the moons had left and put the vent of Pavonis 4,450 km above the young equator (§C, D).
- **An ordinary volcano at Pavonis:** a shallow reservoir about 5 km down, a vent about as wide as the plug, and an overpressure of 10–50 MPa or less (§D.4–D.5).
- **A ring** from the line outward, round and equatorial, whose gaseous inner edge stayed dense enough to hold Phobos until Mars had slowed to a day of about 22 hours (§F).
- **A soft young Mars**, dissipating tidal energy (k_2/Q) 8 to 20 times more than today while Deimos climbed (§G).
- **An explanation of the dark, primitive-looking spectra** of both moons in terms of Martian material altered by water and by long exposure in space.

Contents

Methodological Framework 6

Part I — The Scenario 8

A **The Planetary Puzzle** 8

A.1 Two Moons on Either Side of a Line 8

A.2 Two Distinct Faces 8

A.3 The Martian Water Budget 8

A.4 Conventional Theories and Their Limits 9

B **Young Mars** 10

B.1 The Core Postulate 10

B.2 Initial Physical Properties 10

B.3 Rotation Rate and Moment of Inertia 10

B.4 Rotational Flattening 11

Question-and-answer boxes 1–4

C **The Pear-Shaped Figure** 12

C.1 Rotating Fluid Mechanics 12

C.2 Diagnostic Criteria 13

C.3 Physiography of the Bulge 14

C.4 Rotational Lag and Westward Tilt 15

C.5 Asymmetry of the Topographic Slopes 15

C.6 The Paleotopographic Reconstruction 17

C.7 The Gravitational Signature 19

C.8 Structural Geology 20

C.9 Mars Against the Criteria 20

Question-and-answer boxes 5–7

D **Volcanic Dynamics at the Apex** 21

D.1 Gravity and Sideways Acceleration along the Pear 21

D.2 The Zero-Net-Weight Condition 22

D.3 Why Pavonis 22

D.4 The Magma Chamber Geometry 24

D.5 Ejection Mechanics 24

Question-and-answer boxes 8–12

E **The Ejection Sequence** 25

E.1 The First Release – Deimos 25

E.2 Rotational Deceleration 27

E.3 The Second Release – Phobos 28

Question-and-answer boxes 13–17

F **The Planetary Ring Dynamics** 29
F.1 Ring Formation and Orbital Damping 29
F.2 The Split Interaction 29
F.3 The Velocity Limit 30
F.4 The Gas Leash 30
F.5 The Ring Dissipation 30
F.6 Dating the Dissipation 31
Question-and-answer boxes 18–21
G **Tidal Mechanics** 32
Question-and-answer boxes 22–24
H **The Post-Ejection Drainage** 34
H.1 Relaxation and the Tharsis Bulge 34
H.2 The Neck 34
H.3 The Axial Conduit 35
H.4 The Basin and the Spill 36
H.5 The Dust Provenance 38
H.6 The Residual Volatiles 39
Part II — The Evidence 40
I **The Summit Caldera** 40
I.1 The Caldera Geometry 40
I.2 The Volume Balance 40
I.3 The Caldera Collapse 41
J **The Deimos Record** 41
J.1 The Damped Satellite 41
J.2 The Mass Ablation Profile 42
J.3 The Volatile-Rich Composition 43
K **The Phobos Record** 43
K.1 The Striation Evidence 43
K.2 The Post-Ejection Relaxation 44
K.3 The Roller-Bearing Pits 45
K.4 The Leading Apex 46
K.5 The Intracrater Deformation 46
K.6 Mass Asymmetry and Centre-of-Mass Offset 47
K.7 The Paleo-Orbit Reconstruction 48
K.8 The 3.2 Commensurability 49
L **The Ring Signatures** 50
M **The Hydrological Record** 50

Part III — Empirical Predictions 52

N **The Predictive Matrix** 52

O **System Constraints and Boundary Conditions** 55

Appendix A — Computational Reproducibility 56

Appendix B — Observational Protocols 57

References 59

Methodological Framework

Every number belongs to the young Mars at the time of the two releases unless the row says “today”. The young Mars is treated as homogeneous, as an unsorted, volatile-rich planet would be.

Quantity	Value	Role
The planet		
Mass of Mars	6.4171×10^{23} kg; $GM = 42{,}828$ km^3 s^{-2}	Fixed throughout
Mean radius	young 5,084 km ($1.5\times$); today 3,389.5 km	Expanded, hot and volatile-rich (§B.2)
Mean density	young 1,166 kg m^{-3}; today 3,933	Follows from mass and radius
Day	7.3 h at Deimos’s release; 8.3 h at Phobos’s; today 24.62 h	Mars slows between the two releases (§E.2)
Fastest possible day of a homogeneous young Mars	5.27 h (axisymmetric limit); 5.77 h (onset of a triaxial figure)	Why the day is 7.3–8.3 h and not 5 h (§B.3)
Flattening	0.24 at 7.3 h; 0.18 at 8.3 h	Less spin, less flattening (§B.4)
Equatorial radius	5,561 km at 7.3 h; 5,431 km at 8.3 h	The pear starts from the raised equator; east–west lengths 1.60–1.64 times today’s
Gravity at the mean surface	1.66 m s^{-2} (today 3.73)	
Corotation line	9,083 km at 7.3 h; 9,870 km at 8.3 h; today 20,428 km	Turning with Mars is an orbit there
The pear		
Where it lies today	base 294.7°E 12.5°S, in Valles Marineris; tip 243.9°E 5.4°S	Read on the MOLA topography (§C.3)
Length lying down	2,996 km on today’s Mars; 4,800–4,920 km on the young Mars (8.3–7.3 h)	From the eastern point of the base to the tip; stretched with the equator (§C.6)
Half-width of the base, b	1,340 km today; 1,280–1,450 km on the young Mars	Measured on MOLA, north–south; centre of the base near 272°E, 10°S (§C.6)
Height of the standing pear, $\sqrt{L^2 - b^2}$	2,680 km with today’s measurements; 4,580 km (8.3 h) to 4,740 km (7.3 h) on the young Mars	The length is the hypotenuse, not the height (§C.6)

Quantity	Value	Role
Pavonis on the standing pear	4,450–4,610 km above the equator, 48.9° along the pear and 7.4° off its axis; vent 9,880–10,170 km from the centre	The exit point (§D)
Gravity at the vent	0.41–0.44 $\mathrm{m\,s^{-2}}$, 25–27% of the surface value	Inverse square of the distance
Net weight at the vent	-0.17 (7.3 h) to 0.00 (8.3 h) $\mathrm{m\,s^{-2}}$	Gravity minus the outward effect of the spin: a body there weighs nothing
Sideways speed at the vent	2.43 $\mathrm{km\,s^{-1}}$ (7.3 h); 2.08 $\mathrm{km\,s^{-1}}$ (8.3 h)	Against an orbital speed there of 2.05–2.08 $\mathrm{km\,s^{-1}}$
The volcano and the bodies		
Overpressure at the vent	10–50 MPa	Ordinary volcanic values; no deep chamber, no long conduit (§D.4–D.5)
Exit speed it gives	110–250 $\mathrm{m\,s^{-1}}$	Pushed along the plug's own length
Phobos at release	$1.02\times$ today: 22.9 km across, 1.13×10^{16} kg	Left second (§E.3)
Deimos at release	a little smaller than Phobos then: about 20.7 km, 6.8×10^{15} kg	Left first; ground to 12.4 km (§E.1, F)
The ring and the tide		
Ring	From the corotation line outward, in the equatorial plane; clumps beyond 13,000 km (Roche limit for ice)	Built from steam before the releases (§F)
Tidal friction	$k_2/Q = 1.816 \times 10^{-3}$ today; 8–20 times more in the young Mars	Carries the moons outward beyond the line (§G)
Phobos today	$a = 9{,}375$ km, $e = 0.0151$, $i = 1.09°$	Inside the line, falling
Deimos today	$a = 23{,}463$ km, $e = 0.0003$, $i \approx 1.8°$	Outside the line, receding

Table 1. The working environment. Shading marks the two moments of release.

PART I

The Scenario

A The Planetary Puzzle

A.1 Two Moons on Either Side of a Line

Phobos and Deimos are small, dark and light. Phobos is about 22 km across on average, Deimos about 12. Their bulk densities — 1,853–1,887 $\mathrm{kg\,m^{-3}}$ for Phobos and 1,471 $\mathrm{kg\,m^{-3}}$ for Deimos — mean that a third to a half of each body is empty space (Pätzold et al. 2025; NASA/JPL Solar System Dynamics). Both circle Mars almost exactly above its equator, on almost circular orbits. Phobos, at 9,375 km from the centre of Mars, goes round in 7.65 hours, three times in each Martian day; Deimos, at 23,463 km, takes 30.3 hours.

Between them runs the corotation radius, today 20,428 km: the distance at which a body in free orbit keeps pace with the rotation of the planet itself (the distance at which a body turning with the planet is already moving at orbital speed). The tide acts in opposite directions on the two sides of this line. Inside it a moon outruns the rotation of the planet; the tidal bulge trails behind it and holds it back, and Phobos spirals down. Outside it a moon lags behind the rotation; the tidal bulge runs ahead of it and draws it forward, and Deimos slowly recedes. Any account of the origin of the moons must therefore put Phobos inside the line and Deimos outside it.

A.2 Two Distinct Faces

The surfaces of the two moons are entirely different. Deimos is smooth and rounded: its craters are shallow and worn, and it has no grooves. Phobos is rough: extensive systems of long parallel grooves run across it, and chains of pits run beside them. At its blunt end, toward the leading side of its orbit, opens the depression Stickney, 9 km across (Thomas et al. 1979; Basilevsky et al. 2014). An account of the origin of the moons should say why two bodies of the same planet carry such different surfaces.

A.3 The Martian Water Budget

The surface of Mars records rivers and lakes in its past. Its atmosphere today is enriched in heavy hydrogen (deuterium) about six to seven times relative to the Earth's oceans — the signature of a long loss of water to space; the early planet held at least the equivalent of a global layer 137 m deep (Villanueva et al. 2015).

On the other hand, the seismometer of the InSight lander found that the mid-crust beneath it, between 11.5 and 20 km depth, is best explained by material whose pores are full of liquid water — enough, if the same holds across the planet, for a global layer 1–2 km deep (Wright, Manga & Panning 2024). Mars lost much of its water and kept much of it. An account of the moons that involves water should say where each part went.

A.4 Conventional Theories and Their Limits

Three possible origins of the moons are discussed in the literature today.

Capture of two small asteroids fits the dark colour and the spectrum of the moons, but not their orbits: a captured body arrives on an inclined, eccentric orbit, and no mechanism has been shown to turn two such orbits into equatorial circles.

A giant impact of an external body on Mars throws up a disc of debris from which the moons accrete. Such a disc settles into the plane of the equator and explains the orbits well, and this is the leading account today (Rosenblatt et al. 2016; Hyodo et al. 2017; Canup & Salmon 2018).

Ring–moon cycles also begin with the impact of an external body: its debris forms a ring around Mars, and a moon gathers at the ring's outer edge. The moon, inside the corotation line, is drawn down by the tide until it breaks up at the Roche limit and becomes a ring again, from which a smaller moon forms; the cycle may have repeated three to seven times, each moon about five times smaller than the one before. Phobos is the latest of these moons, and Deimos a survivor of the first disc (Hesselbrock & Minton 2017).

In all three, the different faces of the two moons are explained separately from their origin: by tides, by ejecta from Stickney or by later impacts (Hurford et al. 2016; Ramsley & Head 2019).

The isotopes set the accounts apart:

Capture. A captured asteroid carries the isotopic fingerprint of its own parent body, not that of Mars.

A giant impact of an external body. In impact simulations the disc from which the moons accrete holds a large share of the external body — about 30% of its outer part and up to 65% of the whole (Hyodo et al. 2017) — and its material was melted and partly vaporised, which shifts the isotopes of moderately volatile elements such as zinc and potassium.

Ring–moon cycles. They begin from the disc of an external impact and inherit its mixture.

For the accepted accounts a moon with the isotopes of Mars alone is a difficulty. The account proposed here commits to it: the oxygen, chromium and titanium isotopes of both moons are those of Mars, with no foreign component and no shift by evaporation (N.1). The samples of Phobos that MMX returns in 2031 will decide.

TODAY AND IN THIS PAPER

Today: the moons accreted from debris thrown up by a foreign body, and carry part of it in their isotopes; their different faces have separate, later causes. **Here:** Mars, like every planet with a moon, raised a pear. The moons are two bodies of Martian crust set free, one after the other, from a volcano standing high on it, where the spin of Mars alone gave them their orbits; one ring that Mars had built from its own steam rounded those orbits and gave the two moons their different faces.

B Young Mars

B.1 The Core Postulate

A moon is made of its parent. On the Earth, early on, the spin raised a pear from the equator; its tip crossed the corotation line and became the Moon, and the rest of the pear returned. We take this as a rule: every planet with a moon had a pear in its past. The young Mars falls under this rule too. What differs on Mars is what left the pear: not its tip itself, but two bodies set free gently from a volcano standing high on it.

B.2 Initial Physical Properties

The young Mars was hot, rich in water and gas, and not yet sorted into layers. Its mean radius is taken as 1.5 times today's, 5,084 km, for a mean density of 1,166 $\mathrm{kg\,m^{-3}}$ — the density of the material it later shed and of its two moons. It contracted later, as it lost its volatiles and cooled.

B.3 Rotation Rate and Moment of Inertia

The model. Young Mars turned with a day of seven to nine hours: 7.3 hours when Deimos left and 8.3 hours when Phobos left. Its brake was the material that left through the corotation line (§E.2); this brought it to the 24.6 hours of today.

Support from planet-building studies. Simulations of planet building give young terrestrial planets fast spins: a few large collisions in the last stages of growth set most of the spin, and the typical result is a day of a few hours (Kokubo & Genda 2010). Built from small bodies alone, Mars would turn in about 250 hours, so its spin was set by collisions (Miguel & Brunini 2010).

Support from the theory of rotating bodies. How fast could the young Mars turn? By the classical theory of rotating fluid bodies (Chandrasekhar 1969), a homogeneous body of density 1,166 $\mathrm{kg\,m^{-3}}$ stays an axisymmetric spheroid (a Maclaurin spheroid) only if its day is longer than 5.27 hours, and already below 5.77 hours it prefers an elongated, triaxial figure (a Jacobi ellipsoid). A day of seven to nine hours lies safely within the stable range.

B.4 Rotational Flattening

A spinning homogeneous body flattens (a Maclaurin spheroid), the more so the faster it turns. The slower spin keeps the flattening moderate:

Day	Flattening	Equatorial radius	Polar radius	Line above the equator
7.0 h	0.26	5,618 km	4,164 km	3,214 km
7.3 h	0.24	5,561 km	4,250 km	3,522 km
8.0 h	0.19	5,461 km	4,407 km	4,193 km
8.3 h	0.18	5,431 km	4,455 km	4,439 km
9.0 h	0.15	5,368 km	4,562 km	5,075 km

Table 2. The homogeneous young Mars at five rates of spin; 7.3 and 8.3 hours are the days of the two releases (Deimos and Phobos). The colour fades as the spin and the flattening fall.

Because of the flattening, the equator stands 350–500 km above the mean radius, so the pear starts from raised ground.

QUESTION AND ANSWER 1

Why should every planet with a moon have had a pear?

A moon made of its parent must reach the corotation line, where turning with the planet is already an orbit. On the young Mars the line stood 3,200–5,100 km above the equator (Table 2); no volcano on the ground comes near it. Only a pear carries a vent there. On the Earth the pear's tip became the Moon; on Mars the moons are small and left from a volcano high on the pear, but the pear is what put the volcano at the line.

QUESTION AND ANSWER 2

Why a radius 1.5 times today's?

It is the author's estimate of a hot, expanded planet rich in water and gas. Its mean density, 1,166 kg m^{-3}, lies below those of its moons (1,471–1,860 kg m^{-3}), which are samples of its upper layers. A smaller young Mars would need less mass to leave through the line to slow down (§E.2), so the larger radius is the demanding case.

QUESTION AND ANSWER 3

Is the day of seven to nine hours not simply a choice that suits the model?

No. The early day of Mars has never been measured. Simulations of planet building give days of a few hours (Kokubo & Genda 2010), and nothing known slows Mars from there to 24.6 hours (§B.3). The range is then fixed by the pear itself: the vent of Pavonis stands at the corotation line at 8.3 hours (Table 8), and a moon set free there on a shorter day goes into orbit outside the line (Table 11). The day is read from the topography, not assumed. A day of five hours is ruled out: a homogeneous young Mars cannot turn faster than once in 5.27 hours and keep its axisymmetric figure.

QUESTION AND ANSWER 4

Would so fast a spin not deform Mars beyond recognition?

The flattening is 0.18–0.24, a moderate spheroid whose equator is raised by only 350–500 km above the mean radius (Table 2). It relaxed as Mars slowed and contracted, to today's 1/191.

C The Pear-Shaped Figure

C.1 Rotating Fluid Mechanics

A fluid body that does not spin tends to a sphere; when it spins, it flattens. Maclaurin (1742) showed that a flattened spheroid is a figure of equilibrium for a homogeneous rotating fluid, and that the flattening grows as the spin grows faster. Jacobi (1834) extended this and showed that beyond a certain spin there is a further figure of equilibrium: an ellipsoid with three unequal axes, elongated in one direction in the plane of the equator (Chandrasekhar 1969; Iurato 2014).

Poincaré (1885) found that a new branch leaves the sequence of Jacobi's ellipsoids: the pear-shaped figures, with one end narrow and drawn out and the other broad and rounded. George Darwin saw in this figure the way one body divides into two, and on it he based his explanation of the Moon: it left the Earth through a pear stage (Darwin 1902).




Figure 1. The figures of a rotating fluid body, seen from above its pole. As the spin grows the sphere flattens, elongates and grows a tip (a pear); as the interior settles it returns gradually toward a sphere.

The argument over the stability of the pear lasted about three decades:

Year	Who	What
1742	Maclaurin	The flattened spheroid is a figure of equilibrium
1834	Jacobi	Beyond a certain spin, an elongated ellipsoid
1885	Poincaré	The pear branches off the Jacobi sequence
1902	Darwin	The pear is stable; fission explains the Moon
1905	Lyapunov	The pear is not stable
1917	Jeans	The pear is not stable
1924	Cartan	At the branch point the Jacobi ellipsoid itself loses its stability

Table 3. The classical argument over the stability of the pear (Darwin 1902; Lyapunov 1905; Jeans 1917; Cartan 1924). The colour deepens as the instability is established.

"Not stable" means that the body cannot persist in that shape over time: any small disturbance grows and carries the body on, to division or to a return to a rounder shape. The accepted conclusion is that a pear can form in a rotating fluid body, but it is a passing stage, not a lasting state.

Planetary science knows bodies whose shape today does not fit their spin today, because the shape froze when their crust stiffened while they turned differently:

- **Iapetus**, a moon of Saturn, turns once in 79 days today, but its shape fits a spin of about 16 hours: the shape froze as its crust stiffened while the spin slowed (Castillo-Rogez et al. 2007).
- **The Moon** carries a fossil bulge, the record of an earlier orbit and rate of spin (Garrick-Bethell et al. 2006).
- **Haumea**, a dwarf planet, turns once in 3.9 hours and is strongly elongated: a living example of a body shaped by its spin (Ortiz et al. 2017).

Simulations of planet building give the planets a fast spin in their first stages, a day of a few hours (§B.3). Their interiors were not symmetric: a flow concentrated in one hemisphere could develop in the mantle of the young Mars (Zhong & Zuber 2001), and to this day the centre of mass of Mars does not coincide with its centre of figure but is displaced from it by about 3 km (Zuber et al. 1998).

Sorting began very early: on Mars metal separated from silicate within a few million years (Dauphas & Pourmand 2011). But that is one stage of many. Sorting is a long, graded process that continues today — the Earth's inner core still grows (Labrosse et al. 2001) — fast at first and slower with time.

> THE PRINCIPLE
>
> The spin of the whole planet raises a pear, and the core takes part and moves toward it; the pressure passes through the pear and supports it. As sorting advances, the place of the core is fixed and the pear returns. **The pear always comes before the planet settles**: it was common at the beginning and is rare today. Its instability is the reason it was a passing stage and not a state.

C.2 Diagnostic Criteria

The criteria below set out the marks a pear should leave after it has returned and lain down. They are reasonable, not rigid: a pear that has returned and lain down need not meet each of them in full. Each is tied to what supports it.

	Criterion	Why it marks a pear	Rests on
P1	On the equator	The outward pull of the spin is greatest at the equator; every elongated figure of a spinning body grows in the plane of the equator	Classical theory (Chandrasekhar 1969)

	Criterion	Why it marks a pear	Rests on
P2	One long axis	A pear has one tip; its remnant is elongated in one direction, not spread to all sides	Poincaré's figure
P3	Base and tip	A broad base narrowing to a tip, the highest ground at the tip	Poincaré's figure
P4	Lagging against the spin	Material raised from the ground keeps its angular momentum and falls behind the spin; the tip lies behind the base, on the side opposite to the spin, and the trailing side is steeper. On a planet spinning eastward, as Mars does, the lag is to the west; on one spinning westward it is reversed	Angular momentum in a spinning body
P5	A shape out of step with today's spin	A remnant of a pear is part of the shape that froze and is not explained by the present spin	Castillo-Rogez et al. 2007; Garrick-Bethell et al. 2006
P6	A mass excess	A pear that returned but did not vanish leaves stronger gravity above it	Geodesy
P7	Fractures around it	Rising and sinking stretch the crust around the body; the fractures organise around it, mostly radial	Banerdt et al. 1992
P8	Early	The pear belongs to the fluid, fast period; its remnant is among the oldest structures of the planet	§C.1
P9	A link to the moons	This model holds that if a planet has moons, they left from its pear, for only the pear brings material to the corotation line. A pear that has gone should be tied to the place the moons left from	This model, after Darwin
P10	Return and sorting	As the pear returns, its material is taken back into the planet, in one way or another, and sorted; a planet that froze may leave masses of the pear and traces of the return on its surface	This model

Table 4. Ten criteria for a pear that has gone. Mars is tested against them in §C.9.

C.3 Physiography of the Bulge

On the topography measured by the laser altimeter MOLA (Smith et al. 2001), expanded here in spherical harmonics from the shape model MarsTopo719 (Wieczorek 2015), the pear lies along the equator, just south of it. It runs from its base at 294.7°E,

12.5°S, in Valles Marineris, to its tip at 243.9°E, 5.4°S. Its length is 2,996 km on today's surface, and 4,800–4,920 km on the young, expanded Mars (§C.6; Figure 2).

Beyond the tip, to the west, stand the Tharsis volcanoes. Arsia, just beyond the tip, lies on the continuation of the axis of the pear, 4.1° off it, and its edifice is elongated: above the 9-km contour of MOLA it measures 730 by 415 km (ratio 1.75), its long axis running north-north-east — a body at the tip, drawn out as the pear was flattened and laid down.




Figure 2. Tharsis on the MOLA shape model of Mars (MarsTopo719, to degree 180; contours at 0, 4 and 8 km). Red: the pear as it lies today, from its base in Valles Marineris to its tip. Pavonis stands near the tip, Arsia just beyond it. Triangles: the volcanoes. Mars turns from west to east.

C.4 Rotational Lag and Westward Tilt

While the pear rose, its elongated tip had to be carried faster and faster round the centre of the planet; because angular momentum is conserved, it lagged behind the spin and leaned westward, as the Earth's pear did. The spin of the whole planet had raised it, the core taking part and moving toward it, and the pressure passed through it and supported it. When the moons had left and the vents were open, and as sorting advanced and the core settled, the pear did not fall: it lay down gradually, over tens to hundreds of millions of years, and the lag went on acting on it all that time. It lay down the way it leaned — westward, against the spin.

C.5 Asymmetry of the Topographic Slopes

A body that lay down westward should end in a steep western front, facing a long, gentle slope on the east. Measured from the crest of Tharsis to where the ground falls to zero, along bands of latitude (topography smoothed over about 9°):

Band	Crest	West: distance to 0 km	West slope	East: distance to 0 km	East slope
15–5°S	240°E	1,570 km	7.0 m/km	2,740 km	4.1 m/km
10°S–10°N	244°E	1,360 km	5.8 m/km	3,900 km	2.0 m/km
5°S–5°N	246°E	1,480 km	5.7 m/km	3,790 km	2.2 m/km
0–10°N	252°E	1,710 km	4.2 m/km	3,180 km	2.3 m/km
5–15°N	256°E	1,980 km	4.0 m/km	1,460 km	5.4 m/km

Table 5. The two flanks of Tharsis. Red: the steep western front; blue: the gentle eastern slope. North of 5°N the east falls steeply into Chryse and Kasei Valles.

Along the equator and south of it, the western front is two to three times steeper than the eastern slope (Figure 3), as expected of a body that lay down westward.




Figure 3. Mean elevation in the band 10°S--10°N. The ground falls to zero within 1,360 km to the west of the crest and along a gentle slope of 3,900 km to the east.

QUESTION AND ANSWER 5

How do you know that Tharsis is the pear?

Three findings in the topography point to it. First, it lies on the equator. Second, its crest is the highest point of the large-scale shape of Mars: the long axis of that shape passes through 252°E, the longitude of Tharsis. Third, its two flanks differ along the direction of the spin, steep on the west and gentle on the east (Table 5). This is what a body that leaned back against the spin and lay down westward leaves.

QUESTION AND ANSWER 6

Why does the tip of the pear lie west of its base, and not over it?

Had the pear collapsed quickly, its tip would have come down over the centre of its base. It did not collapse quickly. It lay down slowly, over tens to hundreds of millions of years, and throughout that time the lag acted on it: the tip, far from the axis, trailed the spin of Mars, as the tip of the pear of the Earth did. The slow lying down let the lag work to the end, and the tip came to rest to the west (§C.4). Nor did the spin push it east on the way down: settling 1,000 km in 10 million years gives a sideways (Coriolis) acceleration of about 2×10^{-12} m s^{-2}, which the surrounding material removes entirely.

QUESTION AND ANSWER 7

Why is the crest south of the equator, and why does the band north of 5°N not follow the pattern?

The pear lies along 5–12.5°S, from its base at 12.5°S to its tip at 5.4°S, and its crest follows it. North of 5°N the eastern side is cut by Chryse Planitia and Kasei Valles, which were carved later; the pattern holds fully in the bands that cross the pear.

C.6 The Paleotopographic Reconstruction

The length of 2,996 km is measured on the lying pear, from the eastern point of its base to its tip. When the pear stood, this line was not its height: it runs from the edge of the base to a tip that stands above the centre of the base. It is the hypotenuse of a right triangle whose other two sides are the height h and the half-width of the base b:

$$h = \sqrt{L^2 - b^2}.$$

We measured b on the MOLA topography, across the axis of the pear at its base, smoothed as in Table 5. To the north the rise falls to 0 km 1,280–1,400 km from the axis, at the edge of Kasei Valles and Chryse; to the south no edge is reached, for the axis runs along Valles Marineris and the southern highlands continue beyond it. Taking the northern edge as the half-width of a round base, $b = 1{,}340$ km today; the centre of the base lies near 272°E, 10°S.

With today's measurements the pear would stand

$$h = \sqrt{2{,}996^2 - 1{,}340^2} = 2{,}680 \text{ km}.$$

On the young Mars the measurements were different. Mars was 1.5 times larger and, of low density and turning fast, flattened into a spheroid (Table 2). Along its raised equator, where the pear lies, lengths scale with the equatorial radius a: 1.60 times today's at 8.3 hours, 1.64 times at 7.3 hours. Across the equator, north–south, the flattening compresses them, in proportion to c^2/a. The hypotenuse therefore grows more than the half-width of the base:

	Today's measurements	Young Mars, 8.3 h	Young Mars, 7.3 h
Length lying, L	2,996 km	4,800 km	4,920 km
Half-width of the base, b	1,340 km	1,450 km	1,280 km
Height standing, h	2,680 km	4,580 km	4,740 km

When Phobos left, the pear stood 4,580 km tall (Figure 4). Every point of the pear is carried in the same way: a point at a distance s from the eastern point of the base, along the lying pear, stands $\sqrt{s^2 - b^2}$ above the base when the pear stands. On the lying pear the highest ground is at the tip — Pavonis and Arsia — and the base lies low, in Valles Marineris.




Figure 4. The pear lying and standing. Top: MOLA, the highest point within 400 km of the axis of the pear, from the eastern point of its base to beyond its tip (height exaggerated). The volcanoes at the tip are the highest ground; the base lies low, in Valles Marineris. Bottom: the pear standing, to scale, with today's measurements and on the young Mars at 8.3 hours. The measured length is the hypotenuse; the height is $\sqrt{L^2 - b^2}$.

C.7 The Gravitational Signature

We expanded the GMM-3 gravity field of Mars (Genova et al. 2016) to degree 90.

The long axis of the equator. The equator of Mars is not round. In the broadest part of the field, degree 2, its long axis points to **255°E**: on the pear, between its tip (243.9°E) and the centre of its base (272°E), and close to the long axis of the shape of Mars at 252°E (Question and Answer 5). A body that spins symmetrically about its axis has a round equator; this part of the field is therefore not explained by the present spin. It is a frozen shape, as on Iapetus.

The broad high. Keeping only the broadest parts of the field, degrees 2 and 3 — the large structure, not single mountains — the geoid is highest at **6.5°N, 253.5°E**, between Pavonis and Ascraeus, at the tip of the pear (Figure 5).

Olympus left out. Olympus Mons stands off the pear, a single edifice whose load is its own. With it left out of the full field (within 8°, 10° or 12° of its centre; the result is the same in all three), the highest geoid and the strongest free-air gravity both lie at 11.5°N, 255.5°E, on Ascraeus, the northernmost of the three Tharsis Montes.

Mars is still a mild pear. The first degree of the MOLA shape model, taken about the centre of mass, gives the offset of the centre of figure from the centre of mass. Its north–south part, 3.0 km, is the known offset of the hemispheres (Zuber et al. 1998). Its part in the plane of the equator is 1.37 km, toward **262.7°E** — on the axis of the pear, between its tip (243.9°E) and the centre of its base (272°E). The figure of Mars still bulges toward the pear: a very mild pear, in exactly the direction of the pear of the model. We propose that the core took part and moved toward the pear while it stood; its present position is not measured, and is a prediction for a future seismic network on Mars (Appendix B, B16).

From the base to the tip. The free-air gravity rises steadily along the pear, from the eastern base to the tip:

Place	Free-air gravity	
Eastern point of the base (294.7°E, 12.5°S)	−565 mGal	the canyon (§H.3)
Centre of the base (272°E, 10°S)	+49 mGal	
Tip (243.9°E, 5.4°S)	+393 mGal	
Pavonis	+1,474 mGal	
Arsia	+1,661 mGal	
Ascraeus	+2,307 mGal	

Table 6. Free-air gravity at the surface, GMM-3 degrees 2--90 without the flattening term C_{20}. The colour deepens from the base to the tip.




Figure 5. The geoid of degrees 2--3 (contours, m) on the MOLA topography. Red: the pear. Dashed: the long axis of the equator, 255°E. Star: the broad geoid high. Olympus, off the pear, is left out of the calculation.

C.8 Structural Geology

Anderson et al. (2001) mapped 24,452 grabens and ridges around Tharsis and divided them into five stages, from the Noachian to the Amazonian. In every stage the centre of the radial fractures lies on the high ground of Tharsis: the large early centres in Claritas and Syria, between the base and the tip, and the last stage centred near Pavonis, at the tip of the pear. Most of the mass of Tharsis was in place by the end of the Noachian, in the first hundreds of millions of years of Mars (Phillips et al. 2001), and the tectonic activity around it was greatest at the start and declined with time (Anderson et al. 2001).

C.9 Mars Against the Criteria

These are the findings on Mars against the ten criteria of Table 4:

	Criterion	What Mars shows	
P1	On the equator	The pear lies along 5–12.5°S; the broad geoid high at 6.5°N (§C.3, C.7)	Met
P2	One long axis	2,996 km long against a half-width of 1,340 km at the base (§C.6)	Met
P3	Base and tip	The base low in Valles Marineris; the highest ground at the tip, the Tharsis Montes (§C.3, C.6)	Met

	Criterion	What Mars shows	
P4	Lagging against the spin	Mars spins eastward; the tip west of the base; the western flank two to three times steeper (§C.5)	Met
P5	Frozen shape	The long axis of the equator at 255°E, on the pear, not explained by today's spin; the figure still offset 1.37 km toward 262.7°E: a mild pear even today (§C.7)	Met
P6	Mass excess	Free-air gravity rising from $+49$ to $+1{,}474$–$2{,}307$ mGal from the base to the tip; the geoid high at the tip (§C.7)	Met
P7	Fractures around it	24,452 structures; every stage centred on high Tharsis, the last near Pavonis (§C.8)	Met
P8	Early	Most of Tharsis in place by the end of the Noachian (§C.8)	Met
P9	Link to the moons	Pavonis at the tip; both moons from it (§D, E)	Rests on the model
P10	Return and sorting	Part of the material remains at the surface: Tharsis and its mass excess (P6). On a water-rich Mars, part flowed back in a fluid channel along the axis, unique to Mars: Noctis → Valles Marineris → basin → Eos → chaos → Chryse (§H.2–H.4)	Met

Table 7. The pear of Mars against the ten criteria of Table 4. Green: met. Orange: rests on the model.

Nine of the ten criteria are met by observation; the tenth rests on the model. Tharsis has every mark that a pear which rose, lagged, lay down westward and drained back along its axis would leave.

D Volcanic Dynamics at the Apex

D.1 Gravity and Sideways Acceleration along the Pear

Stood up again, the pear carries each volcano to a height $\sqrt{s^2 - b^2}$ above the base, where s is its distance from the eastern point of the base (§C.6). At these heights gravity falls with the square of the distance from the centre of Mars, while the spin pushes outward in proportion to it. For each volcano there is one day at which the corotation line passes exactly through its vent; on any shorter day the vent stands beyond the line.

Volcano	On the pear	Height above the equator	Distance from the centre	Gravity there	Line through the vent at
Ascraeus	42.1° along, 19.2° off	4,120 km	9,590 km	0.47 m s^{-2} (28%)	7.9 h
Pavonis	48.9° along, 7.4° off	4,450 km	9,880 km	0.44 m s^{-2} (27%)	8.3 h
Arsia	55.0° along (beyond the tip at 50.6°), 4.1° off	4,950 km	10,330 km	0.40 m s^{-2} (24%)	8.9 h

Table 8. The three Tharsis Montes on the standing pear of the young Mars, flattening included and lengths stretched as in §C.6, each at the day on which the line passes through its vent. Positions are measured along the pear from its base and across its axis; percentages are of the gravity at the mean surface. Green: Pavonis.

D.2 The Zero-Net-Weight Condition

At the vent of Pavonis gravity is 0.41–0.44 m s^{-2}, about a quarter of its value at the surface, and the spin almost cancels it: the net weight is -0.17 m s^{-2} at a day of 7.3 hours (slightly outward) and zero at 8.3 hours. A body set free there needs no push to rise. It carries the sideways speed of the vent itself, 2.43 km s^{-1} at 7.3 hours and 2.08 km s^{-1} at 8.3 hours, against a Keplerian orbital speed there of 2.05–2.08 km s^{-1}: the spin alone gives it its orbit.

D.3 Why Pavonis

Pavonis stood high on the pear, near its tip, and beneath it lay the reservoir that fired: its summit pit is the deepest of the four Tharsis calderas for its width, a single circular pit about 5 km deep, the roof of a reservoir that emptied and collapsed (§D.4). Both moons left from Pavonis, one after the other. While the pear stood, every vent on it lay above the centre of its base, at about 10°S; both moons therefore began on orbits tilted by about 10°, and the ring rounded the orbits and brought the tilt down to 1.09° (Phobos) and about 1.8° (Deimos) (§F.1).

QUESTION AND ANSWER 8

Could a pear more than 4,000 km tall stand at all?

The load is least where the pear is tallest: at its top gravity is about a quarter of its surface value, and the spin cancels the rest (Table 1). The spin of the whole planet raised it, the core taking part and moving toward it, and the pressure that passed through it supported it (§C.1); how a pear is raised and held is set out in the papers on the Earth’s pear. Here it is listed as a requirement (§O).

QUESTION AND ANSWER 9

Why Pavonis, and not Arsia, Ascraeus or Olympus?

We tested each of them: a body set free at each vent, on the standing pear, at the two days of the releases:

Volcano	Line through vent	At 7.3 h (Deimos)	At 8.3 h (Phobos)
Ascraeus	7.9 h	orbit, $e = 0.25$	falls back: vent inside the line
Pavonis	8.3 h	orbit, $e = 0.41$	a circle at the line, $e = 0.002$
Arsia	8.9 h	long orbit, to 52,400 km, $e = 0.66$	orbit, $e = 0.19$
Olympus	—	no orbit: off the pear	no orbit

Ascraeus could have set Deimos free at 7.3 hours, on an orbit of eccentricity 0.25. But at 8.3 hours its vent already lies inside the line, and whatever leaves it falls back: it could not have set Phobos free.

Arsia could have set free both: Deimos on a very long orbit ($e = 0.66$, to 52,400 km), and Phobos on an elongated orbit ($e = 0.19$) that the ring would have had to round.

Olympus does not stand on the pear: its summit was about 5,600 km from the centre of the young Mars, against a line at 9,080 km, and no shot from it reached an orbit of its own. It may well have erupted strongly; its material reached the inner edge of the ring, came back, and is part of the dust of Mars (§H.5).

Pavonis alone gives Phobos, at 8.3 hours, a circle at the line ($e = 0.002$) with no push, and it alone carries the mark of a reservoir that fired and emptied: a single deep circular pit, about 5 km deep, of the scale of the two moons together; the caldera of Arsia is broad and shallow. The tilt of the orbits does not choose between them: on the standing pear every vent lay above the centre of the base, at about 10°S, and the ring flattened the tilt (§F.1). The decisive test is gravity: a mass deficit beneath Pavonis larger than beneath Arsia (Appendix B, B10).

QUESTION AND ANSWER 10

The pear is measured on today's surface. How can its length on the young Mars be known?

Its length along today's surface is 2,996 km. The young Mars was 1.5 times larger and flattened by its spin. The pear lies along its equator, which stood 5,430–5,560 km from the centre, so east–west lengths were 1.60–1.64 times today's and the pear was 4,800–4,920 km long; north–south lengths near the equator were compressed by the flattening. Pavonis, near the tip, lay 4,670 km from the eastern point of the base; with the pear standing it stood 4,450 km above the base (§C.6).




Figure 6. Pavonis Mons, the middle of the three Tharsis Montes, between Ascraeus Mons to the north-east and Arsia Mons to the south-west, at 1.48°N, 247.04°E, with its summit caldera 47 km wide. Credit: ACT-REACT QuickMap Mars; MOLA colourised elevation (NASA/GSFC); IAU nomenclature.

D.4 The Magma Chamber Geometry

The model needs no deep magma chamber and no long feeding conduit. Beneath the summit of Pavonis lay a shallow reservoir, about 5 km down: the summit pit of Pavonis, 47 km wide, is about 5 km deep (Crumpler et al. 1996; NASA/JPL PIA25157), and a pit of that depth is the roof of a reservoir that emptied and collapsed.

Before the moons left, the vent above the reservoir was much narrower than the pit is today — about as wide as the plug it held, some 20–23 km — and the plug had to be pushed through it. The pit widened to 47 km only when its roof collapsed after the shots. In this narrow vent Phobos was scored and its boulders rolled (§K.1, K.3).

D.5 Ejection Mechanics

Gas that gathered beneath the plug, at an overpressure of the kind measured in volcanoes on the Earth, pushed it out along its own length. For comparison: before the Vulcanian explosions of Soufrière Hills, Montserrat, in 1997, the dense cap of the conduit needed about 30 MPa to break, and the material beneath it fragmented after a decompression of only a few MPa (Burgisser et al. 2010). Vulcanian explosions throw blocks at up to about 400 $\mathrm{m\,s^{-1}}$ (Fagents & Wilson 1993; Clarke et al. 2015). The speed the gas gives a plug pushed along its own length is $v \approx \sqrt{2\Delta p/\rho}$:

Overpressure	Exit speed	Phobos's orbit at 8.2 h
none (set free)	0	9,890–10,390 km, $e = 0.025$
10 MPa	110 $\mathrm{m\,s^{-1}}$	about $e = 0.06$

Overpressure	Exit speed	Phobos's orbit at 8.2 h
20 MPa	160 m s^{-1}	9,380–11,030 km, $e = 0.08$
50 MPa	250 m s^{-1}	9,020–11,570 km, $e = 0.12$

Table 9. What the volcano adds to the orbit of Phobos (plug density 1,600 kg m^{-3}). The colour fades as the push stretches the orbit.

The pressure lifts nothing; its only role is to open the vent. The orbit comes from the spin, and the gentler the release, the rounder the orbit. The account therefore has a wide margin, below ordinary volcanic pressures.

QUESTION AND ANSWER 11

Can a volcano throw a body 23 km across into orbit?

It does not have to, and it does nothing that volcanoes on the Earth do not do: overpressures of 10–50 MPa and speeds of 110–250 m s^{-1} are the pressures and speeds of ordinary Vulcanian explosions (§D.5). The spin gives the body 2.08–2.43 km s^{-1} (Table 1), the speed of an orbit at the vent; the volcano adds 0.11–0.25 km s^{-1} and only sets the body free. At the vent, on the corotation line, the body weighs nothing.

QUESTION AND ANSWER 12

Would a stronger blast not be safer?

No. A strong push upward, at right angles to the sideways speed that the spin gives, stretches the orbit and lowers its near end (Table 9). The gentler the release, the rounder the orbit.

E The Ejection Sequence

Material left through the pear all the time. Its volcanoes erupted again and again, and most of what they threw fell back to Mars, or reached the inner edge of the ring and returned (§H.5). As ejections, the moons are not unique events. What is unique about them is that they were caught in orbit, or crossed the corotation line, and stayed.

E.1 The First Release – Deimos

Early on, while Mars still turned in 7.3 hours, the vent of Pavonis Mons stood about 1,090 km beyond the corotation line, which then lay 9,080 km from the centre of mass.

The first plug, a little smaller and lighter than Phobos, was set free carrying the sideways speed of the turning vent, 2.43 km s^{-1} — more than a circular orbit needs at that radius. With this excess speed it went onto a long orbit (Figure 7): its near point at 10,170 km, its far point at 23,990 km, eccentricity $e = 0.41$, and an inclination of about 10° to the equator of the young Mars. On every revolution along this orbit it crossed the equatorial ring, whose material moves on circles, at hundreds of metres per second: the ring held it, ground it and in time rounded its orbit (§F).




Figure 7. The two releases, to scale. The pear stands on the equator of the young Mars; the corotation line moves out from 9,080 km to 9,870 km as Mars slows from 7.3 to 8.3 hours. Blue: Deimos's first, long orbit. Green: Phobos's. Lilac: the ring.

QUESTION AND ANSWER 13

What decides that Deimos left first and Phobos second, and not the other way round?

Each volcano on the pear has a window, set by the spin of Mars: while the day is shorter than the day at which the line passes through its vent, a body set free there goes into orbit outside the line. Mars was slowing all the time, and the window of Pavonis Mons closed at 8.3 hours. A body set free at the start of the window, at 7.3 hours, left with excess speed on a long orbit: Deimos. A body set free at the end of the window, near the moment the line passed through the vent, went onto a circle just outside the line: Phobos. A body set free later than 8.3 hours would have started inside the line and fallen back.

QUESTION AND ANSWER 14

Why is Deimos smaller than Phobos, and so much less dense?

Deimos is the first plug, from the top of the vent, and so is made of the youngest material — porous and rich in trapped gas and vapour. Hence its low density, 1,471 kg m^{-3}, against 1,853–1,887 kg m^{-3} for Phobos. It left while Mars was still full of water; in the hour of spin that passed before Phobos left, Mars lost water through the line and slowed (§E.2, J.3). It left about 20.7 km across, a little smaller than Phobos, and drag and grinding in the ring reduced it to its present size, about 12.4 km (§F). Had it not been ground so heavily, Deimos would very likely carry parallel grooves like those of Phobos (§K.1), the marks of a forced passage, under pressure, through a narrow vent. The grooves show that both moons were forced out through the vent under pressure, from the height to which the pear had raised them, into orbit. Other bodies may have left the same source and not survived to an orbit of their own (Appendix B, B9).

E.2 Rotational Deceleration

Between the two releases Mars slowed from 7.3 to about 8.3 hours, and the slowing went on until the day settled at 24.6 hours. The brake is not the moons, whose mass is negligible, but the material that left through the corotation line: water, steam, gas and dust that reached the line carried away the angular momentum of an orbit there. The budget:

Day of the young Mars	Spin, times today's	Share that must leave	Mass through the line
5 h	12.2	92%	11–19% of Mars
7 h	8.7	88%	8–12%
8 h	7.6	87%	7–10%
9 h	6.8	85%	6–8%

Table 10. Angular momentum of a homogeneous young Mars, as a function of the length of its day, against today's ($0.3644MR^2$, 24.62 h). The mass is the amount of material that must cross the corotation line to remove the excess angular momentum; each kilogram carries the angular momentum of a circular orbit at the line, from 9,080 km to 20,428 km. The colour fades with the load.

The moons play no part in this budget: the orbital angular momentum of Phobos is about a millionth of the spin of the young Mars. In the window between 7.3 and 8.3 hours, 6–11% of the mass of Mars must leave this way — about half of what a five-hour Mars would need, and a fifth to a third of the water a young Mars rich in water held. The contraction of Mars, from 5,084 to 3,390 km, makes the load larger, not smaller: had Mars kept its angular momentum while it contracted, it would have spun up by a factor of 2.5, and the brake had to be strong enough to overcome that as well. A young Mars of today's size, turning in 8.5 hours, would need to lose only 2–3% of its mass.

QUESTION AND ANSWER 15

If the mass of the two moons is negligible, what braked Mars so strongly?

The slowing of Mars has nothing to do with its moons; their orbital angular momentum is about a millionth of the spin of the young Mars. Mars was slowed by the water and gas that crossed the line: each kilogram took with it the angular momentum of an orbit at 9,000–20,000 km, four to fifteen times what it held on the ground. A few per cent of the mass of Mars carry off 86–88% of its spin (Table 10).

QUESTION AND ANSWER 16

Is a loss of 6–11% of the mass of Mars in water and gas not far beyond the accepted estimates? The accepted estimate is that the young Mars held water equivalent to a global layer at least 137 m deep, most of it, about 87%, lost to space.

That estimate, about 2×10^{19} kg, measures the slow escape of the atmosphere, which enriched the remaining water in deuterium. Water that left in bulk directly through the corotation line is not sorted by mass, and so leaves no such record. The accepted estimates count only the water that left a record; they include no ring through which water left in bulk. A young Mars rich in water also fits its place in the Solar System: it is the terrestrial planet nearest the asteroid belt, built at the edge of the water-rich region; part of its building material came from the belt (Lunine et al. 2003), and the largest body there, Ceres, is about a quarter water by mass (McCord & Sotin 2005; Park et al. 2016). The account requires a young Mars rich in water — more than 30% of its mass — of which a fifth to a third left this way (§O).

QUESTION AND ANSWER 17

What allowed Mars to slow to 22 hours within its first billion years?

The engine was at its strongest at the start. A hot, expanded planet full of water releases most of its volatiles while it is young; the release fades as it cools and contracts. Most of the slowing came early, and almost all of it before the ring dried: from 8.3 to 22.3 hours, 6–9% of the mass left; from 22.3 to 24.6 hours, only 0.1%. The first billion years is also when, in the accepted view, Mars lost most of its water and atmosphere (Villanueva et al. 2015; Jakosky et al. 2018). How long it took depends on where Phobos stood when the ring dried (Table 13).

E.3 The Second Release – Phobos

When Mars had slowed further, to about 8.3 hours, the corotation line had moved out to 9,870 km, level with the vent of Pavonis Mons. The second plug, Phobos — larger and harder than the first, 1.02 times its present size — was set free there carrying the vent's sideways speed of 2.08 km s^{-1}, almost exactly the speed of a circular orbit at that height:

Day	Line	Orbit from the vent	e	
7.3 h	9,083 km	10,170–23,990 km	0.41	Deimos
7.5 h	9,248 km	10,100–18,850 km	0.30	
7.7 h	9,411 km	10,030–15,420 km	0.21	
7.9 h	9,574 km	9,970–12,970 km	0.13	
8.0 h	9,654 km	9,950–12,000 km	0.09	
8.1 h	9,735 km	9,920–11,140 km	0.06	
8.2 h	9,814 km	9,890–10,390 km	0.03	
8.27 h	9,870 km	9,877–9,919 km	0.002	Phobos
8.4 h	9,973 km	inside the line	0.04	too late

Table 11. The orbits of bodies set free from the vent of Pavonis Mons as Mars slows, with no further push. The window closes at 8.3 hours. Green: the nearly circular orbit in which Phobos was placed, right at the line.

Phobos went directly into a nearly circular orbit just outside the line, at the inner edge of the ring, moving with it at almost its speed.

F The Planetary Ring Dynamics

F.1 Ring Formation and Orbital Damping

The ring was built before the releases, fed directly by Mars itself: steam and fine material from every fissure that reached the corotation line — the height at which turning with Mars is already an orbit — went into orbit, condensed into grains of ice and dust, and settled into an equatorial disc from the line outward. Beyond about 13,000 km the material gathered into clumps.

The material of the ring moves on circles, and a body that lives in it for long takes its orbit from it: circular and equatorial. This is the damping by which discs of gas and debris round the orbits of the bodies within them and flatten them into the equatorial plane. A ring around Mars has been proposed before, on other grounds: from a resonance that tilted Deimos (Ćuk et al. 2020), and as a stage in a cycle of rings and moons (Hesselbrock & Minton 2017). This damping is why the two moons are on such nearly circular orbits today — Deimos at only $e = 0.0003$.

F.2 The Split Interaction

The very different fates of the two moons come from the different conditions in which they met the ring. Three things act the same way:

	Deimos	Phobos
Its orbit in the ring	long ($e = 0.33$): crosses the circling material at hundreds of $\mathrm{m\,s^{-1}}$ on every revolution	nearly circular: moves with the ring, almost no relative speed

	Deimos	Phobos
Where	through the whole ring, into the clumps beyond 13,000 km	the thin inner edge, next to the line
Size against the ring	smaller and lighter: area per unit mass 2.3 times that of Phobos today	larger and denser
Result	held, ground, its orbit rounded; it lost most of its mass	barely touched; held at the line; kept its grooves, pits and Stickney

Table 12. Two bodies, one ring: how each moon responded to it.

On the long orbit of Deimos, the drag of the ring added speed where the body was slowest, at the far end of its orbit, where it moved slower than the ring, and so gradually raised the near end: the orbit rounded outward. With its own angular momentum alone, Deimos would round near 14,300 km; the ring, faster than it at the far end, carries it further out.

F.3 The Velocity Limit

The ring moves on circles at the speed of an orbit. It can bring a slower body up until the speeds are equal — as it did for Deimos, at the far end of its long orbit — but it cannot give a body more than its own speed. Phobos left at almost the speed of an orbit at the line; there was no difference in speed between it and the ring material, and the ring had nothing to give it.

F.4 The Gas Leash

At its innermost edge, next to the line, the ring is mostly thin gas. Gas held up partly by its own pressure moves a little slower than an orbit, so Phobos met there a weak head wind — the same head wind that a gas disc exerts on the solid bodies within it, the basis of the theory of how planets grow (Weidenschilling 1977).

The tide pushed Phobos outward with an acceleration of only about 10^{-11} m s^{-2}; to balance it, a head wind of about 100 m s^{-1} needs a gas density of only 3–8 $\times 10^{-8}$ kg m^{-3}, as thin as the Earth's air at about 115 km. A balance followed: the tide pushed, the gas held back, and Phobos stayed at the inner edge, just outside the line. As Mars slowed, the line moved outward; the inner edge of the ring moved with it and carried Phobos along.

F.5 The Ring Dissipation

The steam that slowed Mars is the steam that fed the inner edge of the ring. The ring was fed as long as the ground was loose. The young Mars — hot, fast-spinning and not yet sorted — had a crust that was not yet compacted, and its volatiles escaped through it freely: everything that drives volatiles out acted at once, and at first the ring was massive. With time the ground tightened. As the spin slowed and the interior went on sorting, the confinement grew stronger; as the crust cooled and thickened,

it became complete. Volcanic outgassing on Mars follows the same course: most efficient before about 4.1 billion years ago, strongly declining through the Hesperian, and insignificant later (Grott et al. 2011).

When the confinement was complete, and Mars had lost its water and approached its present spin, the flow of steam stopped and the ring dried: its material spread, fell back inside the line as frost and dust, or evaporated. Phobos was then near 19,000 km, and the day of Mars about 22 hours.

The drying went from the outside in: the ring was fed only at its inner base, on the line; its distant parts received material only by spreading, and beyond 13,000 km its material gathered into clumps that no longer behaved as a ring.

F.6 Dating the Dissipation

The present orbit of Phobos allows the drying to be dated backwards. From the moment the ring and its band of gas had gone, Phobos fell inward; running its fall backwards at the tidal friction measured today gives the time, and the place it stood gives the day of Mars then:

Phobos when the ring dried	Day of Mars then	Dried, years ago	Time Mars had to slow
17,000 km	18.7 h	1.8 Gyr	2.7 Gyr
18,000 km	20.4 h	2.6 Gyr	1.9 Gyr
18,800 km	21.7 h	3.5 Gyr	1.0 Gyr
19,100 km	22.3 h	3.9 Gyr	0.6 Gyr
19,500 km	23.0 h	4.5 Gyr	none

Table 13. When the ring dried, from the orbit of Phobos today (tidal friction $k_2/Q = 1.816 \times 10^{-3}$, constant). 19,500 km is the upper limit: beyond it the ring would have dried before Mars existed. A softer Mars after the drying would make the fall faster and the drying later.

The most likely reading places the drying of the ring 3.5–3.9 billion years ago, at a day of about 22 hours.

QUESTION AND ANSWER 18

Why did the ring, which carried Deimos out to its present orbit, not push Phobos outward too?

A ring can bring a body up only until its speed equals that of the ring material, not beyond it (§F.3). Deimos left on a long orbit, at the far end of which it was slower than the ring, and so it was pushed outward; Phobos left directly onto an orbit at the line, already at the ring's speed, and there was nothing to push it.

QUESTION AND ANSWER 19

What held Phobos at the line and kept the tide from pushing it outward?

The head wind at the gaseous inner edge of the ring (§F.4). A gas density of a few 10^{-8} kg m^{-3} balances the tide; the gas there was fed by the same steam that left Mars and slowed its spin.

QUESTION AND ANSWER 20

Why are the two orbits so nearly circular and equatorial?

It is the direct result of the time they spent in the ring. The ring was round and lay in the equatorial plane, and a body that lives in it takes its orbit from it (§F.1). Deimos kept this in full, $e = 0.0003$. The present eccentricity of Phobos, 0.015, is not original: the tide raised it later, during its fall inward.

QUESTION AND ANSWER 21

Why should the outer ring have gone first?

The ring was fed from the inside, at the line; its outer parts received material only as it spread. Beyond 13,000 km the material gathered into clumps that no longer behaved as a ring. And as the line moved out, the inner edge was rebuilt on it as long as steam came. When the flow stopped, the outer parts emptied first, and the last of the ring was the gaseous band at the line — the leash.

G Tidal Mechanics

Beyond the corotation line, the tide of the young Mars — hot, soft and less viscous — pushes a moon outward. The climb is faster for a heavier moon and for a larger Mars:

Climb	Mass	8× today's friction	20×
Deimos, 14,300 → 23,463 km	at release, 6.8×10^{15} kg	370 Myr	150 Myr
Deimos, 14,300 → 23,463 km	today, 1.5×10^{15} kg	1.7 Gyr	680 Myr
Phobos, 9,880 → 19,000 km	1.13×10^{16} kg	58 Myr	23 Myr

Table 14. Tidal climbs of the moons on the young Mars (radius 5,084 km), by the mass of the body and the tidal friction of Mars.

Deimos was carried out while it was still heavy, before it was ground, and Mars still large, and passed 20,428 km — the final place of the line — before the line got there; the outer ring shortened the climb. Phobos, the heavier, would have climbed fast, but the leash of the ring's inner edge held it next to the line (§F.4), and it moved outward only as fast as the line did. When the ring dried, near 19,000 km and a day of about 22 hours, the line passed it. From then on Phobos was inside the line, and the tide drew it back — slowly at first, because it was close to the line, where the tide is weak. running

its fall backwards at today's tidal friction places it at 19,100–19,500 km 3.9–4.5 billion years ago.

The fall was slow at first and fast at the end:

Since the drying	Years ago	Phobos	Falling, per century
0	3.9 Gyr	19,100 km	about 4 cm
1 Gyr	2.9 Gyr	18,250 km	
2 Gyr	1.9 Gyr	17,100 km	
3 Gyr	0.9 Gyr	15,250 km	
3.5 Gyr	0.4 Gyr	13,500 km	about 24 cm
3.8 Gyr	0.1 Gyr	10,900 km	
3.9 Gyr	today	9,376 km	1.8 m (measured)

Table 15. The fall of Phobos from the drying of the ring to today, at today's tidal friction. In the first three billion years it fell only 3,800 km; in the last 400 million years, more than 4,000 km.

The leash, then, had to hold until the late stages of the slowing of Mars. Had the ring dried early, while Phobos was still near 9,900 km, the line would have passed it there, and at today's tidal friction it would have fallen to its present orbit in less than a hundred million years and broken up soon after.

QUESTION AND ANSWER 22

Is a fall lasting 3.9 billion years plausible?

Yes. The tide that draws Phobos down weakens with about the 5.5th power of its distance: at 19,100 km it falls fifty times more slowly than today. The rate today is measured — Phobos runs ahead of its predicted place as it falls about 1.8 m a century — and the tidal friction of Mars used here comes from that measurement ($k_2/Q \approx 1.8 \times 10^{-3}$; Lainey et al. 2021). Phobos will reach the Roche limit and break into a ring within some tens of millions of years (Black & Mittal 2015; Hesselbrock & Minton 2017). Most of the fall came in the last few hundred million years (Table 15).

QUESTION AND ANSWER 23

Why did Phobos not fall onto Mars long ago?

Because the leash at the inner edge of the ring held its orbit until Mars had almost finished slowing. Had the ring dried while Phobos was near 9,900 km, the tide would have brought it down to its present orbit in less than a hundred million years and broken up soon after.

QUESTION AND ANSWER 24

Is the tide fast enough to bring Deimos to 23,463 km in time?

Yes, if the tide carried it while it was still heavy and Mars still large and soft: 150–370 million years from 14,300 km (Table 14). It had to pass 20,428 km before the line arrived there; the outer ring, while it lasted, carried it further.

H The Post-Ejection Drainage

H.1 Relaxation and the Tharsis Bulge

With the two bodies gone and the vents open, and as the interior sorted itself and the core settled, the pear returned. It sank and lay down slowly and westward, the way it had leaned (§C.4). Its body is today the Tharsis rise.

The volcanoes that stood on the pear now stand on the ground, Pavonis Mons among them, with the single deep pit from which the two moons left. As the pear returned, the fluid or viscous material inside it drained back from the tip to the base, along its axis and down the slope to the east.

H.2 The Neck

Between the tip and the centre of the base, on the axis of the pear (within 70 km of it, 1,750–2,460 km from the eastern point of the base), lies Noctis Labyrinthus: a network of troughs and pits that cross one another and leave blocks between them, like a broken slab (Figure 8). Toward the east the troughs join into one broad channel — the head of Valles Marineris.

Blocks form when the crust is pulled apart in several directions at once; stretching in one direction gives only parallel cracks. The pits show that material beneath the surface left and the ground above it sank. In the accepted reading the network is a set of grabens at the top of the Tharsis rise, with collapse after material below withdrew.

In this account, at the neck lay a large mass, at the centre of the descent of the pear. As the pear returned, the plates of its surface came down from above. Where the steep slope met the gentle one, the change of angle bent them and broke them into blocks. The fluid or viscous floor beneath them spread sideways, as continents spread but on a small scale, and carried the broken plates apart with it; the troughs are the gaps between them. The labyrinth is thus the head of the drainage: like the tributaries of a river, its troughs gather into one main channel that runs east.

The grooves around the neck. North of the labyrinth, between the equator and about 3°S, and to its south-east near 16°S, the ground is cut by long, straight, parallel grooves (arrows in Figure 8). In the accepted reading they are grabens of the Tharsis fracture system (§C.8). In this account they are the stretching of the same descent: the surface that came down from the pear was pulled toward the spreading floor, and where it did not break into blocks it opened in parallel cracks.




Figure 8. Noctis Labyrinthus, about 5--12°S, 255--265°E (labels −95 to −105°E), and the head of Valles Marineris to the east. Arrows: the long parallel grooves north and south-east of the labyrinth. Credit: ACT-REACT QuickMap Mars; MOLA colour elevation (NASA/GSFC).

H.3 The Axial Conduit

Valles Marineris is the route by which the interior material of the pear flowed east as the pear sank, down the slope. Three findings support this.

It runs along the axis. For about 1,700 km, from within the base to its eastern point, the floor of the canyon runs 100–360 km north of the axis of the pear and nearly parallel to it. The canyon does not cross the pear; it follows it. (The eastern point of the base was itself placed in the canyon; its continuing along the axis for 1,700 km does not follow from that choice.)

Its floor descends to the east. Measured on MOLA expanded to degree 719 (about 15 km) and relative to the areoid computed from GMM-3 — the level along which a fluid flows:

Longitude	Place	Floor
262°E	Noctis Labyrinthus	+4.9 km
270°E		−1.1 km
278°E	Ius / Melas	−4.1 km
288–306°E	Melas, Coprates	−5.0 to −5.2 km
310–320°E	Eos / Capri	−4.3 to −4.5 km

Table 16. The floor of Valles Marineris, the deepest point at each longitude. The colour deepens as the floor falls eastward; deep floors narrower than the resolution are smoothed.

A total fall of about 10 km from west to east, in the direction of the flow.

It lacks mass. The free-air gravity over the canyon is about −500 mGal (Table 6): a channel that material passed through and left.




Figure 9. Valles Marineris from Noctis Labyrinthus (left) to the chaos at its eastern end (right). On its northern side the basins of Ophir, Candor and Hebes Chasmata, and Juventae Chasma further east, open toward the equator (yellow line). Credit: ACT-REACT QuickMap Mars; MOLA colour elevation (NASA/GSFC).

H.4 The Basin and the Spill

The floor does not fall all the way to the eastern exit: from its deepest point it rises about 0.8 km before the eastern exit. This is a basin. Harrison & Chapman (2008) found it on the inner walls of the canyon: scarps and terraces over about 1,500 km, all at one elevation, $-3{,}560$ m, the level of a lake that filled the centre of the canyon. The lake overflowed at the eastern end, in Eos Chasma, into the chaos and the outflow channels; the erosion near the spill point runs east–west. The inner walls kept the level of the flow, and the exit was to the east.

The basins of the canyon open toward the equator. On its northern side Hebes, Ophir, Candor and Juventae Chasmata lie as separate depressions, and from Juventae the channel of Maja Valles runs north (Figure 9). At the eastern end, Eos and Capri Chasmata open into the chaos of Aurorae and Hydraotes, and from there the outflow channels Simud and Tiu run north toward the equator and on to Chryse, with Ares Vallis to their east (Figure 10).

Baker & Milton (1974) described in these channels streamlined hills and longitudinal grooves, like those of the great floods of the Channeled Scabland on Earth; a streamlined hill points its tail downstream. At our resolution the floors of these channels are nearly level (-4.2 to -3.9 km); their direction is shown by their forms, not by their slope.




Figure 10. The eastern end of the canyon: Eos and Capri Chasmata open into the chaos, and from there channels run north toward the equator (yellow line). Credit: ACT-REACT QuickMap Mars; MOLA colour elevation (NASA/GSFC).




Figure 11. The drainage path of the pear. Top: from the neck of the pear along the canyon, over the spill point at Eos, through the chaos and north to Chryse; dashed, the axis of the pear. Bottom: the floor of the canyon and the level of the ponded lake.

THE DRAINAGE OF THE PEAR

Head of the drainage (Noctis Labyrinthus, at the neck) → channel along the axis (Valles Marineris, falling 10 km to the east) → basin filled to one level on the walls ($-3{,}560$ m) → spill to the east (Eos Chasma) → chaos → outflow channels north toward the equator and Chryse, with the forms of flow.

H.5 The Dust Provenance

The ring was not made of steam alone. Besides the enormous amount of volatiles that built it, the volcanoes on the pear threw out solid Martian material — matter of Mars itself — that reached no orbit of its own — shots weaker than those of the two moons, and the debris of the vents. It reached the inner part of the ring, and most of it came back to Mars. While it was in the ring it was ground, against the ring's material and against itself, into small fragments and fine dust.

When the ring dried, its water was broken apart by sunlight and the solar wind, and

the light hydrogen escaped to space; the heavy isotope, deuterium, stayed behind, and the water left on Mars became enriched in it. The fine dust the ring had held came down on Mars, mostly near the equator, together with frost.

In this account that is the source of the dust of Mars: a fine material spread over the whole planet, of much the same composition wherever it has been measured (Berger et al. 2016). The accepted view looks for its source on the ground, in deposits eroded by the wind, the largest being the Medusae Fossae Formation (Ojha et al. 2018). Dust that spent its time in the ring carries what the ring gave it: an age older than 4 billion years, the composition of the Phobos regolith, and grains that carry helium and neon from the solar wind; it should be most abundant in the broad equatorial band (prediction N.8). The loss of the ring left the surface dry.

H.6 The Residual Volatiles

The water inside Mars did not have to come back in when the pear sank: most of it never went out. The steam that fed the ring and the gas beneath the vent, which burst the plugs, were the part of the volatiles that found a way out through fissures and vents.

The rest stayed sealed beneath the crust, and remains there today, in the pores of the mid-crust where InSight found it.

PART II

The Evidence

Part I described the sequence of mechanical and hydrodynamic events in the early history of the system. Part II sets out the geomorphological and geophysical evidence that this sequence left behind, place by place. Each section sets the observation against what the model says about it, and gives the numbers.

I The Summit Caldera

I.1 The Caldera Geometry

Pavonis Mons rises about 14 km above the mean level of Mars on a base about 375 km wide. Its summit caldera is a single circular pit 47 km across and nearly 5 km deep. It is a steep-walled cylinder, not a shallow bowl; to its north-east lies a secondary depression, broader and shallower, about 90 km across (USGS/IAU Gazetteer; NASA/JPL). The ratio of depth to width, about 0.10, is the measure of how much the summit is a barrel.

A comparison of the summits of Tharsis brings out what is particular to Pavonis:

Volcano	Latitude	Caldera width	Caldera depth	Form
Pavonis Mons	1.5°N	47 km	nearly 5 km	single deep circular pit
Olympus Mons	18.7°N	about 70 km (nested)	about 3 km	a complex of overlapping, stepped pits
Arsia Mons	8.4°S	about 110 km	to be measured	one broad pit
Ascraeus Mons	11.8°N	nested pits	to be measured	several overlapping pits

Table 17. The geometry of the four calderas of Tharsis. Green marks Pavonis: a single pit, narrow and deep against those of the other mountains. The depths marked "to be measured" can be read from MOLA topography.

I.2 The Volume Balance

The volume of the Pavonis caldera, taken as a cylinder 47 km wide and 4.8–5.0 km deep, is 8,300–8,700 km^3. The two moons today hold about 5,740 km^3 (Phobos) and

about 1,000 km^3 (Deimos), together about 6,740 km^3.

The pit is therefore of the scale of the two plugs that left it. It is not a mould of the bodies themselves but the space that collapsed after the shots, so it does not set their size.

I.3 The Caldera Collapse

When the plugs had left, the shallow reservoir beneath the vent lost what held up its roof. The roof cannot stand, and it falls vertically inward.

A steep, deep, circular pit is what a reservoir leaves when its roof collapses vertically; its depth, about 5 km, is the depth of the reservoir (§D.4). The model expects a measurement of the gravity beneath the caldera to show the missing mass of a reservoir that fired (prediction N.5).




Figure 12. The Tharsis province, Viking colorised mosaic: Olympus Mons at upper left; Ascraeus, Pavonis and Arsia Montes on the diagonal. Pavonis, in the middle of the row, is the closest to the equator. Credit: ACT-REACT QuickMap Mars; Viking Colorized Global Mosaic (NASA/JPL/USGS).

J The Deimos Record

J.1 The Damped Satellite

Deimos is the smoothest small body yet seen near Mars. Its depressions are shallow and worn, and it has no grooves or long drag marks. Measured on the shape models of Ernst et al. (2023) in the same way for both moons, the depressions wider than 0.7 km on Deimos have a median depth of 1.9% of their width, against 5.5% on Phobos;

84% of them are shallower than 6% of their width, against 56% on Phobos, and only 3% are deeper than 12%, against 13%.

Its orbit is almost a perfect circle (eccentricity 0.0003), close to the equatorial plane, and it lies beyond the corotation radius — the distance at which a body turning with the planet already moves at the speed of a free orbit. A circular, equatorial orbit together with a smooth, worn face are the marks of a body that spent a long time in a ring: the ring carried it into a circular, equatorial orbit, and its particles wore it down.

Deimos goes round more slowly than Mars turns, and it has done so throughout the history of the system: at 14,300 km, where its orbit became round, it went round in 14.4 hours against a Martian day of 7.3–8.3, about 0.5–0.6 times a day; today, at 23,463 km, it goes round in 30.3 hours against 24.6, 0.8 times a day. It has always stood beyond the line. The line moved outward behind it and never caught it — the reverse of Phobos (§K.8).




Figure 13. Deimos, HiRISE (two views; scale bar 2 km). A smooth, rounded body, with shallow, worn craters and no grooves. Credit: NASA/JPL-Caltech/University of Arizona (PIA11826).

J.2 The Mass Ablation Profile

Grinding and drag in the ring remove material. Deimos left the vent of Pavonis considerably larger than it is now, but not larger than Phobos, and it left smooth: as the first plug in the vent it did not rub against its walls. How much it lost during its stay in the ring depends on its diameter when it left:

Diameter on leaving	Volume, km³	Share lost by today	Layer ground away
12.4 km (today)	1,000	—	—
14.0 km	1,430	30%	0.8 km
15.6 km	2,000	50%	1.6 km
16.8 km	2,500	60%	2.2 km
20 km	4,190	76%	3.8 km
22.3 km (as Phobos)	5,800	83%	4.9 km

Table 18. Deimos at departure and in the ring: larger than today, not larger than Phobos. The colour deepens with the share ground away as the diameter at departure grows.

J.3 The Volatile-Rich Composition

Deimos is much lighter than Phobos: 1,471 kg m^{-3}, against 1,853–1,887 kg m^{-3}. The difference supports the order of events in the model directly. The first plug (Deimos) came from the top of the vent, from the youngest part of the crust, the richest in water, steam and trapped gas. The second plug (Phobos) formed later and deeper in the emptied vent, from material that had already dried and compacted.

The order of the densities is the order of the shots, and it is also the order in which Mars dried. Between the two releases Mars lost large amounts of water and gas through the corotation line — that loss is what slowed its spin from 7.3 to 8.3 hours (§E.2). Deimos was made of a planet still full of water; Phobos, an hour of spin later, of a partly dried one, and so it is the denser of the two.

The moons are samples of the crust of the young Mars: a third to a half of each is empty space, a foam that froze, and their densities, 1,471 and about 1,860 kg m^{-3}, are those of the upper layers of a planet whose mean density was only about 1,170 kg m^{-3} — light, porous and rich in water.

K The Phobos Record

K.1 The Striation Evidence

The surface of Phobos is crossed by extensive, coherent families of long, parallel grooves (Thomas et al. 1979; Murray & Heggie 2014). In the model of this paper they are the marks of the walls of the vent. Before the moons left, the vent of Pavonis was much narrower than its pit is today, about as wide as the plug it held, and the plug was pushed through it under high pressure (§D.4).

The steam and fragments that opened the vent did not leave its wall smooth: they left it ribbed with long, thin, hard plates standing out from the wall, like the gills under the cap of a mushroom. A hard, dry plug pushed through such a ribbed vent, and pressed against one side of it by the Coriolis force (a body rising through a spinning planet lags behind the speed of the walls and is deflected westward), is cut by the plates along its whole length, parallel to its motion.

A modern analogue is the solid spines extruded from the crater of Mount St Helens in 2004–2006, which left the conduit scored along their direction of motion (Cashman et al. 2008). The grooves of Phobos survived because it stayed only briefly in the thin edge of a waning ring and lost about 1% of itself, a layer of some 40 m.



Figure 14. Phobos, HiRISE. Centre and left: extensive systems of long parallel grooves and chains of pits; Stickney at the right. Credit: NASA/JPL-Caltech/University of Arizona (PIA10368).

K.2 The Post-Ejection Relaxation

Phobos left the vent long and soft, warm and drawn out along its motion, and its grooves were cut straight along that axis. In space, as it cooled, it settled gradually toward a rounder shape, and the grooves, carried on its surface, bent with it.

Today the grooves form a few large, coherent families across the whole body, curved rather than straight (Figure 15). In the model they are grooves that were straight when cut, on a body that changed its shape afterwards; restoring the present shape to the elongated one should straighten them (prediction N.2).




Figure 15. Groove orientations on Phobos as mapped by Cheng et al. (2022). The grooves form a few large, coherent families that curve along the body. Credit: Cheng, Asphaug, Ballouz, Yu & Baoyin (2022), *Planet. Sci. J.* 3, 249, CC BY 4.0.

K.3 The Roller-Bearing Pits

The chains of pits that run beside the grooves, parallel to them, are a second record of the same passage through the vent, made by a different tool. Boulders caught in the narrow gap between the plug and the wall did not bounce: they were held against the wall and pressed into the soft surface of the plug.

As the plug moved past, they rolled along it like the balls of a bearing, and every bump on a boulder pressed one pit at each turn. This mechanism accounts for the properties of the chains:

- A chain is straight and parallel to the motion.
- Its pits are evenly spaced, at a spacing set by the circumference of the boulder and the number of its bumps.
- It runs toward the rear end of the body, which passed the caught boulders last.
- Grooves and chains were made at the same time and in the same direction, so the chains always run beside the grooves and never cross them.

Pit chains and impact craters. The chains are not chains of secondary impacts. An impact crater that lands on a groove cuts it and erases the part it falls on; such craters exist on Phobos, superposed on grooves (Figure 16, centre). The chains do the opposite: they keep to the direction of the grooves and never cut them. Beside them Phobos carries true impact craters, of three generations: old, shallow, rounded hollows from the time of the ring; small, sharp craters made by material moving in the ring almost in step with Phobos; and sharp, young craters made after the ring had gone (§L).




Figure 16. Phobos, HiRISE detail. Centre: small, later impact craters superposed on grooves, cutting them. Upper right: chains of pits running beside the grooves, in their direction --- made in the same passage through the vent, not later. Credit: NASA/JPL-Caltech/University of Arizona (HiRISE, detail).

K.4 The Leading Apex

Stickney, 9 km across (IAU centre 1°S, 49°W), lies at the blunt end of Phobos. On the shape model of Ernst et al. (2023) its centre lies 79% of the way from the sharp tip to the blunt one, and its rim reaches to within 0.4 km of the blunt tip, at the corner toward the leading side (Figure 18).

In the model the blunt end of the plug led the rise through the vent. At the vent mouth the leading end met the material in its way and pushed it out, and the hollow left by that encounter is Stickney. Stickney therefore formed after the grooves, whose remains are seen on its floor (§K.5).

A second reading. A second reading remains open: Stickney as the mark of a solid body, already moving in the ring before Phobos left, met by a Phobos that was still warm. An impactor of 1.5–2 km at a relative speed of about 1 $\mathrm{km\,s^{-1}}$ suffices; such an impact would have slowed Phobos by about 1 $\mathrm{m\,s^{-1}}$. Both readings place the formation of Stickney on a warm, soft Phobos, and both make it younger than the grooves.

K.5 The Intracrater Deformation

The grooves on the floor of Stickney were cut on the plug before the hollow formed, and were pressed in the encounter. On the 36-m shape model of Ernst et al. (2023) we compared the floor of Stickney, within half its radius and without the later inner crater Limtoc, with a ring around it, from 1.2 to 2 radii (Figure 17). The same filter was applied to both: it picks out long troughs 50–350 m wide in any direction and removes the curvature of the bowl. To judge whether the difference is unusual, the same comparison was repeated at 80 random places on Phobos.

The measurements gave the following values:

Measure	Floor	Outside	Floor / outside	Stickney lower than, of 80 places
Typical groove relief, m	1.56	1.74	0.89	65%
Area with strong grooves (>3 m)	0.9%	5.5%	0.17	76%
Sharpness: depth for width	0.34	0.44	0.77	89%

Table 19. Grooves on the floor of Stickney and around it, measured with the same filter. The colour deepens with how unusual Stickney is among 80 random places on Phobos.

Inside, the grooves keep most of their depth but are wider for it, and the sharp ones have nearly gone; in sharpness the floor is below 89% of the random places. This is the shape of a groove pressed flat, not one cut afresh or covered. No loose material flowed in to fill the floor: the grooves on it are pressed, not buried.

The shape model at this resolution smooths grooves to a few metres of relief, against tens of metres in images, so the contrast it gives is a lower bound; data at higher resolution should show it more sharply (prediction N.2).




Figure 17. Stickney on the 36-m shape model of Phobos (Ernst et al. 2023), shaded relief, in km. Green: the floor compared; dashed: the ring around it. Grooves cross the floor, fainter and broader than outside.


K.6 Mass Asymmetry and Centre-of-Mass Offset

The centre of mass of Phobos is not at its geometric centre: it is displaced toward its blunt end. This distribution was set in the reservoir, by the fast spin of the young Mars,

before the plug left it. At the top of the standing pear the spin outweighed gravity: the net pull on the plug pointed outward, up to 0.17 $\mathrm{m\,s^{-2}}$ (Table 1, §D.2). Under this reversed pull the denser material gathered at the outer end of the plug, and the plug left with that end leading. Gas pressing from below soaked the lower part of the plug and foamed it at the vent mouth, while the front was compacted against the material in its way: a dense front and a light, foamed rear.

As Phobos left the vent, its heavy end led, pointing away from Mars (Figure 18). Today it faces Mars. The spin set the heavy end outward; the gravity of Mars then turned it toward the planet, through 180°. The tide holds the long axis of Phobos toward Mars but does not by itself tell one end from the other; how the turn was completed is set out as a requirement of the model (§O). The prediction stands either way: the centre of mass of Phobos is displaced toward Mars, toward its blunt end, by 300–500 m (prediction N.6).

We examined whether the push of the ring, which held Phobos at the corotation line throughout (§F.4), could have turned its centre of mass. The push acts on the body as a whole, and the centre of mass, displaced by 300–500 m, gives it a lever; the tide, which holds the long axis toward Mars, is stronger by a factor of twenty million to a hundred million (between 19,000 and 9,900 km). The push could turn the heavy end by less than a millionth of a degree. The turning of the centre of mass by the ring is therefore neglected.



Real equatorial outline of Phobos (shape model, Ernst et al. 2023). Red: Stickney (IAU centre 1°S 49°W, 9 km wide). Green: the heavy end, where the centre of mass is displaced.
The same end pointed away from Mars as Phobos left the vent and faces Mars today: it turned through 180° relative to the planet.

Figure 18. The real equatorial outline of Phobos (Ernst et al. 2023). Red: Stickney. Green: the heavy end. It leads, away from Mars, as Phobos leaves the vent; today it faces Mars.

K.7 The Paleo-Orbit Reconstruction

Phobos today goes round Mars in 7.65 hours — 3.2 times in each Martian day — at 2.14 $\mathrm{km\,s^{-1}}$. Its orbit is nearly circular (eccentricity 0.015) and within about a degree of the equatorial plane. Inside the corotation line, every body goes round faster than the planet turns; the tide it raises lags behind it and holds it back, and it spirals down. That is what Phobos is doing, and what it has done since the moving line passed it, at about 19,000 km (§F.5, G).

The back-integration below does not depend on how Phobos reached that height; it fixes only where it stood when its fall began:

Fall began	Distance then	Its orbital period then
3.5 billion years ago	18,807 km	21.8 h
3.9 billion years ago	19,119 km	22.3 h
4.5 billion years ago	19,541 km	23.0 h
Today	9,375 km	7.65 h

Table 20. Phobos run backwards with the tidal friction measured for Mars today ($k_2/Q = 1.816 \times 10^{-3}$). In every case the starting point of the fall lies within a few per cent of the corotation line of a slowed Mars, and it has fallen since.

K.8 The 3.2 Commensurability

When its fall began, Phobos and Mars were nearly in step: at about 19,100 km Phobos went round once in 22.3 hours, and Mars turned once in about 22 hours — close to one orbit per Martian day. The ratio of 3.2 seen today was built afterwards, by two changes acting together in the same direction:

Phobos fell, and falling made it faster. A moon's speed is set by its distance from the centre of the planet: the closer, the faster. The tide holds Phobos back and takes energy from it, and it sinks; as it sinks it speeds up, from about 1.50 km s^{-1} at 19,100 km to 2.14 km s^{-1} at 9,375 km. Its period shortened from 22.3 to 7.65 hours — a factor of 2.9.

Mars slowed. The planet went on losing spin with the water, gas and steam that left it through the corotation line, the ring among them: a ring outside the line carries away the spin it holds, and when it was gone that spin did not return. The Martian day lengthened from about 22 to 24.6 hours — a factor of 1.1.

The two changes together give the ratio of today:

	When the fall began	Today	Change
Distance of Phobos	19,100 km	9,375 km	
Speed of Phobos	1.50 km s^{-1}	2.14 km s^{-1}	
Period of Phobos	22.3 h	7.65 h	faster $\times 2.9$ (the fall)
Martian day	≈ 22 h	24.6 h	slower $\times 1.1$ (Mars)
Orbits per Martian day	≈ 1	3.2	$2.9 \times 1.1 \approx 3.2$

Table 21. From one orbit per day to 3.2. Phobos began nearly in step with Mars; its fall and the slowing of Mars together produced the ratio seen today. The colour shows the growing difference between the speed of the moon and the spin of the planet.

L The Ring Signatures

- **Two equatorial orbits.** Both moons go round Mars within two degrees of the equatorial plane and on nearly circular orbits. This is direct dynamical evidence of an ancient ring: a ring in the equatorial plane draws any body crossing it into that plane and absorbs its excess motion until its orbit is round.
- **A face that follows time and place in the ring.** The surfaces of the two moons reflect where they were in the ring and for how long. Deimos stayed long in the heart of the ring: it is smooth and rounded, and on the more perfect circle (eccentricity 0.0003). Phobos met only the thin inner edge of a waning ring: it was not smoothed, kept the grooved, rough face it was given in the vent, and was left with a small eccentricity (0.015).
- **The inclination of Deimos.** The ring left Deimos on a circle in the equatorial plane; its present tilt of about 1.8° is not original: it was added later, in a passage through a resonance with a massive inner moon. In the accepted literature this tilt has been read, independently, in the accepted literature as the trace of a past ring around Mars (Ćuk et al. 2020). Independent work thus already places a ring around the young Mars.
- **Worn hollows.** The depressions of Deimos are three times shallower than those of Phobos (§J.1), and both moons carry an old generation of worn, rounded hollows beneath a young generation of sharp craters. The model explains this by the state of the moons when they were struck: a moon struck while still warm and soft is not dug into a sharp-rimmed bowl but folds inward, and low-speed contact with bodies moving almost in step with it in the ring leaves broad, shallow hollows. Only a moon that passed through a ring carries them. They were made early, at 10,000–24,000 km, so the near ring held solid bodies — stone and ice thrown from the volcanoes, which survive well inside the 13,000-km limit for loose ice (a solid ice block to about 7,700 km, stone to about 5,300 km). The grinding in the clumpy outer ring came later, and the sharp craters last, on bodies that had cooled and hardened.
- **The Moon never passed through a ring.** No lasting ring could form around the young Earth: the pear that built the Moon swept the equatorial belt in which it would have formed and gathered its material into the forming Moon. At Mars, the time a body spent in the ring sets how much of its first face it lost: Deimos, long in the heart of the ring, is smooth, rounded and reduced; Phobos, briefly at its edge, is grooved and kept its mass; the Moon keeps a saturated crater record and its whole mass.
- **Ages.** Crater counts put the surface of Phobos at about 4.3 billion years in a Mars-orbit chronology, or 3.5 in an asteroid-belt chronology (Schmedemann et al. 2014); the surfaces of both moons are near saturation and of similar crater density. The model expects a direct test of the material to show that Deimos is the older of the two.

M The Hydrological Record

- **Water that left.** Heavy hydrogen in today's atmosphere enriched six to seven times relative to the Earth's oceans, and an early global water layer of at least 137 m (Villanueva et al. 2015): the record of the equatorial ring and of the steam

that crossed the corotation line and went to space.

- **Water that stayed.** A water-saturated mid-crust between 11.5 and 20 km depth beneath InSight, equivalent to a global layer 1–2 km deep (Wright, Manga & Panning 2024), fits the model: it is the water that found no path to the vents or the fractures and stayed sealed beneath the crust. The moons, soft when they were struck in the ring (§L), show that the ground beneath the surface of the young Mars was soft and soaked with water; the mid-crust of today is what remains of it, frozen and stable.
- **One sealed crust.** The same impermeable crust that held the gas under pressure until the plugs moved and the moons left is the crust that holds the water inside Mars today.

PART III

Empirical Predictions

A scientific hypothesis must give quantitative predictions that can be refuted. This part collects the predictions that follow from the LEF model. Each prediction gives what is expected, its value where one can be given, the instrument or data that will measure it and when, and what the accepted explanation expects at the same place — so that each measurement separates the two. The tests will become possible with the data of the coming missions, foremost among them the Martian Moons eXploration (MMX) mission of the Japan Aerospace Exploration Agency (JAXA), which will land on Phobos, fly past Deimos and return samples to the Earth.

N The Predictive Matrix

Prediction	Expected (this paper)	Measured by	When	Accepted explanation expects
N.1 Composition of Phobos	Original Martian crust altered by water: oxygen, chromium and titanium isotopes on the Martian line alone, with no foreign impactor component; moderately volatile elements (zinc, potassium) showing no evaporative shift; hydrated minerals; closed vesicles of a cooled foam	Laboratory analysis of MMX samples, ≥ 10 g of regolith	Return 2031	Impact disc: a mixture of Mars and impactor, much of it melted and re-condensed. Capture: primitive asteroid material

Prediction	Expected (this paper)	Measured by	When	Accepted explanation expects
N.2 Phobos grooves	Cut by the rigid plates of the vent wall: parallel to the direction of motion, concentrated on the side that faced west in the vent; bent with the body as it rounded, so that restoring an elongated shape straightens them; on the floor of Stickney pressed, wider for their depth than outside, more clearly at higher resolution, with no cover of loose regolith; rims sharp or only slightly worn, the surface etched, not smoothed	MMX imaging; HRSC; shape models	Now and 2027 on	Tidal stress or Stickney ejecta; no difference inside Stickney
N.3 Phobos pit chains	Pressed by trapped boulders that rolled between the plug and the wall, as in a ball bearing: straight, parallel to the grooves and never crossing them; pits at equal, regular spacing within each chain; chains running continuously to the rear, sharp end	MMX imaging	2027 on	Secondary impacts or drainage into fractures: irregular spacing, no relation to the ends of the body
N.4 Deimos surface	Ancient, shallow, worn hollows (median depth about 2% of width, against 5.5% on Phobos), the result of long grinding within the ring; same composition as Phobos	MMX flybys of Deimos	After 2027	Accreted rubble; smoothness from later regolith infill

Prediction	Expected (this paper)	Measured by	When	Accepted explanation expects
N.5 Pavonis caldera	The deepest caldera for its width of the four (depth/width ≈ 0.10); a mass deficit beneath it, the mark of the shallow reservoir that emptied	MOLA topography; gravity models (GMM-3)	Now	An ordinary summit caldera of a shield volcano
N.6 The two moons	The same Martian material in both, Deimos the wetter (richer in volatiles, and so lighter and more porous); the centre of mass of Phobos displaced toward Mars, toward its blunt end, by 300–500 m; a light, foamed rear (sharp) end	MMX samples, flybys and gravity field	2027–2031	Separate histories; no prediction for the centre of mass
N.7 Water beneath Tharsis	A water-saturated mid-crust beneath the whole of Tharsis, continuing what was found beneath InSight	Future seismic stations	Future	No link between moons and buried water
N.8 Martian dust	A component older than 4 billion years, from the material of the ring that fell, of the composition of the Phobos regolith, with grains that carry solar-wind helium and neon; most abundant in the broad equatorial band and the dust continents (Tharsis, Arabia Terra, Elysium), mixed with material ground by drought and wind	Returned Perseverance regolith samples; MMX	2031 on	Dust from weathering and wind erosion of surface material alone; no separate ancient component

Table 22. Predictions, instruments and dates of the tests, against the accepted explanation. MMX launches in October 2026, lands on Phobos in 2027 and returns its samples in 2031, with several flybys of Deimos (JAXA; Kuramoto et al. 2022). The full list of proposed measurements, mission by mission, is in Appendix B.

0 System Constraints and Boundary Conditions

The requirements and conditions on which the model rests, collected in one place for the whole paper:

1. **The principle of the pear:** every planet with a moon passed through a pear stage early in its history. On Mars, the Tharsis rise read as the pear lying down, from its base in Valles Marineris to its tip beside Pavonis (§C).
2. **A fast, expanded young Mars:** 1.5 times today's radius, hot and homogeneous (not yet sorted into layers), turning in 7.3 hours at the first release (Deimos) and 8.3 hours at the second (Phobos) (§B, E).
3. **A pear raised by the spin of the whole planet:** the core took part and moved toward it, and the pressure that passed through it supported it; it stood until the moons had left, high enough to put the vent of Pavonis 4,450 km above the young equator (§C.6, D; the length measured today, 2,996 km, is the hypotenuse).
4. **An ordinary volcano** at Pavonis: a shallow reservoir about 5 km down, a vent about as wide as the plug, an overpressure of 10–50 MPa or less; no deep chamber, no long conduit (§D.4–D.5).
5. **The loss of about 87% of the spin** through the corotation line, carried by water and gas: 6–11% of the mass of Mars, from a young Mars of which more than 30% was water, most of it in the first billion years (§E.2).
6. **A ring** from the line outward, built of steam and fine material before the releases, round and in the equatorial plane, which rounded both orbits and flattened their tilt from about 10° (§F.1).
7. **The leash:** the gaseous inner edge of the ring, at a density of a few 10^{-8} kg m^{-3} or more, lasting until Mars had slowed to a day of about 22 hours; its headwind balanced the push of the tide on Phobos and held it at the receding line until it stood near 19,000 km (§F.4–F.6).
8. **A soft young Mars:** dissipating tidal energy 8 to 20 times more than today while Deimos climbed, carrying it past 20,428 km before the line arrived there (§G).
9. **The turn of Phobos:** its heavy end led, away from Mars, as it left the vent, and faces Mars today; what turned it through 180° is left to a focused study (§K.6).
10. **The colour:** the dark, primitive-looking spectra of both moons, which give them the look of asteroids, are not a sign of a foreign origin; they must arise from Martian material altered by water and steam and by long exposure in space.
11. **Sensitivity:** a focused study may give approximate numbers somewhat different from those of this paper; the sequence does not depend on their exact values.

APPENDIX A

Computational Reproducibility

Every number of this paper is computed and printed by the script `mars2_numbers.py` (Python 3, standard library only), kept with the files of this paper.

Input values. $GM = 42{,}828$ km^3 s^{-2}; today's radius 3,389.5 km; day 24.6229 h; today's inertia coefficient 0.3644. The young Mars: radius $1.5\times$ today, homogeneous ($k = 0.4$), its figure a Maclaurin spheroid at each rate of spin (Chandrasekhar 1969).

The geometry of the pear. The pear, from its base at 294.70°E, 12.46°S to its tip at 243.86°E, 5.40°S, is read on the MOLA shape model MarsTopo719 expanded to degree 180 (Smith et al. 2001; Wieczorek 2015). Each candidate volcano stands, on the standing pear, at a height $\sqrt{s^2 - b^2}$ above the equator, where s is its distance from the eastern point of the base and b the half-width of the base (1,340 km today, read on MOLA across the axis of the pear, north edge at 0 km). On the young Mars east–west lengths scale with the equatorial radius a and north–south lengths with c^2/a of the Maclaurin spheroid at the day in question.

Release, tide and headwind. A body is set free at a vent with the sideways speed of the vent and, where stated, a push upward of $\sqrt{2\Delta p/\rho}$ with $\rho = 1{,}600$ kg m^{-3}; Phobos at release 1.13×10^{16} kg, Deimos 6.8×10^{15} kg. Tidal friction is $k_2/Q = 1.816 \times 10^{-3}$ today and 8–20 times more on the young Mars (§G), with $da/dt = 3(k_2/Q)(m/M)(R/a)^5 na$. The headwind at the ring's edge balances the tide on Phobos, $F = \frac{1}{2}\rho_{\mathrm{g}} A v^2$ with $v \approx 100$ m s^{-1}, where ρ_{g} is the gas density and A the cross-section of the moon. The spin removed through the line is carried at the angular momentum of a circular orbit at the receding line, from 9,080 to 20,428 km.

The topography and the shape models of Phobos and Deimos were analysed with separate scripts kept with the code.

APPENDIX B

Observational Protocols

The predictions of this paper can be tested with measurements that current and planned missions are designed to make. We list them by mission, so that each team can see what the account asks of its data. Each row gives what to measure, what the account expects, the result that would refute it, and what the accepted explanation expects.

#	Measure	Expected here	Refuted by	Accepted explanation
MMX (JAXA): Phobos landing 2027, sample return 2031, Deimos flybys				
B1	O, Cr and Ti isotopes; D/H of Phobos samples	On the Martian line alone	Values departing from Mars and matching a primitive asteroid	Capture: asteroid. Impact: Mars plus impactor
B2	Foreign component	None: pure Mars	A significant impactor component	Impact: a large one
B3	Zn and K isotopes	Unshifted: no evaporation and re-condensation	Light isotopes depleted	Impact: depleted
B4	Grain texture	Cooled volcanic foam: closed vesicles, hydrated minerals	Vapour-condensed glass spherules	Impact: melted and re-condensed
B5	Gravity field of Phobos	Centre of mass displaced toward Mars, toward the blunt end, by 300–500 m	Displaced toward the sharp end	No prediction

#	Measure	Expected here	Refuted by	Accepted explanation
B6	Imaging of the grooves	Concentrated on one side; bent with the body, straight on an elongated shape; pressed on the floor of Stickney; rims sharp or slightly worn; surface etched, not smoothed	Grooves on the floor of Stickney as sharp as outside	Tides or Stickney ejecta
B7	Imaging of the pit chains	Straight, beside the grooves and never crossing them; evenly spaced pits; chains ending at the rear, sharp end	Pit chains crossing grooves at random; irregular spacing	Secondary impacts
B8	Deimos flybys: shape and hollows	Shallow, worn hollows; composition as Phobos	A composition unlike Phobos	Accreted rubble; regolith infill
Mars orbiters (MRO, Mars Express, TGO)				
B9	Images east of each of the four Tharsis volcanoes, along their latitude	Chains of large blocks or craters: weak shots that fell back	—	No link to the volcanoes
B10	Gravity beneath the Pavonis caldera, against Arsia	A mass deficit beneath Pavonis, larger than beneath Arsia: the reservoir that fired	A mass excess beneath Pavonis; a larger deficit beneath Arsia (Arsia must then be re-examined)	An ordinary shield caldera
Returned regolith samples of Perseverance				
B11	Age of single dust grains	A component older than 4 billion years (the material of the ring that fell), mixed with locally ground material	—	Weathered material alone
B12	Solar-wind He and Ne in the grains	Ancient grains that carry them, from their time in space in the ring	No such grains	No such grains
B13	Composition of the ancient component	As the Phobos regolith	Unlike it	No prediction
Future seismic stations				

#	Measure	Expected here	Refuted by	Accepted explanation
B14	Mid-crust beneath Tharsis	Water-saturated, as beneath InSight	Dry	No link to the moons
B16	Position of the core	Displaced toward the pear, 255–263°E (§C.7)	Centred, or displaced elsewhere	No prediction
A future mission				
B15	A sample of Deimos	The same Martian fingerprint as Phobos	A different origin	Unrelated histories

Table B1. Proposed measurements and tests of refutation, mission by mission.

References

Anderson, R. C., Dohm, J. M., Golombek, M. P., et al. (2001). Primary centers and secondary concentrations of tectonic activity through time in the western hemisphere of Mars. *Journal of Geophysical Research: Planets*, 106, 20563–20585.

Baker, V. R., & Milton, D. J. (1974). Erosion by catastrophic floods on Mars and Earth. *Icarus*, 23, 27–41.

Banerdt, W. B., Golombek, M. P., & Tanaka, K. L. (1992). Stress and tectonics on Mars. In H. H. Kieffer et al. (Eds.), *Mars* (pp. 249–297). University of Arizona Press.

Basilevsky, A. T., Lorenz, C. A., Shingareva, T. V., Head, J. W., Ramsley, K. R., & Zubarev, A. E. (2014). The surface geology and geomorphology of Phobos. *Planetary and Space Science*, 102, 95–118.

Berger, J. A., Schmidt, M. E., Gellert, R., et al. (2016). A global Mars dust composition refined by the Alpha-Particle X-ray Spectrometer in Gale Crater. *Geophysical Research Letters*, 43, 67–75.

Black, B. A., & Mittal, T. (2015). The demise of Phobos and development of a Martian ring system. *Nature Geoscience*, 8, 913–917.

Burgisser, A., Poussineau, S., Arbaret, L., Druitt, T. H., Giachetti, T., & Bourdier, J.-L. (2010). Pre-explosive conduit conditions of the 1997 Vulcanian explosions at Soufrière Hills Volcano, Montserrat: I. Pressure and vesicularity distributions. *Journal of Volcanology and Geothermal Research*, 194, 27–41.

Canup, R., & Salmon, J. (2018). Origin of Phobos and Deimos by the impact of a Vesta-to-Ceres sized body with Mars. *Science Advances*, 4, eaar6887.

Cartan, É. (1924). Sur la stabilité ordinaire des ellipsoïdes de Jacobi. *Proceedings of the International Mathematical Congress, Toronto*, 2, 9–17.

Cashman, K. V., Thornber, C. R., & Pallister, J. S. (2008). From dome to dust: shallow crystallization and fragmentation of conduit magma during the 2004–2006 dome extrusion of Mount St. Helens, Washington. In *A Volcano Rekindled*, U.S. Geological Survey Professional Paper 1750, 387–413.

Castillo-Rogez, J. C., Matson, D. L., Sotin, C., Johnson, T. V., Lunine, J. I., & Thomas, P. C. (2007). Iapetus' geophysics: rotation rate, shape, and equatorial ridge. *Icarus*,

190, 179–202.

Chandrasekhar, S. (1969). *Ellipsoidal Figures of Equilibrium*. Yale University Press.

Cheng, B., Asphaug, E., Ballouz, R.-L., Yu, Y., & Baoyin, H. (2022). Numerical simulations of drainage grooves in response to extensional fracturing: testing the Phobos groove formation model. *The Planetary Science Journal*, 3, 249.

Clarke, A. B., Esposti Ongaro, T., & Belousov, A. (2015). Vulcanian eruptions. In H. Sigurdsson et al. (Eds.), *The Encyclopedia of Volcanoes* (2nd ed., pp. 505–518). Academic Press.

Crumpler, L. S., Head, J. W., & Aubele, J. C. (1996). Calderas on Mars: characteristics, structure, and associated flank deformation. *Geological Society, London, Special Publications*, 110, 307–348.

Ćuk, M., Minton, D. A., Pouplin, J. L. L., & Wishard, C. (2020). Evidence for a past Martian ring from the orbital inclination of Deimos. *The Astrophysical Journal Letters*, 896, L28.

Darwin, G. H. (1902). On the pear-shaped figure of equilibrium of a rotating mass of liquid. *Philosophical Transactions of the Royal Society A*, 198, 301–331.

Dauphas, N., & Pourmand, A. (2011). Hf–W–Th evidence for rapid growth of Mars and its status as a planetary embryo. *Nature*, 473, 489–492.

Ernst, C. M., Daly, R. T., Gaskell, R. W., et al. (2023). High-resolution shape models of Phobos and Deimos from stereophotoclinometry. *Earth, Planets and Space*, 75, 103.

Fagents, S. A., & Wilson, L. (1993). Explosive volcanic eruptions — VII. The ranges of pyroclasts ejected in transient volcanic explosions. *Geophysical Journal International*, 113, 359–370.

Garrick-Bethell, I., Wisdom, J., & Zuber, M. T. (2006). Evidence for a past high-eccentricity lunar orbit. *Science*, 313, 652–655.

Genova, A., Goossens, S., Lemoine, F. G., et al. (2016). Seasonal and static gravity field of Mars from MGS, Mars Odyssey and MRO radio science. *Icarus*, 272, 228–245.

Grott, M., Morschhauser, A., Breuer, D., & Hauber, E. (2011). Volcanic outgassing of CO_2 and H_2O on Mars. *Earth and Planetary Science Letters*, 308, 391–400.

Harrison, K. P., & Chapman, M. G. (2008). Evidence for ponding and catastrophic floods in central Valles Marineris, Mars. *Icarus*, 198, 351–364.

Hesselbrock, A. J., & Minton, D. A. (2017). An ongoing satellite–ring cycle of Mars and the origins of Phobos and Deimos. *Nature Geoscience*, 10, 266–269.

Hurford, T. A., Asphaug, E., Spitale, J. N., et al. (2016). Tidal disruption of Phobos as the cause of surface fractures. *Journal of Geophysical Research: Planets*, 121, 1054–1065.

Hyodo, R., Genda, H., Charnoz, S., & Rosenblatt, P. (2017). On the impact origin of Phobos and Deimos I: thermodynamic and physical aspects. *The Astrophysical Journal*, 845, 125.

IAU Working Group for Planetary System Nomenclature / USGS Astrogeology Science Center. *Gazetteer of Planetary Nomenclature*, entry “Pavonis Mons”. `https://planetarynames.wr.usgs.gov/`.

Iurato, G. (2014). The dawning of the theory of equilibrium figures: a brief historical account from the 17th through the 20th century. arXiv:1409.3858.

Jakosky, B. M., et al. (2018). Loss of the Martian atmosphere to space: present-day loss rates determined from MAVEN observations and integrated loss through time. *Icarus*, 315, 146–157.

JAXA. *Martian Moons eXploration (MMX)*, mission overview and schedule. `https://www.mmx.jaxa.jp/en/`.

Jeans, J. H. (1917). On the instability of the pear-shaped figure of equilibrium of a rotating mass of liquid. *Philosophical Transactions of the Royal Society A*, 217, 1–34.

Kokubo, E., & Genda, H. (2010). Formation of terrestrial planets from protoplanets under a realistic accretion condition. *The Astrophysical Journal Letters*, 714, L21–L25.

Kuramoto, K., Kawakatsu, Y., Fujimoto, M., et al. (2022). Martian Moons Exploration MMX: sample return mission to Phobos elucidating formation processes of habitable planets. *Earth, Planets and Space*, 74, 12.

Labrosse, S., Poirier, J.-P., & Le Mouël, J.-L. (2001). The age of the inner core. *Earth and Planetary Science Letters*, 190, 111–123.

Lainey, V., et al. (2021). Mars moon ephemerides after 14 years of Mars Express data. *Astronomy & Astrophysics*, 650, A64.

Lunine, J. I., Chambers, J., Morbidelli, A., & Leshin, L. A. (2003). The origin of water on Mars. *Icarus*, 165, 1–8.

Lyapunov, A. M. (1905). Sur un problème de Tchebychef. *Mémoires de l'Académie Impériale des Sciences de St.-Pétersbourg*, 17(3).

McCord, T. B., & Sotin, C. (2005). Ceres: evolution and current state. *Journal of Geophysical Research: Planets*, 110, E05009.

Miguel, Y., & Brunini, A. (2010). Planet formation: statistics of spin rates and obliquities of extrasolar planets. *Monthly Notices of the Royal Astronomical Society*, 406, 1935–1943.

Murray, J. B., & Heggie, D. C. (2014). Character and origin of Phobos' grooves. *Planetary and Space Science*, 102, 119–143.

NASA/JPL Solar System Dynamics. *Planetary Satellite Physical Parameters*. `https://ssd.jpl.nasa.gov/sats/phys_par/`.

Ojha, L., Lewis, K., Karunatillake, S., & Schmidt, M. (2018). The Medusae Fossae Formation as the single largest source of dust on Mars. *Nature Communications*, 9, 2867.

Ortiz, J. L., Santos-Sanz, P., Sicardy, B., et al. (2017). The size, shape, density and ring of the dwarf planet Haumea from a stellar occultation. *Nature*, 550, 219–223.

Park, R. S., Konopliv, A. S., Bills, B. G., et al. (2016). A partially differentiated interior for (1) Ceres deduced from its gravity field and shape. *Nature*, 537, 515–517.

Pätzold, M., Andert, T. P., Cardesín-Moinelo, A., Cicchetti, A., & Ziese, R. (2025). Investigations of the Moon Phobos by Mars Express and implications towards its origin. *Space Science Reviews*, 221, 41.

Phillips, R. J., Zuber, M. T., Solomon, S. C., et al. (2001). Ancient geodynamics and global-scale hydrology on Mars. *Science*, 291, 2587–2591.

Ramsley, K. R., & Head, J. W. (2019). Origin of Phobos grooves: testing the Stickney crater ejecta model. *Planetary and Space Science*, 165, 137–147.

Rosenblatt, P., Charnoz, S., Dunseath, K. M., et al. (2016). Accretion of Phobos and Deimos in an extended debris disc stirred by transient moons. *Nature Geoscience*, 9, 581–583.

Schmedemann, N., Michael, G. G., Ivanov, B. A., Murray, J. B., & Neukum, G. (2014). The age of Phobos and its largest crater, Stickney. *Planetary and Space Science*, 102, 152–163.

Smith, D. E., et al. (2001). Mars Orbiter Laser Altimeter: experiment summary after the first year of global mapping of Mars. *Journal of Geophysical Research*, 106, 23689–23722.

Thomas, P. (1979). Surface features of Phobos and Deimos. *Icarus*, 40, 223–243.

Thomas, P., Veverka, J., Bloom, A., & Duxbury, T. (1979). Grooves on Phobos: their distribution, morphology and possible origin. *Journal of Geophysical Research*, 84, 8457–8477.

Villanueva, G. L., Mumma, M. J., Novak, R. E., et al. (2015). Strong water isotopic anomalies in the martian atmosphere: probing current and ancient reservoirs. *Science*, 348, 218–221.

Weidenschilling, S. J. (1977). Aerodynamics of solid bodies in the solar nebula. *Monthly Notices of the Royal Astronomical Society*, 180, 57–70.

Wieczorek, M. A. (2015). Gravity and topography of the terrestrial planets. In *Treatise on Geophysics*, 2nd ed., vol. 10, Elsevier, 153–193.

Wright, V., Manga, M., & Panning, M. P. (2024). Liquid water in the Martian mid-crust. *Proceedings of the National Academy of Sciences*, 121, e2409983121.

Zhong, S., & Zuber, M. T. (2001). Degree-1 mantle convection and the crustal dichotomy on Mars. *Earth and Planetary Science Letters*, 189, 75–84.

Zuber, M. T., Smith, D. E., Phillips, R. J., et al. (1998). Shape of the northern hemisphere of Mars from the Mars Orbiter Laser Altimeter (MOLA). *Geophysical Research Letters*, 25, 4393–4396.

Declarations

Author: Zohar Duran, Independent Researcher, Ramla, Israel. ORCID 0009-0005-3194-9387.

Funding: This research received no external funding.

Conflicts of interest: The author declares no conflicts of interest.

Data availability: The script that reproduces every number in the paper and its output are provided as supplementary material. The public data used are cited in the text.

Use of AI tools: The author used AI assistants (Claude, Anthropic; ChatGPT, OpenAI;

Google’s AI search assistant) to write and run the computation script, to analyse the shape models, to prepare figures and tables, and for language editing. The account and its conclusions are the author’s. The author checked the content and takes full responsibility for it.