

Functional-coefficient models for nonstationary time series data<sup>1</sup>Zongwu Cai<sup>a,b,c</sup>, Qi Li<sup>d,e</sup>, Joon Y. Park<sup>d</sup><sup>a</sup> Department of Mathematics & Statistics, University of North Carolina at Charlotte, Charlotte, NC 28223, USA<sup>b</sup> Department of Economics, University of North Carolina at Charlotte, Charlotte, NC 28223, USA<sup>c</sup> Wang Yanan Institute for Studies in Economics, Xiamen University, Xiamen, Fujian 361005, China<sup>d</sup> Department of Economics, Texas A&M University, College Station, TX 77843, USA<sup>e</sup> Department of Economics, Tsinghua University, Beijing 100084, China

## article info

## Article history:

Available online 15 October 2008

## Keywords:

Nonstationary

Nonlinearity

Semiparametric estimation

## abstract

This paper studies functional coefficient regression models with nonstationary time series data, allowing also for stationary covariates. A local linear fitting scheme is developed to estimate the coefficient functions. The asymptotic distributions of the estimators are obtained, showing different convergence rates for the stationary and nonstationary covariates. A two-stage approach is proposed to achieve estimation optimality in the sense of minimizing the asymptotic mean squared error. When the coefficient function is a function of a nonstationary variable, the new findings are that the asymptotic bias of its nonparametric estimator is the same as the stationary covariate case but convergence rate differs, and further, the asymptotic distribution is a mixed normal, associated with the local time of a standard Brownian motion. The asymptotic behavior at boundaries is also investigated.

Published by Elsevier B.V.

## 1. Introduction

Nonparametric estimation techniques offer numerous advantages relative to parametric techniques, due mainly to their flexibility and robustness to functional form misspecification, and have been embraced by applied researchers in social, behavioral and economic sciences. Asymptotic theory underlying nonparametric estimators and test statistics for many commonly used models has been well established for independent and identically distributed (iid) data as well as for weakly dependence data. However, little is known about the behavior with nonstationary (in particular, integrated with order one, denoted by I(1)) data, which have predominately been modeled linearly. The early nonparametric asymptotic analyses with nonstationary data include Phillips and Park (1998), Park and Hahn (1999), Chang and Martinez-Chombo (2003) and

Juhl (2005). Phillips and Park (1998) and Juhl (2005) considered nonparametric estimation of regression models when the true data generating process is a linear unit root process, while the others considered the models linearized in the nonstationary variables. More recently, Wang and Phillips (forthcoming, 2008) considered nonparametric estimation of a regression model with an I(1) regressor and Xiao (forthcoming) considered a varying coefficient model with I(1) regressors appearing in the parametric component of the model. Finally, Karlsen et al. (2007) considered nonparametric estimation of a regression model for a different (a more general) type of nonstationary processes, a subclass of the class of null recurrent Markov chains.

In this paper, we tackle a more general set-up for a class of semiparametric models with non-stationary covariates. Specifically, we focus on the popular varying coefficient regression model with some nonstationary covariates

$$Y_t \in \mathbb{R}, Z_t \in \mathbb{R}^d, X_t \in \mathbb{R}^d, 1 \leq t \leq n; \quad (1.1)$$

where  $Y_t$ ,  $Z_t$  and  $X_t$  are scalar,  $X_t \in \mathbb{R}^d$  is a vector of covariates with dimension  $d$ ,  $\beta$  is a  $d \times 1$  column vector function, and the superscript T denotes transpose of a matrix. For ease notation, we assume that  $Z_t$  is univariate case. Extension to multivariate  $Z_t$  involves fundamentally no new ideas but

<sup>1</sup> We thank Peter Robinson for his constructive discussions and comments on an early version of this paper. We also thank an associate editor and two anonymous referees for their helpful comments that greatly improved the paper. Cai's research was supported, in part, by the National Science Foundation grant DMS-0404954 and funds provided by the University of North Carolina at Charlotte. Li's research was partially supported by the Private Enterprise Research Center, Texas A&M University. Park's research is partially supported by the National Science Foundation.

Corresponding author at: Department of Mathematics & Statistics, University of North Carolina at Charlotte, Charlotte, NC 28223, USA.

E-mail addresses: [zca@uncc.edu](mailto:zca@uncc.edu) (Z. Cai), [qi@econmail.tamu.edu](mailto:qi@econmail.tamu.edu) (Q. Li), [jpark@tamu.edu](mailto:jpark@tamu.edu) (J.Y. Park).

<sup>1</sup> The first version of this paper was written independently of these recent works on nonparametric estimation of regression models with non-stationary covariates.

complicated notations. We observe  $\{Y_t; X_t; Z_t\}$  for  $t = 1, \dots, n$ . When  $\{f_t; Z_t\}$  is stationary (denoted by  $I(0)$ ) or iid, various versions of (1.1) have been considered by many authors, including but not limited to, for example, [Chen and Tsay \(1993\)](#), [Hastie and Tibshirani \(1993\)](#), [Cai et al. \(2000\)](#), [Li et al. \(2002\)](#), and among others. When  $f_t$  is stationary and  $Z_t \in \mathbb{R}^d$ , Eq. (1.1) has been tackled by [Robinson \(1989, 1991\)](#), [Cai \(2007\)](#) and [Chen and Hong \(2007\)](#) for stationary  $X_t$ , by [Park and Hahn \(1999\)](#) and [Chang and Martinez-Chombo \(2003\)](#) for nonstationary  $X_t$ , and by [Cai and Wang \(2008\)](#) for nearly integrated  $X_t$ . When  $X_t \in \mathbb{R}^1$  and  $Z_t$  is  $I(1)$ , Eq. (1.1) becomes a standard univariate nonparametric regression model as considered by [Wang and Phillips \(forthcoming, 2008\)](#) and [Karlsen et al. \(2007\)](#). Finally, when  $Z_t$  is  $I(0)$  and  $X_t$  is  $I(1)$ , model (1.1) reduces to the case considered by [Xiao \(forthcoming\)](#).

The advantage of a varying coefficient model specification, compared with an unrestricted nonparametric regression, is that it attenuates the "curse of dimensionality" problem. It also includes many popular semiparametric models as special cases. For example, when  $X_t$  contains a constant, say the first component  $X_{t1} = 1$ , we can write  $X_t^T = (1; \beta_t^T)$ . Further, if the coefficient vector associated with  $\beta_t$  is a vector of constants, say  $\beta$ , then the varying coefficient model reduces to a partially linear model  $E(Y_t|X_t; Z_t) = 1 + \beta^T Z_t$ ; see, e.g., [Robinson \(1988\)](#).

The remainder of the paper is organized as follows: Section 2 discusses the case when  $Z_t$  is stationary. Here, local linear estimators of coefficient functions are developed, and their asymptotic properties are established. A two-step estimation procedure is also proposed when some covariates are nonstationary and the rest are stationary. Section 3 considers the case when  $Z_t$  is nonstationary. Nonparametric kernel smoothing of the coefficient functions is developed and its asymptotic behavior is investigated. Concluding remarks are presented in Section 4. Proofs of the main results of the paper are given in two Appendices.

## 2. Models with stationary $Z_t$

We consider first the case when some or all components of  $X_t$  are  $I(1)$  and  $Z_t$  is strictly stationary. For expositional simplicity, we re-express (1.1) as the following varying coefficient model

$$Y_t = \sum_{i=1}^d Z_{ti} \beta_i + \varepsilon_t, \quad \varepsilon_t \in \mathbb{R}^1, \quad \varepsilon_t \perp Z_t, \quad \varepsilon_t \perp X_t, \quad \varepsilon_t \perp \varepsilon_s, \quad t \neq s, \quad (2.1)$$

where  $X_{t1}$ ,  $Z_t$ , and  $\varepsilon_t$  are stationary,  $X_{t2}$  is an  $I(1)$  vector,  $\beta_i = \beta_i + \beta_i^T Z_t$ ,  $\beta_i^T = \beta_i^T + \beta_i^T Z_t$ , and  $X_t = (X_{t1}; X_{t2}^T)^T$ , where  $X_{ti}$  is a  $d_i = 1$  vector,  $i = 1, 2$ ,  $d_1 \in \mathbb{R}^1$ ,  $d_2 \in \mathbb{R}^d$ , and the first component of  $X_{t1}$  is identically one. In what follows, we assume that  $E(\varepsilon_t | X_t; Z_t) = 0$  which implies that  $X_t$  and  $Z_t$  are uncorrelated with  $\varepsilon_t$ . Note that  $Y_t$  is allowed to be stationary or nonstationary. For example, model (2.1) can be applied to the analysis of purchasing power of parity, in which  $X_{t2}^T = (P_t; P_t; E_t)$  (and no  $X_{t1}$ ), where  $P_t$  and  $P_t$  are the price levels of the domestic and a foreign country,  $E_t$  is the exchange rate between the domestic and the foreign currencies, and  $Z_t = (I_t - I_t)$  is the difference between the domestic interest rate  $I_t$  and the foreign interest rate  $I_t$ . Then if  $Y_t$  is an  $I(0)$  variable, we say that  $P_t$ ,  $P_t$  and  $E_t$  are co-integrated with a varying coefficient co-integration vector  $\beta_i$  which is a vector of smooth functions of  $Z_t$ . This setting is more general than the usual assumption that  $\beta_i$  is a vector of constant parameters in the usual purchasing power of parity analysis.

### 2.1. Local linear estimation

It is well known in the literature; see, e.g., [Fan and Gijbels \(1996\)](#), that a local linear fitting has several nice properties, over the classical Nadaraya–Watson (local constant) method, such as high statistical efficiency in an asymptotic minimax sense,

design-adaptation, and automatic edge correction. We estimate  $\beta_i$  using a local linear fitting from observations  $\{f_t; Z_t; Y_t\}_{t=1}^n$ . We assume throughout the paper that  $f_t$  is twice continuously differentiable, so that for any given grid point  $z$ , we use a local approximation as  $f_t(z) \approx f_t(z) + f_t'(z)(z - z)$  to approximate  $f_t(z)$ , where  $f_t'(z) = df_t/dz$ . Define

$$\hat{\beta}_i = \argmin_{\beta_i} \sum_{t=1}^n \left( Y_t - \sum_{i=1}^d \beta_i^T X_{ti} - \beta_i^T Z_t \right)^2 K_h(Z_t - z), \quad (2.2)$$

where  $K_h(u) = h^{-1}K(u/h)$ ,  $K(\cdot)$  is a kernel function satisfying Assumption A3 below,  $\hat{\beta}_i$  is  $\beta_i$  estimates  $\beta_i$ , and  $\hat{\beta}_i$  is  $\beta_i$  estimates  $\beta_i$ . Then,  $\hat{\beta}_i$  and  $\hat{\beta}_i$  can be expressed as

$$\hat{\beta}_i = \left( \sum_{t=1}^n X_{ti} X_{ti}^T K_h(Z_t - z) \right)^{-1} \sum_{t=1}^n X_{ti} Y_t K_h(Z_t - z), \quad (2.3)$$

where  $A^{-2} = (A^{-1})^T A$  for a vector or matrix  $A$ .

### 2.2. Notations and assumptions

Since  $X_{t2}$  is a vector of  $I(1)$  processes, it can be re-expressed as  $X_{t2} = X_{t2} + \sum_{s=1}^t \varepsilon_{s2}$ , where  $\varepsilon_{s2}$  is an  $I(0)$  process with mean zero and variance  $\sigma^2$ . Then,

$$\frac{1}{n} \sum_{t=1}^n X_{t2} X_{t2}^T = \frac{1}{n} \sum_{t=1}^n X_{t2} X_{t2}^T + \frac{1}{n} \sum_{t=1}^n \varepsilon_{t2} \varepsilon_{t2}^T + \frac{1}{n} \sum_{t=1}^n X_{t2} \varepsilon_{t2}^T + \frac{1}{n} \sum_{t=1}^n \varepsilon_{t2} X_{t2}^T,$$

where  $r = t = n$  and  $|x|$  denotes the integer part of  $x$ . Under some regularity conditions, Donsker's theorem; see, for example, Theorems 14.1 and 19.2 in [Billingsley \(1999\)](#) for iid  $\varepsilon_t$  and  $\varepsilon_t$ -mixing  $\varepsilon_t$ , respectively, generalizes in an obvious way to the multivariate cases and leads to

$$\frac{1}{n} \sum_{t=1}^n X_{t2} X_{t2}^T \Rightarrow W_{22} + \int_0^1 W_2 dW_2, \quad (2.4)$$

where  $W_{22}$  is a  $d_2$ -dimensional Brownian motion on  $[0, 1]$  with covariance matrix  $\sigma^2 I_{d_2}$  and  $\Rightarrow$  represents weak convergence. In particular, it follows from [Merlevède et al. \(2006\)](#) that (2.4) holds if  $\varepsilon_t$  is a stationary strong  $(-)$ -mixing sequence satisfying, for some  $\alpha > 0$ ,

$$E|j|^{2C} < 1; \quad \text{and} \quad \sum_{k=1}^{\infty} k^{2C} \alpha^k < 1; \quad (2.5)$$

where  $\alpha$  is the mixing coefficient; see, e.g., [Hall and Heyde \(1980\)](#) for more discussion on  $\alpha$ -mixing process. Also, for any Borel measurable and totally Lebesgue integrable function  $f$ , one has

$$\frac{1}{n} \sum_{t=1}^n f(X_{t2}) \Rightarrow \int_0^1 f(W_2) dW_2, \quad \text{as } n \rightarrow \infty;$$

where  $\Rightarrow$  denotes the convergence in distribution, so that, for  $I \in \mathbb{R}^1, 2$ ,

$$\frac{1}{n} \sum_{t=1}^n X_{t2} X_{t2}^T \Rightarrow \int_0^1 W_2 W_2^T dr = \int_0^1 W_2 W_2^T dr, \quad \text{as } n \rightarrow \infty. \quad (2.6)$$

see Theorem 1.2 in [Berkes and Horváth \(2006\)](#) for details. Under stronger regularity conditions, (2.4) can be strengthened to the following strong approximation result

$$\sup_{0 \leq r \leq 1} \|X_{t2} - \frac{1}{n} \sum_{t=1}^n W_{22}(r) + \int_0^r W_2 dW_2\| \leq O_p(n^{-1/2} \log n). \quad (2.7)$$

almost surely, where  $\|\cdot\|_k$  is the usual  $L_2$  norm in  $\mathbb{R}^{d_2}$ ,  $D_{1/2} = 2C_1/1 = 2C_1$  and  $D_{1/2} > 0$  is a function of  $\rho$ , provided that  $f_{t|g}$  is a stationary strong mixing sequence satisfying that, for some  $\alpha > 2C_1$  with  $0 < \alpha < 2$ ,

$$E|f_{t|g} - f_{t|g}| < 1; \quad \text{and} \quad \sum_{n \geq 1} n^{\alpha/2} C_1^{1-\alpha} < 1 \quad (2.8)$$

see Theorem 4.1 in Shao and Lu (1987) and Einmahl (1987) for details. It is not hard to see that assumption (2.8) is stronger than (2.5). This is not surprising, since strong approximation in (2.7) usually requires stronger assumptions than weak convergence as in (2.4). Finally, note that if  $f_{t|g}$  is iid and has a finite  $\alpha$ -th moment ( $\alpha > 2$ ), then the right hand side of (2.7) becomes  $O(n^{-\alpha/2})$ , where  $D_{1/2} = 2C_1/1 = 2C_1$ ; see, e.g., Csörgő and Révész (1981, p. 107). We assume throughout the paper that the sequence  $f_{t|g}$  is stationary  $\alpha$ -mixing and satisfies either (2.5) or (2.8). Note that either (2.5) or (2.8) ensures that  $\lim_{n \rightarrow \infty} \text{Var} \cdot n^{-1/2} \sum_{t=1}^n f_{t|g}$  exists and is finite by Davydov's inequality for an  $\alpha$ -mixing process; see, e.g., Corollary A.2 in Hall and Heyde (1980).

Next, we give regularity conditions for the asymptotic distribution of  $b_{1,z}/D_{1/2}$ . We introduce the following notations. Let  $f_{z,z}$  denote the marginal density of  $Z_t$ . Define  $M_{k,z}/D_{1/2} = E \cdot X_{t1}^k \cdot Z_t \cdot D_{1/2} z$  for  $k \in \{1, 2\}$ . Also set, for  $j = 0, 1, 2$ ,  $K_j/D_{1/2} = \int K_j \cdot v \cdot dv$ , and  $J_j \cdot K/D_{1/2} = \int J_j \cdot K^2 \cdot v \cdot dv$ . Further, let

$$S_{z,z}/D_{1/2} = \frac{M_{2,z}/D_{1/2}}{W_{1/2} \cdot M_{1,z}/D_{1/2}} \cdot \frac{M_{1,z}/D_{1/2}}{W_{1/2} \cdot M_{1,z}/D_{1/2}}; \quad (2.9)$$

where  $W_{1/2}$  is defined in (2.6). We make the following assumptions.

#### Assumptions:

- A1.  $f_{z,z}$  is twice continuously differentiable in  $z$  for all  $z \in \mathbb{R}$ .
- A2.  $M_{k,z}/D_{1/2}$  is positive-definite and continuous in a neighborhood of  $z$ .  $f_{z,z}$  is continuously differentiable in a neighborhood of  $z$  and  $f_{z,z} > 0$ .
- A3. The kernel function  $K \cdot /$  is a symmetric and continuous density function, supported by  $[-1, 1]$ .
- A4. The bandwidth  $h$  satisfies  $h \rightarrow 0$  and  $nh \rightarrow \infty$ .
- A5.  $f_{t|g}$  has a finite fourth moment,  $E \cdot f_{t|g}^4 \cdot X_{t1}^2 \cdot Z_t / D_{1/2}^2$  is a positive constant.
- A6.  $f_{t|g}$  is a strictly  $\alpha$ -mixing stationary process with the  $\alpha$ -th moment ( $\alpha > 2$ ).  $E \cdot f_{t|g}^2 \cdot X_{t1}^2 \cdot Z_t^2 / D_{1/2}^2 < \infty$  with  $\alpha > 1$  and  $t/D_{1/2} \rightarrow 0$  for some  $\alpha > \min\{2, 1/\alpha\}$ ;  $2 \cdot \alpha > 2 \cdot \alpha/2$ , where  $\alpha > 4 \cdot 1 = 4 \cdot 1 = 4$  for some  $\alpha$  satisfying  $1 = 1/4 < \alpha < 2$ . Also,  $k \cdot t \cdot k_0 \cdot D_{1/2} [E \cdot f_{t|g}^2]^{1/2} < 1$  with  $q_0 \in [4, 6]$  for some  $1 < \alpha < 4$ . Further,  $\sup_k E \cdot f_{t|g}^2 \cdot Z_{kC1} / D_{1/2} < 1$ .
- A7.  $f_{z,z}$  is continuous at  $x_0$  and  $f_{z,z} > 0$  for  $s = 1$ , where  $f_{z,z}$  is the conditional density of  $Z_0$  given  $X_0 = x_0$ ;  $f_{z,z}$  is the conditional density of  $Z_0$  given  $X_0 = x_0$ ;  $f_{z,z}$  is the conditional density of  $Z_0$  given  $X_0 = x_0$ .
- A8.  $n^{1/2} \cdot h^{1/2} \cdot D_{1/2} > 0$ .

We give some comments on the above conditions. Assumptions A1 and A2 are smoothness conditions. The requirement in Assumption A3 that  $K \cdot /$  be compactly supported is imposed for the sake of brevity of proofs, and can be removed at the cost of lengthier arguments.  $\alpha$ -mixing is one of the weakest mixing conditions for weakly dependent stochastic processes. Stationary linear and nonlinear time series or Markov chains fulfilling certain (mild) conditions are  $\alpha$ -mixing with exponentially decaying coefficients; see discussions and examples in Cai (2002a), Carrasco and Chen (2002) and Chen and Tang (2005). The conditional homoscedastic error Assumption in A5 can be relaxed to allow for conditional heteroscedasticity of the form  $E \cdot f_{t|g}^2 \cdot X_{t1}^2 \cdot Z_t / D_{1/2}$

$\cdot X_{t1}^2 \cdot Z_t / D_{1/2}$ , i.e., the conditional variance is only a function of the stationary covariates  $X_{t1} \cdot Z_t / D_{1/2}$ . However, it is technically difficult to let it also be a function of the nonstationary covariate  $X_{t2}$ . If  $\cdot /$  decays geometrically, then Assumption A6 is fulfilled with some standard moment conditions. Assumption A7 is a standard technical assumption. Clearly, Assumption A8 allows for choosing a wide range of  $h$  and is slightly stronger than the usual condition  $nh \rightarrow \infty$ . For optimal bandwidths selection (i.e.,  $h \rightarrow 0$  for  $0 < \alpha < 1$ ,  $\alpha > 0$ ), A8 is automatically satisfied for  $\alpha > 2 \cdot 1 \cdot C_1/1 = 2 \cdot 1 \cdot C_1/1$  and it is still fulfilled for  $2 < \alpha < 2 \cdot 1 \cdot C_1/1 = 2 \cdot 1 \cdot C_1/1$  if  $\alpha$  satisfies  $\alpha < 2 \cdot 4 \cdot 1 = 2 \cdot 1 \cdot C_1/1 = 2 \cdot 1 \cdot C_1/1$ . Conditions similar to Assumptions A6–A8 are also imposed by Cai et al. (2000) for the stationary data case.

#### 2.3. Asymptotic properties

To establish the asymptotic property of  $b_{1,z}/D_{1/2}$ , we define  $D_n \cdot D_{1/2}$   $\text{diag}\{I_{d_1}, I_{d_2}\}$ , and  $B \cdot z/D_{1/2} \cdot K \cdot / \cdot z/D_{1/2}$ , where  $I_d$  is a  $d \cdot d$  identity matrix. Detailed proof of the following Theorem is provided in Appendix A.

**Theorem 2.1.** Under Assumptions A1–A8, we have

$$\sqrt{nh} \cdot b_{1,z}/D_{1/2} \rightarrow_d N(0, h^2 \cdot B \cdot z/D_{1/2} \cdot MN \cdot z/D_{1/2});$$

where  $MN \cdot z/D_{1/2}$  is a mixed normal distribution with mean zero and conditional covariance matrix given by  $z/D_{1/2} \cdot D_{1/2} \cdot K \cdot S \cdot z/D_{1/2} = f_{z,z}$ .

Here, a mixed normal distribution is defined as follows. Conditional on the random variable that appears at the asymptotic variance, the estimator has an asymptotic normal distribution, see Phillips (1989) and Phillips and Park (1998) for a formal definition of a mixed normal distribution. Clearly, if there is no nonstationary covariate (i.e., removing  $X_{t2}$ ), then Theorem 2.1 reduces to

$$\sqrt{nh} \cdot b_{1,z}/D_{1/2} \rightarrow_d N(0, \frac{1}{2} \cdot f_{z,z} \cdot h^2 \cdot K \cdot / \cdot z/D_{1/2}); \quad (2.10)$$

where  $f_{z,z}/D_{1/2} = K/M_{2,z}/D_{1/2} = f_{z,z}$ , which is non-stochastic and is exactly the same as that in Cai et al. (2000). Further, from Theorem 2.1, we see that  $\text{Var} \cdot b_{1,z}/D_{1/2}$  has a faster rate of convergence than that of  $\text{Var} \cdot b_{2,z}/D_{1/2}$  due to the fact that  $X_{t2}$  is nonstationary ( $\sum_{t=1}^n X_{t2}^2 \cdot D_{1/2} = O(n^2)$  rather than  $O(n)$ ). This is similar to the linear model case. However, the local fitting method renders the asymptotic variance of  $b_{2,z}/D_{1/2}$  to be of the order  $O(n^2 h^{-1})$  rather than the linear model case of  $O(n^2)$ . One can easily derive the integrated asymptotic mean squared error (IAMSE) for  $b_{j,z}/D_{1/2}$

$$\begin{aligned} \text{IAMSE}_j &= \int \text{IAMSE}_j \cdot b_{j,z}/D_{1/2} \\ &= \int \frac{h^4}{4} \cdot \frac{K \cdot K}{2} \cdot \frac{z/D_{1/2}}{j \cdot z/D_{1/2}} \cdot \frac{\text{tr} \cdot z/D_{1/2}}{n \cdot h} \cdot f_{z,z} \cdot dz \end{aligned}$$

for  $j \in \{1, 2\}$ , where  $f_{z,z}$  is a non-negative weight function,  $f_{z,z}$  is the upper-left corner sub-matrix of  $f_{z,z}$ , and  $f_{z,z}$  is the lower-right corner sub-matrix of  $f_{z,z}$ . By minimizing  $\text{IAMSE}_j$  with respect to  $h$ , we obtain the optimal bandwidth for  $b_{j,z}/D_{1/2}$ , which is

$$h_{j,\text{opt}} = \left( \frac{\text{tr} \cdot z/D_{1/2}}{\int \frac{K \cdot K}{2} \cdot \frac{z/D_{1/2}}{j \cdot z/D_{1/2}} \cdot f_{z,z} \cdot dz} \right)^{1/5} \cdot n^{-1/5}; \quad (2.11)$$

With the above choice of  $h_{j,\text{opt}}$ , we see that  $\text{IAMSE}_j$  has an order of  $O(n^{-4/5})$ . Hence,  $\text{IAMSE}_2$  for  $b_{2,z}/D_{1/2}$  has an order  $O(n^{-8/5})$  which is

smaller than  $\text{IAMSE}_1 \propto O(n^{-4/5})$ . Clearly, a single value of  $h$  can not make the estimation of both  $\beta_1(\cdot)$  and  $\beta_2(\cdot)$  optimal (in the sense of minimizing their asymptotic MSE). Therefore, to minimize the asymptotic MSE for each coefficient (function) estimate, an iterative estimation approach is needed. This issue is addressed next.

#### 2.4. A two-step estimation procedure

As discussed in Section 2.3, the one-step estimation procedure based on (2.2) cannot minimize the asymptotic MSE for both  $\beta_1(\cdot)$  and  $\beta_2(\cdot)$ . Therefore, we suggest a two-stage estimation procedure below. The idea is similar to the profile likelihood method; see, e.g., Cai (2002b,c) and Fan and Huang (2005), and is described as follows. If the bandwidth is taken to be of the order  $n^{-2/5}$ ,  $\hat{\beta}_{2,n}(\cdot)$  reaches the optimal convergent rate but  $\hat{\beta}_{1,n}(\cdot)$  is under-smoothed. Therefore, the first step is to estimate  $\beta_2(\cdot)$  with a value of  $h$  that is optimal for estimating  $\beta_2(\cdot)$ . We use  $h_2$  to denote it. For example, we can choose  $h_2 \propto c_2 n^{-2/5}$  for some positive constant  $c_2$ . We know that the resulting estimator  $\hat{\beta}_{2,n}(\cdot)$  has an optimal convergence rate as  $\hat{\beta}_{2,n}(\cdot) \propto O_p(n^{-4/5})$ , and the asymptotic distribution of  $\hat{\beta}_{2,n}(\cdot)$  is given by

$$\sqrt{n^2 h_2} (\hat{\beta}_{2,n}(\cdot) - \beta_2(\cdot)) \xrightarrow{d} MN(0, \Sigma_{\beta_2}(\cdot)); \quad (2.12)$$

where  $\Sigma_{\beta_2}(\cdot)$  is previously defined as the lower-right corner  $d_2 \times d_2$  sub-matrix of  $\Sigma_{\beta}(\cdot)$ .

However, the corresponding estimator of  $\beta_1(\cdot)$  is not optimal with the choice of  $h_2 \propto c_2 n^{-2/5}$  ( $c_2 > 0$ ). Therefore, we suggest that at the second step, one should re-estimate  $\beta_1(\cdot)$  with  $\hat{\beta}_{2,n}(\cdot)$  replaced by  $\hat{\beta}_{2,n}(\cdot)$  obtained at the first step. That is, we replace  $\beta_2(\cdot)$  by  $\hat{\beta}_{2,n}(\cdot)$  in (2.1) to obtain

$$Y_t = \beta_1(\cdot) X_{t1} + \hat{\beta}_{2,n}(\cdot) X_{t2} + \varepsilon_t; \quad (2.13)$$

where  $\varepsilon_t \propto d_2 \times d_2$  CT  $\beta_2(\cdot)$ . Then, we can apply the local linear method to estimate  $\beta_1(\cdot)$ . Let  $\hat{\beta}_{1,2\text{step}}(\cdot)$ ,  $\hat{\beta}_{1,2\text{step}}^{(1)}(\cdot)$  represent the resulting estimator of  $\beta_1(\cdot)$ ,  $\beta_1^{(1)}(\cdot)$ , which is given by

$$\begin{aligned} \hat{\beta}_{1,2\text{step}}(\cdot) &= \frac{\sum_{t \in D_1} X_{t1} Y_t}{\sum_{t \in D_1} X_{t1}^2} \\ \hat{\beta}_{1,2\text{step}}^{(1)}(\cdot) &= \frac{\sum_{t \in D_1} X_{t1} Y_t}{\sum_{t \in D_1} X_{t1}} \end{aligned} \quad (2.14)$$

where  $h_1$  is the bandwidth used at this step for estimating  $\beta_1(\cdot)$ . We will show in Theorem 2.2 below that the asymptotic distribution of  $\hat{\beta}_{1,2\text{step}}(\cdot)$  is the same as the case when  $\beta_2(\cdot)$  were known; that is  $\hat{\beta}_{1,2\text{step}}(\cdot) \propto O_p(n^{-1/2})$  if  $h_1 \propto c_1 n^{-1/5}$  ( $c_1 > 0$ ) and  $h_2$  is as small as possible (see Theorem 2.2 later). Since the right hand side of (2.13) involves only  $\beta_1(\cdot)$ , one can use any data-driven method to select  $h_1$  optimally, such as the nonparametric version of the Akaike information criterion type as in Hurvich et al. (1998) and Cai (2002b,c) or the plug-in method as in Ruppert et al. (1995).

It is clear from (2.12) that  $\hat{\beta}_{2,n}(\cdot) \propto O_p(n^{-4/5})$  if  $h_2 \propto c_2 n^{-2/5}$ , and this pointwise convergence rate is optimal. To establish the asymptotic normality of the estimator given in (2.14), we might need a uniform convergence rate. Therefore, it is assumed that the initial estimator satisfies the following condition

$$\sup_{z \in D} \|\hat{\beta}_{2,n}(\cdot) - \beta_2(\cdot)\| \propto O_p(a_n); \quad (2.15)$$

where  $D$  is a compact support of  $f_z(\cdot)$  (by assuming that  $f_z(\cdot)$  has a compact support  $D$ ) and  $a_n$  satisfies  $a_n \rightarrow 0$  with a certain convergence rate. Note that under some regularity conditions (see, e.g., the assumptions of Theorem 2 in Hansen (2008)), and by following the same arguments as in Hansen (2008), one can show<sup>2</sup> that (2.15) holds with  $a_n \propto n^{-4/5} \log n$ . Alternatively, an assumption similar to (2.15) is also imposed in Linton (2000) for iid samples, and Cai (2002c) for time series data to simplify the proof of the asymptotic results of a two-stage estimator. Now, the asymptotic normality for the proposed two-stage estimator is stated here and its proof is relegated to the Appendix.

**Theorem 2.2.** Under assumption A1–A8 and (2.15), if  $h_2 \propto c_2 n^{-2/5}$ , we have

$$\sqrt{n h_1} (\hat{\beta}_{1,2\text{step}}(\cdot) - \beta_1(\cdot)) \xrightarrow{d} MN(0, \Sigma_{\beta_1}(\cdot));$$

where  $\Sigma_{\beta_1}(\cdot)$  is defined in (2.10).

**Remark 2.1.** Note that the consequences of (2.12) and Theorem 2.2 are that the convergence rates are optimal for estimating each coefficient function at each step. That is, at the first step, the optimal rate is obtained for estimating coefficient functions of non-stationary covariates and the second step is devoted to obtaining the optimal rate for estimating coefficient functions of the stationary covariates. Also, note that the result in Theorem 2.2 is exactly the same as that (see (2.10)) in Cai et al. (2000) for stationary case, which implies that  $\hat{\beta}_{1,2\text{step}}(\cdot)$  is “oracle” in the sense that its asymptotic distribution is the same as the case with a known  $\beta_2(\cdot)$ . Finally, we note that the order of  $h_2$  should be smaller than its optimal value, which means we need to undersmooth the estimate of  $\beta_2(\cdot)$  at the first step. This is a common phenomenon for a two-stage estimation method; see Cai (2002b,c). In practical implementation, we refer to the papers by Cai (2002b,c) for choosing the data-driven fashion bandwidths for a two-stage estimation.

### 3. Models with nonstationary $Z_t$

When  $Z_t$  is a nonstationary I(1) regressor, the asymptotic analysis is much involved. Therefore, we consider only the case that  $X_t$  is stationary in this section. The model is the same as given in (1.1) but now  $Z_t$  is nonstationary; that is,

$$Y_t = \beta_1(\cdot) X_{t1} + \beta_2(\cdot) Z_t + \varepsilon_t; \quad (3.1)$$

where  $X_t$  is a  $d-1$  vector of stationary variables, and  $Z_t$  is a univariate I(1) nonstationary variable. When  $X_t \propto 1$ , model (3.1) was considered by Karlsen et al. (2007) for the case that  $Z_t$  is a nonstationary process from a subclass of the class of null recurrent Markov chains, and by Wang and Phillips (forthcoming, 2008) for the case that  $Z_t$  is I(1), while all of them used the local constant fitting approach to estimate the nonparametric regression function.

We assume that  $\beta_2(\cdot)$  is twice continuously differentiable. Exactly similar to (2.3), we obtain the local linear estimator of  $\beta_2(\cdot)$ , given by

$$\begin{aligned} \hat{\beta}_{2,n}(\cdot) &= \frac{\sum_{t \in D_1} X_{t1} Y_t}{\sum_{t \in D_1} X_{t1}^2} \\ \hat{\beta}_{2,n}^{(1)}(\cdot) &= \frac{\sum_{t \in D_1} X_{t1} Y_t}{\sum_{t \in D_1} X_{t1}} \end{aligned} \quad (3.2)$$

<sup>2</sup> The detailed proof of the uniform convergence result as stated in (2.15) is available from the authors upon request.



<sup>3</sup> Recall that  $Z_t \in \prod_{s \in D_1} u_s \subset Z_0$ , consider the simple case that  $Z_0 \subset \mathbb{R}^n$  and  $u_t$  is iid normal  $N(0, \frac{1}{n}I_n)$ , then  $Z_t \sim N(0, \frac{1}{n}I_n)$ . Here, we consider the case that  $X_t$  and  $Z_t$  are jointly normally distributed.

**Remark 3.1.** The asymptotic properties for  $b_{\cdot}^{1/2} z/\cdot$  can be obtained in a same way as that in Theorem 2.1 and they are omitted. By comparing the results in Theorems 2.1 and 3.1, our new findings are as follows: It is clear that  $h^2 B_{\cdot} z/\cdot$  serves as the asymptotic bias, which is exactly the same as that for stationary case when one uses a local linear estimation method; see Theorem 2.1. This is not surprising since the asymptotic bias term comes from the local linear approximation. However, the asymptotic variance of  $b_{\cdot} z/\cdot$  is of the order  $O_{\cdot} \cdot n^{1/2} h^{-1/2}$ , which is larger than  $O_{\cdot} \cdot n h^{-1/2}$  for the stationary  $Z_t$  case as presented in Theorem 2.2. The integrated asymptotic mean squared error is given by

$$\text{IAMSE} \approx \text{tr} \left( \frac{1}{n} \int_0^1 h^{-1} C \frac{h^4}{4} \frac{2}{2} K \frac{1}{2} z/\cdot^2 M_{\cdot} z/dz \right)$$

Minimizing the IAMSE with respect to  $h$  gives the optimal bandwidth  $h_{\text{opt}} \approx c n^{-1/10}$  for some  $c > 0$ , which is much larger than that for the stationary case; see (2.11).

Now, we consider the asymptotic behavior of  $b_{\cdot} z/\cdot$  at boundaries. When  $Z_t$  is  $I(1)$ , it follows from (3.7) that when  $z \rightarrow a^-$  and  $r \rightarrow d$ ,  $t \rightarrow n$ ,

$$P_{\cdot} Z_t z/\cdot \approx P_{\cdot} Z_t a^- \frac{P_{\cdot} n}{n!} P_{\cdot} W_{n,r}/a^- \approx 1 \cdot a^- \frac{P_{\cdot} n}{r} u/\cdot > 0;$$

where  $\cdot/\cdot$  is the distribution function of the standard normal random variable. This means that there is a great chance for  $|Z_t|$  taking large values. Now the question is how the asymptotic behavior of the estimator looks like when  $z$  is large like  $z \rightarrow a^-$  for any fixed  $a$ . We offer the following asymptotic results at boundary  $z \rightarrow a^-$  for any fixed  $a$ . However, we do not provide the detailed proofs since they follow exactly the same arguments as those used in proving Theorem 3.1.

**Theorem 3.2 (Boundary Behavior).** If Assumptions C1–C5 hold and  $C_{\cdot}/\cdot$  in assumption C3 is bounded as well as  $n^{1/4} h^{5/2} \cdot a^- \frac{P_{\cdot} n}{n!} \rightarrow 0$  for any  $a$ , then, we have

$$P_{\cdot} \frac{b_{\cdot} z/\cdot}{n^{1/2} h} \approx a^- \frac{P_{\cdot} n}{n!} \cdot a^- \frac{P_{\cdot} n}{n!} h^2 B_{\cdot} a^- \frac{P_{\cdot} n}{n!} \rightarrow MN_{\cdot} \cdot a^- \frac{P_{\cdot} n}{n!};$$

where  $MN_{\cdot} \cdot a^- \frac{P_{\cdot} n}{n!}$  is a mixed normal distribution with mean zero and conditional covariance  $\frac{1}{n} \int_0^1 a^- \frac{P_{\cdot} n}{n!} \frac{2}{2} u/\cdot K/E_{\cdot} X_t X_t^T/L \cdot 1; a^- \frac{P_{\cdot} n}{n!}$ .

**Remark 3.2.** Comparing Theorem 3.2 with Theorem 3.1, we observe that the asymptotic variance of  $b_{\cdot} z/\cdot$  at the boundary point differs from that at the interior point. This is different from its stationary counterpart; see Fan and Gijbels (1996) for the stationary case.

#### 4. Discussion

In this paper, we studied the class of varying coefficient models with nonstationary time series data. We suggested using the local linear fitting scheme to estimate the nonparametric coefficient functions and derived the asymptotic properties of the proposed estimators. We would like to mention some interesting future research topics related to this paper. First, it would be very useful and important to discuss how to select data-driven (optimal) bandwidths theoretically and empirically. Secondly, an important extension would be to generalize the asymptotic analysis of this paper to the case where both  $Z_t$  and (or some of the components)  $X_t$  are nonstationary. Further, we conjecture that if some of  $X_t$  in (3.1) are the lagged variables, one might find some regularity conditions to show that  $Y_t$  generated by (3.1) is ergodic, so that it is stationary if it is assumed to be Markovian. We are currently exploring these issues. Finally, it is warranted to consider an extension to other types of nonstationarity such as nearly integrated processes; see, e.g., Torous et al. (2004), Campbell and Yogo (2006) and Polk et al. (2006), which has a potential application in finance, which is under investigation by Cai and Wang (2008).

#### Appendix A. Proofs of Theorems 2.1 and 2.2

In the remaining part of this paper, we denote by  $C$  a generic positive constant, which may take different values at different places. We often need to evaluate (probability) orders of some finite dimensional matrix random variables. Let  $M_t$  be a matrix of finite dimension, and  $b_n$  be a sequence of non-stochastic real numbers, we write  $jM_t \rightarrow O_p \cdot b_n$  to mean that  $jM_{t,ij} \rightarrow O_p \cdot b_n$  for each  $i$  and  $j$ , where  $M_{t,ij}$  is the  $(i, j)$ th component of  $M_t$ . Similarly,  $EjM_t \rightarrow O \cdot b_n$  means that  $EjM_{t,ij} \rightarrow O \cdot b_n$  for all  $i$  and  $j$ ; and  $\sup_t jM_t \rightarrow O \cdot b_n$  means that  $\sup_t jM_{t,ij} \rightarrow O \cdot b_n$  for all  $i$  and  $j$ . Also, we write  $M_t^2 \rightarrow O_p \cdot b_n$  to mean that  $M_{t,ij}^2 \rightarrow O_p \cdot b_n$  for all  $i$  and  $j$ . Finally, we use  $M^{k \times m}$  to denote the real-valued  $k \times m$  matrices and  $\mathcal{D}[0, 1]$  to represent the space of right-continuous with left limits (cadlag) functions on  $[0, 1]$  equipped with the Skorohod metric (as defined in Billingsley (1999, p. 124)).

Before we prove Theorems 2.1 and 2.2, we first give a few lemmas that will be used frequently in the proofs below. First, consider random arrays  $\{U_{nt}, Y_{nt}\}_{n=1}^{\infty}$  where  $U_{nt}$  is a  $k \times m$  matrix and  $Y_{nt}$  is an  $m \times 1$  vector. We transform these arrays into random elements on  $\mathcal{D}[0, 1]$  by  $U_n \cdot s/\cdot \rightarrow U_{n\cdot} s/\cdot$  and  $Y_n \cdot s/\cdot \rightarrow Y_{n\cdot} s/\cdot$  for  $0 \leq s \leq 1$ . Also, define  $W_{nt} = Y_{nt} - Y_{n,t-1}$ . We can then define the stochastic integral

$$Z_s = \int_0^s U_n \cdot r/dY_n \cdot r/\cdot = \int_0^s U_{n\cdot} r/dY_{n\cdot} r/\cdot$$

Let  $BM_{\cdot} z/\cdot$  denote the vector Brownian motion with covariance matrix  $\cdot$ . Finally, we define  $W_t \rightarrow \int_0^t W_{n\cdot} s/\cdot$ , where  $W_{n\cdot} s/\cdot$  satisfies the following assumption:

**Assumption M1:** For some  $p > 2$ ,  $W_{n\cdot} s/\cdot$  is a mean zero and strong mixing sequence with mixing coefficient satisfying  $\cdot/n \cdot O \cdot n^{-p} \cdot$  and  $\sup_t \int_0^1 E|W_{n\cdot} s/\cdot|^p \cdot C < 1$ . In addition,  $E \cdot W_n W_n^T = n \cdot I < 1$  as  $n \rightarrow \infty$ .

Let  $W_n \cdot s/\cdot \rightarrow W_{n\cdot} s/\cdot$  and set  $F_i \rightarrow \cdot U_{nt}; W_t \cdot s/\cdot$  to be the smallest  $\sigma$ -field containing the past history of  $\cdot U_{nt}; W_t/\cdot$  for all  $n$  and  $t \leq i$ , and denote  $E \cdot X_j F_i/\cdot$  by  $E_i \cdot X_j/\cdot$ . Since  $W_n \cdot s/\cdot$  might not be a martingale, we can use a martingale to approximate  $W_n \cdot s/\cdot$ . To this end, define,

$$\tilde{X}_i \rightarrow \int_0^i E_i \cdot W_{iCk}/\cdot E_i \cdot W_{iCk}/\cdot; \text{ and } \tilde{X}_i \rightarrow \int_0^i E_i \cdot W_{iCk}/\cdot$$

Then, it is not hard to verify (see Hansen (1992, p. 492)) that  $W_i \rightarrow \int_0^i \tilde{X}_i$  with  $E_i \cdot \tilde{X}_i/\cdot \rightarrow 0$ , and for  $0 \leq r \leq 1$ ,

$$Z_r = \int_0^r U_n \cdot s/dW_n \cdot s/\cdot = \int_0^r U_n \cdot s/dY_n \cdot s/\cdot - \int_0^r U_n \cdot r/\cdot$$

where  $Y_n \cdot r/\cdot \rightarrow \int_0^r Y_{n\cdot} s/\cdot = \int_0^r \tilde{X}_i$  and

$$U_n \cdot r/\cdot \rightarrow \int_0^r U_{ni} U_{ni}^T \cdot \int_0^r U_n \cdot r/\cdot = \int_0^r U_n \cdot r/\cdot$$

Then,  $\tilde{X}_i; F_i$  is a martingale difference sequence. Therefore, by Theorem 2.1 in Hansen (1992), we have the following result.

**Lemma A.1.** Assume that Assumption M1 holds,  $\sup_{0 \leq r \leq 1} j \cdot n \cdot r/\cdot \rightarrow O_p \cdot 1/\cdot$ , and  $\cdot U_n; W_n/H \cdot \rightarrow U; W/\cdot$  in  $\mathcal{D}_{M^k m} \mathcal{D}[0, 1]$ , then

$$\int_0^1 U_n dY_n \cdot s/\cdot \rightarrow \int_0^1 U dW; \text{ equivalently, } \int_0^1 U_n dW_n \cdot s/\cdot \rightarrow \int_0^1 U dW;$$

where  $U$  is a cadlag process,  $W \cdot s/\cdot \rightarrow BM_{\cdot} z/\cdot$ , and  $\cdot$  is defined in Assumption M1.

**Proof of Theorem 2.1.** First, note that the right hand side of (2.3) has the form of  $A^{-1}B_1$ , define  $H_n = \begin{pmatrix} 1 & 0 \\ 0 & b \end{pmatrix} D_n$ , so that we can

write  $H_n A^{-1} B D H_n A^{-1} H_n H_n^{-1} B D T H_n^{-1} A H_n^{-1} U^{-1} H_n^{-1} B$ . Thus,  $b_{\cdot,z}/$  and  $b_{\cdot,1'}/z/$  can be re-expressed as follows:

$$H_n \frac{b_{\cdot,z}/}{b_{\cdot,1'}/z/} D S_n \cdot z/ \frac{1}{n} \frac{1}{n} \frac{1}{n} K_h \cdot Z_t \cdot z/ \quad (A.6)$$

where  $Z_{t,z,h} D \cdot Z_t \cdot z/h$  and

$$S_n \cdot z/ D \frac{1}{n} \frac{1}{n} K_h \cdot Z_t \cdot z/ \frac{1}{Z_{t,z,h}^2} D_n^{-1} X_t^2$$

with  $j D 0; 1; 2$ ,

$$S_{n,j} \cdot z/ D \frac{1}{n} \frac{1}{n} K_h \cdot Z_t \cdot z/ Z_{t,z,h}^j D_n^{-1} X_t^2$$

Now, to facilitate the analysis of  $S_{n,j} \cdot z/$ , we first express  $S_n \cdot z/$  as

$$S_n \cdot z/ D \frac{F_{n,j,0} \cdot z/}{F_{n,j,1} \cdot z/} \frac{F_{n,j,1} \cdot z/}{F_{n,j,2} \cdot z/}$$

where

$$F_{n,j,0} \cdot z/ D \frac{1}{n} \frac{1}{n} Z_{t,z,h}^j X_{t1} X_{t1}^T K_h \cdot Z_t \cdot z/$$

$$F_{n,j,1} \cdot z/ D \frac{1}{n} \frac{1}{n} K_h \cdot Z_t \cdot z/ Z_{t,z,h}^j X_{t1} X_{t2}^T = \frac{1}{n}$$

and

$$F_{n,j,2} \cdot z/ D \frac{1}{n} \frac{1}{n} Z_{t,z,h}^j K_h \cdot Z_t \cdot z/ X_{t2}^2 = \frac{1}{n}^2$$

For  $l D 1; 2$ , define

$$F_{n,j,l} \cdot z/ D \frac{1}{n} \frac{1}{n} Z_{t,z,h}^j X_{t1}^l K_h \cdot Z_t \cdot z/$$

By noting that  $X_{t1}$  and  $Z_t$  are stationary and using the standard change-of-variable and a Taylor's expansion argument, we have

$$E F_{n,j,l} \cdot z/ D E Z_{t,z,h}^j X_{t1}^l K_h \cdot Z_t \cdot z/ D f_{z,z}/M_l \cdot z/ \cdot j \cdot K/C O(1/)$$

By the kernel theory for the stationary mixing case; see Theorem 1 of Cai et al. (2000) for details, one can easily show that

$$\text{Var } F_{n,j,l} \cdot z/ D O(\cdot n h^{-1}/D O(1/); \quad (A.7)$$

Therefore,

$$F_{n,j,l} \cdot z/ D f_{z,z}/M_l \cdot z/ \cdot j \cdot K/C O_p(1/); \quad (A.8)$$

so that

$$F_{n,j,0} \cdot z/ D F_{n,j,2} \cdot z/ D f_{z,z}/M_2 \cdot z/ \cdot j \cdot K/C O_p(1/); \quad (A.9)$$

Let  $F_i^e D \cdot X_{t1}; Z_t \vee t \cdot i/$  be the smallest  $\sigma$ -field containing the past history of  $\cdot X_{t1}; Z_t/$ , and denote by  $e_t D K_h \cdot Z_t \cdot z/ Z_{t,z,h}^j X_{t1}$

$E[K_h \cdot Z_t \cdot z/ Z_{t,z,h}^j X_{t1}]$ . Then, similar to (A.7), it is easy to verify that

$$\sup_{s=0}^m \text{Var } e_t D O(m=h/); \quad (A.10)$$

for any  $m \cdot 1$ . Define  $U_{nt} D X_{t2} = \frac{1}{n}$  for any  $1 \cdot t \cdot n$  and  $U_{n,r}/ D U_{n, \lfloor nr \rfloor}$  for any  $r \cdot 0; 1/$ . For any small  $0 < \cdot < 1$ , set  $N D \lfloor 1/ \rfloor$ ,  $t_k D \lfloor k n/N \rfloor \subset 1$ ,  $t_k D t_{kC1} \cdot 1$ , and  $t_k D \min\{t_k; n\}$ . Then,

$$\frac{1}{n} \sum_{iD1}^n U_{ni} e_i D \frac{1}{n} \sum_{kD0}^N \sum_{tDt_k}^n U_{nt} e_t$$

$$\frac{1}{n} \sum_{kD0}^N \sum_{tDt_k}^n U_{nt_k} e_t \subset \frac{1}{n} \sum_{kD0}^N \sum_{tDt_k}^n U_{nt} U_{nt_k} e_t$$

$$\frac{1}{n} \sum_{kD0}^N \sum_{tDt_k}^n U_{nt_k} e_t \subset \frac{1}{n} \sum_{kD0}^N \sum_{tDt_k}^n U_{nt} U_{nt_k} e_t$$

$$\sup_{0 \leq s \leq 1} \sum_{j=1}^n U_{n,s}/ \frac{1}{n} \sum_{kD0}^N \sum_{tDt_k}^n e_t$$

$$\subset \sup_{j,r,s,j} \sum_{j=1}^n U_{n,r}/ U_{n,s}/ \frac{1}{n} \sum_{tD1}^n e_t$$

Since  $U_{n,r}/$  converges weakly to a Brownian motion, it is clear that  $\sup_{0 \leq s \leq 1} \sum_{j=1}^n U_{n,s}/ D O_p(1/)$ , while  $\frac{1}{n} \sum_{tD1}^n e_t = \frac{1}{n} D O_p(1/)$  by using the same arguments as those used in the proof of (A.8). Further,

$$E \frac{1}{n} \sum_{kD0}^N \sum_{tDt_k}^n e_t \leq \frac{N}{n} \sup_{k=0}^{N-1} E \sum_{tDt_k}^n e_t$$

$$\sup_{t=n}^n E \frac{1}{n} \sum_{iD1}^n e_i$$

$$\subset \cdot n h^{-1/2} \cdot 0$$

by (A.10) as  $n \cdot 1$ . Therefore, as  $n \cdot 1$ ,

$$\frac{1}{n} \sum_{iD1}^n U_{ni} e_i D O_p(1/); \sup_{j,r,s,j} \sum_{j=1}^n U_{n,r}/ U_{n,s}/ O_p(1/);$$

Note that as  $n \cdot 1$ ,

$$\sup_{j,r,s,j} \sum_{j=1}^n U_{n,r}/ U_{n,s}/ \cdot \sup_{j,r,s,j} \sum_{j=1}^n W_{\cdot,2,r}/ W_{\cdot,2,s}/ \cdot 0$$

as  $\cdot \cdot 0$ . Hence (here we take the sequential limits: first let  $n \cdot 1$ , and then let  $\cdot \cdot 0$ ),

$$\frac{1}{n} \sum_{iD1}^n U_{ni} e_i D O_p(1/);$$

which, by combining (A.9), (2.6), and Lemma A.2, gives that

$$F_{n,j,1} \cdot z/ D E K_h \cdot Z_t \cdot z/ Z_{t,z,h}^j X_{t1} \frac{1}{n} \sum_{tD1}^n U_{nt} \subset \frac{1}{n} \sum_{iD1}^n U_{ni} e_i$$

$$D f_{z,z}/ \cdot j \cdot K/M_1 \cdot z/ W_{\cdot,2}^{-1/2} \subset O_p(1/); \quad (A.11)$$

Similarly,

$$F_{n,j,2} \cdot z/ D f_{z,z}/ \cdot j \cdot K/W_{\cdot,2}^{-2/2} \subset O_p(1/); \quad (A.12)$$

Then, by plugging (A.9), (A.11) and (A.12) into  $S_{n,j} \cdot z/$ , we have

$$S_{n,j} \cdot z/ D f_{z,z}/ \cdot j \cdot K/S \cdot z/ C O_p(1/); \quad (A.13)$$

By noting that  $\cdot_0 \cdot K/D 1$  and  $\cdot_1 \cdot K/D 0$ , we immediately obtain from (A.13) that

$$S_n \cdot z/ D f_{z,z}/ \frac{1}{0} \frac{0}{2 \cdot K/} S \cdot z/ C O_p(1/); \quad (A.14)$$



Let  $R_n(z)^{-1}$  denote the upper-left corner  $d \times d$  sub-matrix of  $S_n(z)^{-1}$ . From (A.14), we immediately obtain that

$$R_n(z)^{-1} D f_z(z)^{-1} S_n(z)^{-1} C o_p(1/\sqrt{n}) \quad (\text{A.15})$$

From (A.6), we have

$$D_n b(z) = z / I_3 C I_4; \quad (\text{A.16})$$

where

$$I_3 D R_n(z)^{-1} B_n(z); \quad (\text{A.17})$$

with

$$B_n(z) D n^{-1} \int_{tD1}^{\infty} K_h \cdot Z_t \cdot z / D_n^{-1} X_t X_t^T f \cdot Z_t / \cdot z / \cdot Z_t \cdot z / \cdot^{1/} \cdot z / g;$$

and

$$I_4 D R_n(z)^{-1} n^{-1} \int_{tD1}^{\infty} K_h \cdot Z_t \cdot z / {}_t D_n^{-1} X_t;$$

Define,

$$G_{n,0}(z) D n^{-1} \int_{tD1}^{\infty} K_h \cdot Z_t \cdot z / X_{t1}^2 f \cdot Z_t / \cdot z / \cdot Z_t \cdot z / \cdot^{1/} \cdot z / g;$$

$$G_{n,1}(z) D n^{-1} \int_{tD1}^{\infty} K_h \cdot Z_t \cdot z / X_{t1} \cdot X_{t2} = \frac{p_-}{n} \int f \cdot Z_t / \cdot z / \cdot Z_t \cdot z / \cdot^{1/} \cdot z / g;$$

$$G_{n,2}(z) D n^{-1} \int_{tD1}^{\infty} K_h \cdot Z_t \cdot z / \cdot X_{t2} = \frac{p_-}{n} \int X_{t1}^T \frac{p_-}{n} f \cdot Z_t / \cdot z / \cdot Z_t \cdot z / \cdot^{1/} \cdot z / g;$$

and

$$G_{n,3}(z) D n^{-1} \int_{tD1}^{\infty} K_h \cdot Z_t \cdot z / \cdot X_{t2} = \frac{p_-}{n} \int \frac{p_-}{n} f \cdot Z_t / \cdot z / \cdot Z_t \cdot z / \cdot^{1/} \cdot z / g;$$

so that

$$B_n(z) D \frac{G_{n,0}(z) / C G_{n,1}(z)}{C_{n,2}(z) / C G_{n,3}(z)}; \quad (\text{A.18})$$

Similar to (A.9), by the kernel theory and an application of Taylor's expansion, it is easy to show that

$$E[G_{n,0}(z)] D h^2 f_z(z) / M_2(z) \cdot \frac{2 \cdot K /}{2} \cdot \frac{2 /}{1} \cdot z / f_1 C o(1/\sqrt{n})$$

and  $\text{Var}[G_{n,0}(z)] D o(1/\sqrt{n})$ , so that

$$G_{n,0}(z) D h^2 f_z(z) / M_2(z) \cdot \frac{2 \cdot K /}{2} \cdot \frac{2 /}{1} \cdot z / f_1 C o_p(1/\sqrt{n});$$

Further, following the proof of (A.11), we can easily show that

$$G_{n,1}(z) D h^2 f_z(z) / M_1(z) \cdot W_{1,2}^{-1/} \cdot \frac{2 \cdot K /}{2} \cdot \frac{2 /}{2} \cdot z / f_1 C o_p(1/\sqrt{n});$$

$$G_{n,2}(z) D h^2 f_z(z) / M_1(z) \cdot W_{1,2}^{-1/} \cdot \frac{p_-}{n} \cdot \frac{2 \cdot K /}{2} \cdot \frac{2 /}{1} \cdot z / f_1 C o_p(1/\sqrt{n});$$

and

$$G_{n,3}(z) D h^2 f_z(z) / W_{1,2}^{-2/} \cdot \frac{p_-}{n} \cdot \frac{2 \cdot K /}{2} \cdot \frac{2 /}{2} \cdot z / f_1 C o_p(1/\sqrt{n});$$

Plugging the above results into (A.18), we obtain

$$B_n(z) D h^2 f_z(z) / S_n(z) D_n \cdot \frac{2 \cdot K /}{2} \cdot \frac{2 /}{1} \cdot z / f_1 C o_p(1/\sqrt{n}); \quad (\text{A.19})$$

Substituting (A.19) into (A.17) and using (A.15) lead to

$$I_3 D D_n h^2 B \cdot z / f_1 C o_p(1/\sqrt{n});$$

Therefore,

$$D_n^{-1} I_3 D h^2 B \cdot z / C o_p(h^2); \quad (\text{A.20})$$

Finally, we consider  $I_4$ . Define

$$T_n(z) D \frac{r}{n} \int_{tD1}^{\infty} K_h \cdot Z_t \cdot z / {}_t D_n^{-1} X_t D \frac{T_{n,1}(z)}{T_{n,2}(z)}$$

with  $T_{n,1}(z) D \frac{q_-}{n} \int_{tD1}^{\infty} K_h \cdot Z_t \cdot z / {}_t X_{t1}$  and  $T_{n,2}(z) D \frac{q_-}{n} \int_{tD1}^{\infty} K_h \cdot Z_t \cdot z / {}_t X_{t2} = \frac{p_-}{n}$ . By combining the above expressions with (A.16) and (A.20), we obtain

$$\frac{p_-}{n h} D_n b(z) \cdot z / h^2 B \cdot z / C o_p(h^2) D R_n(z)^{-1} T_n(z); \quad (\text{A.21})$$

To prove the asymptotic normality of the left hand side of (A.21), it suffices to establish the asymptotic normality of  $T_n(z)$ . Note that  $T_{n,1}$  only involves stationary variables. Hence, by the kernel estimation theory for stationary mixing data; see Theorem 2 of Cai et al. (2000) for details, we have

$$T_{n,1}(z) \xrightarrow{d} N(0; \frac{2}{n} \cdot \frac{0 \cdot K / f_z(z) / M_2(z)}{0 \cdot K / f_z(z) / W_{1,1}(z)}); \quad (\text{A.22})$$

where  $W_{1,1}(z)$  is a  $p_1$ -dimensional Brownian motion on  $[0, 1]$  with covariance matrix  $\frac{2}{n} M_2(z)$ . From (A.22) and note that the first element of  $X_{t1}$  is one, we immediately obtain

$$\frac{p_-}{h=n} \int_{tD1}^{\infty} K_h \cdot Z_t \cdot z / {}_t \xrightarrow{d} N(0; \frac{2}{n} \cdot \frac{0 \cdot K / f_z(z)}{0 \cdot K / f_z(z) / W_{1,1}(z)}); \quad (\text{A.23})$$

where  $W_{1,1}(z)$  is the first element of  $W_{1,1}(z)$ . Note that using the notations of Lemmas A.1–A.3, we have  $U_n(z) D X_{[nr]1} = \frac{p_-}{n}$ ,  $W_n(z) D \int_{sD1}^{[nr]} W_s = \frac{p_-}{n}$  and  $T_{n,2}(z) D \int_0^1 U_n(t) dW_n(t)$ . Hence, (A.22), in conjunction with Lemmas A.1 and A.3

$$T_{n,2}(z) \xrightarrow{d} \frac{p_-}{0 \cdot K / f_z(z)} \int_0^1 W_{1,2}(z) dW_{1,1}(z); \quad (\text{A.24})$$

Therefore, a combination of (A.22) and (A.24) leads to

$$T_n(z) \xrightarrow{d} \frac{p_-}{0 \cdot K / f_z(z)} \int_0^1 \frac{W_{1,1}(z)}{W_{1,2}(z) dW_{1,1}(z)} dW_{1,1}(z);$$

Since  $W_{1,2}(z)$  and  $W_{1,1}(z)$  are uncorrelated (because  $Z_t$  and  $X_t$  are uncorrelated),  $\int_0^1 W_{1,2}(z) dW_{1,1}(z)$  has a mixed normal distribution,

so that the conditional covariance of  $\int_0^1 \frac{W_{1,1}(z)}{W_{1,2}(z) dW_{1,1}(z)}$  is

$$\frac{2}{n} \frac{M_2(z)}{W_{1,2}(z) M_1(z)^T} \frac{M_1(z) W_{1,2}^{-1/}}{W_{1,2}^{-2/}} D \frac{2}{n} S(z); \quad (\text{A.25})$$

Therefore, by Slutsky's theorem, we have



An application of Proposition 11 of Jeganathan (2004) gives

$$b_{j,z}/D_{j,K/L,1;0} = u C_{0,2,1}/$$

as “# By the same token, it is easy to show the case of  $x_n D a$  ( $z D a$ ). For assertion (ii), we have

$$\begin{aligned} E b_{j,z}/D_{j,K/L,1;0} &= E K_{j,h} Z_t z/ \\ &= D_{j,K/L,1;0} \int_0^1 \int_0^1 K_j(t^{1/2}u/h/f_{t,z}, u/du \\ &= D_{j,K/L,1;0} \int_0^1 \int_0^1 K_j(v/f_{t,z}, ht^{1/2}v/dv \\ &= C n^{1/2} \int_0^1 \int_0^1 t^{1/2} D_{0,1}/; \end{aligned}$$

Finally, recall that  $K_{j,h} u/D h^{-1} K_j u=h/$  and  $K_j u/D u^j K u/$ , it can be shown easily by the boundedness of  $f_{t,z}$  that

$$\begin{aligned} E j K_{j,h} Z_t z/ &= D h^p \int_0^1 \int_0^1 K_{j,h} t^{1/2} u=h/ f_{t,z} u/du \\ &= D t^{1/2} h^{1-p} \int_0^1 \int_0^1 K_j v/f_{t,z} t^{1/2} h v/dv = C t^{1/2} h^{1-p}; \end{aligned}$$

This proves the lemma.

**Lemma B.2.** Under Assumptions given in Theorem 3.1, if  $z$  is fixed, we have

$$S_{nj,z}/D E X_t X_t^T / b_{j,z} / C_{0,p,1} = E X_t X_t^T / j,K/L,1;0 = u;$$

**Proof.** Recall that  $V_t D X_t X_t^T$ . By adding and subtracting  $E V_t /$  and  $E V_t j Z_t /$  in  $S_{nj,z}/$ , we decompose  $S_{nj,z}/$  into three terms as follows:

$$\begin{aligned} S_{nj,z}/D B_{1n,1} C B_{1n,2} C B_{1n,3}; \\ \text{where } B_{1n,1} D E V_t / n^{1/2} \int_0^1 K_{j,h} Z_t z/ D E V_t / b_{j,z} /, \\ B_{1n,2} D n^{1/2} \int_0^1 E V_t j Z_t / E V_t / K_{j,h} Z_t z/ \\ D n^{1/2} \int_0^1 g_t Z_t / K_{j,h} Z_t z/; \end{aligned}$$

and

$$\begin{aligned} B_{1n,3} D n^{1/2} \int_0^1 V_t E V_t j Z_t / K_{j,h} Z_t z/ \\ n^{1/2} \int_0^1 t K_{j,h} Z_t z/; \end{aligned}$$

where  $t D V_t E V_t j Z_t /$ . It follows from Lemma B.1 that  $B_{1n,1} \leq j,K/E V_t / L,1;0 = u$ . To show the lemma, it suffices to show that  $B_{1n,2} D_{0,p,1}/$  and  $B_{1n,3} D_{0,p,1}/$ , respectively.

First, we show that  $B_{1n,2} D_{0,p,1}/$ . By (3.5) and the boundedness of  $g_t$  as well as  $t D O(t^{1/2})$  (see Assumption C3), we have

$$\begin{aligned} j B_{1n,2} j &= n^{1/2} \int_0^1 \int_0^1 g_t Z_t / j K_{j,h} Z_t z/ \\ &= C n^{1/2} \int_0^1 \int_0^1 t^{1/2} C Z_t / j K_{j,h} Z_t z/ \\ &= C n^{1/2} \int_0^1 \int_0^1 t^{1/2} C Z_t z/ C z/ j K_{j,h} Z_t z/; \end{aligned}$$

Since  $C u/$  is continuous at  $z$  and  $K_j$  has a finite support, then  $C u/ C_z$  for some  $C_z$  for all  $u$ 's in a neighborhood of  $z$ . Therefore,

$$j B_{1n,2} j = C n^{1/2} \int_0^1 \int_0^1 t^{1/2} j K_{j,h} Z_t z/ = 0$$

by Lemma B.1 and Toeplitz lemma.

Next, we show that  $B_{1n,3} D_{0,p,1}/$ . To do so, it suffices to show that  $E B_{1n,3}^2 \leq O(1)$ . To this end, we have (2 below means  $\int_0^1 \int_0^1$  for any  $i$  and  $j$ , see the discussions below Lemma A.2 for our notation adoption)

$$\begin{aligned} E B_{1n,3}^2 &= E \int_0^1 \int_0^1 K_{j,h}^2 Z_t z/ \\ &= C 2 n^{1/2} \int_0^1 \int_0^1 E \int_0^1 \int_0^1 K_{j,h} Z_t z/ K_{j,h} Z_s z/ \\ &= B_{1n,3,1} C B_{1n,3,2}; \end{aligned}$$

Clearly, by Cauchy–Schwartz inequality, Lemma B.1, and Assumption C1, we have

$$\begin{aligned} j B_{1n,3,1} j &= n^{1/2} \int_0^1 \int_0^1 K_{j,h}^2 Z_t z/ K_{j,h}^2 Z_s z/ \\ &= C n^{1/2} \int_0^1 \int_0^1 t^{1/2} h^{1-p} \\ &= C n h^{4p-2} \int_0^1 \int_0^1 t^{1/2} D_{0,1}/; \end{aligned}$$

since  $n h^{4p-2} \int_0^1 \int_0^1 t^{1/2} D_{0,1}/$  (from Assumption C5). For  $B_{1n,3,2}$ , by (3.5) and (3.6), we have

$$\begin{aligned} E \int_0^1 \int_0^1 Z_t Z_s^T V_s / D E V_t j Z_t Z_s^T V_s / E V_t j Z_t / \\ D \int_0^1 \int_0^1 g_t Z_t Z_s^T V_s / \int_0^1 g_t Z_t /; \end{aligned} \quad (B.1)$$

It is easy to see that

$$\begin{aligned} j B_{1n,3,2} j &= C n^{1/2} \int_0^1 \int_0^1 E f E \int_0^1 \int_0^1 Z_t Z_s^T V_s / s K_{j,h} Z_t z/ K_{j,h} Z_s z/ \\ &= C \int_0^1 \int_0^1 \int_0^1 \int_0^1 E f g_t Z_t Z_s^T V_s / s \\ &= C \int_0^1 \int_0^1 \int_0^1 \int_0^1 E f g_t Z_t Z_s^T V_s / s K_{j,h} Z_t z/ K_{j,h} Z_s z/ \\ &= B_{1n,3,2,1} C B_{1n,3,2,2}; \end{aligned}$$

To evaluate  $B_{1n,3,2,1}$ , first, we consider the following quantity

$$\begin{aligned} E \int_0^1 \int_0^1 g_t Z_t Z_s^T V_s / s K_{j,h} Z_t z/ K_{j,h} Z_s z/ V_s / \\ D \int_0^1 \int_0^1 b_{t,s,z} t^{1/2} h u; s^{1/2} h v / g_{t,s} z/ C h u; z C h v; V_s / \\ \int_0^1 \int_0^1 V_s E V_s / s g_s z/ C h v / K_{j,h} u / K_{j,h} v / t^{1/2} s^{1/2} d u d v \\ D \int_0^1 \int_0^1 K / t^{1/2} s^{1/2} b_{t,s,z} 0; 0 / g_{t,s} z; z; V_s / \\ \int_0^1 \int_0^1 V_s E V_s / s g_s z/ \int_0^1 \int_0^1 C_{0,p,1}/; \end{aligned}$$

Then, using (3.5) and (B.1), we have

$$\begin{aligned} B_{1n,3,2,1} &= C \int_0^1 \int_0^1 \int_0^1 \int_0^1 E f E \int_0^1 \int_0^1 g_t Z_t Z_s^T V_s / s K_{j,h} Z_t z/ \\ &= C \int_0^1 \int_0^1 \int_0^1 \int_0^1 K_{j,h} Z_s z/ V_s / g_j \\ &= C n^{1/2} \int_0^1 \int_0^1 s^{1/2} t^{1/2} b_{t,s,z} 0; 0 / \\ &= E \int_0^1 \int_0^1 g_t Z_t z; z; V_s / f V_s E V_s / s g_s z/ g f 1 C_{0,1}/ j \\ &= O(1) \end{aligned}$$



which, by replacing  $Y_t$  in (3.2) by  $Y_t D X_t^T \cdot Z_t / C_t$ , implies that

$$b.z / \left( \frac{.z / D E.X_t X_t^T / L.1; 0 = u^{-1}}{n^{-1/2} \sum_{t=D1}^n X_t X_t^T \cdot Z_t / \cdot z / \cdot 1 / \cdot z / \cdot Z_t \cdot z /} \right) K_h \cdot Z_t \cdot z / C n^{-1/2} \sum_{t=D1}^n X_t \cdot t K_h \cdot Z_t \cdot z / f1 C o_p.1 / g E.X_t X_t^T / L.1; 0 = u^{-1} B_{2n} C B_{3n} g f1 C o_p.1 / g; \quad (B.2)$$

where  $B_{2n} D n^{-1/2} \sum_{t=D1}^n X_t X_t^T \cdot Z_t / \cdot z / \cdot 1 / \cdot z / \cdot Z_t \cdot z / K_h \cdot Z_t \cdot z /$  and  $B_{3n} D n^{-1/2} \sum_{t=D1}^n X_t \cdot t K_h \cdot Z_t \cdot z /$ . The asymptotic behaviors of  $B_{2n}$  and  $B_{3n}$  are derived in Lemmas B.3 and B.4. Therefore, combining Lemmas B.3 and B.4 with (B.2), we obtain that

$$n^{1/4} h^{1/2} b.z / \cdot z / h^2 B \cdot z / C o_p.h^2 / D u L.1; 0 / E.X_t X_t^T /^{-1} n^{1/4} h^{1/2} B_{3n} f1 C o_p.1 / g \uparrow MN. \cdot 1 /$$

This completes the proof of Theorem 3.1.

## References

- Berkes, I., Horváth, L., 2006. Convergence of integral functionals of stochastic processes. *Econometric Theory* 22, 304–322.
- Billingsley, P., 1999. *Convergence of Probability Measures*, 2nd ed. Wiley, New York.
- Cai, Z., 2002a. Regression quantile for time series. *Econometric Theory* 18, 169–192.
- Cai, Z., 2002b. Two-step likelihood estimation procedure for varying-coefficient models. *Journal of Multivariate Analysis* 82, 189–209.
- Cai, Z., 2002c. A two-stage approach to additive time series models. *Statistica Neerlandica* 56, 415–433.
- Cai, Z., 2007. Trending time varying coefficient time series models with serially correlated errors. *Journal of Econometrics* 137, 163–188.
- Cai, Z., Fan, J., Yao, Q., 2000. Functional-coefficient regression models for nonlinear time series. *Journal of the American Statistical Association* 95, 941–956.
- Cai, Z., Wang, Y., 2008. Instability of predictability of stock returns. Working Paper. Department of Mathematics and Statistics, University of North Carolina at Charlotte.
- Campbell, J.Y., Yogo, M., 2006. Efficient tests of stock return predictability. *Journal of Financial Economics* 81, 27–60.
- Carrasco, M., Chen, X., 2002. Mixing and moment properties of various GARCH and stochastic volatility models. *Econometric Theory* 18, 17–39.
- Chang, Y., Martinez-Chombo, E., 2003. Electricity demand analysis using cointegration and error-correction models with time varying parameters: The Mexican case. Working Paper. Department of Economics, Texas A&M University.
- Chen, B., Hong, Y., 2007. Testing for smooth structural changes in time series models via nonparametric regression. Working Paper. Department of Economics, Cornell University.
- Chen, S.X., Tang, C.Y., 2005. Nonparametric inference of value at risk for dependent financial returns. *Journal of Financial Econometrics* 3, 227–255.
- Chen, R., Tsay, R.S., 1993. Functional-coefficient autoregressive models. *Journal of the American Statistical Association* 88, 298–308.
- Csörgő, M., Révész, P., 1981. *Strong Approximation in Probability and Statistics*. Academic Press, New York.
- Einhärl, U., 1987. A useful estimate in the multidimensional invariance principle. *Probability Theory and Related Fields* 76, 81–101.
- Fan, J., Huang, T., 2005. Profile likelihood inferences on semiparametric varying-coefficient partially linear models. *Bernoulli* 11, 1031–1057.
- Fan, J., Gijbels, I., 1996. *Local Polynomial Modeling and its Applications*. Chapman and Hall, London.
- Hall, P., Heyde, C.C., 1980. *Martingale Limit Theory and its Applications*. Academic Press, New York.
- Hansen, B.E., 1992. Convergence to stochastic integrals for dependent heterogeneous processes. *Econometric Theory* 8, 489–500.
- Hansen, B.E., 2008. Uniform convergence rates for kernel estimation with dependent data. *Econometric Theory* 24, 726–748.
- Hastie, T., Tibshirani, R., 1993. Varying coefficient models (with discussions). *Journal of the Royal Statistical Society, Series B* 55, 757–796.
- Hurvich, C.M., Simonoff, J.S., Tsai, C.L., 1998. Smoothing parameter selection in nonparametric regression using an improved Akaike information criterion. *Journal of the Royal Statistical Society, Series B* 60, 271–293.
- Jeganathan, P., 2004. Convergence of functionals of sums of random variables to local times of fractional stable motions. *The Annals of Probability* 32, 1771–1795.
- Juhl, T., 2005. Functional coefficient models under unit root behavior. *Econometrics Journal* 8, 197–213.
- Karatzas, I., Shreve, S.E., 1991. *Brownian Motion and Stochastic Calculus*, 2nd ed. Springer-Verlag, New York.
- Karlsen, H.A., Myklebust, T., Tjøstheim, D., 2007. Nonparametric estimation in a nonlinear cointegration type model. *Annals of Statistics* 35, 252–299.
- Li, Q., Huang, C., Li, D., Fu, T., 2002. Semiparametric smooth coefficient models. *Journal of Business and Economic Statistics* 20, 412–422.
- Linton, O.B., 2000. Efficient estimation of generalized additive nonparametric regression models. *Econometric Theory* 16, 502–523.
- McLeish, D.L., 1975. A maximal inequality and dependent strong laws. *The Annals of Probability* 3, 829–839.
- Merlevède, F., Peligrad, M., Utev, S., 2006. Recent advances in invariance principles for stationary sequences. *Probability Surveys* 3, 1–36.
- Park, J.Y., Hahn, S.B., 1999. Cointegrating regressions with time varying coefficients. *Econometric Theory* 15, 664–703.
- Phillips, P.C.B., 1989. Partially identified econometric models. *Econometric Theory* 5, 181–240.
- Park, J.Y., Phillips, P.C.B., 1999. Asymptotics for nonlinear transformations of integrated time series. *Econometric Theory* 15, 269–298.
- Phillips, P.C.B., Park, J., 1998. Nonstationary density and kernel autoregression. Unpublished manuscript.
- Polk, C., Thompson, S., Vuolteenaho, T., 2006. Cross-sectional forecasts of the equity premium. *Journal of Financial Economics* 81, 101–141.
- Robinson, P.M., 1988. Root-n consistent semiparametric regression. *Econometrica* 56, 931–954.
- Robinson, P.M., 1989. Nonparametric estimation of time-varying parameters. In: Hackl, P. (Ed.), *Statistical Analysis and Forecasting of Economic Structural Change*. Springer-Verlag, Berlin, pp. 253–264.
- Robinson, P.M., 1991. Time-varying nonlinear regression. In: Hackl, P. (Ed.), *Statistical Analysis and Forecasting of Economic Structural Change*. Springer-Verlag, Berlin, pp. 179–190.
- Ruppert, D., Sheather, S.J., Wand, M.P., 1995. An effective bandwidth selection for local least squares regression. *Journal of the American Statistical Association* 90, 1257–1270.
- Shao, Q., Lu, C., 1987. Strong approximation for partial sums of weakly dependent random variables. *Scientia Sinica* 15, 576–587.
- Torous, W., Valkanov, R., Yan, S., 2004. On predicting stock returns with nearly integrated explanatory variables. *The Journal of Business* 77, 937–966.
- Wang, Q., Phillips, P.C.B., 2006. Asymptotic theory for local time density estimation and nonparametric cointegrating regression. *Econometric Theory* (forthcoming).
- Wang, Q., Phillips, P.C.B., 2008. Structural nonparametric cointegrating regression. Manuscript.
- Xiao, Z., 2007. Functional coefficient co-integration models. *Journal of Econometrics* (forthcoming).