

## Exercise 1

1.  $\subseteq$  is a partial order on any set, to prove it is a lattice, we show that  $\sup\{a, b\}$  and  $\inf\{a, b\}$  exists.  $\sup\{a, b\} = a \cup b$  and clearly  $a \cup b \in Su(A)$ . Since  $a \subseteq a \cup b$  and  $b \subseteq a \cup b$  it is an upper bound for  $\{a, b\}$ . To prove it is the least upper bound, assume that there is some set  $c \in Su(A)$  such that  $a \subseteq c, b \subseteq c$ , and  $c \subseteq a \cup b$ , it means there is some  $k$  in  $a$  or  $b$ , such that  $k \notin c$ , a contradiction.  $\inf\{a, b\} = a \cap b$  can be proved similarly.

2. L1:(a)  $a \vee b = \sup\{a, b\} = \sup\{b, a\} = b \vee a$ .

(b)  $a \wedge b = \inf\{a, b\} = \inf\{b, a\} = b \wedge a$ .

L2: (a) Let  $t = \sup\{a, \sup\{b, c\}\}$  and  $k = \sup\{\sup\{a, b\}, c\}$ , then  $t$  is greater than  $a$  and  $\sup\{b, c\}$ , which means  $t$  is also greater than  $b, c$ , thus  $\sup\{a, b\} \leq t$ , therefore  $\sup\{\sup\{a, b\}, c\} \leq t$ , or  $k \leq t$ .  $k = \sup\{\sup\{a, b\}, c\}$  implies  $c \leq k$  and  $\sup\{a, b\} \leq k$ , which means  $a \leq k$  and  $b \leq k$ . Now we can say that  $\sup\{b, c\} \leq k$ , therefore  $\sup\{a, \sup\{b, c\}\} \leq k$ . From  $t \leq k$  and  $k \leq t$  we conclude  $t = k$ .

(b) Can be proved similarly.

L3: (a) and (b) follows from  $\inf\{a, a\} = \sup\{a, a\} = a$ .

L4: (a) Let  $t = \sup\{a, \inf\{a, b\}\}$  then  $a \leq t$ . Since  $\inf\{a, b\} \leq a$ , we get  $\sup\{a, \inf\{a, b\}\} \leq \sup\{a, a\} = a$ , thus  $t \leq a$  and finally  $a = t$ .

(b) Let  $t = \inf\{a, \sup\{a, b\}\}$ , then  $t \leq a$ , and since  $a \leq \sup\{a, b\}$  we have  $a = \inf\{a, a\} \leq \inf\{a, \sup\{a, b\}\}$ , thus  $a \leq t$ .

3. From L4(b) we have  $x \approx x \wedge (x \vee x)$ , replace  $x \vee x$  for  $y$  in the formula L4(a), we get  $x \approx x \vee [x \wedge (x \vee x)]$ , replacing the first formula in it, we get  $x \approx x \vee x$ . Similarly, replace  $y$  by  $x \wedge x$  in L4(b), we get  $x \approx x \wedge [x \vee (x \wedge x)]$ , from L4(a) we have  $x \approx x \vee (x \wedge x)$ . Therefore  $x \approx x \wedge x$ .

5. Since each item in L1,..., L4 has two symmetric axioms for  $\vee$  and  $\wedge$ , the result trivially follows.

9. (Note: we write  $(a, b) \in R$  instead of  $aRb$ )

Consider the following collection:

$$O = \{\preceq \subseteq A \times A : \preceq \text{ is a partial order on } A\}$$

Clearly,  $(O, \subseteq)$  is a poset and  $\leq \in O$ , we show that every chain in it has an upper bound. Let  $C \subseteq O$  be a chain, clearly  $\bigcup C \subseteq A \times A$ , we show it is a partial order: *reflexive*. Let  $a \in A$ , then for every  $\preceq \in C$  we have  $(a, a) \in \preceq$  (since every partial order is reflexive), therefore  $(a, a) \in \bigcup C$ . *antisymmetric*. Consider  $(a, b), (b, a) \in \bigcup C$ , then for some  $\preceq \in C$  we have  $(a, b) \in \preceq$  and for some  $\preceq' \in C$ ,  $(b, a) \in \preceq'$ . Since  $C$  is a chain we have either  $\preceq \subseteq \preceq'$  or  $\preceq' \subseteq \preceq$ , in either case since they are antisymmetric we get  $a = b$ . *transitive*. Suppose  $(a, b), (b, c) \in \bigcup C$ , then for some  $\preceq, \preceq' \in C$  we have  $(a, b) \in \preceq$  and  $(b, c) \in \preceq'$ , again, since  $C$  is chain one of the orders is superset of another, without loss of generality assume  $\preceq' \subseteq \preceq$ , then by transitivity of  $\preceq$  we get  $(a, c) \in \preceq$ , thus  $(a, c) \in \bigcup C$ . We proved any chain  $C$  of  $O$ ,  $\bigcup C$  is in  $O$ , thus any chain has an upper bound in it. By Zorn's Lemma (see item (e) on page 2)  $O$  has a maximal member, denote it by  $\leq^*$ . Since  $\leq \subseteq \leq^*$  we have if  $a \leq b$  then  $a \leq^* b$ .

Assume that  $\leq^*$  is not a total order, then there are some  $a, b$  such that  $a \not\leq^* b$  and  $b \not\leq^* a$ . Define a relation  $R$  such that  $xRy$  iff  $x \leq^* a$  and  $b \leq^* y$ . You can check that  $\bigcup R$  is partial order on  $A$  (tedious but trivial) and  $\bigcup R \neq \leq^*$  but  $\leq^* \subseteq \bigcup R$ . It contradicts the fact that  $\leq^*$  is a maximal element.

**10.** Let  $S$  be the set of elements  $a$  of  $L$  such that there is no  $a_1 \vee \cdots \vee a_n$  in  $L$  such that  $a = a_1 \vee \cdots \vee a_n$  and each  $a_i$  is join irreducible. This set is non-empty, let  $b$  be a minimal element of it (it exists because  $L$  is finite). Notice that  $b$  is not join irreducible, otherwise we could have  $b = b \vee b$ . Since it is not join-irreducible there must be some element  $c, d \in L$  such that  $b = c \vee d$  but  $b \neq c$  and  $b \neq d$ . On the other hand  $c$  and  $d$  are less than  $b$ , but  $b$  is minimal element so  $c, d \notin S$ , thus we have  $c = a_i \vee \cdots \vee a_n$  and  $d = a'_i \vee \cdots \vee a'_m$  for some  $a_i, i \leq n$  and  $a'_j, j \leq m$  such that each  $a_i$  and  $a_j$  are join irreducible. Now we have  $b = c \vee d = a_i \vee \cdots \vee a_n \vee a'_i \vee \cdots \vee a'_m$  which contradicts the fact that  $b \in S$ .