

Eulerian Polynomials of Digraphs

Sam Spiro, Rutgers University

Joint with Kyle Celano and Nicholas Sieger



Permutation Statistics

Permutation Statistics

A *descent* of a permutation σ on the set $[n] := \{1, 2, \dots, n\}$ is an index $i \in [n-1]$ such that $\sigma(i) > \sigma(i+1)$. An *inversion* is a pair of integers (i, j) with $1 \leq i < j \leq n$ such that $\sigma(i) > \sigma(j)$.

$$\sigma = 23154$$

$$\text{Des}(\sigma) = \{2, 4\}, \text{Inv}(\sigma) = \{(1, 3), (2, 3), (4, 5)\}$$

Permutation Statistics

A *descent* of a permutation σ on the set $[n] := \{1, 2, \dots, n\}$ is an index $i \in [n-1]$ such that $\sigma(i) > \sigma(i+1)$. An *inversion* is a pair of integers (i, j) with $1 \leq i < j \leq n$ such that $\sigma(i) > \sigma(j)$.

$$\sigma = 23154$$

$$\text{Des}(\sigma) = \{2, 4\}, \quad \text{Inv}(\sigma) = \{(1, 3), (2, 3), (4, 5)\}$$

The generating functions



$$A_n(t) = \sum_{\sigma \in \mathfrak{S}_n} t^{\text{des}(\sigma)} \quad M_n(t) = \sum_{\sigma \in \mathfrak{S}_n} t^{\text{inv}(\sigma)}$$

are called Eulerian polynomials and Mahonian polynomials, respectively.

A Common Generalization

A Common Generalization


A *permutation* of an n -vertex digraph $D = (V, E)$ is a bijection $\sigma : V \rightarrow [n]$. A *descent* of such a permutation is an arc $u \rightarrow v$ such that $\sigma(u) > \sigma(v)$.




A Common Generalization

Claim (Foata-Zeilberger)

If $D = \vec{P}_n$, then D -descents are exactly descents. Similarly $D = \vec{K}_n$ corresponds to inversions.



(a) $\text{des}(23154) = 2$




(b) $\text{inv}(23154) = 3$

A Common Generalization

Claim (Foata-Zeilberger)

If $D = \vec{P}_n$, then D -descents are exactly descents. Similarly $D = \vec{K}_n$ corresponds to inversions.



For a digraph $D = (V, E)$, we define

$$A_D(t) = \sum_{\sigma \in \mathfrak{S}_D} t^{\text{des}_D(\sigma)}. \quad (1)$$

In particular, $A_{\vec{P}_n}(t) = A_n(t)$ and $A_{\vec{K}_n}(t) = M_n(t)$.

Main Results

Question

What (if anything) can be said about $A_D(-1)$?

Main Results

Question

What (if anything) can be said about $A_D(-1)$?

$|A_n(-1)| = |A_{\vec{P}_n}(-1)|$ is the number of *alternating permutations*, i.e. those that go

$$\sigma(1) < \sigma(2) > \sigma(3) < \cdots > \sigma(n)$$

Main Results

Question

What (if anything) can be said about $A_D(-1)$?

$|A_n(-1)| = |A_{\vec{P}_n}(-1)|$ is the number of *alternating permutations*, i.e. those that go

$$\sigma(1) < \sigma(2) > \sigma(3) < \cdots > \sigma(n)$$

$|M_n(-1)| = |A_{\vec{K}_n}(-1)|$ also counts something interesting.

Main Results

Question

What (if anything) can be said about $A_D(-1)$?

$|A_n(-1)| = |A_{\vec{P}_n}(-1)|$ is the number of *alternating permutations*, i.e. those that go

$$\sigma(1) < \sigma(2) > \sigma(3) < \cdots > \sigma(n)$$

$|M_n(-1)| = |A_{\vec{K}_n}(-1)|$ also counts something interesting.

and the later being the number of *correct proofs of the Riemann hypothesis*¹.

¹As of the time of writing.

Main Results

Claim

The value of $|A_D(-1)|$ depends only on the underlying graph of G .

Main Results

Claim

The value of $|A_D(-1)|$ depends only on the underlying graph of G .

With this in mind, for any graph G we can define

$$\nu(G) = |A_D(-1)|$$

for any orientation D of G .

Main Results

Claim

The value of $|A_D(-1)|$ depends only on the underlying graph of G .

With this in mind, for any graph G we can define

$$\nu(G) = |A_D(-1)|$$

for any orientation D of G .

Question (Kalai 2002)

What can be said about $\nu(G)$?

Main Results

Claim

The value of $|A_D(-1)|$ depends only on the underlying graph of G .

With this in mind, for any graph G we can define

$$\nu(G) = |A_D(-1)|$$

for any orientation D of G .

Question (Kalai 2002)

What can be said about $\nu(G)$?

For example, $\nu(P_n) = |A_{\vec{P}_n}(-1)|$ is the number of alternating permutations.

Main Results

Definition

Given an n -vertex graph G , we say that an ordering $\pi = (\pi_1, \dots, \pi_n)$ of the vertex set $V(G)$ is an *even sequence* if each of the subgraphs $G[\pi_1, \dots, \pi_i]$ induced by the first i vertices of π have an even number of edges for all $1 \leq i \leq n$.




We claim that $\pi = (3, 1, 2, 5, 4)$ is an even sequence:

Main Results

Definition

Given an n -vertex graph G , we say that an ordering $\pi = (\pi_1, \dots, \pi_n)$ of the vertex set $V(G)$ is an *even sequence* if each of the subgraphs $G[\pi_1, \dots, \pi_i]$ induced by the first i vertices of π have an even number of edges for all $1 \leq i \leq n$.




We claim that $\pi = (3, 1, 2, 5, 4)$ is an even sequence:



Main Results

Definition

Given an n -vertex graph G , we say that an ordering $\pi = (\pi_1, \dots, \pi_n)$ of the vertex set $V(G)$ is an *even sequence* if each of the subgraphs $G[\pi_1, \dots, \pi_i]$ induced by the first i vertices of π have an even number of edges for all $1 \leq i \leq n$.




We claim that $\pi = (3, 1, 2, 5, 4)$ is an even sequence:




Main Results

Definition

Given an n -vertex graph G , we say that an ordering $\pi = (\pi_1, \dots, \pi_n)$ of the vertex set $V(G)$ is an *even sequence* if each of the subgraphs $G[\pi_1, \dots, \pi_i]$ induced by the first i vertices of π have an even number of edges for all $1 \leq i \leq n$.




We claim that $\pi = (3, 1, 2, 5, 4)$ is an even sequence:




Main Results

Definition

Given an n -vertex graph G , we say that an ordering $\pi = (\pi_1, \dots, \pi_n)$ of the vertex set $V(G)$ is an *even sequence* if each of the subgraphs $G[\pi_1, \dots, \pi_i]$ induced by the first i vertices of π have an even number of edges for all $1 \leq i \leq n$.




We claim that $\pi = (3, 1, 2, 5, 4)$ is an even sequence:




Main Results

Definition

Given an n -vertex graph G , we say that an ordering $\pi = (\pi_1, \dots, \pi_n)$ of the vertex set $V(G)$ is an *even sequence* if each of the subgraphs $G[\pi_1, \dots, \pi_i]$ induced by the first i vertices of π have an even number of edges for all $1 \leq i \leq n$.



We claim that $\pi = (3, 1, 2, 5, 4)$ is an even sequence:



Main Results

Let $\eta(G)$ denote the number of even sequences of G .

Main Results

Let $\eta(G)$ denote the number of even sequences of G .

Claim

A permutation π is an even sequence for P_n if and only if π^{-1} is an alternating permutation. In particular, $\nu(P_n) = \eta(P_n)$.

Main Results

Let $\eta(G)$ denote the number of even sequences of G .

Claim

A permutation π is an even sequence for P_n if and only if π^{-1} is an alternating permutation. In particular, $\nu(P_n) = \eta(P_n)$.

Theorem (Celano, Sieger, S. 2023)

We have $\nu(G) = \eta(G)$ whenever G is bipartite

Main Results


Let $\eta(G)$ denote the number of even sequences of G .

Claim

A permutation π is an even sequence for P_n if and only if π^{-1} is an alternating permutation. In particular, $\nu(P_n) = \eta(P_n)$.

Theorem (Celano, Sieger, S. 2023)

We have $\nu(G) = \eta(G)$ whenever G is bipartite, complete multipartite, or a blowup of a cycle.



Main Results

Question

Does every graph satisfy $\nu(G) = \eta(G)$?

Main Results

Question


Does every graph satisfy $\nu(G) = \eta(G)$?

Claim


No!

Main Results

The following graph has $0 = \nu(G) < \eta(G) = 1088$.




(a) C_7^*



(b) An even sequence

Main Results

The following graph has $0 = \nu(G) < \eta(G) = 1088$.



Main Results

Theorem (Celano, Sieger, S. 2023)

We have $\nu(G) = \eta(G)$ whenever G is bipartite, complete multipartite, or a blowup of a cycle.

Main Results

Theorem (Celano, Sieger, S. 2023)

We have $\nu(G) = \eta(G)$ whenever G is bipartite, complete multipartite, or a blowup of a cycle.

Theorem (Celano, Sieger, S. 2023)

If G is a connected graph such that $\nu(G') = \eta(G')$ for all induced subgraphs $G' \subseteq G$, then G is either bipartite, complete multipartite, or a blowup of a cycle.

Extremal Combinatorics!!!!1!!1!!!!

Extremal Combinatorics!!!!1!!1!!!!

Question

What is the maximum/minimum values for $\nu(G)$ (or $\eta(G)$) amongst graphs G with some property?

Extremal Combinatorics!!!!1!!1!!!!

Question



What is the maximum/minimum values for $\nu(G)$ (or $\eta(G)$) amongst graphs G with some property?

Theorem (Celano, Sieger, S. 2023)

If T is a tree on $2n + 1$ vertices, then

$$n!2^n \leq \nu(T) = \eta(T) \leq (2n)!$$

Moreover, equality holds in the lower bound if and only if T is a hairbrush, and equality holds in the upper bound if and only if T is a star.



Main Results

Observe that $\nu(G) = 0$ if and only if -1 is a root of $A_D(t)$.

Main Results

Observe that $\nu(G) = 0$ if and only if -1 is a root of $A_D(t)$.

Question



What can be said about the multiplicity of -1 as a root of $A_D(t)$ in general?

Main Results

Observe that $\nu(G) = 0$ if and only if -1 is a root of $A_D(t)$.

Question

What can be said about the multiplicity of -1 as a root of $A_D(t)$ in general?



$$A_{D_1}(t) = (1+t)^3(1+t+11t^2+t^3+t^4) \quad A_{D_2}(t) = (1+t)(1+5t+16t^2+16t^3+16t^4+5t^5+t^6)$$

Main Results

Theorem (Celano, Sieger, S. 2023)

If D is a tournament on n vertices, then $\text{mult}(A_D(t), -1) = \lfloor \frac{n}{2} \rfloor$.

Main Results

Theorem (Celano, Sieger, S. 2023)

If D is a tournament on n vertices, then $\text{mult}(A_D(t), -1) = \lfloor \frac{n}{2} \rfloor$.

Question

Is $\lfloor \frac{n}{2} \rfloor$ the largest $\text{mult}(A_D(t), -1)$ can be?

Note that tournaments have the largest (potential) degree for $A_D(t)$.

Extremal Combinatorics!!!!1!!!!!!!

Question

Is $\lfloor \frac{n}{2} \rfloor$ the largest $\text{mult}(A_D(t), -1)$ can be?

Extremal Combinatorics!!!!1!!!!!!!

Question

Is $\lfloor \frac{n}{2} \rfloor$ the largest $\text{mult}(A_D(t), -1)$ can be?

Theorem (Celano, Sieger, S. 2023)

If D is an n -vertex digraph, then

$$\text{mult}(A_D(t), -1) \leq n - s_2(n),$$

*where $s_2(n)$ denotes the number of 1's in the binary expansion of n .
Moreover, for all n , there exist n -vertex digraphs D with*

$$A_D(t) = \frac{n!}{2^{n-s_2(n)}} (1+t)^{n-s_2(n)}.$$

The only extremal examples we know are “impartial digraphs”, which is weird.

Proof Ideas

Recall that we want to show $\nu(G) = \eta(G)$ for e.g. bipartite graphs.

Proof Ideas

Recall that we want to show $\nu(G) = \eta(G)$ for e.g. bipartite graphs.

Claim

If G has an odd number of edges, then $\nu(G) = \eta(G) = 0$.

Proof Ideas

Recall that we want to show $\nu(G) = \eta(G)$ for e.g. bipartite graphs.

Claim

If G has an odd number of edges, then $\nu(G) = \eta(G) = 0$.

Claim

If G has an even number of edges, then

$$\eta(G) = \sum_v \eta(G - v).$$

Proof Ideas

Proposition

If G is a graph such that every induced subgraph $G' \subseteq G$ with an even number of edges satisfies

$$\nu(G') = \sum \nu(G' - v),$$

then $\nu(G) = \eta(G)$.

Proof Ideas

Proposition

If G is a graph such that every induced subgraph $G' \subseteq G$ with an even number of edges satisfies

$$\nu(G') = \sum \nu(G' - v),$$

then $\nu(G) = \eta(G)$.

Proof: let G be a smallest counterexample.

Proof Ideas

Proposition

If G is a graph such that every induced subgraph $G' \subseteq G$ with an even number of edges satisfies

$$\nu(G') = \sum \nu(G' - v),$$

then $\nu(G) = \eta(G)$.

Proof: let G be a smallest counterexample. If $e(G)$ is odd then $\nu(G) = \eta(G) = 0$

Proof Ideas

Proposition

If G is a graph such that every induced subgraph $G' \subseteq G$ with an even number of edges satisfies

$$\nu(G') = \sum \nu(G' - v),$$

then $\nu(G) = \eta(G)$.

Proof: let G be a smallest counterexample. If $e(G)$ is odd then $\nu(G) = \eta(G) = 0$, otherwise

$$\nu(G) = \sum \nu(G - v)$$

Proof Ideas

Proposition

If G is a graph such that every induced subgraph $G' \subseteq G$ with an even number of edges satisfies

$$\nu(G') = \sum \nu(G' - v),$$

then $\nu(G) = \eta(G)$.

Proof: let G be a smallest counterexample. If $e(G)$ is odd then $\nu(G) = \eta(G) = 0$, otherwise

$$\nu(G) = \sum \nu(G - v) = \sum \eta(G - v)$$

Proof Ideas

Proposition

If G is a graph such that every induced subgraph $G' \subseteq G$ with an even number of edges satisfies

$$\nu(G') = \sum \nu(G' - v),$$

then $\nu(G) = \eta(G)$.

Proof: let G be a smallest counterexample. If $e(G)$ is odd then $\nu(G) = \eta(G) = 0$, otherwise

$$\nu(G) = \sum \nu(G - v) = \sum \eta(G - v) = \eta(G).$$



Proof Ideas

Corollary

If every bipartite graph with an even number of edges satisfies $\nu(G) = \sum \nu(G - v)$, then $\nu(G) = \eta(G)$ for every bipartite graph.

Proof Ideas

Corollary

If every bipartite graph with an even number of edges satisfies $\nu(G) = \sum \nu(G - v)$, then $\nu(G) = \eta(G)$ for every bipartite graph.

Recall $\nu(G) = |A_D(-1)|$ for any orientation D of G .

Proof Ideas

Corollary

If every bipartite graph with an even number of edges satisfies $\nu(G) = \sum \nu(G - v)$, then $\nu(G) = \eta(G)$ for every bipartite graph.

Recall $\nu(G) = |A_D(-1)|$ for any orientation D of G .

Lemma

For any digraph,

$$A_D(t) = \sum_{v \in V} \frac{t^{\deg_D^+(v)} + t^{\deg_D^-(v)}}{2} A_{D-v}(t)$$

Proof Ideas

Corollary

If every bipartite graph with an even number of edges satisfies $\nu(G) = \sum \nu(G - v)$, then $\nu(G) = \eta(G)$ for every bipartite graph.

Recall $\nu(G) = |A_D(-1)|$ for any orientation D of G .

Lemma

For any digraph,

$$A_D(t) = \sum_{v \in V} \frac{t^{\deg_D^+(v)} + t^{\deg_D^-(v)}}{2} A_{D-v}(t)$$

The corollary will thus apply if we can show for any bipartite graph G , any/some orientation D has the same sign for all of the terms above.

Proof Ideas


Claim

There exists a “natural” orientation D for each bipartite graph G which makes it easy to predict the sign of $A_D(t)$.

Proof Ideas

Claim


There exists a “natural” orientation D for each bipartite graph G which makes it easy to predict the sign of $A_D(t)$.



Proof Ideas

Claim

There exists a “natural” orientation D for each bipartite graph G which makes it easy to predict the sign of $A_D(t)$.




Lemma

For such a digraph we have $A_D(-1) \geq 0$.

Proof Ideas

Claim

There exists a “natural” orientation D for each bipartite graph G which makes it easy to predict the sign of $A_D(t)$.



Lemma


For such a digraph we have $A_D(-1) \geq 0$. In particular, when $e(G)$ is even we have

$$\nu(G) = A_D(-1) = \sum_{v \in V(D)} A_{D-v}(-1) = \sum \nu(G - v).$$

Proof Ideas

Claim


There exists “natural” orientations for complete multipartite graphs/blowup of a cycles which makes it easy to predict the sign of $A_D(t)$.



Proof Ideas

Claim

There exists “natural” orientations for complete multipartite graphs/blowup of a cycles which makes it easy to predict the sign of $A_D(t)$.



Theorem (Celano, Sieger, S. 2023)

We have $\nu(G) = \eta(G)$ whenever G is bipartite, complete multipartite, or a blowup of a cycle.

Proof Ideas


Theorem (Celano, Sieger, S. 2023)

If G is a connected graph such that $\nu(G') = \eta(G')$ for all induced subgraphs $G' \subseteq G$, then G is either bipartite, complete multipartite, or a blowup of a cycle.

Proof Ideas

Theorem (Celano, Sieger, S. 2023)

If G is a connected graph such that $\nu(G') = \eta(G')$ for all induced subgraphs $G' \subseteq G$, then G is either bipartite, complete multipartite, or a blowup of a cycle.



Proposition (Celano, Sieger, S. 2023)

If G is a connected graph, then G is induced odd pan-free if and only if it is either bipartite, complete multipartite, or a blowup of a cycle.

Open Problems

Open Problems

Question

Can one give a combinatorial interpretation for $\nu(G)$ for arbitrary graphs G ?

Open Problems

Question

Can one give a combinatorial interpretation for $\nu(G)$ for arbitrary graphs G ?

Note that $\nu(G) \leq \eta(G)$ in general, so perhaps this is some special subset of even sequences.

Open Problems

Question

Can one give a combinatorial interpretation for $\nu(G)$ for arbitrary graphs G ?

Note that $\nu(G) \leq \eta(G)$ in general, so perhaps this is some special subset of even sequences.

Conjecture

If G is an Eulerian graph, then $\nu(G) = \sum_v \nu(G - v)$.

Open Problems

We gave a non-combinatorial proof that $\nu(G) = \eta(G)$ for all bipartite graphs G .

Open Problems

We gave a non-combinatorial proof that $\nu(G) = \eta(G)$ for all bipartite graphs G .

Problem

For any bipartite graph $G = ([n], E)$ and orientation D of G , construct an explicit involution $\phi : \mathfrak{S}_n \rightarrow \mathfrak{S}_n$ such that

- (a) The set of fixed points \mathcal{F}_ϕ of ϕ is the set of (inverses of) even sequences of G , and
- (b) $(-1)^{\text{des}_D(\sigma)} = -(-1)^{\text{des}_D(\phi(\sigma))}$ for all $\sigma \notin \mathcal{F}_\phi$.

Open Problems

Conjecture

If D is the orientation of a complete multipartite graph which has r parts of odd size, then $\text{mult}(A_D(t), -1) = \lfloor \frac{r}{2} \rfloor$.

Open Problems

Conjecture

If D is the orientation of a complete multipartite graph which has r parts of odd size, then $\text{mult}(A_D(t), -1) = \lfloor \frac{r}{2} \rfloor$.

It is easy to show that $\text{mult}(A_D(t), 0)$ corresponds to the minimum number of arcs one must reverse in order for D to be acyclic.

Open Problems

Conjecture

If D is the orientation of a complete multipartite graph which has r parts of odd size, then $\text{mult}(A_D(t), -1) = \lfloor \frac{r}{2} \rfloor$.

It is easy to show that $\text{mult}(A_D(t), 0)$ corresponds to the minimum number of arcs one must reverse in order for D to be acyclic.

Question

Does there exist a digraph D such that $A_D(t)$ has an integral root which is not equal to either 0 or -1 ?

No such digraph exists on at most 5 vertices, and there exist digraphs with real roots of magnitude larger than 2.