

# Card Guessing with Feedback

Sam Spiro, Rutgers University.

# Feedback Models

# Feedback Models

Start with a deck of  $mn$  cards where there are  $n$  card types each appearing with multiplicity  $m$ .

# Feedback Models

Start with a deck of  $mn$  cards where there are  $n$  card types each appearing with multiplicity  $m$ . For example,  $n = 13$  and  $m = 4$  corresponds to a usual deck of playing cards.

# Feedback Models

Let  $\mathcal{C}_{m,n}^+$  and  $\mathcal{C}_{m,n}^-$  be the maximum and minimum expected scores that the player can get in the complete feedback model.

# Feedback Models

Let  $\mathcal{C}_{m,n}^+$  and  $\mathcal{C}_{m,n}^-$  be the maximum and minimum expected scores that the player can get in the complete feedback model.

Theorem (Diaconis-Graham, 1981)

*For  $n$  fixed,*

$$\mathcal{C}_{m,n}^{\pm} = m \pm c_n \sqrt{m} + o_n(\sqrt{m}).$$

# Feedback Models

# Feedback Models

In the “partial feedback model”, the Guesser guesses the next card and is only told whether their guess was correct or not.



# Feedback Models

In the “partial feedback model”, the Guesser guesses the next card and is only told whether their guess was correct or not. If  $\mathcal{P}_{m,n}^+$  is the maximum expected score in this model,

$$m \leq \mathcal{P}_{m,n}^+$$

# Feedback Models

In the “partial feedback model”, the Guesser guesses the next card and is only told whether their guess was correct or not. If  $\mathcal{P}_{m,n}^+$  is the maximum expected score in this model,

$$m \leq \mathcal{P}_{m,n}^+ \leq \mathcal{C}_{m,n}^+$$

# Feedback Models

In the “partial feedback model”, the Guesser guesses the next card and is only told whether their guess was correct or not. If  $\mathcal{P}_{m,n}^+$  is the maximum expected score in this model,

$$m \leq \mathcal{P}_{m,n}^+ \leq \mathcal{C}_{m,n}^+ = m + o_n(m)$$

# Feedback Models

In the “partial feedback model”, the Guesser guesses the next card and is only told whether their guess was correct or not. If  $\mathcal{P}_{m,n}^+$  is the maximum expected score in this model,

$$m \leq \mathcal{P}_{m,n}^+ \leq \mathcal{C}_{m,n}^+ = m + o_n(m),$$

What happens when  $n$  is large?

# Feedback Models

## Theorem (Diaconis-Graham-He-S., 2020)

*For  $m$  fixed,*

$$\mathcal{C}_{m,n}^+ \sim H_m \log(n),$$

$$\mathcal{C}_{m,n}^- = \Theta(n^{-1/m}),$$

*where  $H_m$  is the  $m$ th harmonic number.*

# Feedback Models

## Theorem (Diaconis-Graham-He-S., 2020)

*For  $m$  fixed,*

$$\mathcal{C}_{m,n}^+ \sim H_m \log(n),$$

$$\mathcal{C}_{m,n}^- = \Theta(n^{-1/m}),$$

*where  $H_m$  is the  $m$ th harmonic number.*

With this we have the trivial bounds

$$m \leq \mathcal{P}_{m,n}^+ \leq \mathcal{C}_{m,n}^+ = O_m(\log n).$$

# Feedback Models

$$m \leq \mathcal{P}_{m,n}^+ \leq O_m(\log n).$$

# Feedback Models

$$m \leq \mathcal{P}_{m,n}^+ \leq O_m(\log n).$$

Theorem (Diaconis-Graham-He-S., 2020+)

*There exist  $c, C > 0$  such that if  $n$  is sufficiently large in terms of  $m$ , we have*

$$m + c\sqrt{m} \leq \mathcal{P}_{m,n}^+$$



# Feedback Models

$$m \leq \mathcal{P}_{m,n}^+ \leq O_m(\log n).$$

Theorem (Diaconis-Graham-He-S., 2020+)

*There exist  $c, C > 0$  such that if  $n$  is sufficiently large in terms of  $m$ , we have*

$$m + c\sqrt{m} \leq \mathcal{P}_{m,n}^+ \leq m + Cm^{3/4} \log m.$$

# A Partial Proof

## Lemma

*Assume that we have played in the partial feedback model for  $t - 1$  rounds such that we have guessed card type  $i$  a total of  $g_i$  times*

# A Partial Proof

## Lemma

*Assume that we have played in the partial feedback model for  $t - 1$  rounds such that we have guessed card type  $i$  a total of  $g_i$  times, and let  $S$  be the total number of points scored.*

# A Partial Proof

## Lemma

*Assume that we have played in the partial feedback model for  $t - 1$  rounds such that we have guessed card type  $i$  a total of  $g_i$  times, and let  $S$  be the total number of points scored. Given this, we have*

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S}.$$

# A Partial Proof

## Lemma

*Assume that we have played in the partial feedback model for  $t - 1$  rounds such that we have guessed card type  $i$  a total of  $g_i$  times, and let  $S$  be the total number of points scored. Given this, we have*

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S}.$$

That is, our upper bound is strongest when  $g_i$  and  $S$  is small.

# A Partial Proof

## Lemma

*Assume that we have played in the partial feedback model for  $t - 1$  rounds such that we have guessed card type  $i$  a total of  $g_i$  times, and let  $S$  be the total number of points scored. Given this, we have*

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S}.$$

That is, our upper bound is strongest when  $g_i$  and  $S$  is small.  
These conditions are necessary

# A Partial Proof

## Lemma

*Assume that we have played in the partial feedback model for  $t - 1$  rounds such that we have guessed card type  $i$  a total of  $g_i$  times, and let  $S$  be the total number of points scored. Given this, we have*

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S}.$$

That is, our upper bound is strongest when  $g_i$  and  $S$  is small. These conditions are necessary: if  $i$  has been guessed incorrectly  $g_i = mn - m$  times, then we know the card must be an  $i$ .

# A Partial Proof

## Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$



# A Partial Proof

## Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all  $i$  and  $t$ , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S}$$

# A Partial Proof

## Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all  $i$  and  $t$ , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S} \approx \frac{m}{mn - g_i},$$

# A Partial Proof

## Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all  $i$  and  $t$ , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S} \approx \frac{m}{mn - g_i},$$

At most one  $i$  is guessed more than  $mn/2$  times.

# A Partial Proof

## Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all  $i$  and  $t$ , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S} \approx \frac{m}{mn - g_i},$$

At most one  $i$  is guessed more than  $mn/2$  times. Every other  $j$  has  $\Pr[\pi_t = j] \leq \frac{2}{n}$  for all  $t$ .

# A Partial Proof

## Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all  $i$  and  $t$ , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S} \approx \frac{m}{mn - g_i},$$

At most one  $i$  is guessed more than  $mn/2$  times. Every other  $j$  has  $\Pr[\pi_t = j] \leq \frac{2}{n}$  for all  $t$ . Thus in expectation at most  $mn \cdot (2/n) = 2m$  cards are guessed correctly from this part

# A Partial Proof

## Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all  $i$  and  $t$ , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S} \approx \frac{m}{mn - g_i},$$

At most one  $i$  is guessed more than  $mn/2$  times. Every other  $j$  has  $\Pr[\pi_t = j] \leq \frac{2}{n}$  for all  $t$ . Thus in expectation at most  $mn \cdot (2/n) = 2m$  cards are guessed correctly from this part, and in total at most  $3m$  are guessed correctly in expectation.  $\square$

# Open Problems

Theorem (Diaconis-Graham-He-S., 2020)

$$m + \Omega(\sqrt{m}) \leq \mathcal{P}_{m,n}^+ \leq m + O(m^{3/4} \log m).$$

# Open Problems

Theorem (Diaconis-Graham-He-S., 2020)

$$m + \Omega(\sqrt{m}) \leq \mathcal{P}_{m,n}^+ \leq m + O(m^{3/4} \log m).$$

Conjecture (Diaconis-Graham-He-S., 2020)

$$\mathcal{P}_{m,n}^+ = m + m^{1/2+o(1)}.$$



# Open Problems

Recall that  $\mathcal{P}_{m,n}^-$  is the minimum number of points one can get in expectation with partial feedback.

# Open Problems

Recall that  $\mathcal{P}_{m,n}^-$  is the minimum number of points one can get in expectation with partial feedback. We trivially have

$$m \geq \mathcal{P}_{m,n}^- \geq \mathcal{C}_{m,n}^- = \Theta(n^{-1/m}).$$

# Open Problems

Recall that  $\mathcal{P}_{m,n}^-$  is the minimum number of points one can get in expectation with partial feedback. We trivially have

$$m \geq \mathcal{P}_{m,n}^- \geq \mathcal{C}_{m,n}^- = \Theta(n^{-1/m}).$$

Theorem (Diaconis-Graham-S., 2020)

$$\mathcal{P}_{m,n}^- \geq 1 - e^{-m} - o_m(1) \geq \frac{1}{2}.$$

# Open Problems

Theorem (Diaconis-Graham-S., 2020)

$$\mathcal{P}_{m,n}^- \geq 1 - e^{-m} - o_m(1) \geq \frac{1}{2}.$$

# Open Problems

Theorem (Diaconis-Graham-S., 2020)

$$\mathcal{P}_{m,n}^- \geq 1 - e^{-m} - o_m(1) \geq \frac{1}{2}.$$

Conjecture (Diaconis-Graham-S., 2020)

*If  $n$  is sufficiently large in terms of  $m$ , then*

$$\mathcal{P}_{m,n}^- \sim m.$$

# Practical Strategies

# Practical Strategies

Theorem (Diaconis-Graham-He-S., 2020)

*There exists a simple strategy showing*

$$\mathcal{P}_{m,n}^+ \geq m + \Omega(\sqrt{m}).$$

# Practical Strategies

Theorem (Diaconis-Graham-He-S., 2020)

*There exists a simple strategy showing*

$$\mathcal{P}_{m,n}^+ \geq m + \Omega(\sqrt{m}).$$

Guess 1 a total of  $mn/2$  times, then do one of two things:



# Practical Strategies

Theorem (Diaconis-Graham-He-S., 2020)

*There exists a simple strategy showing*

$$\mathcal{P}_{m,n}^+ \geq m + \Omega(\sqrt{m}).$$

Guess 1 a total of  $mn/2$  times, then do one of two things:

- (1) If you made less than  $m/2$  correct guesses, guess 1 the rest of the game.

# Practical Strategies

Theorem (Diaconis-Graham-He-S., 2020)

*There exists a simple strategy showing*

$$\mathcal{P}_{m,n}^+ \geq m + \Omega(\sqrt{m}).$$

Guess 1 a total of  $mn/2$  times, then do one of two things:

- (1) If you made less than  $m/2$  correct guesses, guess 1 the rest of the game.
- (2) Else guess 2 the rest of the game.



# Practical Strategies

A very simple strategy is the *safe strategy*, which guesses 1 until  $m$  correct guesses are made, then 2 until  $m$  correct guesses are made, and so on.

# Practical Strategies

A very simple strategy is the *safe strategy*, which guesses 1 until  $m$  correct guesses are made, then 2 until  $m$  correct guesses are made, and so on. Elementary arguments give that the score for this strategy is

$$m + 1 - \frac{1}{m+1} + O(e^{-\beta m})$$

for some  $\beta > 0$ .

# Practical Strategies

A very simple strategy is the *safe strategy*, which guesses 1 until  $m$  correct guesses are made, then 2 until  $m$  correct guesses are made, and so on. Elementary arguments give that the score for this strategy is

$$m + 1 - \frac{1}{m + 1} + O(e^{-\beta m})$$

for some  $\beta > 0$ .

Another simple strategy is the *shifting strategy*, which guesses 1 until a correct guess is made, then 2 until a correct guess is made, and so on.

# Practical Strategies

If  $\pi$  is a word where each letter in  $\{1, 2, \dots, n\}$  exactly  $m$  times, we define  $L(\pi)$  to be the largest integer  $p$  so that  $\pi$  contains a subsequence of the form  $123 \cdots p$ .

# Practical Strategies

If  $\pi$  is a word where each letter in  $\{1, 2, \dots, n\}$  exactly  $m$  times, we define  $L(\pi)$  to be the largest integer  $p$  so that  $\pi$  contains a subsequence of the form  $123 \cdots p$ . For example,

$$\pi = 2345124351 \implies L(\pi) = 3.$$

# Practical Strategies

If  $\pi$  is a word where each letter in  $\{1, 2, \dots, n\}$  exactly  $m$  times, we define  $L(\pi)$  to be the largest integer  $p$  so that  $\pi$  contains a subsequence of the form  $123 \cdots p$ . For example,

$$\pi = 2345124351 \implies L(\pi) = 3.$$

Note that  $L(\pi)$  is (essentially) the score one gets using the shifting strategy if the deck is shuffled according to  $\pi$ .



# Practical Strategies

If  $\pi$  is a word where each letter in  $\{1, 2, \dots, n\}$  exactly  $m$  times, we define  $L(\pi)$  to be the largest integer  $p$  so that  $\pi$  contains a subsequence of the form  $123 \cdots p$ . For example,

$$\pi = 2345124351 \implies L(\pi) = 3.$$

Note that  $L(\pi)$  is (essentially) the score one gets using the shifting strategy if the deck is shuffled according to  $\pi$ .

## Corollary

*If  $n$  is sufficiently large in terms of  $m$ , then*

$$\mathcal{L}_{m,n} := \mathbb{E}[L(\pi)] \leq m + O(m^{3/4} \log m).$$

# Practical Strategies

Conjecture (Diaconis-Graham-He-S., 2020)

*If  $n$  is sufficiently large in terms of  $m$ , then*

$$\mathcal{L}_{m,n} \sim m.$$

# Practical Strategies

Conjecture (Diaconis-Graham-He-S., 2020)

*If  $n$  is sufficiently large in terms of  $m$ , then*

$$\mathcal{L}_{m,n} \sim m.$$

Theorem (Clifton-Deb-Huang-S.-Yoo, 2021)

*We have*

$$\left| \lim_{n \rightarrow \infty} \mathcal{L}_{m,n} - \left( m + 1 - \frac{1}{m+2} \right) \right| \leq O(e^{-\beta m})$$

*for some  $\beta > 0$ .*

# Practical Strategies

More precisely: if  $\alpha_1, \dots, \alpha_m$  are the zeroes of  $\sum_{k=0}^m \frac{x^k}{k!}$ , then

$$\lim_{n \rightarrow \infty} \mathcal{L}_{m,n} = -1 - \sum \alpha_i^{-1} e^{-\alpha_i}.$$

# Practical Strategies

More precisely: if  $\alpha_1, \dots, \alpha_m$  are the zeroes of  $\sum_{k=0}^m \frac{x^k}{k!}$ , then

$$\lim_{n \rightarrow \infty} \mathcal{L}_{m,n} = -1 - \sum \alpha_i^{-1} e^{-\alpha_i}.$$

This implies  $\mathcal{L}_{1,n} \rightarrow e - 1$

# Practical Strategies

More precisely: if  $\alpha_1, \dots, \alpha_m$  are the zeroes of  $\sum_{k=0}^m \frac{x^k}{k!}$ , then

$$\lim_{n \rightarrow \infty} \mathcal{L}_{m,n} = -1 - \sum \alpha_i^{-1} e^{-\alpha_i}.$$

This implies  $\mathcal{L}_{1,n} \rightarrow e - 1$  and that

$$\mathcal{L}_{2,n} \rightarrow e(\cos(1) + \sin(1)) - 1.$$

# Card Guessing

# Card Guessing

The probability of drawing four aces in a row with a deck shuffled uniformly at random is  $1/270725$ .



# Adversarial Card Guessing

The probability of drawing four aces in a row with a deck shuffled uniformly at random is  $1/270725$ .

# Adversarial Card Guessing

The probability of drawing four aces in a row with a deck shuffled uniformly at random is  $1/270725$ .

More precisely, we are now considering a two player game played by Shuffler and Guesser.

# Adversarial Card Guessing

The probability of drawing four aces in a row with a deck shuffled uniformly at random is  $1/270725$ .

More precisely, we are now considering a two player game played by Shuffler and Guesser. Let  $\mathcal{C}_{m,n}(G, S)$  be the expected number of points Guesser scores when the two players follow strategies  $G, S$ .

# Adversarial Card Guessing

$$\Theta_m(n^{-1/m}) \leq \mathcal{C}_{m,n}(G, \text{Uniform}) \leq H_m \log n + o_m(\log n).$$

# Adversarial Card Guessing

$$\Theta_m(n^{-1/m}) \leq \mathcal{C}_{m,n}(G, \text{Uniform}) \leq H_m \log n + o_m(\log n).$$

Theorem (S., 2021)

*There exists a strategy  $S'$  for Shuffler so that*

$$\mathcal{C}_{m,n}(G, S') \leq \log n + o_m(\log n),$$

*and this bound is best possible.*

# Adversarial Card Guessing

$$\Theta_m(n^{-1/m}) \leq \mathcal{C}_{m,n}(G, \text{Uniform}) \leq H_m \log n + o_m(\log n).$$

## Theorem (S., 2021)

*There exists a strategy  $S'$  for Shuffler so that*

$$\mathcal{C}_{m,n}(G, S') \leq \log n + o_m(\log n),$$

*and this bound is best possible.*

This theorem is a first for me, since normally I prove a result, then makes jokes about it during my talk.

# Adversarial Card Guessing

## Theorem

*There exists a strategy  $S'$  for Shuffler so that*  
$$\mathcal{C}_{m,n}(G, S') \leq \log n + o_m(\log n).$$

# Adversarial Card Guessing

## Theorem

*There exists a strategy  $S'$  for Shuffler so that*  
$$\mathcal{C}_{m,n}(G, S') \leq \log n + o_m(\log n).$$

A strategy that gives this is the “greedy strategy”, which is such that if there are  $r$  types of cards remaining in the deck, then Shuffler draws each of these card types with probability  $r^{-1}$  (regardless of how many copies are left in the deck of each type).



# Adversarial Card Guessing

## Theorem

*There exists a strategy  $S'$  for Shuffler so that*  
$$\mathcal{C}_{m,n}(G, S') \leq \log n + o_m(\log n).$$

A strategy that gives this is the “greedy strategy”, which is such that if there are  $r$  types of cards remaining in the deck, then Shuffler draws each of these card types with probability  $r^{-1}$  (regardless of how many copies are left in the deck of each type). E.g. if the deck has a hundred 1's and one 2, we draw a 1 or 2 with probability  $\frac{1}{2}$ .

# Adversarial Card Guessing

## Theorem

*There exists a strategy  $S'$  for Shuffler so that*  
$$C_{m,n}(G, S') \leq \log n + o_m(\log n).$$

A strategy that gives this is the “greedy strategy”, which is such that if there are  $r$  types of cards remaining in the deck, then Shuffler draws each of these card types with probability  $r^{-1}$  (regardless of how many copies are left in the deck of each type). E.g. if the deck has a hundred 1's and one 2, we draw a 1 or 2 with probability  $\frac{1}{2}$ . This gives the desired bound due to a variant of the coupon collector problem. □

# Adversarial Card Guessing

## Theorem (S., 2021)

*The greedy strategy is the unique strategy that minimizes the number of correct guesses if Guesser tries to maximize their score.*

# Adversarial Card Guessing

## Theorem (S., 2021)

*The greedy strategy is the unique strategy that minimizes the number of correct guesses if Guesser tries to maximize their score.*

Interestingly, the greedy strategy is also the “unique” strategy which maximizes the number of correct guesses if Guesser tries to minimize their score.

# Semi-restricted Games

There is a classical game called “Matching Pennies” where two players simultaneously choose one of  $n$  numbers, and if the two match player A gets a point and otherwise player B gets a point.

# Semi-restricted Games

There is a classical game called “Matching Pennies” where two players simultaneously choose one of  $n$  numbers, and if the two match player A gets a point and otherwise player B gets a point.

The adversarial card guessing game can be viewed as a “semi-restricted” version of this game where  $mn$  rounds of Matching Pennies is played and player B must use each number exactly  $m$  times.

# Semi-restricted Games

More generally, one can consider “semi-restricted” versions of any zero sum game.

# Semi-restricted Games

More generally, one can consider “semi-restricted” versions of any zero sum game.

Theorem (S.-Surya-Zeng, 2022)

*In semi-restricted Rock, Paper, Scissors the “greedy strategy” is the unique optimal strategy for the restricted player.*



# Semi-restricted Games

More generally, one can consider “semi-restricted” versions of any zero sum game.

Theorem (S.-Surya-Zeng, 2022)


*In semi-restricted Rock, Paper, Scissors the “greedy strategy” is the unique optimal strategy for the restricted player.*

Theorem (S.-Surya-Zeng, 2022)

*“Almost every” semi-restricted game fails to have an optimal strategy which is greedy.*


# Semi-restricted Games

Given a digraph  $D$ , we define its skew adjacency matrix  $A$  by  $A_{u,v} = +1$  if  $u \rightarrow v$ ,  $A_{u,v} = -1$  if  $v \rightarrow u$ , and  $A_{u,v} = 0$  otherwise.



# Semi-restricted Games

Given a digraph  $D$ , we define its skew adjacency matrix  $A$  by  $A_{u,v} = +1$  if  $u \rightarrow v$ ,  $A_{u,v} = -1$  if  $v \rightarrow u$ , and  $A_{u,v} = 0$  otherwise.



## Question

Which digraphs  $D$  are such that their skew-adjacency matrix  $A$  satisfies  $\text{Null}(A) = \text{span}(\vec{1})$ ?