

Slow Fibonacci Walks

Sam Spiro, UC San Diego.

4/22/20

Joint with Fan Chung and Ron Graham.

Slow Fibonacci Walks

Slow Fibonacci Walks



169



206



I am interested in magic tricks whose explanation requires deep mathematics. The trick should be one that would actually appeal to a layman. An example is the following: the magician asks Alice to choose two integers between 1 and 50 and add them. Then add the largest two of the three integers at hand. Then add the largest two again. Repeat this around ten times. Alice tells the magician her final number n . The magician then tells Alice the next number. This is done by computing $(1.61803398 \dots)n$ and rounding to the nearest integer. The explanation is beyond the comprehension of a random mathematical layperson, but for a mathematician it is not very deep. Can anyone do better?

soft-question

big-list

popularization

[share](#) [cite](#) [edit](#) [follow](#) [flag](#)

edited Jun 8 '17 at 9:58

community wiki

6 revs, 5 users 75%

[Richard Stanley](#)

Slow Fibonacci Walks

Given positive integers a_1, a_2 , we define the (a_1, a_2) -Fibonacci walk to be the sequence $w_k = w_k(a_1, a_2)$ satisfying

$$w_1 = a_1, \quad w_2 = a_2, \quad w_{k+2} = w_{k+1} + w_k.$$

Slow Fibonacci Walks

Given positive integers a_1, a_2 , we define the (a_1, a_2) -Fibonacci walk to be the sequence $w_k = w_k(a_1, a_2)$ satisfying

$$w_1 = a_1, w_2 = a_2, w_{k+2} = w_{k+1} + w_k.$$

For example, if $w_k = w_k(10, 2)$, this gives the sequence

$$10, 2, 12, 14, 26, 40, 66 \dots$$

Slow Fibonacci Walks

Given positive integers a_1, a_2 , we define the (a_1, a_2) -Fibonacci walk to be the sequence $w_k = w_k(a_1, a_2)$ satisfying

$$w_1 = a_1, w_2 = a_2, w_{k+2} = w_{k+1} + w_k.$$

For example, if $w_k = w_k(10, 2)$, this gives the sequence

$$10, 2, 12, 14, 26, 40, 66 \dots$$

We say that w_k is an n -Fibonacci walk if $w_s = n$ for some s . For example, the above w_k is a 40-Fibonacci walk.

Slow Fibonacci Walks

Does there exist an n -Fibonacci walk for all n ?

Slow Fibonacci Walks

Does there exist an n -Fibonacci walk for all n ? Yes, $w_k(x, n)$ is an n -Fibonacci walk for all x , which is kind of boring.

Slow Fibonacci Walks

Does there exist an n -Fibonacci walk for all n ? Yes, $w_k(x, n)$ is an n -Fibonacci walk for all x , which is kind of boring. To make things more interesting, we say that a sequence w_k is an n -slow Fibonacci walk if $w_s = n$ and s is as large as possible.

Slow Fibonacci Walks

Does there exist an n -Fibonacci walk for all n ? Yes, $w_k(x, n)$ is an n -Fibonacci walk for all x , which is kind of boring. To make things more interesting, we say that a sequence w_k is an n -slow Fibonacci walk if $w_s = n$ and s is as large as possible.

For example, the following are all 40-Fibonacci walks.

1024, 40, 1064...

8, 8, 16, 24, 40, 64...

5, 5, 10, 15, 25, 40, 65...

10, 2, 12, 14, 26, 40, 66...

Slow Fibonacci Walks

Does there exist an n -Fibonacci walk for all n ? Yes, $w_k(x, n)$ is an n -Fibonacci walk for all x , which is kind of boring. To make things more interesting, we say that a sequence w_k is an n -slow Fibonacci walk if $w_s = n$ and s is as large as possible.

For example, the following are all 40-Fibonacci walks.

1024, 40, 1064...

8, 8, 16, 24, 40, 64...

5, 5, 10, 15, 25, 40, 65...

10, 2, 12, 14, 26, 40, 66...

However, the first two can't be slow (since the next two achieve 40 with $s = 6$), and one can verify that $w_k(5, 5)$ and $w_k(10, 2)$ are (the unique) 40-slow Fibonacci walks.

Slow Fibonacci Walks

We will say that a pair of positive integers (b, a) is n -good if $w_k(b, a)$ is an n -slow Fibonacci walks.

Slow Fibonacci Walks

We will say that a pair of positive integers (b, a) is n -good if $w_k(b, a)$ is an n -slow Fibonacci walks. Thus understanding n -good pairs is equivalent to understanding slow Fibonacci walks. So what can we say about these pairs?

Slow Fibonacci Walks

We will say that a pair of positive integers (b, a) is n -good if $w_k(b, a)$ is an n -slow Fibonacci walk. Thus understanding n -good pairs is equivalent to understanding slow Fibonacci walks. So what can we say about these pairs? Define $s(n)$ to be the length of any n -slow walk.

Slow Fibonacci Walks

We will say that a pair of positive integers (b, a) is n -good if $w_k(b, a)$ is an n -slow Fibonacci walk. Thus understanding n -good pairs is equivalent to understanding slow Fibonacci walks. So what can we say about these pairs? Define $s(n)$ to be the length of any n -slow walk.

Lemma

$s(n) = 2$ iff $n = 1$, in which case $(x, 1)$ is a 1-good pair for all x .

Slow Fibonacci Walks

We will say that a pair of positive integers (b, a) is n -good if $w_k(b, a)$ is an n -slow Fibonacci walk. Thus understanding n -good pairs is equivalent to understanding slow Fibonacci walks. So what can we say about these pairs? Define $s(n)$ to be the length of any n -slow walk.

Lemma

$s(n) = 2$ iff $n = 1$, in which case $(x, 1)$ is a 1-good pair for all x .

$$w_3(x, y) = x + y \geq 2, w_3(1, n - 1) = n.$$



Slow Fibonacci Walks

Lemma

Assume $s(n) = s > 2$. If (b, a) is n -good, then $a \leq b$.

Slow Fibonacci Walks

Lemma

Assume $s(n) = s > 2$. If (b, a) is n -good, then $a \leq b$.

E.g. instead of

4, 10, 14, ...

Slow Fibonacci Walks

Lemma

Assume $s(n) = s > 2$. If (b, a) is n -good, then $a \leq b$.

E.g. instead of

4, 10, 14, ...

we should start

6, 4, 10, 14, ...

Slow Fibonacci Walks

Lemma

$w_k(b, a) = af_{k-1} + bf_{k-2}$, with f_k the Fibonacci numbers.

Slow Fibonacci Walks

Lemma

$w_k(b, a) = af_{k-1} + bf_{k-2}$, with f_k the Fibonacci numbers.

Lemma

Let $s = s(n) > 2$. If (b, a) is n -good, then (b', a') is n -good iff $a' = a + kf_{s-2} \geq 1$ and $b' = b - kf_{s-1} \geq 1$ for some k .

Slow Fibonacci Walks

Lemma

$w_k(b, a) = af_{k-1} + bf_{k-2}$, with f_k the Fibonacci numbers.

Lemma

Let $s = s(n) > 2$. If (b, a) is n -good, then (b', a') is n -good iff $a' = a + kf_{s-2} \geq 1$ and $b' = b - kf_{s-1} \geq 1$ for some k .

For example, if we know $s(40) = 6$ and $(10, 2)$ is 40-good, then so is $(10 - kf_5, 2 + kf_4) = (10 - 5k, 2 + 3k)$, which only makes sense if $k = 0, 1$.

Slow Fibonacci Walks

Lemma

$w_k(b, a) = af_{k-1} + bf_{k-2}$, with f_k the Fibonacci numbers.

Lemma

Let $s = s(n) > 2$. If (b, a) is n -good, then (b', a') is n -good iff $a' = a + kf_{s-2} \geq 1$ and $b' = b - kf_{s-1} \geq 1$ for some k .

For example, if we know $s(40) = 6$ and $(10, 2)$ is 40-good, then so is $(10 - kf_5, 2 + kf_4) = (10 - 5k, 2 + 3k)$, which only makes sense if $k = 0, 1$.

Proof.

By the above lemma, every n -good pair is a solution to the diophantine equation $n = w_s(b', a') = a'f_{s-1} + b'f_{s-2}$, and the result follows since $\gcd(f_{s-1}, f_{s-2}) = 1$ for $s > 2$. □

Slow Fibonacci Walks

Theorem (Englund, Bicknell-Johnson (1997); Jones, Kiss (1998); Chung, Graham, S. (2019))

For $n > 1$ with $s = s(n)$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = af_{s-1} + bf_{s-2}$ and $1 \leq a \leq b \leq f_{s-1}$. In this case (b, a) is n -good.

Slow Fibonacci Walks

Theorem (Englund, Bicknell-Johnson (1997); Jones, Kiss (1998); Chung, Graham, S. (2019))

For $n > 1$ with $s = s(n)$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = af_{s-1} + bf_{s-2}$ and $1 \leq a \leq b \leq f_{s-1}$. In this case (b, a) is n -good.

Because $w_k(x, y) = yf_{k-1} + xf_{k-2}$, by definition of s there exist a', b' such that $n = w_s(b', a') = a'f_{s-1} + b'f_{s-2}$.

Slow Fibonacci Walks

Theorem (Englund, Bicknell-Johnson (1997); Jones, Kiss (1998); Chung, Graham, S. (2019))

For $n > 1$ with $s = s(n)$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = af_{s-1} + bf_{s-2}$ and $1 \leq a \leq b \leq f_{s-1}$. In this case (b, a) is n -good.

Because $w_k(x, y) = yf_{k-1} + xf_{k-2}$, by definition of s there exist a', b' such that $n = w_s(b', a') = a'f_{s-1} + b'f_{s-2}$. Let k be such that $1 \leq b' - kf_{s-1} \leq f_{s-1}$.

Slow Fibonacci Walks

Theorem (Englund, Bicknell-Johnson (1997); Jones, Kiss (1998); Chung, Graham, S. (2019))

For $n > 1$ with $s = s(n)$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = af_{s-1} + bf_{s-2}$ and $1 \leq a \leq b \leq f_{s-1}$. In this case (b, a) is n -good.

Because $w_k(x, y) = yf_{k-1} + xf_{k-2}$, by definition of s there exist a', b' such that $n = w_s(b', a') = a'f_{s-1} + b'f_{s-2}$. Let k be such that $1 \leq b' - kf_{s-1} \leq f_{s-1}$. Then $(b, a) := (b' - kf_{s-1}, a' + kf_{s-2})$ is n -good.

Slow Fibonacci Walks

Theorem (Englund, Bicknell-Johnson (1997); Jones, Kiss (1998); Chung, Graham, S. (2019))

For $n > 1$ with $s = s(n)$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = af_{s-1} + bf_{s-2}$ and $1 \leq a \leq b \leq f_{s-1}$. In this case (b, a) is n -good.

Because $w_k(x, y) = yf_{k-1} + xf_{k-2}$, by definition of s there exist a', b' such that $n = w_s(b', a') = a'f_{s-1} + b'f_{s-2}$. Let k be such that $1 \leq b' - kf_{s-1} \leq f_{s-1}$. Then $(b, a) := (b' - kf_{s-1}, a' + kf_{s-2})$ is n -good. Thus $a \leq b$, and we have $b \leq f_{s-1}$ by construction.

Slow Fibonacci Walks

Theorem (Englund, Bicknell-Johnson (1997); Jones, Kiss (1998); Chung, Graham, S. (2019))

For $n > 1$ with $s = s(n)$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = af_{s-1} + bf_{s-2}$ and $1 \leq a \leq b \leq f_{s-1}$. In this case (b, a) is n -good.

Because $w_k(x, y) = yf_{k-1} + xf_{k-2}$, by definition of s there exist a', b' such that $n = w_s(b', a') = a'f_{s-1} + b'f_{s-2}$. Let k be such that $1 \leq b' - kf_{s-1} \leq f_{s-1}$. Then $(b, a) := (b' - kf_{s-1}, a' + kf_{s-2})$ is n -good. Thus $a \leq b$, and we have $b \leq f_{s-1}$ by construction. Hence such integers exist, and it's not too hard to prove uniqueness.

Applications

Applications

Let $p(n)$ denote the number of n -good pairs.

Applications

Let $p(n)$ denote the number of n -good pairs.

Corollary

$p(n) \leq 2$ for $n > 1$, with equality iff $a(n) > f_{s-2}$

Applications

Let $p(n)$ denote the number of n -good pairs.

Corollary

$p(n) \leq 2$ for $n > 1$, with equality iff $a(n) > f_{s-2}$

Proof.

Let $1 \leq a \leq b \leq f_{s-1}$ be as in the previous theorem.

Applications

Let $p(n)$ denote the number of n -good pairs.

Corollary

$p(n) \leq 2$ for $n > 1$, with equality iff $a(n) > f_{s-2}$

Proof.

Let $1 \leq a \leq b \leq f_{s-1}$ be as in the previous theorem. Recall that every n -good pair is of the form

$$(b', a') = (b + kf_{s-1}, a - kf_{s-2})$$

for some k such that $b', a' \geq 1$.

Applications

Let $p(n)$ denote the number of n -good pairs.

Corollary

$p(n) \leq 2$ for $n > 1$, with equality iff $a(n) > f_{s-2}$

Proof.

Let $1 \leq a \leq b \leq f_{s-1}$ be as in the previous theorem. Recall that every n -good pair is of the form

$$(b', a') = (b + kf_{s-1}, a - kf_{s-2})$$

for some k such that $b', a' \geq 1$. Because $b \leq f_{s-1}$, we need $k \geq 0$ to have $b' \geq 1$.

Applications

Let $p(n)$ denote the number of n -good pairs.

Corollary

$p(n) \leq 2$ for $n > 1$, with equality iff $a(n) > f_{s-2}$

Proof.

Let $1 \leq a \leq b \leq f_{s-1}$ be as in the previous theorem. Recall that every n -good pair is of the form

$$(b', a') = (b + kf_{s-1}, a - kf_{s-2})$$

for some k such that $b', a' \geq 1$. Because $b \leq f_{s-1}$, we need $k \geq 0$ to have $b' \geq 1$. Because $a \leq f_{s-1} \leq 2f_{s-2}$, we need $k \leq 1$.

Applications

Let $p(n)$ denote the number of n -good pairs.

Corollary

$p(n) \leq 2$ for $n > 1$, with equality iff $a(n) > f_{s-2}$

Proof.

Let $1 \leq a \leq b \leq f_{s-1}$ be as in the previous theorem. Recall that every n -good pair is of the form

$$(b', a') = (b + kf_{s-1}, a - kf_{s-2})$$

for some k such that $b', a' \geq 1$. Because $b \leq f_{s-1}$, we need $k \geq 0$ to have $b' \geq 1$. Because $a \leq f_{s-1} \leq 2f_{s-2}$, we need $k \leq 1$. Thus only the pairs with $k = 0, 1$ can work, and these both work iff $a > f_{s-2}$. □

Applications

Let $T(n) = n^{-1}|\{m \leq n : m \text{ has two } m\text{-slow Fibonacci walks}\}|$.

Applications


Let $T(n) = n^{-1}|\{m \leq n : m \text{ has two } m\text{-slow Fibonacci walks}\}|$.

Theorem (Chung, Graham, S. (2019))

Given n , let c, p be such that $n = \frac{1}{\sqrt{5}}c\phi^p$ with $\frac{1}{\sqrt{5}} \leq c < \frac{1}{\sqrt{5}}\phi$.
Then


$$T(n) = \begin{cases} \frac{1}{2\sqrt{5}\phi^4c} + O(n^{-1/2}) & p \equiv 1 \pmod{2}, \\ \frac{\sqrt{5}}{2}c + \frac{1+\phi^{-5}}{2\sqrt{5}c} - 1 + O(n^{-1/2}) & p \equiv 0 \pmod{2}, c \leq \frac{1+\phi^{-3}}{\sqrt{5}}, \\ 1 - \frac{\sqrt{5}}{2}\phi^{-1}c - \frac{1+\phi^{-2}}{2\sqrt{5}c} + O(n^{-1/2}) & p \equiv 0 \pmod{2}, c \geq \frac{1+\phi^{-3}}{\sqrt{5}}. \end{cases}$$

Applications




(a) Data plot of $T(n)$.

Applications




(a) Data plot of $T(n)$.




(b) Theory plot of $T(n)$.

Applications



(a) Data plot of $T(n)$.



(b) Theory plot of $T(n)$.

Note that this value oscillates.

Applications

Proof Sketch:

Applications

Proof Sketch: Don't count $m \leq n$ which have two pairs, instead count triples $(s(m), a(m), b(m)) = (s, a, b)$ where m has two pairs.

Applications

Proof Sketch: Don't count $m \leq n$ which have two pairs, instead count triples $(s(m), a(m), b(m)) = (s, a, b)$ where m has two pairs. These are exactly the triples satisfying

$$af_{s-1} + bf_{s-2} = m \leq n$$

Applications

Proof Sketch: Don't count $m \leq n$ which have two pairs, instead count triples $(s(m), a(m), b(m)) = (s, a, b)$ where m has two pairs. These are exactly the triples satisfying

$$af_{s-1} + bf_{s-2} = m \leq n,$$

$$f_{s-1} \geq b \geq a$$

Applications

Proof Sketch: Don't count $m \leq n$ which have two pairs, instead count triples $(s(m), a(m), b(m)) = (s, a, b)$ where m has two pairs. These are exactly the triples satisfying

$$af_{s-1} + bf_{s-2} = m \leq n,$$

$$f_{s-1} \geq b \geq a > f_{s-2}.$$

Applications

Proof Sketch: Don't count $m \leq n$ which have two pairs, instead count triples $(s(m), a(m), b(m)) = (s, a, b)$ where m has two pairs. These are exactly the triples satisfying

$$af_{s-1} + bf_{s-2} = m \leq n,$$

$$f_{s-1} \geq b \geq a > f_{s-2}.$$

If s is very small compared to n , then any triple (s, a, b) with $f_{s-1} \geq b \geq a > f_{s-2}$ works (and this is easy to count).

Applications

Proof Sketch: Don't count $m \leq n$ which have two pairs, instead count triples $(s(m), a(m), b(m)) = (s, a, b)$ where m has two pairs. These are exactly the triples satisfying

$$af_{s-1} + bf_{s-2} = m \leq n,$$

$$f_{s-1} \geq b \geq a > f_{s-2}.$$

If s is very small compared to n , then any triple (s, a, b) with $f_{s-1} \geq b \geq a > f_{s-2}$ works (and this is easy to count). Namely, this holds if $n \geq f_{s-1}(f_{s-1} + f_{s-2}) \approx \frac{1}{5}\phi^{2s-1}$.

Applications

Proof Sketch: Don't count $m \leq n$ which have two pairs, instead count triples $(s(m), a(m), b(m)) = (s, a, b)$ where m has two pairs. These are exactly the triples satisfying

$$af_{s-1} + bf_{s-2} = m \leq n,$$

$$f_{s-1} \geq b \geq a > f_{s-2}.$$

If s is very small compared to n , then any triple (s, a, b) with $f_{s-1} \geq b \geq a > f_{s-2}$ works (and this is easy to count). Namely, this holds if $n \geq f_{s-1}(f_{s-1} + f_{s-2}) \approx \frac{1}{5}\phi^{2s-1}$. If s is very large, then no triple (s, a, b) satisfies both inequalities, so no triples exist.

Applications

Proof Sketch: Don't count $m \leq n$ which have two pairs, instead count triples $(s(m), a(m), b(m)) = (s, a, b)$ where m has two pairs. These are exactly the triples satisfying

$$af_{s-1} + bf_{s-2} = m \leq n,$$

$$f_{s-1} \geq b \geq a > f_{s-2}.$$

If s is very small compared to n , then any triple (s, a, b) with $f_{s-1} \geq b \geq a > f_{s-2}$ works (and this is easy to count). Namely, this holds if $n \geq f_{s-1}(f_{s-1} + f_{s-2}) \approx \frac{1}{5}\phi^{2s-1}$. If s is very large, then no triple (s, a, b) satisfies both inequalities, so no triples exist. Namely, this holds if $n < f_{s-2}(f_{s-1} + f_{s-2}) \approx \frac{1}{5}\phi^{2s-2}$ then no such a, b exist.

Applications

Proof Sketch: Don't count $m \leq n$ which have two pairs, instead count triples $(s(m), a(m), b(m)) = (s, a, b)$ where m has two pairs. These are exactly the triples satisfying

$$af_{s-1} + bf_{s-2} = m \leq n,$$

$$f_{s-1} \geq b \geq a > f_{s-2}.$$

If s is very small compared to n , then any triple (s, a, b) with $f_{s-1} \geq b \geq a > f_{s-2}$ works (and this is easy to count). Namely, this holds if $n \geq f_{s-1}(f_{s-1} + f_{s-2}) \approx \frac{1}{5}\phi^{2s-1}$. If s is very large, then no triple (s, a, b) satisfies both inequalities, so no triples exist. Namely, this holds if $n < f_{s-2}(f_{s-1} + f_{s-2}) \approx \frac{1}{5}\phi^{2s-2}$ then no such a, b exist. If n is in between these two values then things are annoying but doable.



Applications

From now on whenever I write w_k I'm assuming it's the n -slow Fibonacci walk $w_k(b(n), a(n))$.

Applications

From now on whenever I write w_k I'm assuming it's the n -slow Fibonacci walk $w_k(b(n), a(n))$.

Corollary

For any n with $s(n) = s$, we have $w_{s+1} = \lfloor \phi n \rfloor$ if s is odd and $w_{s+1} = \lceil \phi n \rceil$ if s is even.

Applications

From now on whenever I write w_k I'm assuming it's the n -slow Fibonacci walk $w_k(b(n), a(n))$.

Corollary

For any n with $s(n) = s$, we have $w_{s+1} = \lfloor \phi n \rfloor$ if s is odd and $w_{s+1} = \lceil \phi n \rceil$ if s is even.

Observe that $w_{s+1}(b, a) = \lfloor \phi n \rfloor$ is equivalent to

$$0 \leq \phi n - w_{s+1}(b, a) < 1.$$

Applications

From now on whenever I write w_k I'm assuming it's the n -slow Fibonacci walk $w_k(b(n), a(n))$.

Corollary

For any n with $s(n) = s$, we have $w_{s+1} = \lfloor \phi n \rfloor$ if s is odd and $w_{s+1} = \lceil \phi n \rceil$ if s is even.

Observe that $w_{s+1}(b, a) = \lfloor \phi n \rfloor$ is equivalent to

$$0 \leq \phi n - w_{s+1}(b, a) < 1.$$

By using $n = w_s(b, a)$ and writing w_k in terms of Fibonacci numbers, this is equivalent to

$$0 \leq (-\phi)^{-s+1}(\phi b - a) < 1.$$

Applications

From now on whenever I write w_k I'm assuming it's the n -slow Fibonacci walk $w_k(b(n), a(n))$.

Corollary

For any n with $s(n) = s$, we have $w_{s+1} = \lfloor \phi n \rfloor$ if s is odd and $w_{s+1} = \lceil \phi n \rceil$ if s is even.

Observe that $w_{s+1}(b, a) = \lfloor \phi n \rfloor$ is equivalent to

$$0 \leq \phi n - w_{s+1}(b, a) < 1.$$

By using $n = w_s(b, a)$ and writing w_k in terms of Fibonacci numbers, this is equivalent to

$$0 \leq (-\phi)^{-s+1}(\phi b - a) < 1.$$

This is true when s is odd for $1 \leq a \leq b \leq f_t$.



Applications

We say that n is a down-integer if $w_{s+1} = \lfloor \phi n \rfloor$, and we define D to be the set of down-integers.

Applications

We say that n is a down-integer if $w_{s+1} = \lfloor \phi n \rfloor$, and we define D to be the set of down-integers. Let $D(n) = n^{-1}|D \cap [n]|$.

Applications

We say that n is a down-integer if $w_{s+1} = \lfloor \phi n \rfloor$, and we define D to be the set of down-integers. Let $D(n) = n^{-1} |D \cap [n]|$.

Theorem (Chung, Graham, S. (2019))

$$D(n) = \begin{cases} \frac{\sqrt{5}n}{2\phi^{q+1}} + \frac{\phi^{q+1}}{10\sqrt{5}n} + O(n^{-1/2}) & \frac{1}{5}\phi^q \leq n < \frac{1}{5}\phi^{q+2}, \quad q \equiv 1 \pmod{4}, \\ 1 - \frac{\sqrt{5}n}{2\phi^{q+1}} - \frac{\phi^{q+1}}{10\sqrt{5}n} + O(n^{-1/2}) & \frac{1}{5}\phi^q \leq n < \frac{1}{5}\phi^{q+2}, \quad q \equiv 3 \pmod{4}. \end{cases}$$

Applications

We say that n is a down-integer if $w_{s+1} = \lfloor \phi n \rfloor$, and we define D to be the set of down-integers. Let $D(n) = n^{-1} |D \cap [n]|$.


Theorem (Chung, Graham, S. (2019))

$$D(n) = \begin{cases} \frac{\sqrt{5}n}{2\phi^{q+1}} + \frac{\phi^{q+1}}{10\sqrt{5}n} + O(n^{-1/2}) & \frac{1}{5}\phi^q \leq n < \frac{1}{5}\phi^{q+2}, \quad q \equiv 1 \pmod{4}, \\ 1 - \frac{\sqrt{5}n}{2\phi^{q+1}} - \frac{\phi^{q+1}}{10\sqrt{5}n} + O(n^{-1/2}) & \frac{1}{5}\phi^q \leq n < \frac{1}{5}\phi^{q+2}, \quad q \equiv 3 \pmod{4}. \end{cases}$$


Proof.

Count triples (s, a, b) with s odd. □

Applications



(a) Data plot of $D(n)$.



(b) Theory plot of $D(n)$.

Applications

This says that roughly half the integers have w_{s+1} rounding down and half have them rounding up.

Applications

This says that roughly half the integers have w_{s+1} rounding down and half have them rounding up. How many n can you have in a row with all of them rounding down?

Applications

This says that roughly half the integers have w_{s+1} rounding down and half have them rounding up. How many n can you have in a row with all of them rounding down?

To this end, write the elements of D in increasing order as d_1, d_2, \dots . The above is equivalent to asking how large $d_{k+1} - d_k$ can be.

Applications

This says that roughly half the integers have w_{s+1} rounding down and half have them rounding up. How many n can you have in a row with all of them rounding down?

To this end, write the elements of D in increasing order as d_1, d_2, \dots . The above is equivalent to asking how large $d_{k+1} - d_k$ can be.

Theorem (Chung, Graham, S. (2019))

$$\{d_{k+1} - d_k : k \geq 1\} = \{1, 2, 3, 5\}.$$

Applications

This says that roughly half the integers have w_{s+1} rounding down and half have them rounding up. How many n can you have in a row with all of them rounding down?

To this end, write the elements of D in increasing order as d_1, d_2, \dots . The above is equivalent to asking how large $d_{k+1} - d_k$ can be.

Theorem (Chung, Graham, S. (2019))

$$\{d_{k+1} - d_k : k \geq 1\} = \{1, 2, 3, 5\}.$$

$$\{d_{k+2} - d_k : k \geq 1\} = \{2, 3, 4, 5, 6, 8, 10\}.$$

Applications

We know that $w_{s+1} = \lfloor \phi n \rfloor$ or $w_{s+1} = \lceil \phi n \rceil$. Intuitively, the smaller $\phi n - \lfloor \phi n \rfloor$ is, the more likely it is that $w_{s+1} = \lfloor \phi n \rfloor$.

Applications

We know that $w_{s+1} = \lfloor \phi n \rfloor$ or $w_{s+1} = \lceil \phi n \rceil$. Intuitively, the smaller $\phi n - \lfloor \phi n \rfloor$ is, the more likely it is that $w_{s+1} = \lfloor \phi n \rfloor$. To this end, we say that n is r -paradoxical if $|\phi n - w_{s+1}| > r$.

Applications

We know that $w_{s+1} = \lfloor \phi n \rfloor$ or $w_{s+1} = \lceil \phi n \rceil$. Intuitively, the smaller $\phi n - \lfloor \phi n \rfloor$ is, the more likely it is that $w_{s+1} = \lfloor \phi n \rfloor$. To this end, we say that n is r -paradoxical if $|\phi n - w_{s+1}| > r$. For $r \geq \frac{1}{2}$, let $P(n, r) = n^{-1} |\{m \leq n : m \text{ is } r\text{-paradoxical}\}|$.

Applications

We know that $w_{s+1} = \lfloor \phi n \rfloor$ or $w_{s+1} = \lceil \phi n \rceil$. Intuitively, the smaller $\phi n - \lfloor \phi n \rfloor$ is, the more likely it is that $w_{s+1} = \lfloor \phi n \rfloor$. To this end, we say that n is r -paradoxical if $|\phi n - w_{s+1}| > r$. For $r \geq \frac{1}{2}$, let $P(n, r) = n^{-1} |\{m \leq n : m \text{ is } r\text{-paradoxical}\}|$.

Theorem (Chung, Graham, S. (2019))

We have $P(n, r) = 0$ if $r \geq 1 - \frac{1}{\sqrt{5}}\phi^{-1} \approx .72$.

Applications


We know that $w_{s+1} = \lfloor \phi n \rfloor$ or $w_{s+1} = \lceil \phi n \rceil$. Intuitively, the smaller $\phi n - \lfloor \phi n \rfloor$ is, the more likely it is that $w_{s+1} = \lfloor \phi n \rfloor$. To this end, we say that n is r -paradoxical if $|\phi n - w_{s+1}| > r$. For $r \geq \frac{1}{2}$, let $P(n, r) = n^{-1} |\{m \leq n : m \text{ is } r\text{-paradoxical}\}|$.

Theorem (Chung, Graham, S. (2019))


We have $P(n, r) = 0$ if $r \geq 1 - \frac{1}{\sqrt{5}}\phi^{-1} \approx .72$. Otherwise, given n , let c, p be such that $n = \frac{1}{\sqrt{5}}c\phi^p$ with $\frac{1}{\sqrt{5}} \leq c < \frac{1}{\sqrt{5}}\phi$. Then $P(n, r)$ satisfies

$$\begin{cases} -\frac{1}{2}\phi^{-1}c + (1-r) + \left(r^2 - r + \frac{1}{2\sqrt{5}}\phi^{-1}\right)c^{-1} + O(n^{-1/2}) & p \text{ odd}, c \leq (1-r)\phi, \\ \frac{\sqrt{5}}{2}\phi \left(r - \frac{1}{\sqrt{5}}\phi\right)^2 c^{-1} + O(n^{-1/2}) & p \text{ odd}, c \geq (1-r)\phi, \\ -\frac{1}{2}c + (1-r) + \left(\phi^{-1}r^2 - \phi^{-1}r + \frac{1}{2\sqrt{5}}\phi^{-2}\right)c^{-1} + O(n^{-1/2}) & p \text{ even}, c \leq 1-r, \\ \frac{\sqrt{5}}{2} \left(r - \frac{1}{\sqrt{5}}\phi\right)^2 c^{-1} + O(n^{-1/2}) & p \text{ even}, 1-r \leq c \leq r \\ \frac{1}{2}c - r + \left(\phi r^2 - \phi r + \frac{1}{2\sqrt{5}}\phi^2\right)c^{-1} + O(n^{-1/2}) & p \text{ even}, c \geq r. \end{cases}$$


Applications




(a) Data plot of $P(n, .5)$.



(b) Theory plot of $P(n, .5)$.



(c) Data plot of $P(n, .6)$.



(d) Theory plot of $P(n, .6)$.

Applications

How does one compute $a(n)$, $b(n)$ in practice?

Applications

How does one compute $a(n), b(n)$ in practice?

- $O(n^2 \log(n))$ algorithm: Try every walk starting a, b, \dots with $a, b \leq n$, pick the pair(s) giving you the slowest walk.

Applications

How does one compute $a(n)$, $b(n)$ in practice?

- $O(n^2 \log(n))$ algorithm: Try every walk starting a, b, \dots with $a, b \leq n$, pick the pair(s) giving you the slowest walk.
- $O(n \log(n))$ algorithm: Try every “reverse walk” ending with \dots, x, n for $x \leq n$ and pick the slowest ones.

Applications

How does one compute $a(n)$, $b(n)$ in practice?

- $O(n^2 \log(n))$ algorithm: Try every walk starting a, b, \dots with $a, b \leq n$, pick the pair(s) giving you the slowest walk.
- $O(n \log(n))$ algorithm: Try every “reverse walk” ending with \dots, x, n for $x \leq n$ and pick the slowest ones.
- $O(\log n)$ algorithm: We know the “standard” slow walk goes $\dots, n, \lfloor \phi n \rfloor$ or $\dots, n, \lceil \phi n \rceil$.

Applications

How does one compute $a(n)$, $b(n)$ in practice?

- $O(n^2 \log(n))$ algorithm: Try every walk starting a, b, \dots with $a, b \leq n$, pick the pair(s) giving you the slowest walk.
- $O(n \log(n))$ algorithm: Try every “reverse walk” ending with \dots, x, n for $x \leq n$ and pick the slowest ones.
- $O(\log n)$ algorithm: We know the “standard” slow walk goes $\dots, n, \lfloor \phi n \rfloor$ or $\dots, n, \lceil \phi n \rceil$. Thus we just have to check these two reverse walks.

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

The Fibonacci sequence isn't special.

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

The Fibonacci sequence isn't special. To this end, we define an n -slow (α, β) -walks to be a sequence with $w_{k+2} = \alpha w_{k+1} + \beta w_k$ such that $w_s = n$ with s as large as possible.

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

The Fibonacci sequence isn't special. To this end, we define an n -slow (α, β) -walks to be a sequence with $w_{k+2} = \alpha w_{k+1} + \beta w_k$ such that $w_s = n$ with s as large as possible. Throughout we assume $\alpha, \beta \geq 1$ and that $\gcd(\alpha, \beta) = 1$.

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

The Fibonacci sequence isn't special. To this end, we define an n -slow (α, β) -walks to be a sequence with $w_{k+2} = \alpha w_{k+1} + \beta w_k$ such that $w_s = n$ with s as large as possible. Throughout we assume $\alpha, \beta \geq 1$ and that $\gcd(\alpha, \beta) = 1$.

Define g_k to be the sequence with

$$g_1 = 1, \quad g_2 = \alpha, \quad g_{k+2} = \alpha g_{k+1} + \beta g_k.$$

Also define

$$\gamma = \frac{1}{2}(\alpha + \sqrt{\alpha^2 + 4\beta}), \quad \lambda = \frac{1}{2}(\alpha - \sqrt{\alpha^2 + 4\beta}).$$

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

The Fibonacci sequence isn't special. To this end, we define an n -slow (α, β) -walks to be a sequence with $w_{k+2} = \alpha w_{k+1} + \beta w_k$ such that $w_s = n$ with s as large as possible. Throughout we assume $\alpha, \beta \geq 1$ and that $\gcd(\alpha, \beta) = 1$.

Define g_k to be the sequence with

$$g_1 = 1, \quad g_2 = \alpha, \quad g_{k+2} = \alpha g_{k+1} + \beta g_k.$$

Also define

$$\gamma = \frac{1}{2}(\alpha + \sqrt{\alpha^2 + 4\beta}), \quad \lambda = \frac{1}{2}(\alpha - \sqrt{\alpha^2 + 4\beta}).$$

Note that when $\alpha = \beta = 1$ we have $g_k = f_k$, $\gamma = \phi$, $\lambda = -\phi^{-1}$.

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

When $\beta = 1$, almost all our proofs from before carry over.

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

When $\beta = 1$, almost all our proofs from before carry over. Let $s(n) = s^{\alpha, \beta}(n)$ be the number of steps it takes an n -slow (α, β) -walk to hit n .

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

When $\beta = 1$, almost all our proofs from before carry over. Let $s(n) = s^{\alpha, \beta}(n)$ be the number of steps it takes an n -slow (α, β) -walk to hit n .

Theorem (S. (2019))

If $s(n) > 2$ and $\beta = 1$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = ag_{s-1} + bg_{s-2}$

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

When $\beta = 1$, almost all our proofs from before carry over. Let $s(n) = s^{\alpha, \beta}(n)$ be the number of steps it takes an n -slow (α, β) -walk to hit n .

Theorem (S. (2019))

If $s(n) > 2$ and $\beta = 1$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = ag_{s-1} + bg_{s-2}$ and $1 \leq a \leq \alpha b$

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

When $\beta = 1$, almost all our proofs from before carry over. Let $s(n) = s^{\alpha, \beta}(n)$ be the number of steps it takes an n -slow (α, β) -walk to hit n .

Theorem (S. (2019))

If $s(n) > 2$ and $\beta = 1$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = ag_{s-1} + bg_{s-2}$ and $1 \leq a \leq \alpha b \leq \alpha g_{s-1}$.

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

When $\beta = 1$, almost all our proofs from before carry over. Let $s(n) = s^{\alpha, \beta}(n)$ be the number of steps it takes an n -slow (α, β) -walk to hit n .

Theorem (S. (2019))

If $s(n) > 2$ and $\beta = 1$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = ag_{s-1} + bg_{s-2}$ and $1 \leq a \leq \alpha b \leq \alpha g_{s-1}$. Moreover, (b, a) is n -good and $w_{s+1}(b, a)$ is either $\lfloor \gamma n \rfloor$ or $\lceil \gamma n \rceil$.

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

When $\beta = 1$, almost all our proofs from before carry over. Let $s(n) = s^{\alpha, \beta}(n)$ be the number of steps it takes an n -slow (α, β) -walk to hit n .

Theorem (S. (2019))

If $s(n) > 2$ and $\beta = 1$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = ag_{s-1} + bg_{s-2}$ and $1 \leq a \leq \alpha b \leq \alpha g_{s-1}$. Moreover, (b, a) is n -good and $w_{s+1}(b, a)$ is either $\lfloor \gamma n \rfloor$ or $\lceil \gamma n \rceil$.

When $\beta \neq 1$ things get a bit more complicated.

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

When $\beta = 1$, almost all our proofs from before carry over. Let $s(n) = s^{\alpha,\beta}(n)$ be the number of steps it takes an n -slow (α, β) -walk to hit n .

Theorem (S. (2019))

If $s(n) > 2$ and $\beta = 1$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = ag_{s-1} + bg_{s-2}$ and $1 \leq a \leq \alpha b \leq \alpha g_{s-1}$. Moreover, (b, a) is n -good and $w_{s+1}(b, a)$ is either $\lfloor \gamma n \rfloor$ or $\lceil \gamma n \rceil$.

When $\beta \neq 1$ things get a bit more complicated.

Theorem (S. (2019))

If $s(n) > 2$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = ag_{s-1} + \beta bg_{s-2}$

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

When $\beta = 1$, almost all our proofs from before carry over. Let $s(n) = s^{\alpha,\beta}(n)$ be the number of steps it takes an n -slow (α, β) -walk to hit n .

Theorem (S. (2019))

If $s(n) > 2$ and $\beta = 1$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = ag_{s-1} + bg_{s-2}$ and $1 \leq a \leq \alpha b \leq \alpha g_{s-1}$. Moreover, (b, a) is n -good and $w_{s+1}(b, a)$ is either $\lfloor \gamma n \rfloor$ or $\lceil \gamma n \rceil$.

When $\beta \neq 1$ things get a bit more complicated.

Theorem (S. (2019))

If $s(n) > 2$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = ag_{s-1} + \beta bg_{s-2}$, $b \leq g_{s-1}$,

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

When $\beta = 1$, almost all our proofs from before carry over. Let $s(n) = s^{\alpha, \beta}(n)$ be the number of steps it takes an n -slow (α, β) -walk to hit n .

Theorem (S. (2019))

If $s(n) > 2$ and $\beta = 1$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = ag_{s-1} + bg_{s-2}$ and $1 \leq a \leq \alpha b \leq \alpha g_{s-1}$. Moreover, (b, a) is n -good and $w_{s+1}(b, a)$ is either $\lfloor \gamma n \rfloor$ or $\lceil \gamma n \rceil$.

When $\beta \neq 1$ things get a bit more complicated.

Theorem (S. (2019))

If $s(n) > 2$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = ag_{s-1} + \beta bg_{s-2}$, $b \leq g_{s-1}$, $1 \leq a \leq \alpha b + (\beta - 1)g_s$

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

When $\beta = 1$, almost all our proofs from before carry over. Let $s(n) = s^{\alpha,\beta}(n)$ be the number of steps it takes an n -slow (α, β) -walk to hit n .

Theorem (S. (2019))

If $s(n) > 2$ and $\beta = 1$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = ag_{s-1} + bg_{s-2}$ and $1 \leq a \leq \alpha b \leq \alpha g_{s-1}$. Moreover, (b, a) is n -good and $w_{s+1}(b, a)$ is either $\lfloor \gamma n \rfloor$ or $\lceil \gamma n \rceil$.

When $\beta \neq 1$ things get a bit more complicated.

Theorem (S. (2019))

If $s(n) > 2$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = ag_{s-1} + \beta bg_{s-2}$, $b \leq g_{s-1}$, $1 \leq a \leq \alpha b + (\beta - 1)g_s$, and $a - \alpha b - \ell g_s$ is not a positive multiple of β for any $\ell \geq 0$.

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

When $\beta = 1$, almost all our proofs from before carry over. Let $s(n) = s^{\alpha, \beta}(n)$ be the number of steps it takes an n -slow (α, β) -walk to hit n .

Theorem (S. (2019))

If $s(n) > 2$ and $\beta = 1$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = ag_{s-1} + bg_{s-2}$ and $1 \leq a \leq \alpha b \leq \alpha g_{s-1}$. Moreover, (b, a) is n -good and $w_{s+1}(b, a)$ is either $\lfloor \gamma n \rfloor$ or $\lceil \gamma n \rceil$.

When $\beta \neq 1$ things get a bit more complicated.

Theorem (S. (2019))

If $s(n) > 2$, there exist unique integers $a = a(n)$, $b = b(n)$ such that $n = ag_{s-1} + \beta bg_{s-2}$, $b \leq g_{s-1}$, $1 \leq a \leq \alpha b + (\beta - 1)g_s$, and $a - \alpha b - \ell g_s$ is not a positive multiple of β for any $\ell \geq 0$. Moreover, (b, a) is n -good and $|w_{s+1}(b, a) - \gamma n| \leq 2\beta^s$.

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

As before we can use this theorem to prove a number of results about slow walks.

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

As before we can use this theorem to prove a number of results about slow walks. Define $p(n)$ to be the number of n -slow (α, β) -walks.

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

As before we can use this theorem to prove a number of results about slow walks. Define $p(n)$ to be the number of n -slow (α, β) -walks.

Theorem (S. (2019))

- If $s(n) > 2$, then

$$p(n) \leq \alpha^2 + 2\beta - 1.$$

Moreover, there always exists an n achieving this.

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

As before we can use this theorem to prove a number of results about slow walks. Define $p(n)$ to be the number of n -slow (α, β) -walks.

Theorem (S. (2019))

- If $s(n) > 2$, then

$$p(n) \leq \alpha^2 + 2\beta - 1.$$

Moreover, there always exists an n achieving this.

- *There exist infinitely many n with*

$$p(n) = \alpha^2 + \beta + \lceil \alpha\beta\gamma^{-1} \rceil - 1,$$

and only finitely many n with

$$p(n) \geq \alpha^2 + \beta + \lceil \alpha\beta\gamma^{-1} \rceil.$$

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

Let S_p denote the set of n with $p(n) > p$.

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

Let S_p denote the set of n with $p(n) > p$.

Theorem (S. (2019))

Given an integer p , let d denote the smallest integer such that $\delta := \beta\gamma^{-1}p - \gamma d \leq \alpha$. If $\beta \leq p \leq \lceil \gamma^2 \rceil - 2$ and $1 \leq c \leq (p - \beta + 1)\gamma/\alpha$, then $n_{c,r}^{-1}|S_p \cap [n_{c,r}]| =$

$$c^{-1} \left(\frac{(2\beta - 2d - 1)\gamma(\alpha - 2\delta + \alpha^{-1}\delta^2)}{2\beta^2(\gamma^2 - 1)} + \frac{\gamma^2}{\gamma^2 - 1} \sum_{q=d+1}^{\beta-1} \frac{\beta - q}{\beta^2} \right) + O(\gamma^{-r} + (\beta\gamma^{-2})^r),$$

$$\text{where } n_{c,r} := \left\lfloor \frac{c\beta}{(\gamma-\lambda)^2} \gamma^{2r+1} \right\rfloor.$$

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

Let S_p denote the set of n with $p(n) > p$.

Theorem (S. (2019))

Given an integer p , let d denote the smallest integer such that $\delta := \beta\gamma^{-1}p - \gamma d \leq \alpha$. If $\beta \leq p \leq \lceil \gamma^2 \rceil - 2$ and $1 \leq c \leq (p - \beta + 1)\gamma/\alpha$, then $n_{c,r}^{-1}|S_p \cap [n_{c,r}]| =$





$$c^{-1} \left(\frac{(2\beta - 2d - 1)\gamma(\alpha - 2\delta + \alpha^{-1}\delta^2)}{2\beta^2(\gamma^2 - 1)} + \frac{\gamma^2}{\gamma^2 - 1} \sum_{q=d+1}^{\beta-1} \frac{\beta - q}{\beta^2} \right) + O(\gamma^{-r} + (\beta\gamma^{-2})^r),$$

$$\text{where } n_{c,r} := \left\lfloor \frac{c\beta}{(\gamma-\lambda)^2} \gamma^{2r+1} \right\rfloor.$$

When $\beta = 1$ the proof is essentially the same as before, otherwise one has to be careful about the divisibility condition.

Generalized Walks: $w_{k+2} = \alpha w_{k+1} + \beta w_k$

$\alpha = 2, \beta = 1, p = 1, 2, 3, 4.$



Slowest Slow Walks

Slowest Slow Walks

How slow is the slowest slow walk?

Slowest Slow Walks

How slow is the slowest slow walk? Define $\mathbf{s}(n) = \max_{(\alpha, \beta)} s^{\alpha, \beta}(n)$, as well as the pairs achieving this $\mathbf{S}(n) = \{(\alpha, \beta) : s^{\alpha, \beta}(n) = \mathbf{s}(n)\}$.

Slowest Slow Walks

How slow is the slowest slow walk? Define $\mathbf{s}(n) = \max_{(\alpha, \beta)} s^{\alpha, \beta}(n)$, as well as the pairs achieving this $\mathbf{S}(n) = \{(\alpha, \beta) : s^{\alpha, \beta}(n) = \mathbf{s}(n)\}$. A priori, any pair (α, β) could be an element of $\mathbf{S}(n)$ for some n .

Slowest Slow Walks

How slow is the slowest slow walk? Define $\mathbf{s}(n) = \max_{(\alpha, \beta)} s^{\alpha, \beta}(n)$, as well as the pairs achieving this $\mathbf{S}(n) = \{(\alpha, \beta) : s^{\alpha, \beta}(n) = \mathbf{s}(n)\}$. A priori, any pair (α, β) could be an element of $\mathbf{S}(n)$ for some n . However, it turns out that only a finite number of pairs have this property.

Slowest Slow Walks

How slow is the slowest slow walk? Define $\mathbf{s}(n) = \max_{(\alpha, \beta)} s^{\alpha, \beta}(n)$, as well as the pairs achieving this $\mathbf{S}(n) = \{(\alpha, \beta) : s^{\alpha, \beta}(n) = \mathbf{s}(n)\}$. A priori, any pair (α, β) could be an element of $\mathbf{S}(n)$ for some n . However, it turns out that only a finite number of pairs have this property.

Theorem (S. (2019))

Let $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (1, 4)\}$.

Slowest Slow Walks

How slow is the slowest slow walk? Define $\mathbf{s}(n) = \max_{(\alpha, \beta)} s^{\alpha, \beta}(n)$, as well as the pairs achieving this $\mathbf{S}(n) = \{(\alpha, \beta) : s^{\alpha, \beta}(n) = \mathbf{s}(n)\}$. A priori, any pair (α, β) could be an element of $\mathbf{S}(n)$ for some n . However, it turns out that only a finite number of pairs have this property.

Theorem (S. (2019))

Let $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (1, 4)\}$.

- For all $n > 1$, we have $\mathbf{S}(n) \subseteq R$.

Slowest Slow Walks

How slow is the slowest slow walk? Define $\mathbf{s}(n) = \max_{(\alpha, \beta)} s^{\alpha, \beta}(n)$, as well as the pairs achieving this $\mathbf{S}(n) = \{(\alpha, \beta) : s^{\alpha, \beta}(n) = \mathbf{s}(n)\}$. A priori, any pair (α, β) could be an element of $\mathbf{S}(n)$ for some n . However, it turns out that only a finite number of pairs have this property.

Theorem (S. (2019))

Let $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (1, 4)\}$.

- For all $n > 1$, we have $\mathbf{S}(n) \subseteq R$.
- For all $(\alpha, \beta) \in R$, there exists an n with $(\alpha, \beta) \in \mathbf{S}(n)$.

Slowest Slow Walks

How slow is the slowest slow walk? Define $\mathbf{s}(n) = \max_{(\alpha, \beta)} s^{\alpha, \beta}(n)$, as well as the pairs achieving this $\mathbf{S}(n) = \{(\alpha, \beta) : s^{\alpha, \beta}(n) = \mathbf{s}(n)\}$. A priori, any pair (α, β) could be an element of $\mathbf{S}(n)$ for some n . However, it turns out that only a finite number of pairs have this property.

Theorem (S. (2019))

Let $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (1, 4)\}$.

- For all $n > 1$, we have $\mathbf{S}(n) \subseteq R$.
- For all $(\alpha, \beta) \in R$, there exists an n with $(\alpha, \beta) \in \mathbf{S}(n)$.
- The set of n with $\mathbf{S}(n) = \{(1, 1)\}$ has density 1.

Slowest Slow Walks

How slow is the slowest slow walk? Define $\mathbf{s}(n) = \max_{(\alpha, \beta)} s^{\alpha, \beta}(n)$, as well as the pairs achieving this $\mathbf{S}(n) = \{(\alpha, \beta) : s^{\alpha, \beta}(n) = \mathbf{s}(n)\}$. A priori, any pair (α, β) could be an element of $\mathbf{S}(n)$ for some n . However, it turns out that only a finite number of pairs have this property.

Theorem (S. (2019))

Let $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (1, 4)\}$.

- For all $n > 1$, we have $\mathbf{S}(n) \subseteq R$.
- For all $(\alpha, \beta) \in R$, there exists an n with $(\alpha, \beta) \in \mathbf{S}(n)$.
- The set of n with $\mathbf{S}(n) = \{(1, 1)\}$ has density 1.

Corollary: the Fibonacci sequence *is* special!

Slowest Slow Walks

Let $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (1, 4)\}$.

Slowest Slow Walks

Let $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (1, 4)\}$.

$\mathbf{S}(32) = \{(1, 1), (1, 2)\}$

Slowest Slow Walks

Let $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (1, 4)\}$.

$\mathbf{S}(32) = \{(1, 1), (1, 2)\}$, $\mathbf{S}(40) = \{(1, 1), (1, 3)\}$

Slowest Slow Walks

Let $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (1, 4)\}$.

$\mathbf{S}(32) = \{(1, 1), (1, 2)\}$, $\mathbf{S}(40) = \{(1, 1), (1, 3)\}$, $\mathbf{S}(3363) = \{(1, 1), (2, 1)\}$

Slowest Slow Walks

Let $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (1, 4)\}$.

$\mathbf{S}(32) = \{(1, 1), (1, 2)\}$, $\mathbf{S}(40) = \{(1, 1), (1, 3)\}$, $\mathbf{S}(3363) = \{(1, 1), (2, 1)\}$,

$\mathbf{S}(5307721328585529) = \{(1, 1), (1, 4)\}$.

Slowest Slow Walks

Let $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (1, 4)\}$.

$\mathbf{S}(32) = \{(1, 1), (1, 2)\}$, $\mathbf{S}(40) = \{(1, 1), (1, 3)\}$, $\mathbf{S}(3363) = \{(1, 1), (2, 1)\}$,

$\mathbf{S}(5307721328585529) = \{(1, 1), (1, 4)\}$.

Can you find n with $\mathbf{S}(n) = \{(\alpha, \beta)\}$?

Slowest Slow Walks

Let $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (1, 4)\}$.

$\mathbf{S}(32) = \{(1, 1), (1, 2)\}$, $\mathbf{S}(40) = \{(1, 1), (1, 3)\}$, $\mathbf{S}(3363) = \{(1, 1), (2, 1)\}$,

$\mathbf{S}(5307721328585529) = \{(1, 1), (1, 4)\}$.

Can you find n with $\mathbf{S}(n) = \{(\alpha, \beta)\}$?

$\mathbf{S}(171) = \{(1, 2)\}$

Slowest Slow Walks

Let $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (1, 4)\}$.

$\mathbf{S}(32) = \{(1, 1), (1, 2)\}$, $\mathbf{S}(40) = \{(1, 1), (1, 3)\}$, $\mathbf{S}(3363) = \{(1, 1), (2, 1)\}$,

$\mathbf{S}(5307721328585529) = \{(1, 1), (1, 4)\}$.

Can you find n with $\mathbf{S}(n) = \{(\alpha, \beta)\}$?

$\mathbf{S}(171) = \{(1, 2)\}$, $\mathbf{S}(11228332) = \{(1, 3)\}$

Slowest Slow Walks

Let $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (1, 4)\}$.

$\mathbf{S}(32) = \{(1, 1), (1, 2)\}$, $\mathbf{S}(40) = \{(1, 1), (1, 3)\}$, $\mathbf{S}(3363) = \{(1, 1), (2, 1)\}$,

$\mathbf{S}(5307721328585529) = \{(1, 1), (1, 4)\}$.

Can you find n with $\mathbf{S}(n) = \{(\alpha, \beta)\}$?

$\mathbf{S}(171) = \{(1, 2)\}$, $\mathbf{S}(11228332) = \{(1, 3)\}$, $\mathbf{S}(22619537) = \{(2, 1)\}$

Slowest Slow Walks

Let $R = \{(1, 1), (1, 2), (1, 3), (2, 1), (1, 4)\}$.

$\mathbf{S}(32) = \{(1, 1), (1, 2)\}$, $\mathbf{S}(40) = \{(1, 1), (1, 3)\}$, $\mathbf{S}(3363) = \{(1, 1), (2, 1)\}$,

$\mathbf{S}(5307721328585529) = \{(1, 1), (1, 4)\}$.

Can you find n with $\mathbf{S}(n) = \{(\alpha, \beta)\}$?

$\mathbf{S}(171) = \{(1, 2)\}$, $\mathbf{S}(11228332) = \{(1, 3)\}$, $\mathbf{S}(22619537) = \{(2, 1)\}$, }

$\mathbf{S}(5000966512101628011743180761388223) = \{(1, 4)\}$.

Slowest Slow Walks

More generally, given a set of relatively prime pairs T , define
 $\mathbf{s}_T(n) = \max_{(\alpha, \beta) \in T} s^{\alpha, \beta}(n)$, $\mathbf{S}_T(n) = \{(\alpha, \beta) : s^{\alpha, \beta}(n) = \mathbf{s}_T(n)\}$.

Slowest Slow Walks

More generally, given a set of relatively prime pairs T , define $\mathbf{s}_T(n) = \max_{(\alpha, \beta) \in T} s^{\alpha, \beta}(n)$, $\mathbf{S}_T(n) = \{(\alpha, \beta) : s^{\alpha, \beta}(n) = \mathbf{s}_T(n)\}$.

Theorem (S. (2019))

There exists a finite set R_T and number n_T such that $\mathbf{S}_T(n) \subseteq R_T$ for all $n \geq n_T$.

Slowest Slow Walks

More generally, given a set of relatively prime pairs T , define $\mathbf{s}_T(n) = \max_{(\alpha, \beta) \in T} s^{\alpha, \beta}(n)$, $\mathbf{S}_T(n) = \{(\alpha, \beta) : s^{\alpha, \beta}(n) = \mathbf{s}_T(n)\}$.

Theorem (S. (2019))

There exists a finite set R_T and number n_T such that $\mathbf{S}_T(n) \subseteq R_T$ for all $n \geq n_T$.

Lemma

For all n with $s^{\alpha, \beta}(n) > 2$, we have

$$\frac{1}{2} \log_{\gamma}(n) - 1 \leq s^{\alpha, \beta}(n) \leq \log_{\gamma}(n) + 2.$$

Slowest Slow Walks

More generally, given a set of relatively prime pairs T , define $\mathbf{s}_T(n) = \max_{(\alpha, \beta) \in T} s^{\alpha, \beta}(n)$, $\mathbf{S}_T(n) = \{(\alpha, \beta) : s^{\alpha, \beta}(n) = \mathbf{s}_T(n)\}$.

Theorem (S. (2019))

There exists a finite set R_T and number n_T such that $\mathbf{S}_T(n) \subseteq R_T$ for all $n \geq n_T$.

Lemma

For all n with $s^{\alpha, \beta}(n) > 2$, we have

$$\frac{1}{2} \log_{\gamma}(n) - 1 \leq s^{\alpha, \beta}(n) \leq \log_{\gamma}(n) + 2.$$

The two extreme cases are $n = g_{s-1} + \beta g_{s-2} \approx \gamma^s$ and $n = \beta g_{s-1} g_s \approx \gamma^{2s}$.



Slowest Slow Walks

Theorem (S. (2019))

There exists a finite set R_T and number n_T such that $\mathbf{S}_T(n) \subseteq R_T$ for all $n \geq n_T$.

Slowest Slow Walks

Theorem (S. (2019))

There exists a finite set R_T and number n_T such that $\mathbf{S}_T(n) \subseteq R_T$ for all $n \geq n_T$.

Proof.

Let $(\alpha', \beta') \in T$ such that $\gamma_{\alpha', \beta'} = \min\{\gamma_{\alpha, \beta} : (\alpha, \beta) \in T\} := \Gamma$.

Slowest Slow Walks

Theorem (S. (2019))

There exists a finite set R_T and number n_T such that $\mathbf{S}_T(n) \subseteq R_T$ for all $n \geq n_T$.

Proof.

Let $(\alpha', \beta') \in T$ such that $\gamma_{\alpha', \beta'} = \min\{\gamma_{\alpha, \beta} : (\alpha, \beta) \in T\} := \Gamma$. If $(\alpha, \beta) \in \mathbf{S}_T(n)$ and $\gamma := \gamma_{\alpha, \beta}$, then

$$\log_{\gamma} n + 2 \geq s^{\alpha, \beta}(n)$$

Slowest Slow Walks

Theorem (S. (2019))

There exists a finite set R_T and number n_T such that $\mathbf{S}_T(n) \subseteq R_T$ for all $n \geq n_T$.

Proof.

Let $(\alpha', \beta') \in T$ such that $\gamma_{\alpha', \beta'} = \min\{\gamma_{\alpha, \beta} : (\alpha, \beta) \in T\} := \Gamma$. If $(\alpha, \beta) \in \mathbf{S}_T(n)$ and $\gamma := \gamma_{\alpha, \beta}$, then

$$\log_{\gamma} n + 2 \geq s^{\alpha, \beta}(n) = \mathbf{s}_T(n)$$

Slowest Slow Walks

Theorem (S. (2019))

There exists a finite set R_T and number n_T such that $\mathbf{S}_T(n) \subseteq R_T$ for all $n \geq n_T$.

Proof.

Let $(\alpha', \beta') \in T$ such that $\gamma_{\alpha', \beta'} = \min\{\gamma_{\alpha, \beta} : (\alpha, \beta) \in T\} := \Gamma$. If $(\alpha, \beta) \in \mathbf{S}_T(n)$ and $\gamma := \gamma_{\alpha, \beta}$, then

$$\log_{\gamma} n + 2 \geq s^{\alpha, \beta}(n) = \mathbf{s}_T(n) \geq s^{\alpha', \beta'}(n)$$

Slowest Slow Walks

Theorem (S. (2019))

There exists a finite set R_T and number n_T such that $\mathbf{S}_T(n) \subseteq R_T$ for all $n \geq n_T$.

Proof.

Let $(\alpha', \beta') \in T$ such that $\gamma_{\alpha', \beta'} = \min\{\gamma_{\alpha, \beta} : (\alpha, \beta) \in T\} := \Gamma$. If $(\alpha, \beta) \in \mathbf{S}_T(n)$ and $\gamma := \gamma_{\alpha, \beta}$, then

$$\log_{\gamma} n + 2 \geq s^{\alpha, \beta}(n) = \mathbf{s}_T(n) \geq s^{\alpha', \beta'}(n) \geq \frac{1}{2} \log_{\Gamma} n - 1$$

Slowest Slow Walks

Theorem (S. (2019))

There exists a finite set R_T and number n_T such that $\mathbf{S}_T(n) \subseteq R_T$ for all $n \geq n_T$.

Proof.

Let $(\alpha', \beta') \in T$ such that $\gamma_{\alpha', \beta'} = \min\{\gamma_{\alpha, \beta} : (\alpha, \beta) \in T\} := \Gamma$. If $(\alpha, \beta) \in \mathbf{S}_T(n)$ and $\gamma := \gamma_{\alpha, \beta}$, then

$$\log_{\gamma} n + 2 \geq s^{\alpha, \beta}(n) = \mathbf{s}_T(n) \geq s^{\alpha', \beta'}(n) \geq \frac{1}{2} \log_{\Gamma} n - 1 > \frac{1}{4} \log_{\Gamma} n + 2.$$

Slowest Slow Walks

Theorem (S. (2019))

There exists a finite set R_T and number n_T such that $\mathbf{S}_T(n) \subseteq R_T$ for all $n \geq n_T$.

Proof.

Let $(\alpha', \beta') \in T$ such that $\gamma_{\alpha', \beta'} = \min\{\gamma_{\alpha, \beta} : (\alpha, \beta) \in T\} := \Gamma$. If $(\alpha, \beta) \in \mathbf{S}_T(n)$ and $\gamma := \gamma_{\alpha, \beta}$, then

$$\log_{\gamma} n + 2 \geq s^{\alpha, \beta}(n) = \mathbf{s}_T(n) \geq s^{\alpha', \beta'}(n) \geq \frac{1}{2} \log_{\Gamma} n - 1 > \frac{1}{4} \log_{\Gamma} n + 2.$$

In particular this implies $\log_{\Gamma} \gamma < 4$.

Slowest Slow Walks

Theorem (S. (2019))

There exists a finite set R_T and number n_T such that $\mathbf{S}_T(n) \subseteq R_T$ for all $n \geq n_T$.

Proof.

Let $(\alpha', \beta') \in T$ such that $\gamma_{\alpha', \beta'} = \min\{\gamma_{\alpha, \beta} : (\alpha, \beta) \in T\} := \Gamma$. If $(\alpha, \beta) \in \mathbf{S}_T(n)$ and $\gamma := \gamma_{\alpha, \beta}$, then

$$\log_{\gamma} n + 2 \geq s^{\alpha, \beta}(n) = \mathbf{s}_T(n) \geq s^{\alpha', \beta'}(n) \geq \frac{1}{2} \log_{\Gamma} n - 1 > \frac{1}{4} \log_{\Gamma} n + 2.$$

In particular this implies $\log_{\Gamma} \gamma < 4$. Since $\gamma_{\alpha, \beta}$ is monotonically increasing in α and β , the set $R_T = \{(\alpha, \beta) : \log_{\Gamma} \gamma_{\alpha, \beta} < 4\} \cap T$ is finite. □

Slowest Slow Walks

It is natural to ask how often a given (α, β) lies in $\mathbf{S}_T(n)$.

Slowest Slow Walks

It is natural to ask how often a given (α, β) lies in $\mathbf{S}_T(n)$. To this end, define $\mathbf{I}_T(\alpha, \beta) = \{n : (\alpha, \beta) \in \mathbf{S}_T(n)\}$.

Slowest Slow Walks

It is natural to ask how often a given (α, β) lies in $\mathbf{S}_T(n)$. To this end, define $\mathbf{I}_T(\alpha, \beta) = \{n : (\alpha, \beta) \in \mathbf{S}_T(n)\}$. Let $\Gamma = \min\{\gamma_{\alpha, \beta} : (\alpha, \beta) \in T\}$.

Slowest Slow Walks

It is natural to ask how often a given (α, β) lies in $\mathbf{S}_T(n)$. To this end, define $\mathbf{I}_T(\alpha, \beta) = \{n : (\alpha, \beta) \in \mathbf{S}_T(n)\}$. Let $\Gamma = \min\{\gamma_{\alpha, \beta} : (\alpha, \beta) \in T\}$.

Proposition

Let T be a set of pairs. Given $(\alpha, \beta) \in T$, let $\gamma = \gamma_{\alpha, \beta}$ and $c = \log_{\Gamma} \gamma$. Then

$$n^{-1} |\mathbf{I}_T(\alpha, \beta) \cap [n]| = O(n^{1-c}).$$

Slowest Slow Walks

It is natural to ask how often a given (α, β) lies in $\mathbf{S}_T(n)$. To this end, define $\mathbf{I}_T(\alpha, \beta) = \{n : (\alpha, \beta) \in \mathbf{S}_T(n)\}$. Let $\Gamma = \min\{\gamma_{\alpha, \beta} : (\alpha, \beta) \in T\}$.

Proposition

Let T be a set of pairs. Given $(\alpha, \beta) \in T$, let $\gamma = \gamma_{\alpha, \beta}$ and $c = \log_{\Gamma} \gamma$. Then

$$n^{-1} |\mathbf{I}_T(\alpha, \beta) \cap [n]| = O(n^{1-c}).$$

In particular, for $(1, 4)$ -walks we have $c \approx 1.95$, which explains why they were so hard to find.

Slowest Slow Walks

Corollary

If T is such that there exists a unique pair $(\alpha', \beta') \in T$ with $\gamma_{\alpha', \beta'} = \Gamma$, then almost every n has $\mathbf{S}_T(n) = \{(\alpha', \beta')\}$.

Slowest Slow Walks

Corollary

If T is such that there exists a unique pair $(\alpha', \beta') \in T$ with $\gamma_{\alpha', \beta'} = \Gamma$, then almost every n has $\mathbf{S}_T(n) = \{(\alpha', \beta')\}$.

Let (α, β) be any other element of R_T .

Slowest Slow Walks

Corollary

If T is such that there exists a unique pair $(\alpha', \beta') \in T$ with $\gamma_{\alpha', \beta'} = \Gamma$, then almost every n has $\mathbf{S}_T(n) = \{(\alpha', \beta')\}$.

Let (α, β) be any other element of R_T . Then

$$n^{-1} |\mathbf{I}_T(\alpha, \beta) \cap [n]| = O(n^{1-c})$$

Slowest Slow Walks

Corollary

If T is such that there exists a unique pair $(\alpha', \beta') \in T$ with $\gamma_{\alpha', \beta'} = \Gamma$, then almost every n has $\mathbf{S}_T(n) = \{(\alpha', \beta')\}$.

Let (α, β) be any other element of R_T . Then

$$n^{-1} |\mathbf{I}_T(\alpha, \beta) \cap [n]| = O(n^{1-c}) = o(1).$$

Slowest Slow Walks

Corollary

If T is such that there exists a unique pair $(\alpha', \beta') \in T$ with $\gamma_{\alpha', \beta'} = \Gamma$, then almost every n has $\mathbf{S}_T(n) = \{(\alpha', \beta')\}$.

Let (α, β) be any other element of R_T . Then

$$n^{-1} |\mathbf{I}_T(\alpha, \beta) \cap [n]| = O(n^{1-c}) = o(1).$$

As $|R_T|$ is finite, the density of $m \leq n$ with some other $(\alpha, \beta) \in \mathbf{S}(m)$ is at most $o(1)$. □

Slowest Slow Walks

Corollary

If T is such that there exists a unique pair $(\alpha', \beta') \in T$ with $\gamma_{\alpha', \beta'} = \Gamma$, then almost every n has $\mathbf{S}_T(n) = \{(\alpha', \beta')\}$.

Slowest Slow Walks

Corollary

If T is such that there exists a unique pair $(\alpha', \beta') \in T$ with $\gamma_{\alpha', \beta'} = \Gamma$, then almost every n has $\mathbf{S}_T(n) = \{(\alpha', \beta')\}$.

What if there isn't such a unique pair?

Slowest Slow Walks


Corollary

If T is such that there exists a unique pair $(\alpha', \beta') \in T$ with $\gamma_{\alpha', \beta'} = \Gamma$, then almost every n has $\mathbf{S}_T(n) = \{(\alpha', \beta')\}$.

What if there isn't such a unique pair? For example, if $T = \{(1, 6), (2, 3)\}$ we have $\gamma_{1,6} = \gamma_{2,3} = 3$.

Slowest Slow Walks

Let $e_{\alpha,\beta}(n) = n^{-1} |\{m : m \leq n, \mathbf{S}_T(m) = \{(\alpha, \beta)\}\}|$ for $T = \{(1, 6), (2, 3)\}$.



Here the open black circles are $e_{2,3}(n)$ and the solid blue dots are $e_{1,6}(n)$.

Open Problems

- Why do $(2, 3)$ -walks tend to be slower than $(1, 6)$ -walks?

Open Problems

- Why do $(2, 3)$ -walks tend to be slower than $(1, 6)$ -walks? Note that in general, $g_s^{2,3} > g_s^{1,6}$, which intuitively should make them faster.

Open Problems

- Why do $(2, 3)$ -walks tend to be slower than $(1, 6)$ -walks?
Note that in general, $g_s^{2,3} > g_s^{1,6}$, which intuitively should make them faster.
- Can you say anything more about how often each set appears in $\mathbf{S}_T(n)$?

Open Problems

- Why do $(2, 3)$ -walks tend to be slower than $(1, 6)$ -walks? Note that in general, $g_s^{2,3} > g_s^{1,6}$, which intuitively should make them faster.
- Can you say anything more about how often each set appears in $\mathbf{S}_T(n)$? For example, do there exist infinitely many n for which $(1, 2)$ is an element of $\mathbf{S}(n)$?

Open Problems

- Why do $(2, 3)$ -walks tend to be slower than $(1, 6)$ -walks? Note that in general, $g_s^{2,3} > g_s^{1,6}$, which intuitively should make them faster.
- Can you say anything more about how often each set appears in $\mathbf{S}_T(n)$? For example, do there exist infinitely many n for which $(1, 2)$ is an element of $\mathbf{S}(n)$? Does there exist an n with $|\mathbf{S}(n)| > 2$?

Open Problems

- Why do $(2, 3)$ -walks tend to be slower than $(1, 6)$ -walks? Note that in general, $g_s^{2,3} > g_s^{1,6}$, which intuitively should make them faster.
- Can you say anything more about how often each set appears in $\mathbf{S}_T(n)$? For example, do there exist infinitely many n for which $(1, 2)$ is an element of $\mathbf{S}(n)$? Does there exist an n with $|\mathbf{S}(n)| > 2$?
- What happens with slow Tribonacci walks, i.e.
 $w_{k+3} = w_{k+2} + w_{k+1} + w_k$?

Open Problems

- Why do $(2, 3)$ -walks tend to be slower than $(1, 6)$ -walks? Note that in general, $g_s^{2,3} > g_s^{1,6}$, which intuitively should make them faster.
- Can you say anything more about how often each set appears in $\mathbf{S}_T(n)$? For example, do there exist infinitely many n for which $(1, 2)$ is an element of $\mathbf{S}(n)$? Does there exist an n with $|\mathbf{S}(n)| > 2$?
- What happens with slow Tribonacci walks, i.e.
 $w_{k+3} = w_{k+2} + w_{k+1} + w_k$?
- What happens if you require a slow walk to hit two prescribed numbers n_1 and n_2 ?

Open Problems

- Why do $(2, 3)$ -walks tend to be slower than $(1, 6)$ -walks? Note that in general, $g_s^{2,3} > g_s^{1,6}$, which intuitively should make them faster.
- Can you say anything more about how often each set appears in $\mathbf{S}_T(n)$? For example, do there exist infinitely many n for which $(1, 2)$ is an element of $\mathbf{S}(n)$? Does there exist an n with $|\mathbf{S}(n)| > 2$?
- What happens with slow Tribonacci walks, i.e.
 $w_{k+3} = w_{k+2} + w_{k+1} + w_k$?
- What happens if you require a slow walk to hit two prescribed numbers n_1 and n_2 ? Note that $w_1 = n_1$, $w_2 = n_2$ works, so this is well defined.

The End

Thank You!