

The Random Turán Problem

Sam Spiro, Rutgers University

Define the Turán number $\text{ex}(n, F)$ to be the maximum number of edges that an F -free graph on n vertices can have.

Define the Turán number $\text{ex}(n, F)$ to be the maximum number of edges that an F -free graph on n vertices can have.

Theorem (Mantel 1907)

$$\text{ex}(n, K_3) = \lfloor n^2/4 \rfloor.$$

Define the Turán number $\text{ex}(n, F)$ to be the maximum number of edges that an F -free graph on n vertices can have.

Theorem (Mantel 1907)

$$\text{ex}(n, K_3) = \lfloor n^2/4 \rfloor.$$

Theorem (Erdős-Stone 1946)

$$\text{ex}(n, F) \sim \left(1 - \frac{1}{\chi(F) - 1}\right) \binom{n}{2}.$$

Let $G_{n,p}$ be the random graph on n vertices where each edge is included independently and with probability p .

Let $G_{n,p}$ be the random graph on n vertices where each edge is included independently and with probability p . Let $\text{ex}(G_{n,p}, F)$ be the maximum number of edges that an F -free subgraph of $G_{n,p}$ can have.

Let $G_{n,p}$ be the random graph on n vertices where each edge is included independently and with probability p . Let $\text{ex}(G_{n,p}, F)$ be the maximum number of edges that an F -free subgraph of $G_{n,p}$ can have. For example,

$$\text{ex}(G_{n,1}, F) = \text{ex}(n, F)$$

Let $G_{n,p}$ be the random graph on n vertices where each edge is included independently and with probability p . Let $\text{ex}(G_{n,p}, F)$ be the maximum number of edges that an F -free subgraph of $G_{n,p}$ can have. For example,

$$\text{ex}(G_{n,1}, F) = \text{ex}(n, F),$$

and with high probability

$$\frac{1}{2}p \binom{n}{2} \lesssim \text{ex}(G_{n,p}, K_3) \lesssim p \binom{n}{2}.$$

Let $G_{n,p}$ be the random graph on n vertices where each edge is included independently and with probability p . Let $\text{ex}(G_{n,p}, F)$ be the maximum number of edges that an F -free subgraph of $G_{n,p}$ can have. For example,

$$\text{ex}(G_{n,1}, F) = \text{ex}(n, F),$$

and with high probability

$$\frac{1}{2}p \binom{n}{2} \lesssim \text{ex}(G_{n,p}, K_3) \lesssim p \binom{n}{2}.$$

The lower bound is tight when $p = 1$.

Let $G_{n,p}$ be the random graph on n vertices where each edge is included independently and with probability p . Let $\text{ex}(G_{n,p}, F)$ be the maximum number of edges that an F -free subgraph of $G_{n,p}$ can have. For example,

$$\text{ex}(G_{n,1}, F) = \text{ex}(n, F),$$

and with high probability

$$\frac{1}{2}p \binom{n}{2} \lesssim \text{ex}(G_{n,p}, K_3) \lesssim p \binom{n}{2}.$$

The lower bound is tight when $p = 1$. The upper bound is tight if p is small enough so that $G_{n,p}$ contains no triangles (i.e. $p \ll n^{-1}$)

Let $G_{n,p}$ be the random graph on n vertices where each edge is included independently and with probability p . Let $\text{ex}(G_{n,p}, F)$ be the maximum number of edges that an F -free subgraph of $G_{n,p}$ can have. For example,

$$\text{ex}(G_{n,1}, F) = \text{ex}(n, F),$$

and with high probability

$$\frac{1}{2}p \binom{n}{2} \lesssim \text{ex}(G_{n,p}, K_3) \lesssim p \binom{n}{2}.$$

The lower bound is tight when $p = 1$. The upper bound is tight if p is small enough so that $G_{n,p}$ contains no triangles (i.e. $p \ll n^{-1}$), or if almost no edges are contained in triangles (i.e. if $p \ll n^{-1/2}$).

Theorem (Frankl-Rödl 1986)

Whp,

$$\text{ex}(G_{n,p}, K_3) \sim \frac{1}{2} p \binom{n}{2} \quad p \gg n^{-1/2}.$$

Theorem (Frankl-Rödl 1986)

Whp,

$$\text{ex}(G_{n,p}, K_3) \sim \frac{1}{2} p \binom{n}{2} \quad p \gg n^{-1/2}.$$

Theorem (Conlon-Gowers, Schacht 2010)

Whp,

$$\text{ex}(G_{n,p}, F) \sim \left(1 - \frac{1}{\chi(F) - 1}\right) p \binom{n}{2} \quad p \gg n^{-1/m_2(F)},$$

where $m_2(F) = \max\{\frac{e(F')-1}{v(F')-2} : F' \subseteq F\}$.

What happens for bipartite graphs?

What happens for bipartite graphs?

Conjecture

If F is a bipartite graph which is not a forest and $\text{ex}(n, F) = \Theta(n^\alpha)$, then whp

$$\text{ex}(G_{n,p}, F) = \begin{cases} \Theta(pn^\alpha) & p \gg n^{-1/m_2(F)}, \\ (1 + o(1))p \binom{n}{2} & p \ll n^{-1/m_2(F)}. \end{cases}$$


What happens for bipartite graphs?

Conjecture

If F is a bipartite graph which is not a forest and $\text{ex}(n, F) = \Theta(n^\alpha)$, then whp

$$\text{ex}(G_{n,p}, F) = \begin{cases} \Theta(pn^\alpha) & p \gg n^{-1/m_2(F)}, \\ (1 + o(1))p \binom{n}{2} & p \ll n^{-1/m_2(F)}. \end{cases}$$

This conjecture turns out to be completely false!



Plot of $\text{ex}(G_{n,p}, C_4)$ (Füredi 1991)

Conjecture (McKinley-S.)

If F is a 2-balanced graph with $\text{ex}(n, F) = \Theta(n^\alpha)$ for some $\alpha \in (1, 2]$, then whp

$$\text{ex}(G_{n,p}, F) = \max\{\Theta(p^{\alpha-1}n^\alpha), n^{2-\frac{v(F)-2}{e(F)-1}}(\log n)^{O(1)}\},$$

provided $p \gg n^{-\frac{v(F)-2}{e(F)-1}}$.

Conjecture (McKinley-S.)

If F is a 2-balanced graph with $\text{ex}(n, F) = \Theta(n^\alpha)$ for some $\alpha \in (1, 2]$, then whp

$$\text{ex}(G_{n,p}, F) = \max\{\Theta(p^{\alpha-1}n^\alpha), n^{2-\frac{v(F)-2}{e(F)-1}}(\log n)^{O(1)}\},$$

provided $p \gg n^{-\frac{v(F)-2}{e(F)-1}}$.

The heart of this problem is in showing $\text{ex}(G_{n,p}, F) = \Theta(p^{\alpha-1}n^\alpha)$ when p is large.

Theorem (Kővari-Sós-Turán 1954)

$$\text{ex}(n, K_{s,t}) = O(n^{2-1/s}).$$

Theorem (Kővari-Sós-Turán 1954)

$$\text{ex}(n, K_{s,t}) = O(n^{2-1/s}).$$

Theorem (Morris-Saxton 2013)

$$\text{ex}(G_{n,p}, K_{s,t}) = O(p^{1-1/s} n^{2-1/s}) \text{ for } p \text{ large.}$$

Theorem (Kővari-Sós-Turán 1954)

$$\text{ex}(n, K_{s,t}) = O(n^{2-1/s}).$$

Theorem (Morris-Saxton 2013)

$$\text{ex}(G_{n,p}, K_{s,t}) = O(p^{1-1/s} n^{2-1/s}) \text{ for } p \text{ large.}$$

Moreover, this bound is tight whenever $\text{ex}(n, K_{s,t}) = \Theta(n^{2-1/s})$.

Theorem (Bondy-Simonovits 1974)

$$\text{ex}(n, C_{2b}) = O(n^{1+1/b}).$$

Theorem (Bondy-Simonovits 1974)

$$\text{ex}(n, C_{2b}) = O(n^{1+1/b}).$$

Theorem (Morris-Saxton 2013)

$$\text{ex}(G_{n,p}, C_{2b}) = O(p^{1/b} n^{1+1/b}) \text{ for } p \text{ large.}$$

Theorem (Bondy-Simonovits 1974)

$$\text{ex}(n, C_{2b}) = O(n^{1+1/b}).$$


Theorem (Morris-Saxton 2013)

$$\text{ex}(G_{n,p}, C_{2b}) = O(p^{1/b} n^{1+1/b}) \text{ for } p \text{ large.}$$

Moreover, this is tight whenever $\text{ex}(n, \{C_3, C_4, \dots, C_{2b}\}) = \Theta(n^{1+1/b})$.


Theorem (Faudree-Simonovits 1974)

$$\text{ex}(n, \theta_{a,b}) = O(n^{1+1/b}).$$



Theorem (Faudree-Simonovits 1974)

$$\text{ex}(n, \theta_{a,b}) = O(n^{1+1/b}).$$




Theorem (McKinley-S. 2023++)

$$\text{ex}(G_{n,p}, \theta_{a,b}) = O(p^{1/b} n^{1+1/b}) \text{ for } p \text{ large.}$$

Theorem (Faudree-Simonovits 1974)

$$\text{ex}(n, \theta_{a,b}) = O(n^{1+1/b}).$$






Theorem (McKinley-S. 2023++)

$$\text{ex}(G_{n,p}, \theta_{a,b}) = O(p^{1/b} n^{1+1/b}) \text{ for } p \text{ large.}$$

Moreover, this bound is tight whenever a is sufficiently large in terms of b .




Theorem (Bukh-Conlon 2015)

If T^ℓ is the " ℓ th power of a balanced tree with density b/a ", then $\text{ex}(n, T^\ell) = \Omega(n^{2-a/b})$ if ℓ is sufficiently large.



Theorem (Bukh-Conlon 2015)

If T^ℓ is the " ℓ th power of a balanced tree with density b/a ", then $\text{ex}(n, T^\ell) = \Omega(n^{2-a/b})$ if ℓ is sufficiently large.



Theorem (S. 2022)

$$\text{ex}(G_{n,p}, T^\ell) = \Omega(p^{1-a/b} n^{2-a/b}).$$


Theorem (Jiang-Longbreak 2022)

If F satisfies “mild conditions”, then


$$\text{ex}(G_{n,p}, F) = O(p^{1-m_2^*(F)(2-\alpha)} n^\alpha) \text{ for } p \text{ large,}$$

where $m_2^*(F) = \max\{\frac{e(F')-1}{v(F')-2} : F' \subsetneq F, e(F') \geq 2\}$.

Hypergraphs



Hypergraphs




We define the *loose cycle* C_ℓ^r to be the r -uniform hypergraph obtained by inserting $r - 2$ distinct vertices into each edge of the graph cycle C_ℓ .

Theorem (Nie-S.-Verstaëte 2020; Nie 2023)

For $r \geq 3$, if $p \gg n^{-r+3/2}$ then whp

$$\text{ex}(G_{n,p}^r, C_3^r) = \max\{p^{\frac{1}{2r-3}} n^{2+o(1)}, pn^{r-1+o(1)}\}.$$



Picture due to Jiaxi Nie.

Theorem (Mubayi-Yepremyan 2020; Nie 2023)

For $r \geq 4$, if $p \gg n^{-r+1+\frac{1}{2\ell-1}}$ then whp

$$\text{ex}(G_{n,p}^r, C_{2\ell}^r) = \max\{n^{1+\frac{1}{2\ell-1}}, pn^{r-1}\}.$$


Theorem (Mubayi-Yepremyan 2020; Nie 2023)

For $r \geq 4$, if $p \gg n^{-r+1+\frac{1}{2\ell-1}}$ then whp


$$\text{ex}(G_{n,p}^r, C_{2\ell}^r) = \max\{n^{1+\frac{1}{2\ell-1}}, pn^{r-1}\}.$$

It's suspected that this continues to hold for $r = 3$, but there is a gap for medium values of p .


We say that F is a Berge C_ℓ if it has edges e_1, \dots, e_ℓ and distinct vertices v_1, \dots, v_ℓ with $v_i \in e_i \cap e_{i+1}$ for all i .



We say that F is a Berge C_ℓ if it has edges e_1, \dots, e_ℓ and distinct vertices v_1, \dots, v_ℓ with $v_i \in e_i \cap e_{i+1}$ for all i .



Let \mathcal{B}_ℓ^r denote the set of r -uniform Berge C_ℓ 's.



Plot of $\text{ex}(G_{n,p}^3, \mathcal{B}_4^3)$ (S.-Verstraëte 2021)

Theorem (S.-Verstraëte 2021)


Let K_{s_1, \dots, s_r}^r denote the complete r -partite r -graph with parts of sizes s_1, \dots, s_r . There exist constants $\beta_1, \beta_2, \beta_3, \gamma$ depending on s_1, \dots, s_r such that, for s_r sufficiently large in terms of s_1, \dots, s_{r-1} , we have whp

$$\text{ex}(G_{n,p}^r, K_{s_1, \dots, s_r}^r) = \begin{cases} \Theta(pn^r) & n^{-r} \ll p \leq n^{-\beta_1}, \\ n^{r-\beta_1+o(1)} & n^{-\beta_1} \leq p \leq n^{-\beta_2}(\log n)^\gamma, \\ \Theta(p^{1-\beta_3} n^{r-\beta_3}) & n^{-\beta_2}(\log n)^\gamma \leq p \leq 1. \end{cases}$$

Lower Bound Techniques

Lower Bound Techniques

For the bipartite case, lower bounds for $\text{ex}(n, F)$ often imply lower bounds for $\text{ex}(G_{n,p}, F)$ through “random homomorphisms.”




$\text{ex}(t, F):$



Lower Bound Techniques

For the bipartite case, lower bounds for $\text{ex}(n, F)$ often imply lower bounds for $\text{ex}(G_{n,p}, F)$ through “random homomorphisms.”




$\text{ex}(t, F):$




Lower Bound Techniques

For the bipartite case, lower bounds for $\text{ex}(n, F)$ often imply lower bounds for $\text{ex}(G_{n,p}, F)$ through “random homomorphisms.”



$\text{ex}(t, F):$



Upper Bound Techniques

Upper Bound Techniques

Let $N_m(n, F)$ denote the number of F -free graphs with n vertices and m edges.

Upper Bound Techniques

Let $N_m(n, F)$ denote the number of F -free graphs with n vertices and m edges.

Theorem (Füredi, 1991; Morris-Saxton, 2013)

If $m \geq n^{4/3}(\log n)^2$, then

$$N_m(n, C_4) \leq e^{cm}(\log n)^m \left(\frac{n^{3/2}}{m} \right)^{2m}$$

Upper Bound Techniques

Corollary

If $p \geq n^{-1/3}(\log n)^3$, then a.a.s.

$$\text{ex}(G_{n,p}, C_4) \leq O\left(p^{1/2} n^{3/2} \log n\right).$$

Upper Bound Techniques

Corollary

If $p \geq n^{-1/3}(\log n)^3$, then a.a.s.

$$\text{ex}(G_{n,p}, C_4) \leq O\left(p^{1/2}n^{3/2}\log n\right).$$

Observe that $\text{ex}(G_{n,p}, C_4) \geq m$ if and only if $G_{n,p}$ contains at least one C_4 -free subgraph on m edges.

Upper Bound Techniques

Corollary

If $p \geq n^{-1/3}(\log n)^3$, then a.a.s.

$$\text{ex}(G_{n,p}, C_4) \leq O\left(p^{1/2}n^{3/2}\log n\right).$$

Observe that $\text{ex}(G_{n,p}, C_4) \geq m$ if and only if $G_{n,p}$ contains at least one C_4 -free subgraph on m edges. Define the random variable X to be the number of C_4 -free subgraphs of $G_{n,p}$ on m edges.

Upper Bound Techniques

Corollary

If $p \geq n^{-1/3}(\log n)^3$, then a.a.s.

$$\text{ex}(G_{n,p}, C_4) \leq O\left(p^{1/2}n^{3/2}\log n\right).$$

Observe that $\text{ex}(G_{n,p}, C_4) \geq m$ if and only if $G_{n,p}$ contains at least one C_4 -free subgraph on m edges. Define the random variable X to be the number of C_4 -free subgraphs of $G_{n,p}$ on m edges. Then

$$\Pr[\text{ex}(G_{n,p}, C_4) \geq m] = \Pr[X \geq 1]$$

Upper Bound Techniques

Corollary

If $p \geq n^{-1/3}(\log n)^3$, then a.a.s.

$$\text{ex}(G_{n,p}, C_4) \leq O\left(p^{1/2}n^{3/2}\log n\right).$$

Observe that $\text{ex}(G_{n,p}, C_4) \geq m$ if and only if $G_{n,p}$ contains at least one C_4 -free subgraph on m edges. Define the random variable X to be the number of C_4 -free subgraphs of $G_{n,p}$ on m edges. Then

$$\Pr[\text{ex}(G_{n,p}, C_4) \geq m] = \Pr[X \geq 1] \leq \mathbb{E}[X]$$

Upper Bound Techniques

Corollary

If $p \geq n^{-1/3}(\log n)^3$, then a.a.s.

$$\text{ex}(G_{n,p}, C_4) \leq O\left(p^{1/2} n^{3/2} \log n\right).$$

Observe that $\text{ex}(G_{n,p}, C_4) \geq m$ if and only if $G_{n,p}$ contains at least one C_4 -free subgraph on m edges. Define the random variable X to be the number of C_4 -free subgraphs of $G_{n,p}$ on m edges. Then

$$\Pr[\text{ex}(G_{n,p}, C_4) \geq m] = \Pr[X \geq 1] \leq \mathbb{E}[X] = p^m \cdot N_m(n, C_4)$$

Upper Bound Techniques

Corollary

If $p \geq n^{-1/3}(\log n)^3$, then a.a.s.

$$\text{ex}(G_{n,p}, C_4) \leq O\left(p^{1/2} n^{3/2} \log n\right).$$

Observe that $\text{ex}(G_{n,p}, C_4) \geq m$ if and only if $G_{n,p}$ contains at least one C_4 -free subgraph on m edges. Define the random variable X to be the number of C_4 -free subgraphs of $G_{n,p}$ on m edges. Then

$$\Pr[\text{ex}(G_{n,p}, C_4) \geq m] = \Pr[X \geq 1] \leq \mathbb{E}[X] = p^m \cdot N_m(n, C_4) \approx p^m \cdot \left(\frac{n^{3/2}}{m}\right)^{2m}$$

Summary

- Relatively little is known about this problem (especially for hypergraphs).

Summary

- Relatively little is known about this problem (especially for hypergraphs).
- There are a number of tried and true techniques to employ.

Summary

- Relatively little is known about this problem (especially for hypergraphs).
- There are a number of tried and true techniques to employ.
- Plenty of problems left to solve!