



# Extremal Problems for Random Objects

Sam Spiro, Rutgers University






# Part I: Card Guessing with Feedback



In the “Complete Feedback Model,” we start with a deck of  $mn$  cards where there are  $n$  card types each appearing with multiplicity  $m$ .

In the “Complete Feedback Model,” we start with a deck of  $mn$  cards where there are  $n$  card types each appearing with multiplicity  $m$ .



Let  $\mathcal{C}_{m,n}^+$  and  $\mathcal{C}_{m,n}^-$  be the maximum and minimum expected scores that the player can get in the complete feedback model.

Let  $\mathcal{C}_{m,n}^+$  and  $\mathcal{C}_{m,n}^-$  be the maximum and minimum expected scores that the player can get in the complete feedback model.

### Theorem (Diaconis-Graham, 1981)

For  $n$  fixed,

$$\mathcal{C}_{m,n}^\pm = m \pm c_n \sqrt{m} + o_n(\sqrt{m}).$$

# Another Game

## Another Game

In the “partial feedback model”, the Guesser guesses the next card and is only told whether their guess was correct or not.

## Another Game

In the “partial feedback model”, the Guesser guesses the next card and is only told whether their guess was correct or not. If  $\mathcal{P}_{m,n}^+$  is the maximum expected score in this model,

$$m \leq \mathcal{P}_{m,n}^+$$

## Another Game

In the “partial feedback model”, the Guesser guesses the next card and is only told whether their guess was correct or not. If  $\mathcal{P}_{m,n}^+$  is the maximum expected score in this model,

$$m \leq \mathcal{P}_{m,n}^+ \leq \mathcal{C}_{m,n}^+$$

## Another Game

In the “partial feedback model”, the Guesser guesses the next card and is only told whether their guess was correct or not. If  $\mathcal{P}_{m,n}^+$  is the maximum expected score in this model,

$$m \leq \mathcal{P}_{m,n}^+ \leq \mathcal{C}_{m,n}^+ = m + o_n(m)$$

## Another Game

In the “partial feedback model”, the Guesser guesses the next card and is only told whether their guess was correct or not. If  $\mathcal{P}_{m,n}^+$  is the maximum expected score in this model,

$$m \leq \mathcal{P}_{m,n}^+ \leq \mathcal{C}_{m,n}^+ = m + o_n(m).$$

What happens when  $n$  is large?

## Theorem (Diaconis-Graham-He-S., 2020)

For  $m$  fixed,

$$\begin{aligned}\mathcal{C}_{m,n}^+ &\sim H_m \log(n), \\ \mathcal{C}_{m,n}^- &= \Theta(n^{-1/m}),\end{aligned}$$

where  $H_m$  is the  $m$ th harmonic number.

## Theorem (Diaconis-Graham-He-S., 2020)

For  $m$  fixed,

$$\begin{aligned}\mathcal{C}_{m,n}^+ &\sim H_m \log(n), \\ \mathcal{C}_{m,n}^- &= \Theta(n^{-1/m}),\end{aligned}$$

where  $H_m$  is the  $m$ th harmonic number.

With this we have the trivial bounds

$$m \leq \mathcal{P}_{m,n}^+ \leq \mathcal{C}_{m,n}^+ = O_m(\log n).$$

$$m \leq \mathcal{P}_{m,n}^+ \leq O_m(\log n).$$

$$m \leq \mathcal{P}_{m,n}^+ \leq O_m(\log n).$$

### Theorem (Diaconis-Graham-He-S., 2020)

*There exist  $c, C > 0$  such that if  $n$  is sufficiently large in terms of  $m$ , we have*

$$m + c\sqrt{m} \leq \mathcal{P}_{m,n}^+$$

$$m \leq \mathcal{P}_{m,n}^+ \leq O_m(\log n).$$

### Theorem (Diaconis-Graham-He-S., 2020)

*There exist  $c, C > 0$  such that if  $n$  is sufficiently large in terms of  $m$ , we have*

$$m + c\sqrt{m} \leq \mathcal{P}_{m,n}^+ \leq m + Cm^{3/4} \log m.$$

$$m \leq \mathcal{P}_{m,n}^+ \leq O_m(\log n).$$

### Theorem (Diaconis-Graham-He-S., 2020)

*There exist  $c, C > 0$  such that if  $n$  is sufficiently large in terms of  $m$ , we have*

$$m + c\sqrt{m} \leq \mathcal{P}_{m,n}^+ \leq m + Cm^{3/4} \log m.$$

### Theorem (Z. Nie, 2022)

*If  $n \gg m$ , then*

$$\mathcal{P}_{m,n}^+ = m + \Theta(\sqrt{m}).$$

# A Partial Proof

# A Partial Proof

## Lemma

*Assume that we have played in the partial feedback model for  $t - 1$  rounds such that we have guessed card type  $i$  a total of  $g_i$  times*

# A Partial Proof

## Lemma

Assume that we have played in the partial feedback model for  $t - 1$  rounds such that we have guessed card type  $i$  a total of  $g_i$  times, and let  $S$  be the total number of points scored.

# A Partial Proof

## Lemma

Assume that we have played in the partial feedback model for  $t - 1$  rounds such that we have guessed card type  $i$  a total of  $g_i$  times, and let  $S$  be the total number of points scored. Given this, we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S}.$$

# A Partial Proof

## Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

# A Partial Proof

## Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all  $i$  and  $t$ , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S}$$

# A Partial Proof

## Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all  $i$  and  $t$ , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S} \approx \frac{m}{mn - g_i},$$

# A Partial Proof

## Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all  $i$  and  $t$ , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S} \approx \frac{m}{mn - g_i},$$

At most one  $i$  is guessed more than  $mn/2$  times.

# A Partial Proof

## Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all  $i$  and  $t$ , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S} \approx \frac{m}{mn - g_i},$$

At most one  $i$  is guessed more than  $mn/2$  times. Every other  $j$  has  $\Pr[\pi_t = j] \leq \frac{2}{n}$  for all  $t$ .

# A Partial Proof

## Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all  $i$  and  $t$ , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S} \approx \frac{m}{mn - g_i},$$

At most one  $i$  is guessed more than  $mn/2$  times. Every other  $j$  has  $\Pr[\pi_t = j] \leq \frac{2}{n}$  for all  $t$ . Thus in expectation at most  $mn \cdot (2/n) = 2m$  cards are guessed correctly from this part

# A Partial Proof

## Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all  $i$  and  $t$ , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S} \approx \frac{m}{mn - g_i},$$

At most one  $i$  is guessed more than  $mn/2$  times. Every other  $j$  has  $\Pr[\pi_t = j] \leq \frac{2}{n}$  for all  $t$ . Thus in expectation at most  $mn \cdot (2/n) = 2m$  cards are guessed correctly from this part, and in total at most  $3m$  are guessed correctly in expectation. □

# Card Guessing

## Card Guessing

The probability of drawing four aces in a row with a deck shuffled uniformly at random is  $1/270725$ .

# Adversarial Card Guessing

The probability of drawing four aces in a row with a deck shuffled uniformly at random is  $1/270725$ .

## Adversarial Card Guessing

The probability of drawing four aces in a row with a deck shuffled uniformly at random is  $1/270725$ .

More precisely, we are now considering a two player game played by Shuffler and Guesser.

## Adversarial Card Guessing

The probability of drawing four aces in a row with a deck shuffled uniformly at random is  $1/270725$ .

More precisely, we are now considering a two player game played by Shuffler and Guesser. Let  $\mathcal{C}_{m,n}(G, S)$  be the expected number of points Guesser scores when the two players follow strategies  $G, S$ .

$$\Theta_m(n^{-1/m}) \leq \mathcal{C}_{m,n}(G, \text{Uniform}) \leq H_m \log n + o_m(\log n).$$

$$\Theta_m(n^{-1/m}) \leq \mathcal{C}_{m,n}(G, \text{Uniform}) \leq H_m \log n + o_m(\log n).$$

### Theorem (S., 2021)

*There exists a strategy  $S'$  for Shuffler so that*

$$\mathcal{C}_{m,n}(G, S') \leq \log n + o_m(\log n),$$

*and this bound is best possible.*

## Theorem

*There exists a strategy  $S'$  for Shuffler so that*  
 $\mathcal{C}_{m,n}(G, S') \leq \log n + o_m(\log n).$

## Theorem

*There exists a strategy  $S'$  for Shuffler so that*  
 $\mathcal{C}_{m,n}(G, S') \leq \log n + o_m(\log n).$

A strategy that gives this is the “greedy strategy”, which is such that if there are  $r$  types of cards remaining in the deck, then Shuffler draws each of these card types with probability  $r^{-1}$ .

## Theorem

*There exists a strategy  $S'$  for Shuffler so that*  
 $\mathcal{C}_{m,n}(G, S') \leq \log n + o_m(\log n).$

A strategy that gives this is the “greedy strategy”, which is such that if there are  $r$  types of cards remaining in the deck, then Shuffler draws each of these card types with probability  $r^{-1}$ . E.g. if the deck has a hundred 1's and one 2, we draw a 1 or 2 with probability  $\frac{1}{2}$ .

## Theorem

*There exists a strategy  $S'$  for Shuffler so that*  
 $\mathcal{C}_{m,n}(G, S') \leq \log n + o_m(\log n).$

A strategy that gives this is the “greedy strategy”, which is such that if there are  $r$  types of cards remaining in the deck, then Shuffler draws each of these card types with probability  $r^{-1}$ . E.g. if the deck has a hundred 1’s and one 2, we draw a 1 or 2 with probability  $\frac{1}{2}$ . This gives the desired bound due to a variant of the coupon collector problem. □

## Theorem (S., 2021)

*The greedy strategy is the unique strategy that minimizes the number of correct guesses if Guesser tries to maximize their score.*

## Theorem (S., 2021)

*The greedy strategy is the unique strategy that minimizes the number of correct guesses if Guesser tries to maximize their score.*

Interestingly, the greedy strategy is also the “unique” strategy which maximizes the number of correct guesses if Guesser tries to minimize their score.

# Future Problems

# Future Problems

## Problem

Prove non-trivial bounds for the partial feedback model with adversarial shufflings.

# Future Problems


## Problem

Prove non-trivial bounds for the partial feedback model with adversarial shufflings.

## Conjecture

*The minimum expected score one can get with partial feedback is asymptotic to  $m$ .*

## Part II: Turán's Problem in Random Graphs




Define the Turán number  $\text{ex}(n, F)$  to be the maximum number of edges that an  $F$ -free graph on  $n$  vertices can have.

Define the Turán number  $\text{ex}(n, F)$  to be the maximum number of edges that an  $F$ -free graph on  $n$  vertices can have.

### Theorem (Mantel 1907)


$$\text{ex}(n, K_3) = \lfloor n^2/4 \rfloor.$$



Define the Turán number  $\text{ex}(n, F)$  to be the maximum number of edges that an  $F$ -free graph on  $n$  vertices can have.

Theorem (Mantel 1907)

$$\text{ex}(n, K_3) = \lfloor n^2/4 \rfloor.$$



Theorem (Erdős-Stone 1946)

$$\text{ex}(n, F) = \left(1 - \frac{1}{\chi(F) - 1} + o(1)\right) \binom{n}{2}.$$

Let  $G_{n,p}$  be the random graph on  $n$  vertices where each edge is included independently and with probability  $p$ .

Let  $G_{n,p}$  be the random graph on  $n$  vertices where each edge is included independently and with probability  $p$ . Let  $\text{ex}(G_{n,p}, F)$  be the maximum number of edges that an  $F$ -free subgraph of  $G_{n,p}$  can have.

Let  $G_{n,p}$  be the random graph on  $n$  vertices where each edge is included independently and with probability  $p$ . Let  $\text{ex}(G_{n,p}, F)$  be the maximum number of edges that an  $F$ -free subgraph of  $G_{n,p}$  can have. For example,

$$\text{ex}(G_{n,1}, F) = \text{ex}(n, F)$$

Let  $G_{n,p}$  be the random graph on  $n$  vertices where each edge is included independently and with probability  $p$ . Let  $\text{ex}(G_{n,p}, F)$  be the maximum number of edges that an  $F$ -free subgraph of  $G_{n,p}$  can have. For example,

$$\text{ex}(G_{n,1}, F) = \text{ex}(n, F),$$

and with high probability

$$p \cdot \text{ex}(n, F) \lesssim \text{ex}(G_{n,p}, F) \lesssim p \binom{n}{2}.$$

Let  $G_{n,p}$  be the random graph on  $n$  vertices where each edge is included independently and with probability  $p$ . Let  $\text{ex}(G_{n,p}, F)$  be the maximum number of edges that an  $F$ -free subgraph of  $G_{n,p}$  can have. For example,

$$\text{ex}(G_{n,1}, F) = \text{ex}(n, F),$$

and with high probability

$$p \cdot \text{ex}(n, F) \lesssim \text{ex}(G_{n,p}, F) \lesssim p \binom{n}{2}.$$

The lower bound is tight when  $p = 1$ .

Let  $G_{n,p}$  be the random graph on  $n$  vertices where each edge is included independently and with probability  $p$ . Let  $\text{ex}(G_{n,p}, F)$  be the maximum number of edges that an  $F$ -free subgraph of  $G_{n,p}$  can have. For example,

$$\text{ex}(G_{n,1}, F) = \text{ex}(n, F),$$

and with high probability

$$p \cdot \text{ex}(n, F) \lesssim \text{ex}(G_{n,p}, F) \lesssim p \binom{n}{2}.$$

The lower bound is tight when  $p = 1$ . The upper bound is tight if  $p$  is “small.”

$$\frac{1}{2}p\binom{n}{2} \lesssim \text{ex}(G_{n,p}, K_3) \lesssim p\binom{n}{2},$$

with the lower bound tight for  $p = 1$  and the upper bound tight for  $p \ll n^{-1/2}$ .

$$\frac{1}{2}p\binom{n}{2} \lesssim \text{ex}(G_{n,p}, K_3) \lesssim p\binom{n}{2},$$

with the lower bound tight for  $p = 1$  and the upper bound tight for  $p \ll n^{-1/2}$ .

### Theorem (Frankl-Rödl 1986)

Whp,

$$\text{ex}(G_{n,p}, K_3) \sim \frac{1}{2}p\binom{n}{2} \quad p \gg n^{-1/2}.$$

$$\frac{1}{2}p \binom{n}{2} \lesssim \text{ex}(G_{n,p}, K_3) \lesssim p \binom{n}{2},$$

with the lower bound tight for  $p = 1$  and the upper bound tight for  $p \ll n^{-1/2}$ .

### Theorem (Frankl-Rödl 1986)

Whp,

$$\text{ex}(G_{n,p}, K_3) \sim \frac{1}{2}p \binom{n}{2} \quad p \gg n^{-1/2}.$$

### Theorem (Conlon-Gowers, Schacht 2010)

Whp,

$$\text{ex}(G_{n,p}, F) = p \cdot \left(1 - \frac{1}{\chi(F) - 1} + o(1)\right) \binom{n}{2} \quad p \gg n^{-1/m_2(F)},$$

where  $m_2(F) = \max\left\{\frac{e(F') - 1}{v(F') - 2} : F' \subseteq F\right\}$ .

What happens for bipartite graphs?

What happens for bipartite graphs?

### Conjecture

*If  $F$  is a bipartite graph which is not a forest, then whp*

$$\text{ex}(G_{n,p}, F) = \begin{cases} \Theta(p \cdot \text{ex}(n, F)) & p \gg n^{-1/m_2(F)}, \\ (1 + o(1))p \binom{n}{2} & p \ll n^{-1/m_2(F)}. \end{cases}$$


What happens for bipartite graphs?

### Conjecture

*If  $F$  is a bipartite graph which is not a forest, then whp*

$$\text{ex}(G_{n,p}, F) = \begin{cases} \Theta(p \cdot \text{ex}(n, F)) & p \gg n^{-1/m_2(F)}, \\ (1 + o(1))p \binom{n}{2} & p \ll n^{-1/m_2(F)}. \end{cases}$$

This conjecture turns out to be completely false!



Plot of  $\text{ex}(G_{n,p}, C_4)$  (Füredi 1991)

## Conjecture (McKinley-S.)


If  $F$  is a graph with  $\text{ex}(n, F) = \Theta(n^\alpha)$  for some  $\alpha \in (1, 2]$ , then whp

$$\text{ex}(G_{n,p}, F) = \max\{\Theta(p^{\alpha-1} n^\alpha), n^{2-1/m_2(F)} (\log n)^{O(1)}\},$$

provided  $p \gg n^{-1/m_2(F)}$ .


## Theorem (Kővari-Sós-Turán 1954)

$$\text{ex}(n, K_{s,t}) = O(n^{2-1/s}).$$



Theorem (Kővari-Sós-Turán 1954)

$$\text{ex}(n, K_{s,t}) = O(n^{2-1/s}).$$




Theorem (Morris-Saxton 2013)

$$\text{ex}(G_{n,p}, K_{s,t}) = O(p^{1-1/s} n^{2-1/s}) \text{ for } p \text{ large.}$$

Theorem (Kővari-Sós-Turán 1954)

$$\text{ex}(n, K_{s,t}) = O(n^{2-1/s}).$$




Theorem (Morris-Saxton 2013)

$$\text{ex}(G_{n,p}, K_{s,t}) = O(p^{1-1/s} n^{2-1/s}) \text{ for } p \text{ large.}$$

Moreover, this bound is tight whenever  $\text{ex}(n, K_{s,t}) = \Theta(n^{2-1/s})$ .


## Theorem (Bondy-Simonovits 1974)

$$\text{ex}(n, C_{2b}) = O(n^{1+1/b}).$$



Theorem (Bondy-Simonovits 1974)

$$\text{ex}(n, C_{2b}) = O(n^{1+1/b}).$$




Theorem (Morris-Saxton 2013)

$$\text{ex}(G_{n,p}, C_{2b}) = O(p^{1/b} n^{1+1/b}) \text{ for } p \text{ large.}$$

Theorem (Bondy-Simonovits 1974)

$$\text{ex}(n, C_{2b}) = O(n^{1+1/b}).$$




Theorem (Morris-Saxton 2013)

$$\text{ex}(G_{n,p}, C_{2b}) = O(p^{1/b} n^{1+1/b}) \text{ for } p \text{ large.}$$

Moreover, this is tight whenever  $\text{ex}(n, \{C_3, C_4, \dots, C_{2b}\}) = \Theta(n^{1+1/b})$ .


## Theorem (Faudree-Simonovits 1974)

$$\text{ex}(n, \theta_{a,b}) = O(n^{1+1/b}).$$



Theorem (Faudree-Simonovits 1974)

$$\text{ex}(n, \theta_{a,b}) = O(n^{1+1/b}).$$




Theorem (McKinley-S. 2023)

For  $a \geq 100$ ,

$$\text{ex}(G_{n,p}, \theta_{a,b}) = O(p^{1/b} n^{1+1/b}) \text{ for } p \text{ large.}$$

Theorem (Faudree-Simonovits 1974)

$$\text{ex}(n, \theta_{a,b}) = O(n^{1+1/b}).$$



Theorem (McKinley-S. 2023)

For  $a \geq 100$ ,



$$\text{ex}(G_{n,p}, \theta_{a,b}) = O(p^{1/b} n^{1+1/b}) \text{ for } p \text{ large.}$$

Moreover, this bound is tight whenever  $a$  is sufficiently large in terms of  $b$ .

## Theorem (Bukh-Conlon 2015)

If  $T^\ell$  is the “ $\ell$ th power of a balanced tree” and  $\ell$  is sufficiently large, then



$$\text{ex}(n, T^\ell) = \Omega(n^{2-\rho(T)}).$$



## Theorem (Bukh-Conlon 2015)

If  $T^\ell$  is the “ $\ell$ th power of a balanced tree” and  $\ell$  is sufficiently large, then

$$\text{ex}(n, T^\ell) = \Omega(n^{2-\rho(T)}).$$




## Theorem (S. 2022)

$$\text{ex}(G_{n,p}, T^\ell) = \Omega(p^{1-\rho(T)} n^{2-\rho(T)}),$$

provided  $\ell$  is sufficiently large.

# Hypergraphs



## Theorem (S.-Verstraëte 2021)

Let  $K_{s_1, \dots, s_r}^r$  denote the complete  $r$ -partite  $r$ -graph with parts of sizes  $s_1, \dots, s_r$ . There exist constants  $\beta_1, \beta_2, \beta_3, \gamma$  depending on  $s_1, \dots, s_r$  such that, for  $s_r$  sufficiently large in terms of  $s_1, \dots, s_{r-1}$ , we have whp

$$\text{ex}(G_{n,p}^r, K_{s_1, \dots, s_r}^r) = \begin{cases} \Theta(pn^r) & n^{-r} \ll p \leq n^{-\beta_1}, \\ n^{r-\beta_1+o(1)} & n^{-\beta_1} \leq p \leq n^{-\beta_2}(\log n)^\gamma, \\ \Theta(p^{1-\beta_3} n^{r-\beta_3}) & n^{-\beta_2}(\log n)^\gamma \leq p \leq 1. \end{cases}$$

## Question

Does the McKinley-Spiro conjecture extend to hypergraphs?

## Question


Does the McKinley-Spiro conjecture extend to hypergraphs?

## Theorem (Nie-S. 2023 (Informal))

*Any hypergraph which is not Sidorenko fails to have a flat middle range.*


## Other Hypergraph Results


- ① Solved for loose triangles (Nie-S.-Verstraëte 2020; Nie 2023)
- ② Solved for loose even cycles of uniformity  $r \geq 4$  (Mubayi-Yepremyan 2020; Nie 2023)
- ③ (Non-optimal) bounds for Berge cycles (S.-Verstraëte 2021; Nie 2023)
- ④ \*Improved lower bound for non-Sidorenko hypergraphs (Nie-S. 2023)
- ⑤ \*Lifting upper bounds from graphs to hypergraphs (Nie-S. 20XX++)




# Lower Bound Techniques

# Lower Bound Techniques




$$Ex(P_h, F) =$$


# Lower Bound Techniques




$\mathcal{P}_X(P_h, F) =$


A hand-drawn square representing the domain  $\mathcal{P}_X(P_h, F)$ . The square has vertices at the corners and is drawn with a single continuous line.



# Lower Bound Techniques



$G_{h,\rho} =$




$\rho_x(p_h, F) =$  

# Future Problems

# Future Problems

## Problem


Prove tight bounds for the 3-uniform loose 4-cycles.



# Future Problems

## Problem

Prove tight bounds for the 3-uniform loose 4-cycles.



## Problem

Prove tight bounds for subdivisions of complete bipartite graphs.

# Thanks!