

Methods in Extremal Combinatorics

Sam Spiro

May 23, 2024

Contents

I	Basic Probabilistic Methods	6
1	Introduction	7
2	Some Random Examples	12
2.1	Graphs with Small and Large Chromatic Numbers	12
2.2	Random Permutations and Extremal Set Theory	13
2.3	The Crossing Lemma and Incidence Geometry	16
3	The Lovász Local Lemma	18
3.1	Applications	19
3.2	Related Lemmas	21
4	The Second Moment Method	22
4.1	Concentration Inequalities	22
4.2	The Rödl Nibble	23
4.3	Steiner Systems	26
II	Further Probabilistic Methods	28
5	Dependent Random Choice	29
6	Strong Concentration Inequalities	34
6.1	The Chernoff Bound	34

6.2	Martingales	35
6.3	Talagrand's Inequality	37
7	Coupling	40
7.1	Graphs and Digraphs	40
7.2	Graphs and Hypergraphs	42
8	Random Algebraic Constructions	48
8.1	Random Multilinear Maps	48
8.2	Random Polynomial Graphs	50
8.3	Multicolor Ramsey Numbers	52
III	Spread Hypergraphs	55
9	The Spreadness Theorem	55
9.1	Applications	55
9.2	Proof of the Spreadness Theorem	57
9.3	Losing Logarithms	59
10	TODOThe Park-Pham Theorem	61
11	Spread Approximations	62
11.1	Further Results	65
IV	Entropy	66
12	Introduction	66
12.1	The Main Properties	66
12.2	Basic Application 1: Binomial Coefficients	68
12.3	Basic Application 2: Walks in Graphs	69
12.4	Further Applications	71
13	Shearer's Lemma	72
14	Random Chain Rules	76

15 The Union-Closed Sets Conjecture	80
V Hypergraph Containers	84
16 Introduction	85
17 Graph Containers	86
17.1 Regular Graphs	88
17.2 A Randomized Sperner's Theorem	90
17.3 Counting Sidon Sets	93
18 A Proof of an r-Uniform Container Lemma	94
18.1 An Informal Discussion	94
18.2 A Formal Algorithm	95
18.3 Proof of The Formal Result	98
19 Hypergraph Containers and Triangle-Free Graphs	100
19.1 Mantel's Theorem in Random Graphs	102
19.2 Maximal Triangle-Free Graphs	104
20 TODOCounting F-free graphs	108
VI Linear Algebra Methods	109
21 Modular Intersections	110
22 Pseudo-Adjacency Matrices	115
22.1 Huang's Theorem	115
22.2 Hoffman's Bound and Erdős-Ko-Rado	117
23 The Polynomial Method I: Slice Rank	122
23.1 Slice Rank	122
23.2 The Capset Problem	123
23.3 Nonuniform Sunflowers	125
24 The Polynomial Method II: Combinatorial Nullstellensatz	127

24.1	Cauchy-Davenport	128
24.2	Covering the Hypercube	129
24.3	Regular Subgraphs mod p	130
VII	Other Methods	131
25	The Delta-system Method	132
25.1	Proof of the Semilattice Intersection Lemma	136
25.2	Other Hypergraph Approximations	137
26	Expansion and α-maximality	138
VIII	Bibliography	145
IX	Appendix: Spectral Graph Theory TODO	151

Preface

This is a work in progress. There are typos, missing references, and so on scattered throughout. **Please let me know if you notices such errors, find anything confusing, or if you have any other suggestions!** If you prefer, you can let me know about any of this anonymously through [this link](#).

The following is a set of lecture notes for a graduate level course in extremal combinatorics. These notes focus on standard methods that have been used to solve a large number of problems in extremal combinatorics. Throughout I assume basic knowledge of asymptotic analysis, probability theory, and linear algebra.

Due to the sheer scope of extremal combinatorics, there are many methods which I am not able to cover at all (and there is no topic which I am able to cover in complete depth). Below is a small list of methods and topics ***not*** covered by this text (which may be written up at some point), as well as some sources for thorough treatments of the topics.

- Extremal Combinatorics in general: see books of Lovasz [80] or Bollobás [16]; surveys by Simonovits and Szemerédi [102] and Füredi and Simonovits [53]; and online courses by [Morris](#) and [Gowers](#) [60].
- The Regularity Lemma: see the excellent book by Zhao [112] (as well as his corresponding [video lectures](#)).
- Additive combinatorics and discrete Fourier analysis: again Zhao [112] is a good introductory text, see also the book by Tao and Vu [107] and the online course by [Prendiville](#).
- Discrete geometry: see the books by Sheffer [101] and Matoušek [84], as well as the online minicourse on finite geometry and Ramsey theory by [Bishnoi](#).
- Statistical mechanics: see notes by [Will Perkins](#).
- The discharging method: see the survey by Cranston and West [30].
- Absorption [need to find a reference](#).

Part I

Basic Probabilistic Methods

This part is based heavily off of the book by Alon and Spencer [\[6\]](#) (which goes into much more depth on the topic), as well as lecture notes by Verstraëte.

1 Introduction

One of the most exciting developments in extremal combinatorics over the past century has been the incorporation of ideas and tools from probability theory into solving combinatorial problems. The first such use was by Erdős who proved an exponential lower bound for Ramsey numbers. We recall that the *Ramsey number* $R(s, t)$ is the smallest integer N such that any 2-coloring of the edges of K_N contains a monochromatic clique.

Theorem 1.1 ([40]). *For all n , we have*

$$R(n, n) \geq (1 + o(1)) \frac{n}{e\sqrt{2}} 2^{n/2}.$$

This is essentially the best known lower bound (though we prove a slightly stronger bound in Theorem 3.3). The best known upper bound is roughly 4^n , so there's still quite a gap!

For this proof and throughout the text, we make heavy use of the union bound: if A, B are events in a probability space, then $\Pr[A \cup B] \leq \Pr[A] + \Pr[B]$. Often we will use an equivalent version: $\Pr[\bar{A} \cap \bar{B}] \leq 1 - \Pr[A] - \Pr[B]$, which follows from De Morgan's laws.

Proof. Let G be a **random** coloring of K_N with N to be determined later¹. That is, for each edge of K_N , we independently and uniformly choose the edge to be colored either red or blue. The key observation is that if $\Pr[G \text{ contains no monochromatic } K_n] > 0$, then there exists a coloring of K_N with no monochromatic K_n (since otherwise the probability would be zero), proving the desired lower bound.

If S is a set of n vertices, we let A_S be the event that G contains a monochromatic K_n on S . With this we have

$$\Pr[G \text{ contains a monochromatic } K_n] = \Pr \left[\bigcup_{S \in \binom{[N]}{n}} A_S \right] \leq \sum_{S \in \binom{[N]}{n}} \Pr[A_S] = \binom{N}{n} \cdot 2^{1 - \binom{n}{2}}.$$

If this quantity is less than 1, then we can conclude that $\Pr[G \text{ contains no monochromatic } K_n] > 0$, so our goal is to choose N as large as possible so that this happens. By using the bound $\binom{N}{n} \leq (eN/n)^n$ (which we will use many times throughout the text), we see that it suffices to have²

$$1 > (eN/n) 2^{1 - \binom{n}{2}} = 2(eN/n 2^{(n-1)/2})^n.$$

Solving this shows that the desired bound holds if $N < 2^{1/n} \cdot \frac{n}{e\sqrt{2}} 2^{n/2}$, proving the result³. \square

¹When trying to prove results in extremal and probabilistic combinatorics, one often uses a method that depends on some parameter such as N or p . Typically it is best to proceed through the argument without deciding what N, p is ahead of time, and only in the end do you optimize your parameter to give you the best bounds possible.

²Finding the “right” way to bound expressions like this takes time and practice. A reasonable strategy for these sorts of problems is try and get all of the main terms to have the same form (e.g. x^n in this example). Much more about the art of asymptotic analysis can be found in the book *Asymptopia* by Spencer [104].

³In fact, a closer analysis of this proof shows that asymptotically, almost every coloring of K_N with $N = (2 - \epsilon)^{n/2}$ contains no monochromatic K_n . Despite almost every coloring working, we know of no explicit coloring that gives more than a polynomial lower bound for $R(n, n)$. Thus the probabilistic method gives us a way to find [the hay in the haystack](#).

The proof of Theorem 1.1 implicitly used the following general principle, which is at the heart of the probabilistic method.

- (*) Let T be an object chosen randomly from a set \mathcal{T} (in some way) and P some property that objects in \mathcal{T} could have. If $\Pr[T \text{ has property } P] > 0$, then there exists some $T' \in \mathcal{T}$ with this property.

We now turn to another classical extremal problem with a slick probabilistic proof. Recall that $\alpha(G)$ denotes the largest independent set of a graph G , i.e. the largest set of vertices I such that there exist no edge contained in I .

Theorem 1.2 (Caro-Wei Bound). *Let G be an n -vertex graph with degrees d_1, \dots, d_n . Then*

$$\alpha(G) \geq \sum \frac{1}{d_i + 1}.$$

Moreover, equality holds if and only if G is a disjoint union of cliques.

Here and throughout the text we make heavy use of the principle of linearity of expectation: for two (possibly dependent) real-valued random variables, we have $\mathbb{E}[X + Y] = \mathbb{E}[X] + \mathbb{E}[Y]$.

Proof. For π a bijection from $V(G)$ to $[n]$, we define

$$I(\pi) = \{v \in V(G) : \pi(v) < \pi(u) \forall u \in N(v)\}.$$

That is, $I(\pi)$ is the set of vertices which are smaller than all of their neighbors under π . Observe that $I(\pi)$ is an independent set (if u, v are adjacent we must have, say $\pi(v) < \pi(u)$, in which case $u \notin I(\pi)$), so in particular $\alpha(G) \geq |I(\pi)|$ for all π .

Let π be a random bijection chosen uniformly amongst all bijections from $V(G)$ to $[n]$, and let 1_v be the indicator variable which is 1 if $v \in I(\pi)$ and 0 otherwise. Note that regardless of what π is, we have $\alpha(G) \geq |I(\pi)| = \sum 1_v$, so by linearity of expectation we have

$$\alpha(G) \geq \mathbb{E}[|I(\pi)|] = \sum \mathbb{E}[1_v] = \sum \Pr[1_v = 1]. \quad (1)$$

Observe that $1_v = 1$ if and only if $\pi(v) = \min_{u \in \{v\} \cup N(v)} \pi(u)$. Since π was chosen uniformly at random, each $u \in \{v\} \cup N(v)$ is equally likely to achieve this minimum, so $\Pr[1_v = 1] = \frac{1}{d(v)+1}$, and plugging this into (1) gives the result.

Note that equality holds in (1) if and only if $I(\pi)$ is an independent set of maximum size for all bijections π . It is not too difficult to show that this holds if and only if G is a disjoint union of cliques, and we leave this as an exercise to the reader. \square

Theorem 1.2 implies Turán's theorem, which is essentially the result that jump started the entire field of extremal combinatorics¹ (though the original proof was not probabilistic).

¹The first theorem in extremal combinatorics is typically attributed to Mantel, which is the $r = 3$ case of Turán's Theorem. However, it wasn't until Turán's result 30 years later that the field really took off.

To state this result, we define $\text{ex}(n, F)$ to be the largest number of edges that an n -vertex F -free graph can have¹ which is called the *Turán number* or *extremal number* of F . We define the *Turán graph* $T_r(n)$ to be the complete n -vertex r -partite graph with parts of sizes as equal as possible. We let $t_r(n) = e(T_r(n))$. For example, $T_2(n) = K_{\lfloor n/2 \rfloor, \lceil n/2 \rceil}$ and $t_2(n) = \lfloor n/2 \rfloor \cdot \lceil n/2 \rceil = \lfloor n^2/4 \rfloor$. More generally we have

$$t_r(n) \leq \binom{r}{2} (n/r)^2 = \frac{r-1}{r} \cdot \frac{n^2}{2} = \left(1 - \frac{1}{r}\right) \frac{n^2}{2},$$

with equality holding if $r|n$ and otherwise $t_r(n)$ is the floor of this upper bound.

Corollary 1.3 (Turán's Theorem). *For all $r \leq n$ we have*

$$\text{ex}(n, K_r) = t_{r-1}(n).$$

Moreover, $T_{r-1}(n)$ is the unique n -vertex K_r -free graph with $t_{r-1}(n)$ edges.

Proof. The lower bound $\text{ex}(n, K_r) \geq t_{r-1}(n)$ follows by considering $T_{r-1}(n)$. Let G be an n -vertex K_r -free graph with degrees d_1, \dots, d_n . Observe that the complement \overline{G} contains no independent set of size r , so by Theorem 1.2 we have

$$r-1 \geq \alpha(\overline{G}) \geq \sum \frac{1}{n-d_i}. \quad (2)$$

Observe that if x, y are positive numbers, then²

$$\frac{1}{x} + \frac{1}{y} \geq \frac{1}{\frac{1}{2}(x+y)} + \frac{1}{\frac{1}{2}(x+y)}$$

with equality holding if and only if $x = y$. In view of this inequality, we see that (2) is minimized when all of the d_i are as close together as possible. Because $\sum d_i = 2e(G)$, we have

$$r-1 \geq n \cdot \frac{1}{n-2e(G)/n} = \frac{n^2}{n^2-e(G)} \implies e(G) \leq \left(1 - \frac{1}{r-1}\right) n^2/2,$$

so $e(G) \leq t_{r-1}(n)$ as desired. Moreover, to have equality, \overline{G} must be a union of cliques with sizes as close as possible to each other, i.e. G must be a complete r -partite graph with parts having sizes as close as possible to each other, i.e. G must be the Turán graph. \square

In addition to using the probabilistic method to get an upper bound for $\text{ex}(n, K_n)$ as in Corollary 1.3, one can also use it to give a general lower bound for $\text{ex}(n, F)$.

Theorem 1.4. *Let F be a graph with v vertices and $e \geq 2$ edges. If $e \geq v$, then*

$$\text{ex}(n, F) = \Omega_v(n^{2-\frac{v-2}{e-1}}).$$

¹Throughout the text, a graph being F -free means that it contains no subgraph which is isomorphic to F (and we don't care whether this subgraph is induced or not).

²By multiplying both sides of the above expression by $xy(x+y)$, we see that this is equivalent to saying $y(x+y) + x(x+y) \geq 4xy$, which is equivalent to saying $x^2 - 2xy + y^2 = (x-y)^2 \geq 0$.

For this proof we use an object that is fundamental to probabilistic and extremal combinatorics. This is the *Erdős-Rényi random graph* $G_{n,p}$, which is the random graph on n vertices that contains each edge $e \in E(K_n)$ independently and with probability p . For example, $G_{n,1} = K_n$ and $G_{n,1/2}$ is equally likely to be any labeled graph on n vertices. The random graph is an incredibly fascinating object in its own right. We will not discuss it in too much depth in this text, see the book by Frieze and Karoński [51] for a thorough treatment of it.

Proof. Let $G_{n,p}$ be the random graph with p a quantity to be determined later. Let X denote the number of copies of F in $G_{n,p}$. For S a set of v vertices, let 1_S be the indicator variable which is 1 if S contains a copy of F in $G_{n,p}$ and which is 0 otherwise. With this,

$$\sum 1_S \leq X \leq v! \sum 1_S,$$

since each set of v vertices contains at most $v!$ copies of F . To have $1_S = 1$, we in particular need S to contain at least e edges, so

$$\Pr[1_S = 1] \leq \sum_{k \geq e} \binom{\binom{v}{2}}{k} p^k (1-p)^{\binom{v}{2}-k} \leq v^2 2^{v^2} p^e \leq 4^{v^2} p^e.$$

In total this gives

$$\mathbb{E}[X] \leq v! \binom{n}{v} \cdot 4^{v^2} p^e \leq (4^v n)^v p^e.$$

Observe that when $p \gg n^{v/e}$, the calculation above suggests that $G_{n,p}$ will contain copies of F (at least in expectation), so $G_{n,p}$ will not work as an F -free graph for this range of p . However, we can get around this by using the following trick known as the method of alterations. Let G be any subgraph of $G_{n,p}$ obtained by deleting an edge from each copy of F in $G_{n,p}$. By definition G will be F -free. Moreover, the number of edges that G has is at least $e(G_{n,p}) - X$ since at most X of the original edges from $G_{n,p}$ are deleted. Using linearity of expectation gives

$$\mathbb{E}[e(G)] \geq \mathbb{E}[e(G_{n,p}) - X] \geq p \binom{n}{2} - (4^v n)^v p^e \geq \frac{1}{4} p n^2 - (4^v n)^v p^e. \quad (3)$$

At this point we want to choose p so that the above expression is roughly maximized. Intuitively this will happen when both terms on the rightside of (3) are roughly equal to each other, i.e. when $p n^2 \approx n^v p^e$. This suggests taking $p \approx n^{\frac{2-v}{e-1}}$. And indeed, after playing around for a bit, one sees that, for example, taking $p = \frac{1}{20 \cdot 16^v} n^{\frac{2-v}{e-1}}$ and plugging it into (3) gives¹ $\mathbb{E}[e(G)] \geq \frac{1}{160 \cdot 16^v} n^{2-\frac{2-v}{e-1}}$. Because G is a (random) F -free graph, by (*) there exists some deterministic graph G' which is F -free with this many edges, proving the result. \square

For many F , there are known constructions which give much better lower bounds for $\text{ex}(n, F)$ than Theorem 1.4. However, this is the best known lower bound which works for arbitrary F .

The method used in this proof is known as the method of alterations. Typically this works by defining some initial random set A (e.g. a set of edges of a graph) which contains some bad

¹Here we use $4^{v^2} \leq 4^{ve}$ and that $e \geq 2$.

subsets B (e.g. subsets of edges forming a forbidden graph F). We then define a random set A' by deleting an element from each bad subset B , giving that $|A'| \geq |A| - |B|$ and that A' has no bad subsets. At this point we win provided

$$\mathbb{E}[|A'|] = \mathbb{E}[|A|] - \mathbb{E}[|B|]$$

is large. Typically the expectations $\mathbb{E}[|A|], \mathbb{E}[|B|]$ depend on some common parameter p , and we often optimize this expression by finding p such that $\mathbb{E}[|A|] \approx \mathbb{E}[|B|]$, and then ultimately choosing p to be a bit smaller than this so that, say, $\mathbb{E}[|B|] \leq \frac{1}{2}\mathbb{E}[|A|]$.

(**) The method of alterations detailed above is often very useful.

The last core tenant of the probabilistic method that we have implicitly used throughout this section is the following.

(***) If one is trying to find a nice object, one should always try and see how well a random object does (possibly after applying alterations).

For example, the most straightforward random coloring gave the bound of Theorem 1.1, and the random graph together with alterations gave Theorem 1.4.

Lastly, we note that in principle many of these results could be proven without needing to use probability. However, for certain problems a probabilistic perspective is genuinely useful since it allows one to use powerful tools from probability theory (e.g. martingales and concentration inequalities). Even when it isn't strictly needed, probability often provides for a much clearer perspective on a problem.

2 Some Random Examples

This section consists of an assorted collection of examples which provides both practice with the general principles of the probabilistic method, as well as proofs of many fundamental results from extremal combinatorics.

2.1 Graphs with Small and Large Chromatic Numbers

We start with a very simple example that will be used throughout the text (often without reference).

Lemma 2.1. *If G is an n -vertex graph, then there exists a bipartite subgraph $G' \subseteq G$ such that $e(G') \geq \frac{1}{2}e(G)$. Moreover, we can choose G' such that its partition classes U, V have sizes $\lfloor n/2 \rfloor, \lceil n/2 \rceil$.*

Given this lemma, if you want to prove a statement of the form “any graph G with $\Omega(m)$ edges has some monotone graph property”, then you only need to consider graphs which are (balanced) bipartite.

Proof. The first part is very easy: let $U \subseteq V(G)$ be obtained by including each vertex independently and with probability $\frac{1}{2}$, and let $V = V(G) \setminus U$. Let G' be the graph which consists of every edge $e \in E(G)$ with one vertex in U and one vertex in V . It is easy to check that $\mathbb{E}[e(G')] = \frac{1}{2}e(G)$, so such a (bipartite) subgraph exists.

The second part is conceptually easy but computationally a little tedious. Let $U \subseteq V(G)$ be a set of size $\lfloor n/2 \rfloor$ chosen uniformly at random and let $V = V(G) \setminus U$. Let G' be the graph which consists of every edge $e \in E(G)$ with one vertex in U and one vertex in V . Observe that the probability that a given edge $xy \in E(G)$ is in G' is exactly

$$1 - \frac{\lfloor n/2 \rfloor \cdot (\lfloor n/2 \rfloor - 1)}{n(n-1)} - \frac{\lceil n/2 \rceil \cdot (\lceil n/2 \rceil - 1)}{n(n-1)} \geq \frac{1}{2},$$

with the last step following from a case analysis based on whether n is even or odd. Thus in expectation G' has at least $\frac{1}{2}e(G)$ edges, so such a balanced bipartite subgraph of G must exist. \square

A graph G is said to have *girth* ℓ if its smallest cycle is of size ℓ , and we say that it has infinite girth if G has no cycles. Observe that graphs of large girth locally look like a tree, i.e. if you pick any vertex v , then the graph induced by every vertex within distance ℓ of v is a tree. In particular, “locally” graphs of large girth can be properly colored using few colors, but does this necessarily hold globally as well? That is, does there exist graphs with girth at least ℓ and chromatic number at least k for all ℓ, k ? A clever (random) argument of Erdős shows that such a graph does indeed exist.

Theorem 2.2 (Erdős). *For all ℓ, k there exist graphs of girth at least ℓ and chromatic number at least k .*

For this proof we use Markov's inequality: if X is a non-negative real-valued random variable, then $\Pr[X \geq x] \leq \mathbb{E}[X]/x$ for $x > 0$.

Proof. Consider $G_{n,p}$ with n, p to be determined later. Let $X_{\leq \ell}$ denote the number of cycles in $G_{n,p}$ of size at most ℓ . Linearity of expectation gives

$$\mathbb{E}[X_{\leq \ell}] \leq \sum_{t=3}^{\ell} n^t \cdot p^t \leq \ell(pn)^\ell.$$

Thus if we wanted $G_{n,p}$ to have girth smaller than ℓ with high probability, by Markov's inequality it would suffice to take $p \ll n^{-1}$. Unfortunately this naive approach is too weak since in this case $G_{n,p}$ will have very small chromatic number. To get around this, we will take p slightly larger than n^{-1} and then use alterations to delete a vertex from every small cycle of $G_{n,p}$. With some foresight¹ we will take $p = n^{-1+1/2\ell}$. With this we see that

$$\Pr[X_{\leq \ell} \geq n/2] \leq \mathbb{E}[X_{\leq \ell}]/(n/2) \leq 2\ell n^{-1/2}. \quad (4)$$

We now turn to the chromatic number of $G_{n,p}$, which is a slightly trickier quantity to get a handle on. To do this we use the inequality $\chi(G) \geq |V(G)|/\alpha(G)$, which follows from the fact that a k -coloring of G is a partition of $V(G)$ into independent sets. Thus for $G_{n,p}$ to have large chromatic number, it suffices to show that all of its independent sets are small. For m an integer we let Y_m be the number of independent sets of size m in $G_{n,p}$. Using linearity of expectation and $(1-x) \leq e^{-x}$ gives for $m \geq 2$

$$\mathbb{E}[Y_m] = \binom{n}{m} \cdot (1-p)^{\binom{m}{2}} \leq n^m \cdot (e^{-p(m-1)/2})^m \leq (ne^{-pm/4})^m.$$

By Markov's inequality and our choice of $p = n^{-1+1/2\ell}$, we find for $m = n/2k$ and n sufficiently large in terms of k, ℓ that

$$\Pr[Y_{n/2k} \geq 1] \leq (ne^{-n^{1/2\ell}/8k})^m < \frac{1}{2}. \quad (5)$$

By combining (4) and (5), we see for n sufficiently large that $X_{\leq \ell} < n/2$ and $Y_{n/2k} = 0$ both occur with positive probability, i.e. there exists a graph G such that both of these events occur. Let G' be G after deleting a vertex from each cycle of length at most ℓ in G . This deletes at most half the vertices of G by assumption of $X_{\leq \ell}$, and we have $\alpha(G') \leq \alpha(G) \leq n/2k$. Thus

$$\chi(G') \geq |V(G')|/\alpha(G') \geq k,$$

proving the result. □

2.2 Random Permutations and Extremal Set Theory

In this subsection, we use random permutations (similar to the proof of Theorem 1.2) to prove two famous results from extremal set theory, which is roughly speaking the study of

¹The exact choice of p doesn't matter here, the important thing is to take $p = n^{-1+\alpha}$ with $0 < \alpha < 1/\ell$.

extremal problems for hypergraphs. We only scratch the surface of this topic, see Frankl and Tokushige [47] for a more thorough treatment.

We start with the most fundamental theorem in extremal set theory: the Erdős-Ko-Rado theorem.

Theorem 2.3 (Erdős-Ko-Rado Theorem). *Let $\mathcal{F} \subseteq \binom{[n]}{k}$ be an intersecting family, i.e. $F \cap F' \neq \emptyset$ for any $F, F' \in \mathcal{F}$. If $n \geq 2k$, then*

$$|\mathcal{F}| \leq \binom{n-1}{k-1}.$$

This bound is sharp by taking \mathcal{F} to consist of every set containing the element 1 (and in fact, up to isomorphism this is the unique extremal construction when $n > 2k$). Note that if $n < 2k$, then $\mathcal{F} = \binom{[n]}{k}$ is an intersecting family, so we need $n \geq 2k$ for us to be able to prove a non-trivial bound.

Proof. The proof uses what is known as Katona's circle method, which involves choosing a random cyclic ordering $\pi : [n] \rightarrow \mathbb{Z}_n$, where \mathbb{Z}_n is the integers mod n . Given such a π and a set $A \in \mathcal{F}$, we let 1_A be the indicator variable with $1_A = 1$ if $A = \{\pi(i), \pi(i) + 1, \dots, \pi(i) + k - 1\}$ for some $i \in [n]$. We claim that $1_A = 1$ for at most k sets A .

Indeed, if $1_A = 0$ for all A then there is nothing to prove, so assume $1_A = 1$ for some A , say with $A = \{\pi(i), \pi(i) + 1, \dots, \pi(i) + k - 1\}$. Let $S_j = \{\pi(i) + j, \pi(i) + j + 1, \dots, \pi(i) + j + k - 1\}$, and observe that if $B \in \mathcal{F}$ has $1_B = 1$, then we must have¹ $B = S_j$ for some $-k < j < k$. Moreover, for each pair $\{S_{-k+\ell}, S_\ell\}$ with $0 \leq \ell < k$, at most one $B \in \mathcal{F}$ is equal to one of these sets since $S_{-k+\ell}, S_\ell$ are disjoint, so in total we conclude that $1_A = 1$ for at most k different $A \in \mathcal{F}$.

Observe² that $\Pr[1_A = 1] = n \binom{n}{k}^{-1}$, and this together with the claim above implies

$$k \geq \mathbb{E}[\sum_{A \in \mathcal{F}} 1_A] = \sum_{A \in \mathcal{F}} \Pr[1_A = 1] = |\mathcal{F}| \cdot n \binom{n}{k}^{-1},$$

and rearranging gives the desired bound. □

There are many, many proofs of the Erdős-Ko-Rado theorem, as well as many generalizations and applications. Again, we refer the reader to [47] for more on this. Our second result related to extremal set theory is the following.

Theorem 2.4 (Bollobás Set Pairs Inequality). *Let $\mathcal{A} = \{A_1, \dots, A_m\}$ and $\mathcal{B} = \{B_1, \dots, B_m\}$ be set systems such that $A_i \cap B_i = \emptyset$ for all i and $A_i \cap B_j \neq \emptyset$ for all $i \neq j$. Then*

$$\sum_{i=1}^m \left(\frac{|A_i| + |B_i|}{|A_i|} \right)^{-1} \leq 1$$

¹Here we use that each $B \in \mathcal{F}$ intersects A and that $n \geq 2k$ implies S_k is disjoint from A

²This follows because for any cyclic ordering π there are exactly n sets S which have $1_S = 1$

Pairs of families as in Theorem 2.4 are called *cross-intersecting*.

Proof. Let π be a random permutation of the underlying ground set (the size of which is irrelevant for the conclusion/proof). Let 1_i be the indicator variable with $1_i = 1$ if $\pi(x) < \pi(y)$ for all $x \in A_i$ and $y \in B_i$. That is, 1_i is the indicator for the event that A_i appears completely before B_i under π . A simple counting argument shows that $\Pr[1_i = 1] = \binom{|A_i| + |B_i|}{|A_i|}^{-1}$ (where here we implicitly use that $A_i \cap B_i = \emptyset$, as otherwise $\Pr[1_i = 1] = 0$).

We claim that there is at most one i such that $1_i = 1$. Indeed, say $1_i = 1$. Then for any $j \neq i$, by hypothesis there is some $x \in A_j \cap B_i \subseteq A_j$ and $y \in A_i \cap B_j \subseteq B_j$, and since $1_i = 1$, we have $\pi(x) > \pi(y)$. Thus $1_j = 0$ for all $j \neq i$. With this claim we have

$$1 \geq \mathbb{E}\left[\sum_i 1_i\right] = \sum \Pr[1_i = 1] = \sum \binom{|A_i| + |B_i|}{|A_i|}^{-1}.$$

□

Theorem 2.4 has many applications. One such application involves *antichains*, which are collections of sets \mathcal{F} such that there exist no distinct $A, B \in \mathcal{F}$ with $A \subseteq B$.

Corollary 2.5 (LYM Inequality). *If $\mathcal{F} \subseteq 2^{[n]}$ is an antichain, then*

$$\sum_{A \in \mathcal{F}} \binom{n}{|A|}^{-1} \leq 1.$$

Proof. Let $\mathcal{F} = \{A_1, \dots, A_m\}$ and define $B_i = [n] \setminus A_i$. It is not difficult to check that since \mathcal{F} is an antichain, $A_i \cap B_j = \emptyset$ if and only if $i = j$. The bound then follows from Theorem 2.4. □

We note that the proof of Corollary 2.5 is a nice simplification of the proof of Theorem 2.4: now $1_i = 1$ if and only if $A_i = \{\pi(1), \dots, \pi(|A_i|)\}$.

Corollary 2.6 (Sperner's Theorem). *If $\mathcal{F} \subseteq 2^{[n]}$ is an antichain, then*

$$|\mathcal{F}| \leq \binom{n}{\lfloor n/2 \rfloor}.$$

This result is sharp, as can be seen by taking $\mathcal{F} = \binom{[n]}{\lfloor n/2 \rfloor}$ or $\binom{[n]}{\lceil n/2 \rceil}$.

Proof. We have $\binom{n}{k} \leq \binom{n}{\lfloor n/2 \rfloor}$ for all k , so by the LYM inequality

$$1 \geq \sum_{A \in \mathcal{F}} \binom{n}{|A|}^{-1} \geq |\mathcal{F}| \binom{n}{\lfloor n/2 \rfloor}^{-1},$$

and moving things around gives the desired result. □

2.3 The Crossing Lemma and Incidence Geometry

Our final result concerns drawings of graphs. Without being too precise with our definitions, we define the *crossing number* of a graph G to be the minimum number of crossings that an embedding $\phi(G)$ in the plane will have. For example, a graph is planar if and only if $cr(G) = 0$.

Lemma 2.7. *If G is an n -vertex graph with m edges, then $cr(G) \geq m - 3n$.*

Sketch of Proof. Let $\phi(G)$ be an embedding of G with $cr(G)$ crossings. By deleting an edge from each crossing, we obtain a planar graph G' with n vertices and at least $m - cr(G)$ edges. A simple consequence of Euler's formula shows that this means $m - cr(G) \leq 3n$, giving the result. \square

We will use the probabilistic method to “amplify” the elementary bound of Lemma 2.7 and give a bound that is effective for dense graphs.

Lemma 2.8 (Crossing Lemma). *If G is an n -vertex graph with $m \geq 4n$ edges, then*

$$cr(G) \geq \frac{m^3}{64n^2}.$$

Proof. Let $\phi(G)$ be an embedding of G which has $cr(G)$ crossings. Let $V_p \subseteq V(G)$ be obtained by keeping each vertex of $V(G)$ independently and with probability p , and let $G_p = G[V_p]$. Observe that there is a natural embedding of G_p , namely the restriction of ϕ to G_p .

Let X denote the number of crossings in $\phi(G_p)$, and note that $\mathbb{E}[X] = p^4 cr(G)$ since a crossing survives if and only if all four of its relevant vertices lie in V_p . Using Lemma 2.7, we see that

$$p^4 cr(G) = \mathbb{E}[X] \geq \mathbb{E}[e(G') - 3|V_p|] = p^2 m - 3pn \implies cr(G) \geq p^{-2}m - 3p^{-3}n.$$

This lower bound will roughly be optimized when $p^{-2}m = p^{-3}n$, i.e. when $p = n/m$. More precisely, taking $p = 4n/m$ gives the desired bound. However, implicitly this argument requires that $0 \leq p \leq 1$, i.e. that $m \geq 4n$, and this holds by hypothesis. \square

In addition to being interesting in its own right, the crossing lemma gives a short proof of a fundamental result in incidence geometry.

Theorem 2.9 (Szemerédi-Trotter Theorem). *Let \mathcal{P} be a set of n points and \mathcal{L} a set of m lines in the plane, and let $\mathcal{I} \subseteq \mathcal{P} \times \mathcal{L}$ denote their set of incidences, i.e. pairs (p, ℓ) with $p \in \ell$. Then*

$$|\mathcal{I}| = O(m^{2/3}n^{2/3} + m + n).$$

This bound is essentially best possible, though we omit the details of the (not too difficult) construction.

Proof (due to Székely). Without loss of generality, we can assume every point and line is in at least one incidence (otherwise we can delete these points/lines). Let G be the graph on \mathcal{P} which makes two points p_1, p_2 adjacent if there exists a line $\ell \ni p_1, p_2$ such that there is no third point

q on the line segment p_1p_2 . In other words, G is the graph obtained by drawing the points and lines on the plane, and then erasing the rays of lines which go off to infinity.

If $i(\ell)$ denotes the number of points incident to ℓ , then it is not difficult to see that $e(G) = \sum i(\ell) - 1 = |\mathcal{I}| - m$, where here we implicitly used that $i(\ell) \geq 1$ for all ℓ . If $|\mathcal{I}| \leq 2m$, then in particular $|\mathcal{I}| = O(m)$ and the result follows, so we can assume $e(G) \geq \frac{1}{2}|\mathcal{I}|$, and similarly we can assume $|\mathcal{I}| \geq 8m$ and hence $e(G) \geq 4n$. Thus by the crossing lemma we have

$$cr(G) \geq \frac{|\mathcal{I}|^3}{2^9 n^2}.$$

The critical observation is that $cr(G) \leq \binom{m}{2}$ since each crossing corresponds to two lines of \mathcal{L} intersecting. Plugging this into the expression above gives the desired result. \square

As a brief aside, we note that this idea of taking a weak result (Lemma 2.7) and amplifying it to a stronger result (Lemma 2.8) shows up in many other places in extremal combinatorics. For example, it is easy to prove a weak version of the Szemerédi-Trotter theorem with a bound of roughly $O(mn^{1/2} + n)$ by observing that there exist no points p_1, p_2 and ℓ_1, ℓ_2 such that all of the incidences (p_i, ℓ_j) are present, i.e. the “incidence graph” on $\mathcal{P} \cup \mathcal{L}$ contains no C_4 . One can then use the method of polynomial partitioning to dissect \mathbb{R}^2 into small regions where this bound is effective. For much more on incidence geometry and polynomial partitioning, we refer the reader to the excellent book by Sheffer [101].

3 The Lovász Local Lemma

We say that an event A_i is mutually independent of a set of events $\{A_j : j \in J\}$ if for any $J' \subseteq J$, we have $\Pr[A_i \cap \bigcap_{j \in J'} A_j] = \Pr[A_i] \cdot \Pr[\bigcap_{j \in J'} A_j]$. We say that A_1, \dots, A_n are mutually independent events if A_i is mutually independent of $\{A_j : j \in [n] \setminus \{i\}\}$ for all i . Note that in this case we have $\Pr[\bigcap A_i] = \prod \Pr[A_i]$. In this section we consider a result which roughly says that if the A_i 's are “almost independent”, then we have $\Pr[\bigcap A_i] \approx \prod \Pr[A_i]$.

Theorem 3.1. [Lovász Local Lemma] *Let A_1, \dots, A_n be events and let $D_1, D_2, \dots, D_n \subseteq [n]$ be such that A_i is mutually independent of $\{A_j : j \notin D_i \cup \{i\}\}$ for all i . If there exist real numbers $\gamma_i \in [0, 1)$ such that $\Pr[A_i] \leq \gamma_i \prod_{j \in D_i} (1 - \gamma_j)$ for all i , then*

$$\Pr[\bigcap \overline{A_i}] \geq \prod (1 - \gamma_i) > 0.$$

This result is often just referred to as “the local lemma”. Note that if the A_i were all mutually independent, then we could take $D_i = \emptyset$ and $\gamma_i = \Pr[A_i]$ for all i and conclude from the local lemma that $\Pr[\bigcap \overline{A_i}] \geq \prod \Pr[\overline{A_i}]$.

Proof. We claim that for all i and $S \subseteq [n]$, we have

$$\Pr[\overline{A_i} \mid \bigcap_{j \in S} \overline{A_j}] \geq 1 - \gamma_i.$$

This will give the result since then

$$\Pr[\bigcap_i \overline{A_i}] = \prod_i \Pr[\overline{A_i} \mid \bigcap_{j \in [i-1]} \overline{A_j}] \geq \prod_i (1 - \gamma_i).$$

We prove this claim by induction¹ on $|S|$. The base case $|S| = 0$ is equivalent to saying $\Pr[A_i] \leq \gamma_i$ for all i , and this follows from $\Pr[A_i] \leq \gamma_i \prod_{j \in D_i} (1 - \gamma_j) \leq \gamma_i$. Now consider any set S , and in particular assume we have proven the result for all $S' \subsetneq S$. If $i \in S$ then the result is trivial, so we can assume $i \notin S$. Observe that

$$\begin{aligned} \Pr[A_i \mid \bigcap_{j \in S} \overline{A_j}] &= \frac{\Pr[A_i \cap \bigcap_{j \in S} \overline{A_j}]}{\Pr[\bigcap_{j \in S} \overline{A_j}]} \leq \frac{\Pr[A_i \cap \bigcap_{j \in S \setminus D_i} \overline{A_j}]}{\Pr[\bigcap_{j \in S \setminus D_i} \overline{A_j}] \cdot \Pr[\bigcap_{k \in S \cap D_i} \overline{A_k} \mid \bigcap_{j \in S \setminus D_i} \overline{A_j}]} \\ &= \frac{\Pr[A_i]}{\Pr[\bigcap_{k \in S \cap D_i} \overline{A_k} \mid \bigcap_{j \in S \setminus D_i} \overline{A_j}]}, \end{aligned} \tag{6}$$

where the first inequality used that we are taking a product over fewer events, and the second equality used that A_i is mutually independent of events not in D_i . Let $S \cap D_i = \{k_1, \dots, k_p\}$. Then we can rewrite the probability in the denominator of (6) as

$$\prod_{q=1}^p \Pr[\overline{A_{k_q}} \mid \bigcap_{j \in (S \setminus D_i) \cup \{k_1, \dots, k_{q-1}\}} \overline{A_j}] \geq \prod_{q=1}^p (1 - \gamma_{k_q}) \geq \prod_{j \in D_i} (1 - \gamma_j),$$

¹It is perhaps more natural to try and prove the result by induction on n rather than on this somewhat weird looking claim. However, if one plays around with this problem, one quickly sees that one needs to prove something like the stated claim.

where the first inequality used the inductive hypothesis and the last step used $k_q \in S \cap D_i \subseteq D_i$ for all q . This together with (6) and the hypothesis $\Pr[A_i] \leq \gamma_i \prod_{j \in D_i} (1 - \gamma_j)$ implies that $\Pr[A_i | \bigcap_{j \in S} \overline{A_j}] \leq \gamma_i$, which is equivalent to saying $\Pr[\overline{A_i} | \bigcap_{j \in S} \overline{A_j}] \geq 1 - \gamma_i$. This proves the inductive hypothesis of our claim, and hence proves the result. \square

The following version of the local lemma is often sufficient for most applications (and again this is often referred to as “the local lemma”).

Corollary 3.2 (Symmetric Lovász Local Lemma). *Let A_1, \dots, A_n be events and let $D_1, D_2, \dots, D_n \subseteq [n]$ be such that A_i is mutually independent of $\{A_j : j \notin D_i \cup \{i\}\}$ for all i . If $\Delta \geq 1$ is such that $|D_i| \leq \Delta$ and $\Pr[A_i] \leq \frac{1}{e(\Delta+1)}$ for all i , then $\Pr[\bigcap \overline{A_i}] > 0$.*

Proof. Observe that for all i we have

$$\frac{1}{\Delta+1} \prod_{j \in D_i} \left(1 - \frac{1}{\Delta+1}\right) \geq \frac{1}{\Delta+1} \left(1 - \frac{1}{\Delta+1}\right)^\Delta \geq \frac{1}{e(\Delta+1)} \geq \Pr[A_i],$$

where the second to last inequality used that $(1 - 1/x)^{x-1} > 1/e$ for $x \geq 2$. Thus the (asymmetric) local lemma applies with $\gamma_i = \frac{1}{\Delta+1}$ for all i , proving the result. \square

We note that this result is essentially best possible. Indeed, consider rolling a fair $(\Delta + 1)$ -sided dice and let A_i be the event that the dice rolls i . In this case A_i is dependent on all of $D_i = [\Delta + 1] \setminus \{i\}$ and we have $\Pr[A_i] = \frac{1}{\Delta+1} > \frac{1}{e(\Delta+1)}$, so the local lemma does not apply (which is good since we have $\Pr[\bigcap \overline{A_i}] = 0$). In particular, this example shows that we can not improve the requirement $\Pr[A_i] \geq \frac{1}{e(\Delta+1)}$ in the symmetric local lemma to $\Pr[A_i] \geq \frac{1}{\Delta+1}$ in general. Thus the hypothesis in the symmetric local lemma is sharp up to a factor of e , and in fact Shearer proved that this factor of e is necessary [100].

3.1 Applications

Our first application of the local lemma will be an asymptotic improvement to our lower bound for Ramsey numbers from Theorem 1.1.

Theorem 3.3 (Spencer [103]). *For all n we have*

$$R(n, n) \geq (1 + o(1)) \frac{\sqrt{2}n}{e} 2^{n/2}.$$

Proof. Uniformly at random color the edges of K_N . For $S \in \binom{[N]}{n}$, let A_S be the event that G contains a monochromatic K_n on S , and as before we note that $\Pr[A_S] = 2^{1-\binom{n}{2}}$. Let D_S consist of all the sets $T \in \binom{[N]}{n} \setminus \{S\}$ such that $|S \cap T| \geq 2$. It is not difficult to see that A_S is mutually independent of $\{A_T : T \notin D_S \cup \{S\}\}$ since the color given to each pair of S is independent of these events. A weak bound gives $|D_S| \leq \binom{n}{2} \binom{N}{n-2} - 1 \leq n^2(eN/(n-2))^{n-2} - 1$, so by the (symmetric) local lemma we have that $\Pr[\bigcap \overline{A_S}] > 0$ provided $2^{1-\binom{n}{2}} < \frac{1}{en^2}(eN/n-2)^{2-n}$, i.e. if

$$(2en^2)^{1/n-2} \cdot 2^{\binom{n}{2}/n-2} \cdot \frac{n-2}{eN} = (2en^2)^{1/n-2} \cdot 2^{n/2+1/2-1/(n-2)} \cdot \frac{n-2}{eN} < 1,$$

and this happens if $N = (1 - \epsilon) \frac{\sqrt{2n}}{e} 2^{n/2}$ for any $\epsilon > 0$ provided n is sufficiently large, giving the desired result. \square

The local lemma works best if there are few dependencies between events. As such, it performs much better for off-diagonal Ramsey numbers.

Theorem 3.4. *For all n we have*

$$R(3, n) = \Omega(n^2 / \log^2 n).$$

Proof. Randomly color each edge of K_N red with probability p and blue otherwise. Given a set $S \in \binom{[N]}{3}$, we let R_S be the event that the vertices of S form a red triangle, and similarly for $T \in \binom{[N]}{n}$ we define B_T . Observe that $\Pr[A_S] = p^3$ and $\Pr[B_T] = (1 - p)^{\binom{n}{2}}$.

Given $S \in \binom{[N]}{3} \cup \binom{[N]}{n}$, we define D_S to be the sets of sizes 3 and n which intersect S in at least two vertices. Observe that if $|S| = 3$, then D_S contains at most $3N$ set of size 3 and at most $\binom{N}{n}$ sets of size n , and if $|S| = n$, we have that D_S contains at most $N\binom{n}{2}$ sets of size 3 and at most $\binom{N}{n}$ sets of size n . Our goal now is to choose some parameters γ_S, γ_T so that the (asymmetric) local lemma applies to the R_S, B_T events.

At this point there's a lot of undetermined variables floating around: N, p, γ_S . Let's think about reasonable guesses for how to optimize things. First of all, it seems clear that we probably want two parameters γ_3, γ_n such that we set $\gamma_S = \gamma_{|S|}$ when applying the local lemma. With this we in particular need

$$p^3 \leq \gamma_3(1 - \gamma_3)^{3N}(1 - \gamma_n)^{\binom{N}{n}}. \quad (7)$$

In particular we need $\gamma_3 \geq p^3$, so let's naively take $\gamma_3 = Cp^3$ for some large constant C . Given this, we also need $\gamma_n \leq c\binom{N}{n}^{-1}$ in order to have the $(1 - \gamma_n)^{\binom{N}{n}}$ term be no larger than a constant. If we take $\gamma_n = c\binom{N}{n}^{-1}$, we see that (7) is satisfied provided $p = o(N^{-1/3})$ and c, C are chosen appropriately.

The other condition we need to satisfy is

$$(1 - p)^{\binom{n}{2}} \leq \gamma_n(1 - \gamma_3)^{N\binom{n}{2}}(1 - \gamma_n)^{\binom{N}{n}},$$

and by plugging in our choices for γ_3, γ_n and the assumption that p must be fairly small, we essentially need to have

$$e^{-p\binom{n}{2}} \leq (n/N)^n \cdot e^{-p^3 N\binom{n}{2}},$$

and for this to hold we in particular need something like $p\binom{n}{2} \geq p^3 N\binom{n}{2}$, i.e. $p = O(N^{-1/2})$. Taking $p = c'N^{-1/2}$, we see that we also need roughly

$$p\binom{n}{2} \approx c'N^{-1/2}n^2 \geq n \log(N/n).$$

Assuming $N \geq n^{1+\epsilon}$ for some small $\epsilon > 0$, this reduces to $N^{1/2} \leq n / \log n$, i.e. $N = n^2 / (\log n)^2$.

Thus in total, a heuristic argument suggests that we can apply the local lemma with $N = \Theta(n^2 / (\log n)^2)$ by taking $p = \Theta(N^{-1/2})$, $\gamma_3 = \Theta(N^{-3/2})$, and $\gamma_n = \Theta(\binom{N}{n})$. And indeed, a careful analysis shows that this will work out for n sufficiently large. \square

We note that the bound of $n^2/\log^2 n$ is the best one can do using this approach. However, it turns out that $R(3, n) = \Theta(n^2/\log n)$. This improved lower bound was originally proved by Kim [73]. The idea of their proof was to start with a K_N which is entirely colored blue, and then to iteratively randomly pick an edge of K_N and color it red if it does not create a red triangle. A careful analysis shows that with positive probability the final graph at the end contains no large blue clique, and it contains no red clique by construction. We will see a shorter proof of this lower bound in [a later section \(expanders\)](#).

3.2 Related Lemmas

While the local lemma is very powerful, there are certain circumstances where it doesn't give you quite what you want. Fortunately there are many other lemmas which allow one to prove bounds on $\Pr[\bigcap \overline{A_i}]$ even when the A_i depend on each other in some way. For example, it is not too difficult to generalize the local lemma as follows (and as an exercise the reader should convince themselves that they can prove this result).

Theorem 3.5. *Let A_1, \dots, A_n be events. Assume there exists partitions $D_i \cup E_i = [n] \setminus \{i\}$ for all i and real numbers $0 \leq \delta, \gamma \leq 1$ such that $\gamma(1 - \gamma)^{|D_i|} \geq \delta$ and for all $E \subseteq E_i$ we have $\Pr[A_i | \bigcap_{e \in E} \overline{A_e}] \leq \delta$ and . Then*

$$\Pr[\bigcap \overline{A_i}] \geq (1 - \gamma)^n > 0.$$

Note that when $\delta = \gamma$ we more or less recover Theorem 3.1 when $\gamma_i = \gamma$ for all i . The power here is that we allow each A_i to possibly be dependent of every event, but it is not “very dependent” on the events of E_i .

Another result in a similar spirit as the local lemma is Janson's inequality. Given a set $S \subseteq X$ and \vec{p} , let \mathcal{A}_S be the set containing

Theorem 3.6 (Janson's inequality). *Let H be a hypergraph on a set V , and let V_p be the set obtained by including each vertex of V independently and with probability p . Let A_i denote the event that V_p contains the i th edge of H and define*

$$\mu = \sum \Pr[A_i], \quad \Delta = \sum_{(S_i, S_j): S_i \cap S_j \neq \emptyset} \Pr[A_i \cap A_j].$$

Then

$$\prod_i \Pr[\overline{A_i}] \leq \Pr[\bigcap_i \overline{A_i}] \leq e^{-\mu + \frac{\Delta}{2}}.$$

Note that if all of the edges of H are disjoint, then these bounds are roughly $e^{-\mu} \leq \Pr[\bigcap_i \overline{A_i}] \leq e^{-\mu/2}$. Again there are many variants of Theorem 3.6 which are useful in different situations.

4 The Second Moment Method

In the previous section we saw several results that can be used to bound the probability of some event happening. In this section we look at what is perhaps the most broadly applicable result of this form: Chebyshev's inequality.

4.1 Concentration Inequalities

Roughly speaking, concentration inequalities are results which say that under reasonable circumstances, a random variable is likely to be close to its expectation.

Perhaps the most famous (one-sided) concentration inequality is Markov's inequality. We already saw this around the proof of Theorem 2.2, but for good measure we'll formally state it here.

Lemma 4.1 (Markov's inequality). *If X is a non-negative real-valued random variable, then for all $\lambda > 0$ we have*

$$\Pr[X \geq \lambda] \leq \frac{\mathbb{E}[X]}{\lambda}.$$

In particular, if X is integer-valued, then

$$\Pr[X \neq 0] \leq \mathbb{E}[X].$$

Proof. For simplicity we only prove the result when X is integer valued. In this case we have

$$\Pr[X \geq \lambda] = \sum_{k \geq \lambda} \Pr[X = k] \leq \sum_{k \geq \lambda} \frac{k}{\lambda} \cdot \Pr[X = k] = \frac{\mathbb{E}[X]}{\lambda}.$$

The second statement follows by taking $\lambda = 1$. □

The “in particular” part of this lemma is probably the most common usage of Markov's inequality. To reiterate, this says that $\mathbb{E}[X] \rightarrow 0$ implies $X = 0$ with high probability, and this application of Markov's inequality is often known as *the first moment method*.

Unfortunately it is not true in general that $\mathbb{E}[X] \rightarrow \infty$ implies $X > 0$ with high probability (e.g. take $X = n$ with probability $n^{-1/2}$ and $X = 0$ otherwise). However, for many reasonable examples this implication does hold. Often one can show this by utilizing Chebyshev's inequality. We recall that the variance of a random variable is $\text{Var}(X) = \mathbb{E}[(X - \mathbb{E}[X])^2]$.

Lemma 4.2 (Chebyshev's inequality). *Let X be a real-valued random variable with $\text{Var}(X) = \sigma^2$. Then for all $\lambda > 0$, we have*

$$\Pr[|X - \mathbb{E}[X]| \geq \lambda\sigma] \leq \frac{1}{\lambda^2}.$$

Proof. We have

$$\Pr[|X - \mathbb{E}[X]| \geq \lambda\sigma] = \Pr[(X - \mathbb{E}[X])^2 \geq \lambda^2\sigma^2] \leq \frac{1}{\lambda^2},$$

where this last step used Markov's inequality applied to the (non-negative) random variable $Y := (X - \mathbb{E}[X])^2$ after noting that $\mathbb{E}[Y] = \sigma^2$ by definition. \square

Morally speaking, Chebyshev's inequality says that if $\sigma = o(\mathbb{E}[|X|])$, then X is close to its expectation with high probability. The usage of Chebyshev's inequality is often referred to as *the second moment method*.

4.2 The Rödl Nibble

Our approach in this subsection borrows heavily from Alon and Spencer [6].

We say that a set of edges $E \subseteq E(K_n^r)$ is a *k-covering* if every k -set $S \subseteq [n]$ is contained in some edge of E , and we will simply call this a *covering* if $k = 1$. It isn't hard to see that every k -covering needs at least $\binom{n}{k} / \binom{r}{k}$ edges, with equality holding if and only if every k -set is contained in exactly one edge of E . S Erdős and Hannini asked whether one could find k -coverings with asymptotically this many edges, and this was answered positively by Rödl.

Theorem 4.3 (Rödl [96]). *For all fixed $k \leq r$, there exists a k -covering $E \subseteq E(K_n^r)$ with*

$$|E| = (1 + o(1)) \frac{\binom{n}{k}}{\binom{r}{k}},$$

where the $o(1)$ term tends to 0 as n tends towards infinity.

The first step of this argument is to reduce the problem of finding k -coverings to simply finding coverings. To this end we define $H_n^{r,k}$ to be the $\binom{r}{k}$ -uniform hypergraph whose vertex set is $\binom{[n]}{k}$ and whose edge set is $\{\binom{S}{k} : S \in \binom{[n]}{r}\}$. That is, the hyperedges of $H_n^{r,k}$ are all the sets of size k covered by an edge $S \in E(K_n^r)$. It is not too difficult to check that Theorem 4.3 is equivalent to saying that there exists a covering of $H_n^{r,k}$ of the stated size.

Our approach for proving that $H_n^{r,k}$ has a small covering will more generally show that any “nice” r -uniform hypergraph H has a small covering, with the approach roughly being the following:

- Randomly choose $\epsilon n / r$ edges E_1 from H for some small $\epsilon > 0$. With high probability E_1 will cover about $e^{-\epsilon} n$ vertices.
- Delete vertices covered by E_1 to get a new hypergraph H_2 . With high probability H_2 is also “nice”, so we can iterate the procedure above and pick some random set of edges E_2 .
- We keep doing this until ϵn vertices remain uncovered, and at this point we can trivially cover them using at most ϵn additional edges.

Broadly speaking this approach is known as the Rödl nibble or the semirandom method. To reiterate, the core idea is that you iteratively do something to a small chunk of vertices in such a way that the structure of the rest of your hypergraph remains roughly the same with high

probability, which allows one to keep iterating this procedure until one is left with a very small problem to solve.

It turns out that for our purposes, “nice” hypergraphs are those which are roughly D -regular for some D and such that every pair of vertices is in $o(D)$ edges, i.e. the hypergraph is almost linear. These conditions are reasonable for trying to prove Theorem 4.3 since $H_n^{r,k}$ is $\binom{n-k}{r-k}$ -regular and every pair of vertices is in at most $\binom{n-k-1}{r-k-1}$ edges. Our main technical lemma in this direction is the following, where throughout this section we write $c = 1 \pm \delta$ if $c \in [1 - \delta, 1 + \delta]$.

Lemma 4.4. *For every $r \geq 2$ and reals $K \geq 1$ and $\epsilon, \delta' > 0$, there are $\delta = \delta(r, K, \epsilon, \delta') > 0$ and $D_0 = D_0(r, K, \epsilon, \delta')$ such that for every $n \geq D \geq D_0$ the following holds.*

Let $H = (V, E)$ be an n -vertex r -graph such that

- (i) For all but at most δn vertices $x \in V$, we have $d(x) = (1 \pm \delta)D$,*
- (ii) For all $x \in V$ we have $d(x) < KD$, and*
- (iii) For any two distinct $x, y \in V$, we have $d(x, y) < \delta D$.*

In this case there exist a set of edges $E' \subseteq E$ such that

- (a) $|E'| = (1 \pm \delta')(\epsilon n/r)$,*
- (b) The set $V' := V - \bigcup_{e \in E'} e$ has $|V'| = (1 \pm \delta')e^{-\epsilon}n$, and*
- (c) For all but at most $\delta'|V'|$ vertices $x \in V'$, the degree $d'(x)$ of x in the induced hypergraph $H[V']$ satisfies $d'(x) = (1 \pm \delta')De^{-\epsilon(r-1)}$.*

Again, (i) and (ii) say that H is close to D -regular and (iii) says it has small codegrees. The main point of the conclusion is that the number of uncovered vertices and their degrees shrink in a predictable way.

Proof. We only give a sketch of the proof from [6] (their full version is complicated enough to involve 20 named constants!). Throughout the proof we’ll introduce various constants δ_i which we always assume to be sufficiently small in terms of our relevant parameters.

Randomly choose a subset $E' \subseteq E$ such that each edge of E appears in E' independently and with probability $p = \epsilon/D$. Roughly speaking, our goal will be to show that with this choice of E' , each of (a),(b),(c) occur in expectation, and then we will use Chebyshev to show that each of these occur with high probability.

To start, because H is essentially D -regular, we have $|E| = (1 \pm \delta_1)Dn/r$, so

$$\mathbb{E}[|E'|] = p|E| = (1 \pm \delta_1)\epsilon n/r.$$

We also have

$$\text{Var}(|E'|) = p(1-p)|E| \leq 2\epsilon n/r.$$

Because $\text{Var}(|E'|) = o(\mathbb{E}[|E'|])$, Chebyshev should be able to show that E' is close to $\mathbb{E}[E']$ with high probability. More precisely, Chebyshev's inequality implies

$$\Pr[||E'| - \mathbb{E}[|E'|]| \geq \delta_1 \sqrt{2\epsilon n/r} \cdot \sqrt{2\epsilon n/r}] \leq \frac{r}{2\delta_1^2 \epsilon n} \leq .01,$$

with this last step holding for n sufficiently large. Thus with probability at least .99,

$$|E'| = \mathbb{E}[|E'|] \pm 2\delta_1 \epsilon n/r = (1 \pm 3\delta_1) \epsilon n/r.$$

This shows that (a) occurs with high probability. To deal with (b), let us first get a grasp on $\mathbb{E}[|V'|]$. For $x \in V$, let $1_x = 1$ if $x \notin \bigcup_{e \in E'} e$ and $1_x = 0$ otherwise. With this we see $|V'| = \sum 1_x$, so by linearity of expectation it suffices to bound each of $\mathbb{E}[1_x]$.

We will say that a vertex x is *good* if $d(x) = (1 \pm \delta)D$ and that it is *bad* otherwise. If x is bad we will simply use the trivial estimates $0 \leq \mathbb{E}[1_x] \leq 1$. If x is good we have

$$\mathbb{E}[1_x] = (1 - p)^{d(x)} = (1 - \epsilon/D)^{(1 \pm \delta)D} = (1 \pm \delta_3) e^{-\epsilon}, \quad (8)$$

where this last step used that $1 - p$ is within a constant factor of e^{-p} for p sufficiently small and that δ is chosen to be sufficiently small in terms of ϵ (e.g. we can make sure that it's smaller than ϵ^{-1}).

Having at most δn bad vertices together with (8) implies $\mathbb{E}[|V'|] = (1 \pm \delta_4) n e^{-\epsilon}$. To compute the variance, we observe that

$$\text{Var}[|V'|] = \sum_x \text{Var}[1_x] + \sum_x \sum_{y \neq x} \mathbb{E}[1_x 1_y] - \mathbb{E}[1_x] \mathbb{E}[1_y]. \quad (9)$$

Because each 1_x is an indicator random variable, we have

$$\sum_x \text{Var}[1_x] \leq \sum_x \mathbb{E}[1_x] = \mathbb{E}[|V'|].$$

For the mixed terms of (9), we have for any x, y that

$$\begin{aligned} \mathbb{E}[1_x 1_y] - \mathbb{E}[1_x] \mathbb{E}[1_y] &= (1 - p)^{d(x) + d(y) - d(x, y)} - (1 - p)^{d(x) + d(y)} \\ &\leq (1 - p)^{-d(x, y)} - 1 \leq (1 - \epsilon/D)^{-\delta D} - 1 \leq e^{\epsilon \delta} - 1 \leq \delta_5. \end{aligned}$$

In total we find

$$\text{Var}[|V'|] \leq \mathbb{E}[|V'|] + \delta_5 n^2 \leq \delta_6 (\mathbb{E}[|V'|])^2,$$

where this last step used that $\mathbb{E}[|V'|] = \Theta_\epsilon(n)$. By Chebyshev we can guarantee with probability at least .99 that

$$|V'| = (1 \pm \delta_7) \mathbb{E}[|V'|] = (1 \pm \delta_8) n e^{-\epsilon}.$$

Proving that condition (c) holds with high probability is a little complicated, so we'll omit the full details¹ (which can be found in [6]). Let us instead give a heuristic argument as to why

¹The argument is similar in spirit to that of (b): you define $1_e = 1$ if e survives in $H[V']$ and $1_e = 0$ otherwise. Then $d'(x)$ is just the sum of some of these indicator random variables, so one has to bound terms of the form $\mathbb{E}[1_e]$ and $\mathbb{E}[1_e 1_f]$. If e, f are "typical" edges then the computation of $\mathbb{E}[1_e]$ and $\mathbb{E}[1_e 1_f]$ are straightforward to estimate, and there are few terms involving e which are not typical.

(c) holds in expectation. We first condition on the event $x \in V'$, which means that no edge containing x is in E' . An edge $e \ni x$ survives in $H[V']$ only if every edge f with $e \cap f \neq \emptyset$ has $f \notin E'$. Because H is roughly linear and D -regular, there are about rD such edges f , but D of these (namely those containing x) are automatically not in E' since we conditioned on $x \in V'$. The remaining $(r-1)D$ edges are each included independently and with probability ϵ/D , so the probability that none are included is $(1 - \epsilon/D)^{(r-1)D} \approx e^{-(r-1)\epsilon}$, and summing this over all of the roughly D edges containing x gives the result.

Once we have shown that each of (a),(b),(c) holds with probability at least .99, then the probability that all of them hold is at least .97, so in particular some choice of E' exists which satisfies these conditions. \square

By repeatedly applying this lemma with carefully chosen values of ϵ, δ , one can prove the following result (and again, we omit the details of this, see [6]).

Theorem 4.5 (Pippenger). *For every $r \geq 2$ and reals $K \geq 1$ and $a > 0$, there are $\delta = \delta(r, K, a) > 0$ and $D_0 = D_0(r, K, a)$ such that for every $n \geq D \geq D_0$ the following holds.*

Let $H = (V, E)$ be an n -vertex r -graph such that

- (i) For all but at most δn vertices $x \in V$, we have $d(x) = (1 \pm \delta)D$,*
- (ii) For all $x \in V$ we have $d(x) < KD$, and*
- (iii) For any two distinct $x, y \in V$, we have $d(x, y) < \delta D$.*

Then there exists a cover of H using at most $(1 + a)(n/r)$ edges.

In particular, $H_n^{r,k}$ satisfies the conditions of the theorem, proving Theorem 4.3.

4.3 Steiner Systems

We say that a hypergraph is a Steiner system $S(n, r, k)$ if it has n -vertices, is r -uniform, and every k -set of its vertex set is contained in exactly one edge. Note that Steiner systems are k -coverings with exactly $\binom{n}{k} / \binom{r}{k}$ edges, so Theorem 4.3 shows that “approximate” Steiner systems exist, but when do actual Steiner systems exist?

The simplest non-trivial case is $S(n, 3, 2)$, which are also known as Steiner triple systems. It is not difficult to see that if a Steiner triple system on n vertices exists, then $3 \mid \binom{n}{2}$ (each edge covers 3 pairs and each of the $\binom{n}{2}$ pairs are covered exactly once) and $2 \mid (n-1)$ (for any given vertex v , each edge contains 2 pairs containing v and there are exactly $n-1$ such pairs). Equivalently, this argument says that if a Steiner triple system exists, then it is necessary that $n \equiv 1, 3 \pmod{6}$. It turns out that this condition is also sufficient due to certain constructions involving quasigroups and latin squares.

In general for an $S(n, r, k)$ to exist, there are certain “obvious” divisibility conditions that must be satisfied, but in general these are not sufficient. In fact, as of 2014, it wasn’t even known if,

say, any $S(n, k, 6)$ Steiner systems existed, let alone if there were infinitely many n for which such a Steiner system existed. In a major breakthrough, it was shown by Keevash [71] and independently by Glock, Kühn, Lo, and Osthus [58] that if n is sufficiently large in terms of r, k , then $S(n, r, k)$ systems exist if and only if n satisfies the obvious divisibility conditions. The core of Keevash's proof was a variant of the Rödl nibble in an algebraic setting, but the proof is very, very complicated!

Part II

Further Probabilistic Methods

5 Dependent Random Choice

The following is all based off of the excellent survey by Fox and Sudakov [44]. Throughout this section we denote the common neighborhood of a set of vertices S by $N(S)$, i.e. $N(S) = \{u : u \in N(v) \forall v \in S\}$. Before we explain what dependent random choice is, let's first see an example of it in action.

Lemma 5.1. *Let G be an n -vertex graph with average degree at least d . For any choice of integers m, r, t , there exists a set $U \subseteq V(G)$ such that every r -subset of U has at least m common neighbors, and such that*

$$|U| \geq \frac{d^t}{n^{t-1}} - \binom{n}{r} \left(\frac{m}{n}\right)^t.$$

Proof. The statement of the result suggests how we should prove it: we'll randomly pick a set W which will have expected size at least d^t/n^{t-1} , and then we'll use the method of alterations to delete from W a set of bad vertices, which in expectation will have size at most $\binom{n}{r}(m/n)^t$. The key twist is that we don't start by, say, defining W to include each vertex independently and with probability $p = d^t/n^t$, but instead W will end up depending on a different random set T .

To this end, let T be the random set obtained by uniformly at random selecting t vertices with repetition (i.e. each vertex is equally likely to be the i th vertex added to T , and in total T has size at most t), and define $W = N(T)$. The probability that a given vertex v is included in W is exactly $(d(v)/n)^t$, so by linearity of expectation and convexity we find that

$$\mathbb{E}[|W|] = \sum (d(v)/n)^t \geq d^t/n^{t-1}.$$

We say that a set of vertices $S \subseteq V(G)$ of size r is bad if $|N(S)| \leq m$. The probability that W contains a given bad set S is at most $(m/n)^t$ (since $S \subseteq W$ iff $T \subseteq N(S)$). Thus the expected number of bad sets of W is at most $\binom{n}{r}(m/n)^t$. If we let U be the set obtained by deleting a vertex from each bad set of W , then it has the desired properties by construction and in expectation it has the desired size, so such a choice of U exists. \square

Again, the key idea of this proof is that instead of defining W by including each vertex independently and with probability $p = d^t/n^t$, we instead formed it so that, on average, each vertex has probability at least p of being added, but the vertices are added in a very dependent way. In particular, the dependent way that W was generated made it more likely to have our desired property (i.e., we generated W by taking a common neighborhood, which made it less likely for W to contain sets of vertices with small common neighborhoods).

We can use Lemma 5.1 to prove some bounds on Turán numbers by using the following embedding lemma.

Lemma 5.2. *Let F be a bipartite graph on $A \cup B$ with $|A| = a$, $|B| = b$ such that the vertices in B all have degree at most r . If G is a graph which contains a set U such that $|U| = a$ and such that any subset of U of size r contains at least $a + b$ common neighbors, then G contains F as a subgraph.*

Proof. We define an injective homomorphism ϕ from $V(F)$ to $V(G)$ as follows. Choose $\phi|_A$ to be an arbitrary bijection onto U . For each $v \in B$ that has yet to be assigned, choose $\phi(v)$ to be any common neighbor of $\phi(N_F(v))$ which has yet to be assigned by ϕ . Note that there exist at least $a + b$ common neighbors of $\phi(N_F(v))$, so there certainly exists one which has yet to be assigned. This mapping gives the result. \square

With this we can quickly prove the following.

Theorem 5.3 (Füredi [52]; Alon, Krivelevich, Sudakov [5]). *If F is a bipartite graph on $A \cup B$ such that the vertices of B all have degree at most r , then*

$$\text{ex}(n, F) < 3(a + b)n^{2-1/r}.$$

Observe that this result generalizes Kővari-Sós-Turán, at least in terms of order of magnitude.

Proof. Assume G is an n -vertex F -free graph with average degree $d = 6(a + b)n^{1-1/r}$. By Lemma 5.2, we would be done if we could find a set U of size at least a such that every subset of size r had at least $m = a + b$ common neighbors. By Lemma 5.1, for any t we can find a set U with these properties of size at least

$$\frac{d^t}{n^{t-1}} - \binom{n}{r} \left(\frac{m}{n}\right)^t \geq (6a + 6b)^t n^{1-t/r} - (e/r)^r (a + b)^t n^{r-t}.$$

We see that taking $t = r$ makes the powers of n on both sides equal, and in total this gives a set of size at least

$$(6a + 6b)^r - (e(a + b)/r)^r.$$

Note that $6(a + b) \geq \frac{1}{2}(e(a + b)/r)$, so this is at least $\frac{1}{2}(6a + 6b)^r \geq a$. We have thus found our desired set U , which together with Lemma 5.2 gives a copy of F in G , a contradiction. \square

Another application of this method is to subdivisions. We define the 1-subdivision H^* of a graph H to be the graph obtained by replacing each edge of H by a P_2 (i.e. by inserting a new vertex in the middle of each edge). Note that subdivisions are bipartite graphs with all of its $e(H)$ new vertices having degree 2. Thus the previous theorem gives $\text{ex}(n, K_a^*) = O(a^2 n^{3/2})$. It turns out that one can significantly improve upon this dependency of a .

Theorem 5.4 (Alon, Krivelevich, Sudakov [5]). *For all a we have*

$$\text{ex}(n, K_a^*) = O(an^{3/2}).$$

Note that this only gives a reasonable bound when $a = O(n^{1/2})$, which makes sense since K_a^* has about a^2 vertices and thus can always be avoided by an n -vertex graph if $a \gg n^{1/2}$.

Unfortunately Lemma 5.1 on its own is not enough to prove Theorem 5.3, essentially because the size of U that we're guaranteed is too small. We can increase the size of U by demanding slightly weaker conditions for it to have, i.e. we only need that most pairs have many common neighbors¹. More precisely, we use the following.

¹This is a common situation that happens in applications of dependent random choice, though the exact way you weaken the conditions of Lemma 5.1 depends on the particular problem at hand.

Lemma 5.5. *Let G be an n -vertex graph with $an^{3/2}$ edges. Then G contains a subset of vertices U with $|U| = a$ such that for all $1 \leq i \leq \binom{a}{2}$, there are less than i pairs of vertices in U with fewer than i common neighbors in $V(G) \setminus U$.*

For example, this says that every pair of vertices of U has at least one common neighbor outside of U , and that there is at least one pair which has at least a common neighbors outside of U .

Add more intuition for the proof.

Proof. For simplicity we assume n is even, and by losing at most half of our edges we can assume that G is bipartite on $V_1 \cup V_2$ with $|V_1| = |V_2| = n/2$. Without loss of generality we can assume $\sum_{v \in V_1} d(v)^2 \leq \sum_{v \in V_2} d(v)^2$.

Let T be a random set obtained by including two vertices uniformly at random from V_1 with replacement. Let $W = N(T)$ and $X = |W|$. Similar to our computation before, we find

$$\mathbb{E}[X] = \sum_{v \in V_2} (d(v)/(n/2))^2 \geq 4n^{-2} \cdot (n/2)(an^{1/2})^2 = 2a^2.$$

Given distinct vertices $x, y \in V_2$, we define $f(x, y) = \frac{1}{|N_{V_1}(x, y)|}$ and we let $Y = \sum_{x, y \in W} f(x, y)$. Observe that

$$\begin{aligned} \mathbb{E}[Y] &= \sum_{x, y \in V_2} f(x, y) \cdot \Pr[x, y \in W] = \sum_{x, y \in V_2} \frac{1}{|N_{V_1}(x, y)|} \cdot \left(\frac{|N_{V_1}(x, y)|}{n/2} \right)^2 = 4n^{-2} \sum_{x, y \in V_2} |N_{V_1}(x, y)| \\ &= 4n^{-2} \sum_{z \in V_1} \binom{d(z)}{2} \leq 2n^{-2} \sum_{z \in V_1} d(z)^2 \leq 2n^{-2} \sum_{z \in V_2} d(z)^2 = \frac{1}{2} \mathbb{E}[X]. \end{aligned}$$

With this we see $\mathbb{E}[X - \mathbb{E}[X]/2 - Y] \geq 0$, and thus there exists a choice of T such that $X \geq Y$ and $X \geq \mathbb{E}[X]/2 \geq a^2$.

The trick now is to take $U \subseteq W$ a set of size exactly a uniformly at random, and let $Y' = \sum_{x, y \in U} f(x, y)$. In this case

$$\mathbb{E}[Y'] = \sum_{x, y \in W} f(x, y) \cdot \Pr[x, y \in U | x, y \in W] \leq Y \cdot \frac{a(a-1)}{X(X-1)} \leq X \cdot (a/X)^2 \leq 1.$$

Thus there exists a choice of U such that $Y' \leq 1$. We claim that such a U has the desired properties. Indeed, if there existed i pairs with fewer than i common neighbors, then this would immediately imply $Y' \geq i \cdot \frac{1}{i-1} > 1$, a contradiction. \square

Theorem 5.4 follows almost immediately from Lemma 5.5, and we omit its proof.

For our last result, we say that a graph F is r -degenerate if every subgraph of F contains a vertex of degree at most r . In this setting we can prove an embedding lemma analogous to Lemma 5.2.

Lemma 5.6. *Let G be a graph with vertex sets U_1, U_2 such that, for $k = 1, 2$, every subset of at most r vertices in U_k contains at least m common neighbors in U_{3-k} . Then G contains every r -degenerate bipartite graph H on m vertices.*

Proof. Let F_1 be an m -vertex r -degenerate bipartite graph on $V_1 \cup V_2$. By definition this means that there exists a vertex $v_1 \in F_1$ such that $d_{F_1}(v_1) \leq r$, and that there is some $v_2 \in F_2 := F_1 - v_1$ with $d_{F_2}(v_2) \leq r$ and so on. We now define a map $\phi : V_1 \cup V_2 \rightarrow U_1 \cup U_2$ with $\phi(V_i) \subseteq U_i$ as follows. Iteratively assume we have defined $\phi(v_m), \phi(v_{m-1}), \dots, \phi(v_{q+1})$ and that $v_q \in V_i$. Since $S := N(v_q) \cap \{v_m, \dots, v_{q+1}\}$ has at most r vertices by assumption, the set $\phi(S) \subseteq U_{3-i}$ has at least m common neighbors, so choose $\phi(v_q)$ to be any of these vertices that has yet to be assigned. It is not difficult to see that this gives the desired embedding. \square

Motivated by this lemma, we prove the following variant of Lemma 5.1.

Lemma 5.7. *Let $r, m \geq 2$ and let G be an n -vertex graph with at least $mn^{1-1/6r}$ edges. Then G contains two subsets U_1, U_2 such that, for $k = 1, 2$, every subset of r vertices in U_k has at least m common neighbors in U_{3-k} .*

Proof. The rough strategy of the proof is as follows. We will first apply Lemma 5.1 directly to obtain a large set U_1 such that every q -subset of U_1 (with $q > r$) has at least m common neighbors. We then mimic the proof of Lemma 5.1 by choosing a random set $T \subseteq U_1$ of size t and letting $U_2 = N(U_1)$. By choosing an appropriate value of t , the set U_2 will satisfy the condition. Moreover, if $q - t \geq r$, then for any r -subset $S \subseteq U_1$, the set $S \cup T$ has at least m common neighbors, all of which in particular lie in $N(T) = U_2$, so U_1 will also have the desired property.

We now begin the formal argument. Apply Lemma 5.1 using $q = 3r$ instead of r, t to get a set U_1 such that every subset of size $3r$ has at least m common neighbors and such that

$$|U_1| \geq \frac{d^{3r}}{n^{3r-1}} - \binom{n}{3r} (m/n)^{3r} \geq m^{3r} n^{1/2} - m^r / (3r)! \geq mn^{1/2}.$$

Now let T be a set obtained by including $t = 2r$ vertices uniformly at random from U_1 with replacement, and let $U_2 = N(T)$. The probability that U_2 contains a set of r vertices which have fewer than m common neighbors in U_1 is at most

$$\binom{n}{r} (m/|U_1|)^{2r} \leq \frac{1}{r!} < 1,$$

and in particular there exists a choice of T such that no r -subset of U_2 has fewer than m common neighbors. Note that for any r -subset $S \subseteq U_1$, the set $S \cup T$ has size at most $3r$ vertices, so by construction S has at least m common neighbors which lie in $N(T) = U_2$. Thus U_1, U_2 gives the desired result. \square

Combining these two lemmas immediately gives the following.

Theorem 5.8. *If F is an m -vertex r -degenerate graph, then*

$$\text{ex}(n, F) < mn^{2-1/6r}.$$

We note that one can optimize the proof of Lemma 5.7 to improve the exponent of this theorem slightly (by using $(3 - 2\sqrt{2})r$ instead of $3r$ throughout). However, the end result is still weaker

than the best known bound of $\text{ex}(n, F) \leq m^{1/2r} n^{2-1/4r}$ due to Alon, Krivelevich, and Sudakov [5], with their proof more or less being a slight refinement of the argument we gave.

As all of these examples illustrate: if you have a problem that could be magically solved if you had a large set of vertices U such that every r -set of U had many common neighbors, then a variant of dependent random choice might be worth trying out!

6 Strong Concentration Inequalities

As we have already seen in Section 4, there are many instances where one would like to show a random variable is concentrated around its expectation. Chebyshev's inequality is one tool which achieves this, and for arbitrary random variables this is essentially the best one can do. However, there are many specific kinds of random variables where one can get significantly stronger bounds. Here we present three results of this form in roughly increasing order of power.

We will omit the majority of the proofs of these results, focusing mostly on their applications. We refer the reader to the book of Dubhashi and Panconesi [37] for complete proofs. We note that the appendix of [37] consists of a very nice summary of these inequalities and many of their generalizations.

6.1 The Chernoff Bound

The Chernoff bound says that binomial random variables have exponential concentration around their means. **This version is incorrect, need to adjust.**

Theorem 6.1. *Let X_1, \dots, X_n be independent Bernoulli random variables each with probability of success p , and let $X = \sum X_i$. Then for all $\lambda > 0$,*

$$\Pr[|X - pn| \geq \lambda pn] < 2e^{-\lambda^2 pn/2}.$$

Sketch of Proof. Observe that for all $\lambda, t > 0$, we have

$$\Pr[X \geq (1 + \lambda)pn] = \Pr[e^{tX} \geq e^{t(1+\lambda)pn}] \leq \mathbb{E}[e^{tX}]e^{-t(1+\lambda)pn},$$

with this last step using Markov's inequality. We note that $e^{tX} = \sum \frac{t^m \mathbb{E}[X^m]}{m!}$ is the *moment generating function* of X , and it is a common trick in probability to rephrase inequalities in terms of e^{tX} . And indeed, because the X_i are all independent, we have

$$\mathbb{E}[e^{tX}] = \prod \mathbb{E}[e^{tX_i}] = (e^t p + (1 - p))^n.$$

Thus we are left with the problem of choosing t so that $\frac{e^t p + (1 - p)}{e^{-t(1+\lambda)p}}$ is minimized. One can do this using calculus, and this will give $\Pr[X \geq (1 + \lambda)pn] < e^{-\lambda^2 pn/2}$. The same argument gives $\Pr[X \leq (1 - \lambda)pn] < e^{-\lambda^2 pn/2}$, and combining these inequalities gives the desired result. \square

The Chernoff bound can be generalized, for example, by replacing the bernoulli random variables with any bounded random variable, see [37].

Given a hypergraph H and partition $V(H) = R \sqcup B$, define the discrepancy of the partition by

$$\text{disc}(H, R, B) = \max_{e \in E(H)} ||e \cap R| - |e \cap B||,$$

and define the discrepancy of the hypergraph by $\text{disc}(H) = \min_{R, B} \text{disc}(H, R, B)$. In other words, $\text{disc}(H)$ measures how well one can partition the vertex set so that each edge has about the same number of vertices from each part.

Theorem 6.2. *If H is an r -uniform hypergraph with m edges, then $\text{disc}(H) \leq 2\sqrt{r \log(2m)}$.*

If H is a clique on $2r - 1$ vertices, then $\text{disc}(H) = r$ and $m \approx 4^r$, so this result is essentially best possible for general H .

Proof. Assign each vertex of H to R or B independently and with probability $\frac{1}{2}$. For $e \in E(H)$, let A_e be the event that

$$\left| |e \cap R| - \frac{1}{2}r \right| \geq \sqrt{r \log(2m)} = 2\sqrt{\frac{\log(2m)}{r}} \cdot \frac{1}{2}r.$$

Because $|e \cap R|$ has a binomial distribution, the Chernoff bound gives $\Pr[A_e] < 2e^{-\log(2m)} = m^{-1}$, and by a union bound we have $\Pr[\bigcup A_e] < 1$. Thus with positive probability, there exists a partition R, B such that none of the A_e occur. This means $\text{disc}(H, R, B) \leq 2\sqrt{r \log 2m}$, proving the result. \square

Much more can be said about discrepancy problems, see [6, Chapter 13].

6.2 Martingales

We say that a sequence of real-valued random variables X_0, X_1, \dots is a *martingale* if $\mathbb{E}[X_{i+1}|X_i] = X_i$ for all i . One important class of Martingales, called Doob martingales, are defined as follows. Given random variables Y_1, \dots, Y_m and a real-valued function f , let

$$X_i = \mathbb{E}[f(Y_1, \dots, Y_m) | Y_1, \dots, Y_i].$$

It is not too difficult to show that any sequence of random variables X_i defined in this way is indeed a martingale.

One of the most common classes of (Doob) martingales in probabilistic combinatorics are the edge-exposure martingales. In this case, Y_i denotes the indicator random variable which is 1 if the i th pair of vertices in $G_{n,p}$ is an edge (where the pairs are ordered in some arbitrary way). Intuitively in this situation we think of revealing the edges of $G_{n,p}$ one at a time, and X_i denotes the value that we expect f to be after we reveal all of the remaining edges.

Let us look at the very concrete case of the edge-exposure martingale when $n = 3$ and f is the number of triangles in $G_{3,p}$. With this we have

$$X_0 = \mathbb{E}[f] = p^3, \quad X_1 = \mathbb{E}[f|Y_1] = p^2 Y_1,$$

$$X_2 = \mathbb{E}[f|Y_1, Y_2] = p Y_1 Y_2, \quad X_3 = \mathbb{E}[f|Y_1, Y_2, Y_3] = Y_1 Y_2 Y_3.$$

The main concentration result for martingales is Hoeffding's inequality.

Theorem 6.3 (Hoeffding's inequality). *Let X_0, \dots be a martingale with $|X_i - X_{i-1}| \leq \alpha_i$ for all i . Then for all $\lambda > 0$, we have*

$$\Pr[|X_m - X_0| \geq \lambda] < 2e^{\frac{-2\lambda^2}{\sum \alpha_i^2}}.$$

Sketch of Proof. Let $Y_i = X_i - X_{i-1}$. Similar to the proof of the Chernoff bound, we have

$$\Pr[X_m - X_0 \geq \lambda] = \Pr[e^{t(X_m - X_0)} \geq e^{t\lambda}] \leq \mathbb{E}[e^{t \sum_{i=1}^m Y_i}] e^{-t\lambda}.$$

We claim that this expectation is at most $e^{\frac{1}{8}t^2 \sum_{i=1}^m \alpha_i^2}$. Indeed, we can use conditional expectations to write

$$\mathbb{E}[e^{t \sum_{i=1}^m Y_i}] = \mathbb{E} \left[\mathbb{E}[e^{t \sum_{i=1}^m Y_i} | X_0, \dots, X_{m-1}] \right] = \mathbb{E}[e^{t \sum_{i=1}^{m-1} Y_i} \cdot \mathbb{E}[e^{t Y_m} | X_0, \dots, X_{m-1}]] ,$$

where this last step used that Y_i with $i < m$ is fixed given X_0, \dots, X_{m-1} . Observe that conditional on X_0, \dots, X_{m-1} , we have $\mathbb{E}[Y_m] = 0$ (due to the martingale property) and $|Y_m| \leq \alpha_m$ (due to the hypothesis of the theorem). One can show that for random variables of this form, the expected value of its moment generating function is at most $e^{\alpha_m^2 t^2 / 8}$. One gets the claim by repeating this argument inductively on the remaining terms.

In total, we have for any $t > 0$ that

$$\Pr[X_m - X_0 > \lambda \sqrt{m}] \leq e^{\frac{1}{8}t^2 \sum \alpha_i^2 - t\lambda}.$$

Taking $t = 4\lambda / \sum \alpha_i^2$ gives $\Pr[X_m - X_0 \geq \lambda] < e^{\frac{-2\lambda^2}{\sum \alpha_i^2}}$. A symmetric argument shows $\Pr[X_m - X_0 \leq -\lambda] < e^{\frac{-2\lambda^2}{\sum \alpha_i^2}}$ (this can also be seen by considering the martingale $X'_i := -X_i$ and applying the first inequality), which gives the result. \square

In the special case where $\alpha_i = 1$ for all i , this result is referred to as Azuma's inequality¹.

Corollary 6.4 (Azuma's inequality). *Let X_0, \dots be a martingale which satisfies $|X_i - X_{i-1}| \leq 1$ for all i . Then for all $\lambda > 0$, we have*

$$\Pr[|X_m - X_0| \geq \lambda \sqrt{m}] < 2e^{-2\lambda^2}.$$

There are many generalizations of the Hoeffding's inequality which weakens the hypothesis that $|X_{i+1} - X_i| \leq \alpha_i$. For example, it suffices to have that this difference holds in expectation, or that it holds with high probability. Again, see [37] for details.

One application of Azuma's inequality is the following.

Proposition 6.5. *We have*

$$\Pr[|\chi(G_{n,p}) - \mathbb{E}[\chi(G_{n,p})]| \geq \lambda \sqrt{n}] < 2e^{-\lambda^2/2}.$$

We note that this result tells us that $\chi(G_{n,p})$ is concentrated around its expectation, but it gives no indication of what this expectation is. This is a common phenomenon when applying concentration inequalities.

¹The naming convention for these inequalities are all over the place: some people call these Hoeffding's inequalities, others Azuma (which is probably the most popular name in the combinatorics community), some Azuma-Hoeffding, and yet others Hoeffding-Azuma.

Proof. We consider a vertex-exposure martingale, i.e. a Doob martingale $f(Y_1, \dots, Y_n)$ where Y_i is the set of vertices $j > i$ in $G_{n,p}$ which are adjacent to i . In particular, taking $f = \chi$ and $X_i = \mathbb{E}[\chi | Y_1, \dots, Y_i]$ gives $X_0 = \mathbb{E}[\chi(G_{n,p})]$ and $X_n = \chi(G_{n,p})$. It is clear that each time we reveal a set Y_i that the expected chromatic number changes by at most 1, i.e. $|X_{i+1} - X_i| \leq 1$ for all i . Thus Azuma's inequality applies, giving the result. \square

We note that one could try and prove this result using an edge-exposure martingale instead of a vertex-exposure martingale, but this approach gives essentially trivial bounds. In general, when using martingales you want to reveal information in as few rounds as possible, while also making it so that the information you reveal can't dramatically change your function each round.

While Proposition 6.5 says nothing about $\mathbb{E}[\chi(G_{n,p})]$, it is well known that this value is asymptotic to $\frac{n}{2 \log_{1/(1-p)} n}$ for any fixed p . This was first proven by Bollobás using a clever martingale argument. Much more can be said about $\chi(G_{n,p})$, see for example the paper by Heckel and Riordan [64] which, in addition to surveying many of the known results on $\chi(G_{n,p})$, shows that the concentration in Proposition 6.5 is in some sense close to best possible.

There are a number of variants of all of the concentration inequalities stated in this chapter. One particular version of Azuma that we will need at some point is the following.

Lemma 6.6 ([78]). *Let X_0, \dots be a martingale which satisfies $|X_i - X_{i-1}| \leq \alpha$ for all i . Then for all $\delta \in [0, 1]$, we have*

$$\Pr[|X_m - X_0| \geq \delta \alpha m] < e^{-\delta^2 \alpha m / 6c}.$$

Note that $|X_m - X_0| \leq \alpha m$ deterministically, so $\delta \alpha m$ is at least a δ fraction of the mean of $X_m - X_0$, and as such this is referred to as the “multiplicative Azuma inequality” (since its error term is multiplicative relative to the expectation as opposed to additive).

6.3 Talagrand's Inequality

Let $\Omega = \Omega_1 \times \dots \times \Omega_n$ be a product of probability spaces. For $\alpha = (\alpha_1, \dots, \alpha_n)$ a vector of non-negative real numbers, we define the *weighted Hamming distance* d_α on Ω by $d_\alpha(x, y) = \sum_{i: x_i \neq y_i} \alpha_i$. For example, $\alpha = (1, \dots, 1)$ gives the usual Hamming distance on product spaces.

Given α as above, a set $A \subseteq \Omega$, and a non-negative number t , we define

$$A_{\alpha,t} = \{x : \exists y \in A, d_\alpha(x, y) \leq t\}.$$

That is, $A_{\alpha,t}$ is the set of points in Ω which are within distance t of A . We also define $\overline{A} = \Omega \setminus A$. Our goal is to prove “isoperimetric” inequalities which state that, for any $A \subseteq \Omega$, we have

$$\Pr[A] \cdot \Pr[\overline{A_{\alpha,t}}] \leq f(t),$$

where f is some rapidly shrinking function. Isoperimetric inequalities are intimately related to concentration inequalities. For example, a corollary of an inequality as above is that if $\Pr[A] \geq \frac{1}{2}$, then $\Pr[\overline{A_{\alpha,t}}] \leq 2f(t)$ (i.e., most of Ω is concentrated around A). On the other hand, one can prove isoperimetric inequalities by using concentration inequalities.

Proposition 6.7. *For Ω a product space, $A \subseteq \Omega$, and α such that $\sum \alpha_i^2 = 1$, we have for all t that*

$$\Pr[A] \Pr[\overline{A_{\alpha,t}}] \leq 4e^{-t^2}$$

Proof. Define the function $f : \Omega \rightarrow \mathbb{R}$ by $f(y) = d_\alpha(y, A)$. Let $Y = (Y_1, \dots, Y_n)$ be chosen according to the probability distribution on Ω and let $X_i = \mathbb{E}[f(Y)|Y_1, \dots, Y_i]$. Observe that $X_n = 0$ iff $Y \in A$ and $X_n > t$ iff $Y \in \overline{A_{\alpha,t}}$, and also that $|X_i - X_{i-1}| \leq \alpha_i$. Thus Hoeffding's inequality implies

$$\begin{aligned} \Pr[A] \Pr[\overline{A_{\alpha,t}}] &= \Pr[X_n = 0] \Pr[X_n > t] \\ &\leq \Pr[|X_n - X_0| \geq X_0] \Pr[|X_n - X_0| > t - X_0] \\ &\leq 4e^{-2X_0^2 - 2(t-X_0)^2} \leq 4e^{-t^2}, \end{aligned}$$

where this last step used that the exponent is maximized when $X_0 = \frac{1}{2}t$. \square

A remarkable result of Talagrand shows that Proposition 6.7 essentially holds even when comparing A with the set of points which are at least distance t from A for *some* choice of α .

Theorem 6.8 (Talagrand's inequality). *For all $A \subseteq \Omega$ and $t \geq 0$, we have*

$$\Pr[A] \Pr \left[\bigcap_{\alpha} \overline{A_{\alpha,t}} \right] \leq e^{-t^2/4},$$

where the intersection ranges over all α with $\sum \alpha_i^2 = 1$.

Again we emphasize that $\bigcap_{\alpha} \overline{A_{\alpha,t}}$ can be much larger than $\overline{A_{\alpha,t}}$ for any given α , but still essentially the same bound as in Proposition 6.7 holds. We omit the proof of Theorem 6.8, and we refer the reader to [6] for a direct proof, and to [37] for a longer, but perhaps more enlightening argument.

We note that Talagrand's inequality is often stated in the following equivalent form: Given $x \in \Omega$ and $A \subseteq \Omega$, define $d'(x, A) = \min_{y \in A} \max_{\alpha} d_\alpha(x, y)$, where the maximum ranges over all α with $\sum \alpha_i^2 = 1$. Note that having $d'(x, A) \leq t$ is equivalent to saying that for all α there exist $y \in A$ with $d_\alpha(x, y) \leq t$, which is equivalent to saying $x \in \bigcap_{\alpha} A_{\alpha,t}$. Thus Theorem 6.8 can be seen as an isoperimetric inequality with respect to the pseudo-distance d' .

Talagrand's inequality has a number of applications to concentration of random variables. One particular application is for certifiable functions. For a function $s : \mathbb{R} \rightarrow \mathbb{N}$, we say that a real-valued function f defined on a product space Ω is *s-certifiable* if having $f(x) \geq c$ implies that there exists a set $I \subseteq [n]$ of size $s(c)$ such that $f(y) \geq c$ whenever $y_i = x_i$ for all $i \in I$ (that is, the values in position I “certify” that $f(x) \geq c$).

For example, if $f(x) = |\{i : x_i \neq 0\}|$, then f is *s-certifiable* with $s(c) = c$, since $f(x) \geq c$ implies there exist c coordinates with $x_i \neq 0$, and any y which agrees with x on these coordinates satisfies $f(y) \geq c$. Lastly, we say that a function f is Lipschitz if $|f(x) - f(y)| \leq 1$ whenever x, y differ in at most one coordinate.

Corollary 6.9. *If f is an s -certifiable Lipschitz function on the product space Ω and X is chosen according to the probability space Ω , then for all m and $t > 0$ we have*

$$\Pr[f(X) < m - t\sqrt{s(m)}] \Pr[f(X) \geq m] \leq e^{-t^2/4}.$$

Proof. Let $A = \{x : f(x) < m - t\sqrt{s(m)}\}$. We claim that $\bigcap_{\alpha} A_{\alpha,t} \subseteq \{y : f(y) < m\}$.

Assume for contradiction that $y \in \bigcap_{\alpha} A_{\alpha,t}$ and $f(y) \geq m$. Because f is s -certifiable, there exists a set of $s(m)$ indices I which certifies $f(y) \geq m$. Let $\alpha'_i = \frac{1}{\sqrt{s(m)}}$ if $i \in I$ and $\alpha'_i = 0$ otherwise. Because $y \in \bigcap_{\alpha} A_{\alpha,t} \subseteq A_{\alpha',t}$, there exists some $x \in A$ such that $d_{\alpha'}(x, y) \leq t$, i.e. such that restricted to I , the vectors x, y differ in at most $t\sqrt{s(m)}$ coordinates. Let z be defined by $z_i = y_i$ if $i \in I$ and $z_i = x_i$ otherwise. Then $f(z) \geq m$ by definition of I , and f being Lipschitz implies $f(x) \geq f(z) - t\sqrt{s(m)} \geq m - t\sqrt{s(m)}$, contradicting $x \in A$. This proves the claim.

The contrapositive of the claim implies $\{y : f(y) \geq m\} \subseteq \overline{\bigcap_{\alpha} A_{\alpha,t}}$, so the first result follows from Talagrand's inequality. \square

We emphasize that Proposition 6.7 is too weak to prove Corollary 6.9: we genuinely have to make use of the fact that Talagrand's inequality allows us to choose a different distance function for each choice of y .

In most applications, one applies Corollary 6.9 where either m is a median, i.e. $\Pr[f(X) \geq m] = \frac{1}{2}$, or where $m - t\sqrt{s(m)}$ is a median. While medians are hard to estimate directly, a concentration result like that of Corollary 6.9 can usually be used to show that the median and expectation must be close to each other, see for example [37, Problem 11.4].

As an application, let $X = (X_1, \dots, X_n)$ be a random vector with each X_i distributed uniformly on $[0, 1]$. Let $f(X)$ denote the length of a longest increasing subsequence, i.e. the largest k such that there exist indices with $X_{i_1} < X_{i_2} < \dots < X_{i_k}$. Note that f is Lipschitz and is s -certifiable with $s(c) = c$, so if m is a median we conclude $\Pr[f(X) < m - t\sqrt{m}] \leq 2e^{-t^2/4}$. It is well known that $\mathbb{E}[f(X)] \sim 2\sqrt{n}$, so at least heuristically, this argument suggests $f(X)$ is highly concentrated around $2\sqrt{n} + \Theta(n^{1/4})$ (and it's not hard to make this more precise). In contrast, if one attempted to get concentration results for $f(X)$ by utilizing martingales, one would conclude that $f(X)$ is highly concentrated around $\Theta(n^{1/2})$, which is significantly weaker. One can literally dedicate an entire book to the longest increasing subsequence problem, see Romik [97] for more on this topic.

7 Coupling

It is often the case that one can understand a random variable X by comparing it to a “similar” random variable Y which is easier to do calculations for. One way to do this to form a *coupling*, i.e. a pair of random variables (X', Y') such that X', Y' have the same distribution as X, Y , respectively, and such that X', Y' have some (nice) relation between them. In particular, we will consider two beautiful couplings around random graphs, one relating $G_{n,p}$ to random digraphs, and the other to random hypergraphs.

7.1 Graphs and Digraphs

Define the random digraph $D_{n,p}$ as the digraph on n vertices obtained by including each ordered pair (u, v) as an arc independently and with probability p . Note that $D_{n,p}$ may have directed 2-cycles, i.e. it will contain both arcs (u, v) and (v, u) with probability p^2 .

Our goal for this subsection is to use coupling arguments to show that $D_{n,p}$ and $G_{n,p}$ exhibit similar “behaviors”, which will allow one to lift results from one setting to the other. Our approach can be generalized to a very large class of properties of graphs/digraphs, but for ease of presentation, we focus only on the property of Hamiltonicity. To this end, we say a digraph D is *Hamiltonian* if one can order the vertices as v_1, \dots, v_n such that $v_i v_{i+1} \in E(D)$ for all $1 \leq i \leq n$ (with indices being written cyclicly), and we will call such an ordering of its vertices a *Hamiltonian cycle*.

As a warmup, we prove the following basic coupling result.

Proposition 7.1. *For all $p \in [0, 1]$, we have*

$$\Pr[D_{n,p} \text{ is Hamiltonian}] \leq \Pr[G_{n,q} \text{ is Hamiltonian}],$$

where $q := 2p - p^2$.

Proof. Our main goal is to construct (correlated) random variables D, G such that (1) D and G have the same distributions as $D_{n,p}$ and $G_{n,q}$, respectively, and such that (2) G is Hamiltonian whenever D is. From this the result will quickly follow. And in this case the path forward is relatively easy: start with a random digraph $D \sim D_{n,p}$, then take G to be the graph obtained from D by “forgetting” the orientations of each arc. One can easily check that conditions (1) and (2) are satisfied here. For completeness, we consider a more formal argument below.

Let $\{X_{u,v} : u, v \in [n]\}$ be a collection of iid Bernoulli random variables with success probability p . Define the random digraph D on $[n]$ by including the arc (u, v) iff $X_{u,v} = 1$ and define the random graph G on $[n]$ by including the edge uv if and only if $\max\{X_{u,v}, X_{v,u}\} = 1$. It is not difficult to see that D being Hamiltonian implies that G is Hamiltonian (since in particular, v_1, \dots, v_n being a Hamiltonian cycle in D implies it is also a Hamiltonian cycle in G), so in particular we find

$$\Pr[D \text{ is Hamiltonian}] \leq \Pr[G \text{ is Hamiltonian}].$$

The result follows since (as is easy to check), D, G are random variables distributed according to $D_{n,p}, G_{n,q}$ respectively. \square

We now wish to reverse Proposition 7.1 by lowering bounding the probability that $D_{n,p}$ is Hamiltonian in terms of that for $G_{n,p}$ by using a very nice coupling by McDiarmid [85].

Theorem 7.2 ([85]). *For all $p \in [0, 1]$, we have*

$$\Pr[G_{n,p} \text{ is Hamiltonian}] \leq \Pr[D_{n,p} \text{ is Hamiltonian}].$$

Proof. Naively, one might try to perform the same approach as in Proposition 7.1 by starting with $G \sim G_{n,p}$ and then replacing each edge of G with some number of arcs. However, one can quickly work out that such a scheme will not recover a digraph distributed according to $D_{n,p}$, since in particular one expects the number of (unordered) pairs of vertices in $D_{n,p}$ connected by some arc to be about twice as many as the number of edges in $G_{n,p}$.

To remedy the situation, we use a clever idea of McDiarmid. Roughly speaking, instead of considering just the two models $G_{n,p}$ and $D_{n,p}$, we will instead consider a sequence of models D_0, D_1, \dots, D_N which “interpolate” between $G_{n,p}$ and $D_{n,p}$ in the sense that $D_0 = G_{n,p}$ and $D_N = D_{n,p}$ and are such that D_{i-1} and D_i differ only on how they include edges/arcs on a single pair of vertices e_i . Because there is such a small difference between the models D_{i-1} and D_i , we will easily be able to compare the probabilities of each being Hamiltonian, and by iterating this we will obtain our desired comparison for $D_0 = G_{n,p}$ and $D_N = D_{n,p}$.

As a small technical aside, it will be slightly more convenient in this argument to work exclusively with digraphs. To this end, we define a new random digraph model $D_{n,p}^*$ which independently for each unordered pair $\{u, v\}$ includes both arcs $(u, v), (v, u)$ with probability p and excludes both arcs with probability $1 - p$. That is, $D_{n,p}^*$ is equivalent to the digraph obtained from $G_{n,p}$ by replacing each edge uv with the 2-cycle $(u, v), (v, u)$, and in particular the probability that $G_{n,p}$ is Hamiltonian is equal to the probability that $D_{n,p}^*$ is Hamiltonian. As such, it suffices to work with this latter model instead.

We now turn to the formal details. Let $e_1, \dots, e_{\binom{n}{2}}$ be an arbitrary ordering of the unordered pairs of $[n]$, say with $e_j = \{u_j, v_j\}$. For all $0 \leq i \leq \binom{n}{2}$, define the random digraph D_i on $[n]$ as follows: for all $j > i$, independently and with probability p include both of the arcs (u_j, v_j) and (v_j, u_j) in D_i ; and for all $j \leq i$, independently and with probability p include the arc (u_j, v_j) , and independently and with probability p include the arc (v_j, u_j) . Observe that $D_0 \sim D_{n,p}^*$ and that $D_{\binom{n}{2}} \sim D_{n,p}$ (with more generally D_i having exactly i pairs behaving like those in $D_{n,p}$ and the rest like $D_{n,p}^*$), so we will be done if we can show for all $i \geq 1$ that

$$\Pr[D_{i-1} \text{ is Hamiltonian}] \leq \Pr[D_i \text{ is Hamiltonian}].$$

It is not difficult to couple D_{i-1} and D_i in such a way that they agree outside of the pair e_i , and we let D denote this common digraph. Let A denote the event that D is not Hamiltonian but adding at least one arc of e_i makes D Hamiltonian. Observe that

$$\Pr[D_{i-1} \text{ is Hamiltonian} | A^c] = \Pr[D_i \text{ is Hamiltonian} | A^c],$$

since either D is Hamiltonian (in which case D_{i-1}, D_i will both be Hamiltonian with probability 1), or D plus both arcs of e_i is not Hamiltonian (in which case D_{i-1}, D_i will both be Hamiltonian with probability 0). Conditional on A , we have that $\Pr[D_{i-1} \text{ is Hamiltonian} | A^c] = p$, as we will be Hamiltonian iff we include both arcs on e_i . Crucially, we also have $\Pr[D_i \text{ is Hamiltonian} | A^c] \geq$

p , as p is the probability of including some given arc on e_i which will make D_i Hamiltonian (with it being possible for two such arcs to exist, in which case the probability becomes higher). Thus regardless of the event A we see that D_i is at least as likely to be Hamiltonian as D_{i-1} , proving the claim and hence the proof. \square

This coupling approach is very flexible and has found applications in many other settings

Discuss variants of this method, e.g. to rainbow setting as well as perturbed random setting. See e.g. “Rainbow subgraphs of uniformly coloured randomly perturbed graphs”, “Rainbow Hamilton cycles in random graphs and hypergraphs”, and “Spanning cycles in random directed graphs,”

7.2 Graphs and Hypergraphs

Here we discuss a beautiful coupling due to Riordan and Heckel allowing us to translate facts about random hypergraphs to random graphs. As a partial warmup to the argument, we consider a coupling result in a completely unrelated setting (namely that of random walks), whose proof is spiritually similar to theirs while having the advantage of being substantially simpler.

Let S^n denote a simple random walk of length n , i.e. S^n is a random vector $(S_0^n, S_1^n, \dots, S_n^n)$ where $S_0^n = 0$, and $\Pr[S_i^n = S_{i-1}^n + 1] = \Pr[S_i^n = S_{i-1}^n - 1] = \frac{1}{2}$. For n even, let T^n denote a random walk after conditioning on having $T_n^n = 0$ (i.e. we uniformly at random pick a walk which returns to 0 at the end of the walk). It is easy to show via Chernoff bounds that S_t^n is likely to be within roughly \sqrt{t} of 0 for any given value t . While Chernoff bounds don't apply to the random variables T_t^n , intuitively the same conclusion should also hold for T_t^n , since the condition of $T_n^n = 0$ should force T_t^n to be closer to the origin than S_t^n in general. It is possible to make this intuition rigorous, allowing one to bootstrap bounds of S_t^n to T_t^n .

Proposition 7.3. *For n even and all s, t , we have $\Pr[|T_t^n| \geq s] \leq \Pr[|S_t^n| \geq s]$.*

Proof. Our goal is to define a new random vector R^n such that (1) R^n has the same distribution as T^n , and (2) $|R_t^n| \leq |S_t^n|$ for all t . From this the result will quickly follow. Intuitively, we will define R_t^n in rounds by flipping biased coins. If the t th coin lands heads, then R_{t+1}^n moves towards 0, and if it lands tails, it moves towards/away from 0 if and only if S_{t+1}^n moves towards/away from 0. Such a process will always satisfy (2), and it will satisfy (1) by choosing the probability of our biased coins appropriately.

To this end, set $R_0^n = 0$. Given R_t^n , we define a random variable Y_t (which will be our biased coin flips) that equals 1 with probability $\frac{|R_t^n|}{|S_t^n|}$ and is 0 otherwise. If $Y_t = 1$, we set $R_{t+1}^n = R_t^n \pm 1$ such that $|R_t^n| > |R_{t+1}^n|$ (i.e. such that R^n moves towards 0; note that this is well defined since $Y_t = 1$ implies $R_t^n \neq 0$). If $Y_t = 0$ and $R_t^n \neq 0$, then we set $R_{t+1}^n = R_t^n \pm 1$ such that $|R_t^n| > |R_{t+1}^n|$ if and only if $|S_t^n| > |S_{t+1}^n|$ (i.e. R^n move away/towards 0 if S^n moves away/towards 0). If $R_t^n = 0$ then we set $R_t^n = \pm 1$ with equal probability.

It is straightforward to see that (2) is achieved from this process¹. It is not difficult to prove that

¹Any time S^n moves towards 0, R^n does as well, except when $R_t^n = 0$. In this case S_t^n must be an even distance away from 0, so after one step R_t^n is still at least as close to 0.

for T^n , we have that $|T_t^n| > |T_{t+1}^n|$ happens with probability¹ $\frac{\frac{1}{2}(n-t+|T_t^n|)}{n-t}$. One can check that R^n has $|R_t^n| > |R_{t+1}^n|$ with probability $\frac{\frac{1}{2}(n-t+|R_t^n|)}{n-t}$, so we conclude (1) and hence the result. \square

Recall that an F -factor in a graph G is a collection of vertex disjoint copies of F such that every vertex is in one of these copies of F . By a similar argument as in Theorem 9.2, one can show that $G_{n,p}$ contains a K_r -factor provided $r|n$ and $p \gg n^{-1/\binom{r}{2}} \log n$. Intuitively, it seems reasonable that the set of K_r 's in $G_{n,p}$ should be distributed like the hyperedges of $G_{n,\pi}^r$ where $\pi = p^{\binom{r}{2}}$ (at the very least, the expected number of K_r 's in $G_{n,p}$ is equal to the expected number of hyperedges in $G_{n,\pi}^r$). If this were true, then $G_{n,p}$ would contain a K_r -factor when $G_{n,\pi}^r$ contains a perfect matching, and Theorem 9.2 says this should happen when $p^{\binom{r}{2}} \approx \pi \gg n^{-1} \log n$, which implies that taking $p \gg n^{-1/\binom{r}{2}} \log^{1/\binom{r}{2}} n$ should suffice. And indeed, Johansson, Kahn, and Vu [68] proved that this is the threshold for K_r -factors in $G_{n,p}$ using a somewhat involved argument. A nice coupling result of Riordan [95] will allow us to conclude the result in a much easier way.

Let $(V_1, E_1), \dots, (V_{\binom{n}{r}}, E_{\binom{n}{r}})$ be an arbitrary ordering of all the K_r 's in K_n . To prove our desired coupling, we would like to construct a pair of random variables (G, H) such that (1) $G \sim G_{n,p}$ and $H \sim G_{n,\pi}^r$ with $\pi \approx p^{\binom{r}{2}}$, and such that (2) every hyperedge in H is a K_r in G . Note that (2) means that H containing a perfect matching implies that G has a K_r -factor. Let us first consider the following (very, very) naive attempt at this coupling.

Algorithm 1. Generate a random graph $G \sim G_{n,p}$. Let H be an initially empty r -graph on $[n]$. For each i with $E_i \subseteq G$, add V_i as a hyperedge to H . Output (G, H) .

This algorithm definitely satisfies (2), but it completely fails at (1). Indeed, let A_i denote the event that $E_i \subseteq G$ (i.e. the event that V_i is a hyperedge in H), and assume V_1, V_2 have at least two vertices in common. Then $\Pr[A_2|A_1] \geq p^{\binom{r}{2}-1}$ and $\Pr[A_2|\overline{A_1}] < p^{\binom{r}{2}}$. But to have $H \sim G_{n,\pi}^r$ we would, in particular, need these two probabilities to equal each other. Thus we'll need to consider a somewhat more complicated algorithm. As before, let (V_i, E_i) be the K_r 's in K_n , and let π be a parameter which will be approximately $p^{\binom{r}{2}}$.

Algorithm 2. Generate a random graph $G \sim G_{n,p}$ and an initially empty hypergraph H on $[n]$. We proceed in $\binom{n}{r}$ rounds as follows. For the i th round, let π_i be the conditional probability of having $E_i \subseteq G$ given all the information from the previous rounds.

- If $\pi_i < \pi$, then with probability π we add V_i to H .
- If $\pi_i \geq \pi$, then with probability $\frac{\pi}{\pi_i}$ we test whether $E_i \subseteq G$, and if so, we add V_i to H . Otherwise² we declare this hyperedge to be absent in H .

¹Given T_t^n , we still need to make $\frac{1}{2}(n-t+|T_t^n|)$ steps in the direction of 0 and $\frac{1}{2}(n-t-|T_t^n|)$ in the direction away from 0.

²Note that with probability $1 - \frac{\pi}{\pi_i}$ we do not reveal any additional information about E_i . When working with random objects, it is usually best to reveal as little information as possible in order to “preserve” the randomness of your object.

We note that the $\pi_i \geq \pi$ case of Algorithm 2 is similar in spirit to the proof of Proposition 7.3: each round we flip a coin which is biased based off of the current information we have. If the coin lands heads we do something to H independent of G , and otherwise we have H behave “in the same way” as G .

For Algorithm 2, it is not difficult to see that $H \sim G_{n,\pi}^r$. Unfortunately, if $\pi_i < \pi$, then it is possible that H contains edges which are not K_r ’s in G , i.e. the coupling could fail to satisfy (2). The key insight is that for applications, it suffices to have (2) be satisfied with high probability, which will turn out to be the case.

To try and convince ourselves that this algorithm has a chance of winning even when $\pi_i < \pi$, let’s consider the most dangerous situation, namely that $\pi_i = 0$. It is not too hard to see that $\pi_i = 0$ if and only if there exists some $j < i$ such that (a) we revealed that $E_j \not\subseteq E(G)$ and (b) every edge of $E_j \setminus E_i$ has been revealed to be in G . If this situation happens and if the algorithm adds V_i to H , then the coupling fails to satisfy (2). However, when this happens, every graph edge of E_j is contained in a hyperedge of H (by (b) and $V_i \in E(H)$) and V_j is not a hyperedge of H (by (a)). Thus the probability of this situation happening is at most the probability of $H \sim G_{n,\pi}^r$ containing such a configuration. These configurations can essentially be described as follows.

Lemma 7.4. *If H is an r -graph with $r \geq 4$ which contains a set of r vertices $V \notin E(H)$ such that every pair of V is contained in a hyperedge of H , then H contains a subgraph F which has $e(F) \leq \binom{r}{2}$ and $|V(F)| \leq (r-1)e(F) - 1$.*

This statement is false for $r = 3$. Indeed, one could take H to be the loose triangle with edges $\{1, 2, 4\}, \{2, 3, 5\}, \{1, 3, 6\}$ which satisfies the hypothesis of Lemma 7.4 with $V = \{1, 2, 3\}$ but which fails to satisfy the conclusion.

Proof. Let V_1, \dots, V_t be hyperedges such that every pair of V is contained in some V_i . By throwing away redundant hyperedges, we can assume that $|V_i \cap V| \geq 2$ for all i and that $t \leq \binom{r}{2}$. Let $F \subseteq H$ be the hypergraph with hyperedges V_1, \dots, V_t .

First assume $|V_1 \cap V_2| \geq 2$. Then $V_1 \cup V_2$ consists of at most $2r - 2$ vertices, and it is not difficult to see that it is possible to order the remaining sets so that $|V_i \setminus \bigcup_{j < i} V_j| \leq r - 1$ and that $|V_t \setminus \bigcup_{j < t} V_j| \leq r - 2$. In total this implies that F has the desired properties.

Thus we can assume that $|V_i \cap V_j| \leq 1$ for all i, j . This means every pair of V is covered by some unique V_i , so $t = e(F) = \binom{r}{2}$ and the number of vertices of F is at most $r + (r-2)e(F) = (r-1)e(F) - (e(F) - r) \leq (r-1)e(F) - 1$ since $\binom{r}{2} - r \geq 1$ for $r \geq 4$. \square

Lemma 7.5. *For $r \geq 4$, if $H \sim G_{n,\pi}^r$ and $\pi \leq n^{-(r-1)+o(1)}$, then a.a.s. H does not contain a set V as in Lemma 7.4.*

Proof. If H did contain such a set V , then it must contain a subgraph F as in Lemma 7.4. Up to isomorphism, there are only finitely many subgraphs that F could be, and for each of these the expected number of copies of F in H is at most

$$O(\pi^{e(F)} n^{|V(F)|}) = O(\pi^{e(F)} n^{(r-1)e(F)-1}) = o(1).$$

We conclude the result by Markov’s inequality. \square

We note that for $\pi \approx p^{\binom{r}{2}}$ this lemma applies when $p \approx n^{-2/r}$. Thus when p is about this value, none of the “bad” configurations of Lemma 7.4 are likely to appear, and in this regime we have the following.

Theorem 7.6 ([95]). *For $r \geq 4$ and $p \leq n^{-2/r+o(1)}$, there exists some $\pi \sim p^{\binom{r}{2}}$ such that Algorithm 2 produces a pair (G, H) with $G \sim G_{n,p}$, $H \sim G_{n,\pi}^r$, and such that a.a.s. every hyperedge of H is the vertex set of a K_r in G .*

We emphasize that the theorem as stated does not cover the case $r = 3$. However, Heckel [63] showed that the same conclusion does hold for $r = 3$ by using a slightly different coupling.

For the proof of Theorem 7.6, we will need a standard result known as Harris’ inequality (also referred to as Kleitman’s inequality).

Lemma 7.7 (Harris’ Inequality). *Let $f, g, h : \mathbb{R}^n \rightarrow \mathbb{R}$ be functions such that f, g are non-decreasing and h is non-increasing. Let $X = (X_1, \dots, X_n)$ be a random vector such that the X_i ’s are mutually independent. Then*

$$\mathbb{E}[f(X)g(X)] \geq \mathbb{E}[f(X)]\mathbb{E}[g(X)],$$

$$\mathbb{E}[f(X)h(X)] \leq \mathbb{E}[f(X)]\mathbb{E}[h(X)].$$

Proof. For $n = 1$, we deterministically have

$$(f(y) - f(z))(g(y) - g(z)) \geq 0 \geq (f(y) - f(z))(h(y) - h(z)).$$

Thus if Y, Z are independent random variables with the same distribution as $X = X_1$, the first inequality implies

$$0 \leq \mathbb{E}[f(Y)g(Y) + f(Z)g(Z) - f(Y)g(Z) - f(Z)g(Y)] = 2\mathbb{E}[f(X)g(X)] - 2\mathbb{E}[f(X)]\mathbb{E}[g(X)].$$

This gives the first bound, and the second bound follows from an identical argument.

Assume the result has been proven up to some $n > 1$. By the inductive hypothesis and the $n = 1$ case applied to $f'(X_1) := \mathbb{E}[f(X)|X_1]$ and $g'(X_1) = \mathbb{E}[g(X)|X_1]$, we find

$$\begin{aligned} \mathbb{E}[f(X)g(X)] &= \mathbb{E}[\mathbb{E}[f(X)g(X)|X_1]] \geq \mathbb{E}[\mathbb{E}[f(X)|X_1] \cdot \mathbb{E}[g(X)|X_1]] \\ &= \mathbb{E}[f'(X_1)g'(X_1)] \geq \mathbb{E}[f'(X_1)]\mathbb{E}[g'(X_1)] = \mathbb{E}[f(X)]\mathbb{E}[g(X)]. \end{aligned}$$

This proves the first inequality, and the second follows from an identical argument. \square

The main application of Harris’ inequality is when f is an indicator function. More precisely, we say that a set system $\mathcal{A} \subseteq 2^{[n]}$ is an *upset* if $A \in \mathcal{A}$ implies $B \in \mathcal{A}$ for all $B \supseteq A$, and we similarly define what it means for \mathcal{A} to be a *downset*.

Corollary 7.8. *Let \mathcal{A}, \mathcal{B} be upsets and \mathcal{C} a downset of $[n]$, and let $S \subseteq [n]$ be obtained by including each element i independently and with probability p_i . Then*

$$\Pr[S \in \mathcal{A} \cap \mathcal{B}] \geq \Pr[S \in \mathcal{A}] \Pr[S \in \mathcal{B}],$$

$$\Pr[S \in \mathcal{A} \cap \mathcal{C}] \leq \Pr[S \in \mathcal{A}] \Pr[S \in \mathcal{C}].$$

Proof. Define $f : \mathbb{R}^n \rightarrow \mathbb{R}$ by having $f(x) = 1$ if $\{i : x_i > 0\} \in \mathcal{A}$ and $f(x) = 0$ otherwise. Similarly define g, h with respect to \mathcal{B}, \mathcal{C} . The result follows from Harris' inequality by letting $X = (X_1, \dots, X_n)$ with the X_i being independent Bernoulli random with probability p_i . \square

Proof of Theorem 7.6. Again, the only cases where the algorithm can fail is when π_i is small, so let us try and lower bound this quantity in terms of the (random) information we have at the i th step. Let Y be the set of “yes” indices j such that we have revealed that $E_j \subseteq E(G)$ and let N be the set of “no” indices such that we have revealed $E_j \not\subseteq E(G)$. Let $R = \bigcup_{j \in Y} E_j$ be the set of revealed edges, and let G' be the random graph which contains all of the edges of R , and which contains any $e \notin R$ independently and with probability p . Let $E'_j = E_j \setminus R$, and let A'_j be the event that $E'_j \subseteq G'$. It is not too hard to see that in total we have

$$\pi_i = \Pr[A'_i | \bigcap_{j \in N} \overline{A'_j}].$$

Define

$$D_0 = \bigcap_{j \in N, E_j \cap E_i = \emptyset} \overline{A'_j}, \quad D_1 = \bigcap_{j \in N, E_j \cap E_i \neq \emptyset} \overline{A'_j}.$$

Intuitively D_0 shouldn't really influence π_i , and we can prove this using Harris' inequality. First note that

$$\pi_i = \Pr[A'_i | D_0 \cap D_1] \geq \Pr[A'_i \cap D_1 | D_0] = \Pr[A'_i | D_0] - \Pr[A'_i \cap \overline{D_1} | D_0] = \Pr[A'_i] - \Pr[A'_i \cap \overline{D_1} | D_0],$$

where this last step used that A'_i and D_0 are independent. Observe that A'_j is an upset for all j (i.e. A'_j is achieved precisely when the random set $E(G')$ is an element of an appropriately defined upset), D_0 is a downset (since complements of upsets are downsets, and downsets/upsets are preserved under intersection), and $A'_i \cap \overline{D_1}$ is an upset. Thus by Harris' inequality, we have¹

$$\Pr[A'_i] - \Pr[A'_i \cap \overline{D_1} | D_0] = \Pr[A'_i] - \frac{\Pr[A'_i \cap \overline{D_1} \cap D_0]}{\Pr[D_0]} \geq \Pr[A'_i] - \Pr[A'_i \cap \overline{D_1}].$$

Now let $N_1 = \{j \in N : E_j \cap E_i \neq \emptyset\}$. Note that $\overline{D_1} = \bigcup_{j \in N_1} A'_j$, so by a union bound we have

$$\pi_i \geq \Pr[A'_i] - \sum_{j \in N_1} \Pr[A'_j \cap A'_i] = p^{|E_i|} - \sum_{j \in N_1} p^{|E'_i \cup E'_j|} = p^{|E'_i|} (1 - Q_i) \geq p^{\binom{r}{2}} (1 - Q_i),$$

where

$$Q_i := \sum_{j \in N_1} p^{|E_j \setminus (E_i \cup R)|}.$$

Let Δ denote the maximum degree of G . We next prove a (somewhat imprecise) claim.

Claim 7.9. *Either $\Delta > n^{o(1)}$, or for all i , either $Q_i = o(1)$ or $V_i \in E(H)$ implies H contains a configuration as in Lemma 7.4.*

¹This same sort of argument is essentially what you need to do prove Janson's inequality Theorem 3.6.

Proof. Assume $\Delta \leq n^{o(1)}$ and consider some index i . Given j , let K_j denote the graph on V_j with edge set $E_i \cup R$, and let C_1, \dots, C_{k+1} with $k \geq 0$ denote the connected components of K_j , say with $|V(C_\ell)| = r_\ell$ for all ℓ . Observe that $|E_j \setminus (E_i \cup R)|$ is at least the number of edges which aren't contained in any K_j component, i.e.

$$|E_j \setminus (E_i \cup R)| \geq \binom{r}{2} - \sum \binom{r_\ell}{2} \geq \binom{r}{2} - \binom{r-k}{2},$$

where this last inequality holds since if there are two terms with $r_\ell \geq 2$, then one can adjust these two terms to get a stronger bound. Because K_j is a graph using edges of $R \cup E_i \subseteq G$, we have that the number of $j \in N_1$ such that K_j has $k+1$ components is at most $rn^k \Delta^{r-k-1} = n^{k+o(1)}$, where the factor of r comes from the fact that $j \in N_1$ implies that K_j contains at least one vertex of V_i since E_i, E_j intersect in at least one edge.

In total then, the contribution to Q_i coming from j such that K_j has $k+1 \geq 2$ components is at most

$$\sum_{k=1}^{r-2} n^{k+o(1)} p^{\binom{r}{2} - \binom{r-k}{2}} = o(1),$$

where the equality follows from a simple calculation. Thus it remains to show that the contribution from terms with K_j connected is small. Because there are only $r\Delta^{r-1} = n^{o(1)}$ such terms, a similar argument shows that the contribution is negligible for terms with $e(K_j) < \binom{r}{2}$. The only non-trivial case then is when $e(K_j) = \binom{r}{2}$, i.e. when every edge of V_j is contained in $R \cup E_i$. In this case, $V_i \in E(H)$ implies that H contains a configuration as in Lemma 7.4. Thus for all i , either this happens or $Q_i = o(1)$, proving the result. \square

Since $H \sim G_{n,\pi}^r$, we have that the expected degree of every vertex is roughly $\pi n^{r-1} = n^{o(1)}$. Thus if \mathcal{B}_1 is the “bad” event that $\Delta > n^{o(1)}$, then by the Chernoff bound we have $\Pr[\mathcal{B}_1] = o(1)$. Similarly if \mathcal{B}_2 is the event that H contains one of the configurations as in Lemma 7.4, then $\Pr[\mathcal{B}_2] = o(1)$ by Lemma 7.5.

We are now ready to complete the proof. Recall that the theorem claims the result holds for *some* $\pi \sim p^{\binom{r}{2}}$, so it suffices to prove it for $\pi = p^{\binom{r}{2}}(1 - o(1))$ where the $o(1)$ term is the upper bound for Q_i from the claim. Now all we have to do is verify that with this choice, a.a.s. every hyperedge of H is a K_r in G . The only way this can fail is if there exists an i with $\pi_i < \pi$ such that V_i is added as a hyperedge to H . By the previous claim and our choice of π , this is only possible if $\mathcal{B}_1 \cup \mathcal{B}_2$ occurs. As these occur with probability $o(1)$, we conclude the result. \square

As noted previously, the proof of Theorem 7.6 does not go through for $r = 3$ due to the existence of loose triangles, but Heckel [63] managed to get around this issue. Essentially the idea of her proof is to first do a coupling on edges of H and G which are in loose triangles and then to run Riordan's argument.

It is also proven in [95] that one can to some extent generalize this approach to finding F -factors for sufficiently nice F . In this setting, H is not exactly a uniform hypergraph, but instead a collection of copies of F in K_n chosen with some probability π .

8 Random Algebraic Constructions

One can easily extend our general lower bound for graph Turán numbers Theorem 1.4 to the setting of hypergraphs as follows.

Theorem 8.1. *Let F be an r -graph with v vertices and $e \geq r$ edges. If $e \geq v$, then*

$$\text{ex}(n, F) = \Omega_v(n^{r - \frac{v-r}{e-1}}).$$

Sketch of Proof. Consider $G_{n,p}^r$, which in expectation has about pn^r edges and $p^{e(F)}n^{|V(F)|}$ copies of F . At $p = Cn^{-\frac{v-r}{e-1}}$ for some large constant C this first quantity is much larger than the second, so we can delete an edge from each copy of F to give the result. \square

One way you could try and improve upon this argument is to delete edges which are in many copies of F . In $G_{n,p}^r$ this is too much to ask for, but it is possible to do this in other random hypergraph models. In particular, if our random model contains some algebraic structure, then it is often the case that edges will either be in many copies of F or almost none. We look at a few examples of this phenomenon.

8.1 Random Multilinear Maps

The problem of determining the Turán number of $K_{2,\dots,2}^r$, the complete r -partite r -graph with each part having size 2, is called the Erdős box problem. Theorem 8.1 gives a lower bound of $n^{r - \frac{r}{2^r-1}}$, and for certain values of r this lower bound was improved by Gunderson, Rödl, and Sidorenko [62]. This result was significantly improved by Conlon, Pohoata, and Zakharov [29] who gave a polynomial improvement to the bound of Theorem 8.1 for all values of r .

Theorem 8.2 ([29]). *For all $r \geq 2$, we have*

$$\text{ex}(n, K_{2,\dots,2}^r) = \Omega(n^{r - \lceil \frac{2^r-1}{r} \rceil^{-1}}).$$

Note that r never¹ divides $2^r - 1$, so this does always give a polynomial improvement to Theorem 1.4.

We prove this result by considering a random hypergraph based off of multilinear maps. Recall that if V_1, \dots, V_r are vector spaces over \mathbb{F}_q , then a map $T : V_1 \times \dots \times V_r \rightarrow \mathbb{F}_q$ is said to be *multilinear* if the one dimensional function $f(x) = T(v_1, \dots, v_{i-1}, x, v_{i+1}, \dots, v_r)$ is linear for all i and any choice of v_j . Note that there are only finitely many such maps over \mathbb{F}_q if V_1, \dots, V_r are finite dimensional, so in this setting we can talk about choosing such a T uniformly at random.

Let $s = \lceil \frac{2^r-1}{r} \rceil$, and let V_1, \dots, V_r be copies of \mathbb{F}_q^s with q a large prime power. Given a multilinear map T , let H_T denote the r -partite r -graph on $V_1 \cup \dots \cup V_r$ with $\{v_1, \dots, v_r\} \in E(H_T)$ if and only if $T(v_1, \dots, v_r) = 1$ (here and throughout we assume $v_i \in V_i$ for all i). The proof relies on the following three results.

¹If r is prime then $2^r - 1 \equiv 2 - 1 \pmod r$ by Fermat's little theorem **though I don't see why this holds otherwise**

Lemma 8.3. *Let T be a uniformly random multilinear map and assume q is sufficiently large in terms of r . Then the following hold:*

- (a) *We have $\mathbb{E}[e(H_T)] = (q^s - 1)^r q^{-1} \approx q^{rs-1}$.*
- (b) *Let \mathcal{F} denote the set of tuples $(v_1^0, v_1^1, \dots, v_r^0, v_r^1)$ with $v_i^j \in V_i$ and $v_i^0 \neq v_i^1$ such that $T(v_1^{j_1}, \dots, v_r^{j_r}) = 1$ (i.e. such that this forms a $K_{2, \dots, 2}^r$ in H_T). Then $\mathbb{E}[|\mathcal{F}|] \sim q^{2rs-2^r}$.*
- (c) *Let \mathcal{B} denote the set of edges $\{v_1, \dots, v_r\}$ such that $(v_1, v_1', \dots, v_r, v_r') \in \mathcal{F}$ for some $\{v_1', \dots, v_r'\}$. Then $\mathbb{E}[|\mathcal{B}|] \leq (1 + o(1))q^{-r}\mathbb{E}[|\mathcal{F}|]$.*

We note that $G_{n,p}^r$ with p, n chosen appropriately already roughly satisfy (a) and (b), so the crucial thing we gain here is (c), which says that there are not many edges that are contained in some $K_{2, \dots, 2}^r$, i.e. the copies of $K_{2, \dots, 2}^r$ are all clumped together. This is the key fact that we acquire from using a random algebraic construction.

Let us briefly observe that this lemma gives the result. Indeed, we can form a $K_{2, \dots, 2}^r$ -free hypergraph H'_T by deleting every edge of \mathcal{B} . The expected number of edges for this will be asymptotically at least $q^{rs-1} - q^{2rs-2^r-r}$. Because $s = \lceil \frac{2^r-1}{r} \rceil$, we have $s < \frac{2^r-1}{r} + 1$, which is equivalent to saying $rs - 1 > 2rs - 2^r - r$, and hence the number of edges is roughly q^{rs-1} . Since H'_T has $rq^s := n$ vertices, this gives $\text{ex}(n, K_{2, \dots, 2}^r) = \Omega(n^{r-1/s})$ as desired. Thus it remains to prove the lemma.

Proof of Lemma 8.3. For (a), note that $T(v_1, \dots, v_r) = 0$ if $v_i = 0$ for some i . For any other tuple, let $U_i \subseteq V_i$ be the one-dimensional subspace containing v_i and 0. It is not too hard to argue that T restricted to $U_1 \times \dots \times U_r$ is still a uniform multilinear map. Further, every multilinear map on this space is uniquely determined by the value of $T(1, \dots, 1)$, and it is not hard to see that exactly one of these q maps has $T(v_1, \dots, v_r) = 1$. Thus such a tuple is an edge with probability q^{-1} and the result follows from linearity of expectation.

For (b), observe that the only tuples that can be in \mathcal{F} are those such that $v_i^0 \neq \lambda v_i^1$ for any i , as otherwise

$$\lambda = \lambda T(v_1^0, \dots, v_i^0, \dots, v_r^0) = T(v_1^0, \dots, v_i^1, \dots, v_r^0) = 1,$$

which means $\lambda = 1$, contradicting $v_i^0 \neq v_i^1$. The number of such tuples with this property is asymptotic to q^{2rs} . For such a tuple, let U_i be the span of v_i^0, v_i^1 , which is a 2-dimensional subspace. Again T restricted to $U_1 \times \dots \times U_r$ is uniform, and it is not too hard to see that there are q^{2^r} choices for T with exactly one of these placing the tuple in \mathcal{F} . The result follows from linearity of expectation.

It remains to deal with (c). Given affine lines ℓ_1, \dots, ℓ_r in V_1, \dots, V_r , let $P(\ell_1, \dots, \ell_r)$ denote the set of tuples $(v_1, v_1', \dots, v_r, v_r')$ such that $v_i, v_i' \in \ell_i$. It is not difficult to show the following:

- The sets $P(\ell_1, \dots, \ell_r)$ are disjoint for distinct choices of lines.
- We have $|P(\ell_1, \dots, \ell_r)| = q^r(q-1)^r$.
- Every element of \mathcal{F} is contained in some $P(\ell_1, \dots, \ell_r)$.

- If $P(\ell_1, \dots, \ell_r) \cap \mathcal{F} \neq \emptyset$ then $P(\ell_1, \dots, \ell_r) \subseteq \mathcal{F}$, i.e. $T(u_1, \dots, u_r) = 1$ for any $u_i \in \ell_i$.

If \mathcal{L} denotes the set of tuples (ℓ_1, \dots, ℓ_r) with $P(\ell_1, \dots, \ell_r) \cap \mathcal{F} \neq \emptyset$, then the above implies that

$$|\mathcal{L}|q^r(q-1)^r = |\mathcal{F}|.$$

Further, we have

$$|\mathcal{B}| = \left| \bigcup_{(\ell_1, \dots, \ell_r) \in \mathcal{L}} \ell_1 \times \dots \times \ell_r \right| \leq q^r |\mathcal{L}| = (q-1)^{-r} |\mathcal{F}|,$$

so taking expectations gives the result. \square

We note that one can get a slightly stronger result by not just considering one multilinear map T , but a family of (random) multilinear maps T_1, \dots, T_ℓ and then defining H_{T_1, \dots, T_ℓ} by having a hyperedge if and only if $T_i(v_1, \dots, v_r) = 1$ for all i . The analysis here is mostly the same, but for ease of presentation we only considered a single map.

8.2 Random Polynomial Graphs

Somewhat more complicated constructions can be made by utilizing random polynomials as opposed to random multilinear maps. This approach was first popularized by Bukh [20], and since then Bukh and Conlon have developed a lot of theory surrounding it.

To set things up, given a field \mathbb{F}_q , we define $\mathcal{P}_{d,b}$ to be the set of polynomials over \mathbb{F}_q in t variables with degree at most d . We will say that f is a random polynomial from $\mathcal{P}_{d,b}$ if it is chosen uniformly at random from $\mathcal{P}_{d,b}$, which can be done, for example, by uniformly at random choosing the coefficient of each possible monomial. With a little bit of linear algebra one can show the following, which says that a random polynomial has the same distribution as a random function when evaluated on a few number of points.

Lemma 8.4 ([22] Lemma 2.3). *If $q > \binom{m}{2}$ and $d \geq m-1$, then if $f \in \mathcal{P}_{d,b}$ is uniformly random and x_1, \dots, x_m are m distinct points of \mathbb{F}_q^b , then*

$$\Pr[f(x_i) = 0 \ \forall i] = q^{-m}.$$

Maybe include proof.

The next lemma requires just a smidge of terminology from algebraic geometry. A *variety* is any set of the form $X = \{x \in \mathbb{F}_q^b : f_1(x) = \dots = f_a(x) = 0\}$ where $f_1, \dots, f_a : \mathbb{F}_q^t \rightarrow \mathbb{F}_q$ are polynomials. The variety X is said to have complexity at most M if a, b and the degrees of the f_i are bounded by M . One can prove the following using standard results from algebraic geometry.

Lemma 8.5 ([22] Lemma 2.7). *Let X, D be varieties over \mathbb{F}_q of complexity at most M . If q is sufficiently large in terms of M , then either $|X \setminus D| \geq q/2$ or $|X \setminus D| \leq c$ for some c depending only on M .*

The actual lemma statement involves the algebraic closure, which I think is an artifact of the proof and isn't necessary in the statement. Please let me know if you think I'm wrong (or right) about this point.

One can think of this lemma as being analogous to the fact that if f is a degree d polynomial in one variable which is 0 on at least $d + 1$ points, then it must in fact be 0 on an entire line. With these two results we can prove the following.

Theorem 8.6. *For all $s \geq 2$, there exists some $t_0 = t_0(s)$ such that for all $t \geq t_0$, we have*

$$\text{ex}(n, K_{s,t}) = \Theta(n^{2-1/s}).$$

Proof. The upper bound follows from the Kővári-Sós-Túran theorem. For the lower bound, let q be a sufficiently large prime power, and with some foresight we define

$$r = s^2 + 1, \quad d = rs + 1, \quad N = q^s.$$

Let $f \in \mathcal{P}_{d,2s}$ be a polynomial chosen uniformly at random. Let G be the (random) graph with vertex set $\mathbb{F}_q^s \times \mathbb{F}_q^s$ where vertices $x^1 \in \mathbb{F}_q^s, x^2 \in \mathbb{F}_q^s$ form an edge of G if and only if $f(x^1, x^2) = 0$.

Fix vertices $x^1, \dots, x^s \in \mathbb{F}_q^s \cup \mathbb{F}_q^s$. Let C be the set of vertices y such that $x^i \sim y$ for all i (noting that $C = \emptyset$ if the x^i don't all belong to the same copy of \mathbb{F}_q^s , and otherwise this means e.g. $f(x^i, y) = 0$ for all i). Observe that the number of $K_{s,r}$'s of G which has the x^i as its set of size s is equal to $\binom{|C|}{r}$, and motivated by this we will attempt to bound the r th moment $\mathbb{E}[|C|^r] = \mathbb{E}[|C^r|]$ (which will be slightly easier to work with compared to the r th falling moment). To this end, we observe that if a given tuple (y^1, \dots, y^r) with k distinct elements lies in C^r , then the corresponding copy of $K_{s,k}$ lies in G . By Lemma 8.4, the probability that any given copy of $K_{s,k}$ appears in G is exactly q^{-sk} (provided q is sufficiently large in terms of s, r). Moreover, the number of tuples with k distinct elements is $O_r(N^k)$. In total we conclude that

The exposition here can probably be cleaned up

$$\mathbb{E}[|C^r|] \leq \sum_{k=1}^r q^{-sk} \cdot O_r(N^k) = O_r(1).$$

Note that C is an algebraic variety by definition. By Lemma 8.5, there exists some constant c such that either $|C| \leq c$ or $|C| \geq q/2$. Thus

$$\Pr[|C| > c] = \Pr[|C| \geq q/2] = \Pr[|C|^r \geq (q/2)^r] \leq \frac{\mathbb{E}[|C|^r]}{(q/2)^r} = O_r(q^{-r}),$$

with the last step using the previous inequality.

Call a sequence (x^1, \dots, x^s) *bad* if there are more than c vertices y such that $x^i \sim y$ for all i , and let B_i denote the number of i -bad sequences. Our analysis above gives

$$\mathbb{E}[B_i] \leq 2N^s \cdot O_r(q^{-r}) = O_r(q^{s^2-r}) = o_r(1). \quad (10)$$

Now let $G' \subseteq G$ be defined by deleting a vertex from each bad sequence. Because each vertex is in at most $N = q^s$ edges in G , by (??) and (10) we find

$$\mathbb{E}[e(G')] \geq \mathbb{E}[e(G)] - \mathbb{E}[B] \cdot q^s = \Omega(q^{2s-1}),$$

where this last step used the previous inequality and Lemma 8.4 to deduce $\mathbb{E}[e(G)] = q^{2s} \cdot q^{-1}$. By definition G' contains no copy of $K_{s,c}$, so for $t \geq t_0 := c$, we have shown that there exists a graph G' on at most q^{2s} vertices such that it contains at least $\Omega(q^{2s-1})$ edges and no copy of $K_{s,t}$. This gives the desired lower bound when n is a sufficiently large prime power. Using Bertrand's postulate gives the desired bound for all n . \square

We will admit that on its own Theorem 8.6 is not particularly groundbreaking. Indeed, there exist explicit constructions showing $\text{ex}(n, K_{s,t}) = \Theta(n^{2-1/s})$ when $s > (t-1)!$, and this dependency on t is far better than the implicit constant t_0 one gets from the proof. However, the proof of Theorem 8.6 gives us a new idea for constructing F -free graphs.

And indeed, many developments have been made on this method since Bukh's original construction of this form. In particular, by using more sophisticated tools from algebraic geometry, Bukh [21] showed that Theorem 8.6 holds with $t_0 = C^s$ for some absolute constant C , which stands as the best known bounds for this problem. On the other hand, by carefully modifying the current proof of Theorem 8.6, Conlon [26] was able to construct large graphs avoiding theta graphs $\theta_{a,b}$, which we recall denotes the graph consisting of a internally disjoint paths of length b between two fixed vertices.

Theorem 8.7 ([26]). *For all $b \geq 2$, there exists some $a_0 = a_0(b)$ such that for all $a \geq a_0$, we have*

$$\text{ex}(n, \theta_{a,b}) = \Theta(n^{1+1/b}).$$

Proof Sketch. The upper bound is a result of Faudree and Simonovits [43]. For the lower bound, the key idea is to consider *multiple* random polynomials $f_1, \dots, f_a \in \mathcal{P}_{d,2b}$ chosen independently, with us defining our graph G on $\mathbb{F}_q^b \cup \mathbb{F}_q^b$ by having $x \sim y$ if and only if $f_i(x, y) = 0$ for all i .

Fix two vertices $x^1, x^{b+1} \in \mathbb{F}_q^b \cup \mathbb{F}_q^b$ and define C to be the tuples of distinct vertices (x^2, \dots, x^{b-1}) such that $x^1 \dots x^{b+1}$ is a path in G . As before (but with a somewhat more difficult analysis), one can show $\mathbb{E}[|C|^r] = O(1)$. However, in this case C is not quite an algebraic variety because of us requiring C to use distinct vertices. However, one can write C as $X \setminus D$ for two varieties X, D , so Lemma 8.5 still applies and the rest of the proof goes through. \square

One can further interpolate between this theorem and Theorem 8.6 to give effective lower bounds on $\text{ex}(n, F)$ whenever F is a “large power of a rooted tree” (e.g. $K_{s,t}$ is just many copies of a star $K_{s,1}$, and $\theta_{a,b}$ is just many copies of a path P_b). This was done by Bukh and Conlon [22] in order to show that for every rational number $r \in [1, 2]$, there exists a finite set of graphs \mathcal{F} such that $\text{ex}(n, \mathcal{F}) = \Theta(n^r)$. The rational exponents conjecture, which says that one can achieve this with \mathcal{F} consisting of a single graph, remains a major open problem.

8.3 Multicolor Ramsey Numbers

Let $r(t; \ell)$ denote the smallest number N such that every ℓ -coloring of $E(K_N)$ contains a monochromatic clique of size t . **Inset history and connection with the earlier Ramsey results proven in the text. Also sketch the proof of the bound you get with the naive method for comparison.**

The following observation will be the key towards going further. The initial idea for this lemma can be seen in Conlon and Ferber [27], though it was first really used by Wigderson [109] and then generalized by Sawin [99].

Lemma 8.8. *Let G be graph with no clique of size t , and let p be the probability that vertices $v_1, \dots, v_t \in V(G)$ chosen independently and uniformly at random form an independent set. Then for all $\ell \geq 2$, we have*

$$r(t; \ell) \geq p^{-(\ell-2)/t} 2^{(t-1)/2}.$$

Note that when $\ell = 2$ this recovers the usual lower bound for Ramsey numbers from the random coloring.

Proof. Let N be an integer to be determined later, and let $f_1, \dots, f_{\ell-2} : V(K_N) \rightarrow V(G)$ be chosen independently and uniformly at random. Define a coloring $\chi : E(K_N) \rightarrow [\ell]$ in the following way: for distinct $x, y \in V(K_N)$, if there exists i such that $f_i(x)f_i(y) \in E(G)$, then set $\chi(xy)$ to be the minimum i with this property. Otherwise, set $\chi(xy)$ to be $\ell - 1$ or ℓ with probability $1/2$ each. That is (as Wigderson notes in his paper), this coloring comes from covering K_N with $\ell - 2$ randomly permuted blowups of G and then randomly using two colors to deal with any uncovered vertices.

We first observe that there is no monochromatic K_t in any color $i \leq \ell - 2$. Indeed, if $\{x_1, \dots, x_t\}$ were such a clique then this would imply $\{f_i(x_1), \dots, f_i(x_t)\}$ forms a clique in G (since $\chi(x_j x_k) = i$ implies $f_i(x_j)f_i(x_k) \in E(G)$). Thus it remains to show that, with positive probability, there is no monochromatic K_t in color $i \in \{\ell - 1, \ell\}$. Observe that a clique K_t in K_N has all of its edges colored by $\ell - 1$ or ℓ if and only if each f_i maps K_t to an independent set of G , and the probability that this happens is exactly $p^{\ell-2}$ by hypothesis, and from there this K_t will be monochromatic with probability $2^{1-\binom{t}{2}}$. In total then, the expected number of monochromatic cliques will equal $\binom{N}{t} p^{\ell-2} 2^{1-\binom{t}{2}}$, and this will be less than 1 provided $N \leq p^{-(\ell-2)/t} 2^{(t-1)/2}$. Thus there exists a coloring of this size with no monochromatic clique, giving the desired result. \square

Observe that the p in Lemma 8.8 roughly corresponds to the number of independent sets of size at most t in G , so we need to find a graph with small clique number and not too many small independent sets. To this end, let $V \subseteq \mathbb{F}_2^t$ be the set of vectors v with $v \cdot v = 0$ (i.e. vectors with even Hamming weight), and let G be the graph on V where two vectors u, v are adjacent if and only if $u \cdot v = 1$.

Lemma 8.9. *If t is even, then the graph G contains no clique of size t .*

Proof. Assume for contradiction that there exist distinct vectors $v_1, \dots, v_t \in V$ with $v_i \cdot v_j = 1$ for all $i \neq j$ (and $= 0$ for $i = j$ by definition of V). We claim that these vectors are linearly independent. Indeed, if there exists $\alpha_i \in \{0, 1\}$ with $\sum \alpha_i v_i = 0$, then by taking the dot product of v_j on both sides we find $\sum_{i \neq j} \alpha_i \equiv 0$ for all j , and it is not difficult to show that this implies $\alpha_i = 0$ for all i (here we need that t is even, else $\alpha_i = 1$ for all i would work). However, V is a $t - 1$ dimensional subspace, so it contains no set of t linearly independent vectors, proving the result. \square

Lemma 8.10. *The probability p that a uniformly random tuple $(v_1, \dots, v_t) \in V^t$ is such that $\{v_1, \dots, v_t\}$ is independent in G is at most $2^{-3t^2/8 + o(t^2)}$.*

Proof. Let X be the set of tuples $(v_1, \dots, v_t) \in V^t$ such that $v_i \cdots v_j = 0$ for all i, j , so our goal is to upper bound $|X|/|V|^t = |X|2^{-t^2}$. Define the rank of a tuple in X to be the rank of the smallest subspaces containing every vertex of the tuple. We claim that the number of tuples in X of rank r is at most

$$t! \left(\prod_{i=0}^{r-1} 2^{t-i} \right) \cdot 2^{(t-r)r} = 2^{tr - \binom{r}{2} + tr - r^2}. \quad (11)$$

Indeed, possibly by reordering the tuple (giving us the factor of $t!$) we can assume the first r vectors are linearly independent, and given v_1, \dots, v_i with $0 \leq i < r$, the number of choices for a v_{i+1} which is linearly independent of v_1, \dots, v_i is exactly q^{t-i} . After this every vector must lie in the span of v_1, \dots, v_r , giving exactly q^r choices for the remaining $t - r$ vectors.

We next claim that there exists no tuple in X of dimension larger than $t/2$. Indeed, if S is the span of the vectors in a tuple of X , then note that $S \subseteq S^\perp$ since $v_i \cdot v_j = 0$ for all i, j . From linear algebra we have $t = \dim S + \dim S^\perp \geq 2 \dim S$, proving the claim.

It is not hard to prove that (11) is increasing for $r \leq t/2$, so plugging in $r = t/2$ gives an upper bound for $|X|/(t/2)$ of the form $2^{5t^2/8 + o(t^2)}$, giving the desired bound on $|X|/|V|^t$. \square

Putting all these lemmas together gives the following.

Corollary 8.11. *For $\ell \geq 3$ we have*

$$r(t; \ell) \geq \left(2^{\frac{3\ell}{8} - \frac{1}{4}} \right)^{t - o(t)}.$$

This bound stood as the best for about a year until Sawin [99] realized one could do somewhat better by replacing the algebraic graph G described above with a purely random graph, namely $G_{n,p}$ with $p \approx .455$. Thus, although the initial breakthrough for multicolor Ramsey numbers came from a random algebraic approach, the method was later subsumed by a simpler random model. This sort of thing happens somewhat often with proofs using the random algebraic method. Because of this, some mathematicians are of the opinion that any time the random algebraic method is used, there exists a simpler random model which gives better results. I don't personally believe that this is true, and even if it were, the fact that random algebraic methods consistently give initial breakthroughs to longstanding open problems makes them worth considering in my eyes.

Maybe comment on other ways random homomorphisms are useful.

Part III

Spread Hypergraphs

9 The Spreadness Theorem

Throughout this section we consider hypergraphs \mathcal{H} which may have repeated edges, and we will typically denote the edges of \mathcal{H} by S . We recall that $d(A)$ denotes the degree of a set of vertices A in \mathcal{H} , i.e. the number of edges of \mathcal{H} containing A .

We say that a hypergraph \mathcal{H} is r -bounded if all of its edges have size at most r . We say that a hypergraph \mathcal{H} is q -spread¹ for some $0 < q < 1$ if \mathcal{H} is non-empty and if $d(A) \leq q^{|A|} |\mathcal{H}|$ for all sets of vertices A . The main result for q -spread hypergraphs is the following.

Theorem 9.1 ([7, 49]). *Let \mathcal{H} be an r -bounded q -spread hypergraph on V . There exists an absolute constant K_0 such that if W is a set of size $Cq \log r \cdot |V|$ chosen uniformly at random from V with $C \geq K_0$, then*

$$\Pr[W \text{ contains an edge of } \mathcal{H}] \geq 1 - 8C^{-1}.$$

We note that better quantitative versions of Theorem 9.1 exist, see e.g. [Tao's reformulation below?](#), but as stated this theorem already does a lot. Let's start by looking at some applications before turning it's short (though very dense!) proof.

9.1 Applications

Our first application is the following.

Theorem 9.2. *Let $G_{n,m}^r$ be the r -graph chosen uniformly at random amongst all r -graphs with n vertices and m edges. Then there exists a constant C such that if $m \geq Cn \log n$ and n is a multiple of r , then $G_{n,m}^r$ contains a perfect matching a.a.s.²*

It is not too difficult to show that this bound on m is essentially best possible. We note that morally speaking, $G_{n,m}^2$ acts the same as $G_{n,p}$ where $p = m/\binom{n}{2}$. In particular, one can use Theorem 9.2 to prove that $G_{n,p}$ contains a perfect matching a.a.s. if $p = \Omega(\log n/n)$. Proving Theorem 9.2 for $r = 2$ is not hard, but the result for general r was thought to be very difficult, with its first proof due to Johansson, Kahn, and Vu [68] using a rather involved argument. We will prove Theorem 9.2 in just a few lines with Theorem 9.1.

Proof. Let \mathcal{H} be the hypergraph on $E(K_n^r)$ where each hyperedge S is a perfect matching of

¹Some texts would say that such an \mathcal{H} is q^{-1} -spread.

²This means “asymptotically almost surely”, i.e. the probability of this event happening tends to 1 as n tends to infinity.

K_n^r . Observe that for any set $A \subseteq E(K_n^r)$, we have

$$\begin{aligned}
d(A) \cdot |\mathcal{H}|^{-1} &= \frac{(n - r|A|)!}{(r!)^{n/r - |A|} (n/r - |A|)!} \cdot \frac{(r!)^{n/r} (n/r)!}{n!} \\
&= (r!)^{|A|} \binom{n/r}{|A|} \binom{n}{r|A|}^{-1} \frac{|A|!}{(r|A|)!} \\
&\leq (r!)^{|A|} (en/r|A|)^{|A|} \cdot (n/r|A|)^{-r|A|} \cdot (|A|)^{|A|} \cdot (r|A|/e)^{-r|A|} \\
&= (r!)^{|A|} e^{(r+1)|A|} n^{-(r-1)|A|} \leq (n/re^3)^{-(r-1)|A|}.
\end{aligned}$$

Thus \mathcal{H} is $(n/re^3)^{-r+1}$ -spread. It is also (n/r) -uniform and has a ground set $V = E(K_n^r)$ of size $\binom{n}{r}$. By Theorem 9.1, we see that if m is at least as large as in our hypothesis, then with high probability a random m -subset of \mathcal{H} will contain a hyperedge, i.e. $H_{n,m}^r$ will contain a perfect matching with high probability. \square

Another basic example is the following.

Proposition 9.3. *Let F be an r -graph and define $t(F) = \max\{|E(F')|/|V(F')| : F' \subseteq F\}$. Let $G_{n,m}^r$ be as in Theorem 9.2. There exists a constant $C(F)$ such that if $m \geq C(F)n^{r-1/t(F)}$, then $G_{n,m}^r$ contains a copy of F a.a.s.*

A simple first moment argument shows that this bound is tight. One can prove Proposition 9.3 using a standard but somewhat tedious second moment argument, but using Theorem 9.1 gives a shorter proof.

Proof. Let \mathcal{H} be the hypergraph on $E(K_n^r)$ whose hyperedges correspond to copies of F . Observe that \mathcal{H} being q -spread is equivalent to having $(d(A)/|\mathcal{H}|)^{1/|A|} \leq q$ for all $A \subseteq V = E(K_n^r)$. Any set $A \subseteq E(K_n^r)$ of positive degree in \mathcal{H} forms a subgraph $F' \subseteq F$ with $|E(F')| = |A|$, and in this case

$$\left(\frac{d(A)}{|\mathcal{H}|} \right)^{1/|A|} \leq \left(\frac{n^{|V(F)| - |V(F')|}}{\binom{n}{|V(F')|}} \right)^{1/|A|} \leq |V(F)|^{|V(F')|} \cdot n^{-|V(F')|/|E(F')|}.$$

Thus we see that \mathcal{H} is q -spread with

$$q = \max\{|V(F)|^{|V(F')|} \cdot n^{-|V(F')|/|E(F')|} : F' \subseteq F\} = |V(F)|^{|V(F)|} \cdot n^{-1/t(F)}.$$

Plugging this into Theorem 9.1 gives the result. \square

The study of q -spread hypergraphs was initiated by Alweiss, Lovett, Wu, and Zhang [7] where they proved a slightly weaker version of Theorem 9.1. Their motivation came from the Erdős sunflower conjecture. A k -sunflower is a hypergraph with edges S_1, \dots, S_k such that there exists a set K called the *kernel* which has $S_i \cap S_j = K$ for all $i \neq j$.

Theorem 9.4. *There exists an absolute constant $C > 0$ such that if \mathcal{H} is an r -graph with at least $(Ck \log r)^r$ edges, then \mathcal{H} contains a k -sunflower.*

We note that [7] was the first to prove bounds of the form $(\log r)^{r+o(1)}$ for fixed k , with [92, 14] later giving better bounds in terms of k . Prior to [7], the best known bounds were of the form $r^{r-o(1)}$. It is a famous conjecture of Erdős that one can prove a bound of the form $c_k^{r+o(1)}$.

Proof. We prove the result by induction on r , the $r = 1$ case being trivial. Let \mathcal{H} be an r -graph with at least $(Ck \log r)^r$ edges. If \mathcal{H} is not q -spread with $q = (Ck \log r)^{-1}$, then there exists some $A \subseteq V(H)$ such that $d(A) \geq (Ck \log r)^{r-|A|}$. This means that the link hypergraph $\mathcal{H}_A = \{S \setminus A : S \in \mathcal{H}, A \subseteq S\}$ has size at least $(Ck \log r)^{r-|A|}$. Since \mathcal{H}_A is an $(r - |A|)$ -uniform hypergraph, by induction \mathcal{H}_A contains a k -sunflower, say with edges $S_1 \setminus A, \dots, S_k \setminus A \in \mathcal{H}_A$. It is not difficult to check that $S_1, \dots, S_k \in \mathcal{H}$ forms a k -sunflower in \mathcal{H} . We conclude that any \mathcal{H} with at least $(Ck \log r)^r$ edges which is not q -spread contains a k -sunflower, so from now on we may assume \mathcal{H} is q -spread.

Possibly by adding isolated vertices to \mathcal{H} , we can assume that the size of the vertex set V of \mathcal{H} is a multiple of $2k$. Let V_1, \dots, V_{2k} be a random partition of V such that each $V_i \subseteq V$ has size $(2k)^{-1}|V|$. This means that each V_i is a uniformly chosen set of V of size $(2k)^{-1}|V| = \frac{1}{2}C(\log r)q|V|$. Let 1_i be the indicator variable for the event that V_i contains an edge of \mathcal{H} . By Theorem 9.1, we have $\Pr[1_i = 1] \geq \frac{1}{2}$ provided C is sufficiently large. In this case, $\mathbb{E}[\sum 1_i] \geq k$, and hence there exists some partition V_1, \dots, V_{2k} such that $\sum 1_i \geq k$, which in particular means there exist k disjoint edges of \mathcal{H} . This is a k -sunflower in \mathcal{H} , proving the result. \square

9.2 Proof of the Spreadness Theorem

A maybe simpler proof of this/Park-Pham can be found in Rao's talk on "The Sunflower Lemma and Monotone Thresholds", see also the link commented out below

There are by now a number of proofs of Theorem 9.1, though most of them maintain the same core set of ideas. The proof we present here is based off of a proof due to Rao [93] which gives weaker quantitative bounds. We emphasize that while the proof itself is very short, it is also very dense in content, so we'll spend some time trying to build up some intuition for it.

Recall that \mathcal{H} is an r -bounded q -spread hypergraph on V , and that we want to show that a uniformly random set W of size $Cq \log r \cdot |V|$ contains an edge of \mathcal{H} with high probability. In order to use an iterative approach, we consider a uniform random vector of disjoint sets $(W_1, \dots, W_{\log r})$ each of size $Cq|V|$. Let $W_{\leq i} = \bigcup_{j \leq i} W_j$, and note that W has the same distribution as $W_{\log r}$, so it suffices to work with these random sets.

A super ideal situation for our iterative approach would be if for all $S \in \mathcal{H}$, we have $|S - W_{\leq i}| < 2^{-i}r$. Indeed, with this at $i = \log r$, we would get that every edge is contained in $W_{\leq \log r}$. Of course, this is far too much to hope for. However, since we only need $W_{\leq \log r}$ to contain a single edge, it would suffice to have this work out for *some* S . As such it perhaps make sense to say that an edge S "succeeds" at step i if $|S - W_{\leq i}| < 2^{-i}r$, and then to argue that with high probability some edge succeeds at each step. Unfortunately this notion of success is too restrictive to work. The key insight is that we can loosen our condition by saying that an edge S "succeeds" if there exists some edge $S' \subseteq S \cup W_{\leq i}$ (or equivalently $S' - W_{\leq i} \subseteq S - W_{\leq i}$) such that $|S' - W_{\leq i}| < 2^{-i}r$. The point is that (1) this condition is easier to achieve (and in particular will be achieved by "most" S at each step), and (2) if some S succeeds at each step, then the S' it points to for step $\log r$ will be contained in $W_{\leq \log r}$. **Need to fact check that this is really what's going on, and in particular we maybe need S' also to have not failed at this point.**

With this in mind, given W_1, \dots, W_i , we iteratively define "failure" hypergraph \mathcal{F}_i to be those

$S \notin \mathcal{F}_{\leq i-1} := \bigcup_{j \leq i-1} \mathcal{F}_j$ (i.e. which haven't failed at any previous step) such that for all $S' \in \mathcal{H} - \mathcal{F}_{\leq i-1}$ satisfying $S' \subseteq S \cup W_{\leq i}$, we have $|S' - W_{\leq i}| \geq 2^{-i}r$. The key claim is the following.

Lemma 9.5. *Given $W_{\leq i-1}$, we have $\mathbb{E}[|\mathcal{F}_i|] \leq 2(C/4)^{-2^{-i}r} |\mathcal{H}|$.*

To upper bound $|\mathcal{F}_i|$, it will help to instead upper bound the size of an auxiliary hypergraph defined as follows. Given W_1, \dots, W_i , we define the *fragment* $T(S, W_{\leq i})$ of an edge $S \in \mathcal{H} - \mathcal{F}_{i-1}$ to be a set of minimum size in $\{S' - W_{\leq i} : S' \in \mathcal{H} - \mathcal{F}_{\leq i-1}, S' \subseteq S \cup W_{\leq i}\}$ (say the lexicographically smallest set if there are multiple of minimum size). We let \mathcal{G}_i be the hypergraph where T is an edge if $T = T(S, W_{\leq i})$ for some $S \in \mathcal{F}_i$. Note that by definition this means $|T| \geq 2^{-i}r$, and that for every $S \in \mathcal{F}_i$, there exists some $T \in \mathcal{G}_i$ with $T \subseteq S$. This last condition says \mathcal{G}_i is an *undercover* of \mathcal{F}_i , which will also be a key condition in our upcoming proof of the Park-Pham Theorem.

Proof. Let $w := Cq|V|$, which we recall is the size of W_i , and let $n_i = |V - W_{\leq i-1}|$. Let \mathcal{P} consist of all pairs (S, W) with $S \in \mathcal{H}$ and $W \in \binom{V - W_{\leq i-1}}{w}$ such that $S \in \mathcal{F}_i$ whenever $W_i = W$. Similarly given an integer $a \geq 2^{-i}r$, let \mathcal{P}_a consist of all pairs (T, W) with $|T| = a$ and $W \in \binom{V - W_{\leq i-1}}{w}$ such that $T \in \mathcal{G}_i$ whenever $W_i = W$. We claim that. **Probably use t instead of a**

$$\mathbb{E}[|\mathcal{F}_i|] = |\mathcal{P}| \binom{n_i}{w}^{-1} \leq \sum_{a \geq 2^{-i}r} q^a |\mathcal{H}| |\mathcal{P}_a| \binom{n_i}{w}^{-1}. \quad (12)$$

Indeed, the equality is straightforward. Because \mathcal{G}_i is an undercover of \mathcal{F}_i , for every pair $(S, W) \in \mathcal{P}$ there exists a pair $(T, W) \in \bigcup_a \mathcal{P}_a$ such that $T \subseteq S$. Moreover, for each set T of size a , the number of $S \in \mathcal{H}$ with $T \subseteq S$ is at most $q^a |\mathcal{H}|$ by the definition of \mathcal{H} being q -spread. This gives the stated inequality

It remains to count the number of elements $(T, W) \in \mathcal{P}_a$. We will identify such a pair by first specifying the set $T \cup W$, and then specifying T (which uniquely determines W). We first note that $T \cup W$ is a set of size $a + w$, so the number of choices for this step is at most

$$\binom{n_i}{a+w} \leq (n_i/w)^a \cdot \binom{n_i}{w} = (Cq)^{-a} \binom{n_i}{w}.$$

Given $T \cup W$, choose any $S' \in \mathcal{H} - \mathcal{F}_{\leq i-1}$ with $S' - W_{\leq i-1} \subseteq T \cup W$. Crucially, we must have $T \subseteq S' - W_{\leq i-1}$, as otherwise if $T = T(S, W_{\leq i-1} \cup W)$ for some S , then taking $T' = S' - (W_{\leq i-1} \cup W) \subsetneq T$ (with the inclusion holding because $S' - W_{\leq i-1} \subseteq T \cup W$, and the strictness holding if $T \not\subseteq S' - W_{\leq i-1}$), we find that T cannot be the fragment of S (since T' is a smaller set than T satisfying the same properties). Note that $S' \notin \mathcal{F}_{\leq i-1}$ implies $|S' - W_{\leq i-1}| \leq 2^{-i+1}r$, so the number of choices for T is at most $2^{2^{-i+1}r} = 4^{2^{-i}r}$.

In total we conclude that $|\mathcal{P}_a| \leq (Cq)^{-a} 4^{2^{-i}r} \binom{n_i}{w}$. Plugging this into (12), we find

$$\mathbb{E}[|\mathcal{F}_i|] \leq \sum_{a \geq 2^{-i}r} q^a |\mathcal{H}| \cdot (Cq)^{-a} 4^{2^{-i}r} = 4^{2^{-i}r} |\mathcal{H}| \sum_{a \geq 2^{-i}r} C^{-a} \leq 2(C/4)^{-2^{-i}r} |\mathcal{H}|,$$

with this last step holding for C sufficiently small. \square

With this lemma, we have

$$\mathbb{E}[|\mathcal{F}_{\leq \log r}|] \leq \sum_{i=1}^r 2(C/4)^{-2^{-i}r} |\mathcal{H}| \leq 16C |\mathcal{H}|.$$

By Markov, the probability that $|\mathcal{F}_{\leq \log r}| = |\mathcal{H}|$ is at most $\frac{1}{16C}$, so $\mathcal{F}_{\leq \log r} \neq \mathcal{H}$ is at least $1 - \frac{1}{16C}$. As noted above, if there exists $S \in \mathcal{H} - \mathcal{F}_{\leq \log r}$ then S points to an edge which is contained in $W_{\leq \log r}$, so we conclude that this random set $W_{\leq \log r}$ of size $Cq \log(r)|V|$ contains an edge with probability at least $1 - \frac{1}{16C}$.

9.3 Losing Logarithms

As we noted earlier, the bound of Theorem 9.2 is best possible. In particular, the $\log r$ term of Theorem 9.1 is necessary in general. However, under certain conditions one can remove this logarithmic term. This was first observed by Kahn, Narayanan, and Park [70] where they found tight bounds on the threshold of a square of a Hamiltonian cycle in $G_{n,p}$. Here we briefly outline how, under special circumstances, one can modify the previous proof to get rid of the $\log r$ factor.

The main idea is that instead of setting our cutoff points for our fragments to be $r/2, r/4, \dots$, we instead set them to be k_1, k_2, \dots for some suitable sequence k_i with significantly fewer than $\log r$ terms. In this setup, one could try to naively go through Lemma 9.5 and replace $2^{-i+1}r$ with k_{i-1} and $2^{-i}r$ with k_i , which will roughly give us

$$\mathbb{E}[|\mathcal{F}_i|] \leq 2^{k_{i-1}} C^{-k_i} |\mathcal{H}|,$$

but this will be terrible unless k_{i-1} differs from k_i by a multiplicative constant depending on C .

One way we can get around this is if we impose that for all sets A and integers j with $k_{i-1} \geq |A| \geq j \geq k_i$, we have that the number of edges S' with $|A \cap S'| = j$ is at most $q^j |\mathcal{H}|$. Note that this is stronger than spreadness since, when taking $j = |A|$, the condition $|A \cap S'| = j$ just says S' is an edge containing A , so this bound exactly says $\deg_{\mathcal{H}}(A) \leq q^{|A|} |\mathcal{H}|$, which is the spreadness condition. Assuming this condition holds, we will count the pairs $(S, W) \in \mathcal{P}$ in a more subtle way.

Let $T = T(S, W_{\leq i-1} \cup W)$, which we note has $|T| \geq k_i$ and $T \subseteq S$. We first specify $T \cup W$, which as before can be done in roughly $(Cq)^{-a} \binom{n_i}{w}$ ways. We then pick some edge $S' \notin \mathcal{F}_{\leq i-1}$ such that $A := S' - W_{\leq i-1} \subseteq T \cup W$, where as before we have $T \subseteq A$. Since $T \subseteq A \cap S$, we have $j := |A \cap S| \geq a$. Given j , the number of S with $|A \cap S| = j$ is at most $q^j |\mathcal{H}|$ by our condition (and we have $|A| \leq k_{i-1}$ as otherwise we would have $S' \in \mathcal{F}_{\leq i-1}$). Since we now know S and $T \cup W$, we also know $S \cup W$ (since $T \subseteq S$), and hence T (since T is purely a function of the set $S \cup W$ and $W_{\leq i-1}$ **Need to double check this; in any case there are trivially at most 2^j choices for $T \subseteq A \cap S$**), and hence $W = (T \cup W) \setminus T$ (since T is disjoint from W by definition). With this, we see that the total number of choices is at most

$$\sum_{a \geq k_i} C^{-a} \binom{n_i}{w} \sum_{j \geq a} q^j |\mathcal{H}| \approx C^{-k_i} \binom{n_i}{w} |\mathcal{H}|,$$

giving the desired result. A more formal theorem/proof can be found in [105], though the approach used there is an older and more complicated version of the one presented here. See also [51] for a proof in the specific case of getting rid of the logarithm for the square of a Hamiltonian cycle.

10 **TODO**The Park-Pham Theorem

11 Spread Approximations

Motivated by our success in proving bounds on hypergraphs which don't contain sunflowers in Theorem 9.4, we consider some additional applications of spread hypergraphs to extremal set theory. For this it will be convenient to define a slightly different notion of spreadness which was introduced by Kupavskii and Zakharov [77].

Given a set of vertices A , we define the link hypergraph $\mathcal{H}(A) = \{e \setminus A : e \in \mathcal{H}, A \subseteq e\}$. We say that an n -vertex r -graph \mathcal{H} is τ -homogeneous if

$$d(A) = |\mathcal{H}(A)| \leq \tau^{|A|} \frac{\binom{n-|A|}{r-|A|}}{\binom{n}{r}} |\mathcal{H}|.$$

Intuitively, τ -homogeneous hypergraphs should be thought of as (τ/n) -spread hypergraphs. Formally, we have the following.

Lemma 11.1. *If \mathcal{H} is an n -vertex r -graph which is τ -homogeneous, then it is q -spread with $q = \frac{\tau r}{n}$.*

Proof. Note that $\binom{n-1}{r-1} = \frac{r}{n} \binom{n}{r}$. By repeating this logic and using that $\frac{r-i}{n-i} \leq \frac{r}{n}$ whenever $i \leq r \leq n$, we find

$$d(A) \leq \tau^{|A|} \frac{\binom{n-|A|}{r-|A|}}{\binom{n}{r}} |\mathcal{H}| \leq \left(\frac{\tau r}{n}\right)^{|A|} |\mathcal{H}|,$$

proving the result. □

The main motivation for this definition is the following.

Theorem 11.2 ([77]). *Fix $k \in \mathbb{Z}_{\geq 1}$ and $\tau \in \mathbb{R}_{\geq 1}$. For every n -vertex r -graph \mathcal{H} , there exists an “approximation” hypergraph \mathcal{S} with edges of size at most k , and a “remainder” $\mathcal{H}' \subseteq \mathcal{H}$ with the following properties:*

- *For every $e \in \mathcal{H} \setminus \mathcal{H}'$, there exists $S \in \mathcal{S}$ such that $S \subseteq e$.*
- *$|\mathcal{H}'| \leq \tau^{-k-1} \binom{n}{r}$.*
- *For every $S \in \mathcal{S}$ there exists $\mathcal{H}_S \subseteq \mathcal{H}$ such that the link hypergraph $\mathcal{H}_S(S)$ is τ -homogeneous.*

This first condition says that \mathcal{S} “approximates” $\mathcal{H} \setminus \mathcal{H}'$ in the sense that it's an undercover, while the second condition says that “most” of \mathcal{H} is approximated. The third condition gives us the very useful property that each edge S of our approximating hypergraph \mathcal{S} has a well spread link, which will imply that a random partition of our vertex set will give a sunflower whose kernel contains S .

Add more intuition for this proof, possibly via recalling the proof of the improved sunflower bounds (namely you go by picking a largest S such that $\mathcal{F}(S)$ isn't spread).

Proof. If $n \leq k$, then we can take $\mathcal{S} = \mathcal{H}$, $\mathcal{H}_S = \{S\}$, and $\mathcal{H}' = \emptyset$ to get the result, so we may assume $n > k$. For this proof we define $\mathcal{H}[S] = \{e : e \in \mathcal{H}, S \subseteq e\}$, i.e. this is just the link hypergraph $\mathcal{H}(S)$ after adding the set S back into each edge.

Let $\mathcal{H}^1 = \mathcal{H}$. Given \mathcal{H}^i , let S_i be a maximal set of vertices with $d_{\mathcal{H}^i}(S_i) \geq \tau^{|S_i|} \frac{\binom{n-|S_i|}{r-|S_i|}}{\binom{n}{r}} |\mathcal{H}^i|$. If $|S_i| > k$ (which happens automatically if $\mathcal{H}^i = \emptyset$ and $n > k$), then stop the procedure. Otherwise set $\mathcal{H}^{i+1} = \mathcal{H}^i \setminus \mathcal{H}^i[S_i]$.

Say we stop this procedure at $i = m+1$. Set $\mathcal{H}' = \mathcal{H}^{m+1}$, $\mathcal{S} = \{S_i : i \leq m\}$, and $\mathcal{H}_{S_i} = \mathcal{H}^i[S_i]$. Observe that by construction each edge of \mathcal{S} has size at most k and that the first condition of the theorem is satisfied (since $e \in \mathcal{H} \setminus \mathcal{H}'$ implies $e \in \mathcal{H}^i$ for some $i \leq m$, and taking $S = S_i$ works). Again by construction, we have

$$|\mathcal{H}'| = |\mathcal{H}^{m+1}| \leq d_{\mathcal{H}^{m+1}}(S_{m+1}) \tau^{-|S_{m+1}|} \frac{\binom{n}{r}}{\binom{n-|S_{m+1}|}{r-|S_{m+1}|}} \leq \tau^{-k-1} \binom{n}{r},$$

where this last step used the trivial bound $d_{\mathcal{H}^{m+1}}(S_{m+1}) \leq \binom{n-|S_{m+1}|}{r-|S_{m+1}|}$ and that $|S_{m+1}| \geq k+1$ by assumption of us stopping the procedure here. This establishes the second condition.

For the third condition, note that for any set T disjoint from S_i , we have

$$d_{\mathcal{H}_{S_i}}(T) = d_{\mathcal{H}^i}(S \cup T) < \tau^{|S_i \cup T|} \frac{\binom{n-|S_i \cup T|}{r-|S_i \cup T|}}{\binom{n}{r}} |\mathcal{H}^i|,$$

where this last step used the maximality of S_i . We also have

$$|\mathcal{H}^i| \leq \tau^{-|S_i|} \frac{\binom{n}{r}}{\binom{n-|S_i|}{r-|S_i|}} d_{\mathcal{H}^i}(S_i) = \tau^{-|S_i|} \frac{\binom{n}{r}}{\binom{n-|S_i|}{r-|S_i|}} |\mathcal{H}_{S_i}|.$$

Combining these two inequalities gives the third condition, proving the result. \square

It turns out that when τ is small, the \mathcal{S} given by Theorem 11.2 inherits intersection properties of the original hypergraph \mathcal{H} . Recall that a hypergraph is t -intersecting if every two edges intersect in at least t vertices.

Lemma 11.3. *There exists an absolute constant C such that the following holds. Let \mathcal{H} be an n -vertex r -graph, and let \mathcal{S} be the hypergraph guaranteed by Theorem 11.2 with parameters k, τ . If $n \geq C\tau r \max\{\log r, k\}$, then*

$$\{S \cap T : S, T \in \mathcal{S}\} \subseteq \{e \cap f : e, f \in \mathcal{H}\}.$$

We emphasize that we allow $S = T$ in the lemma statement.

Proof. Let $S, T \in \mathcal{S}$, and note that Lemma 11.1 implies $\mathcal{H}_S(S)$ is $\frac{\tau r}{n}$ -spread. Let $\mathcal{H}'_S \subseteq \mathcal{H}_S(S)$ be the hypergraph obtained by deleting all of the vertices $x \in T \setminus S$. Then

$$|\mathcal{H}'_S| \geq |\mathcal{H}_S(S)| - \sum_{x \in T \setminus S} d_{\mathcal{H}_S(S)}(x) \geq (1 - \frac{\tau r}{n} \cdot k) |\mathcal{H}_S(S)| \geq \frac{1}{2} |\mathcal{H}_S(S)|,$$

where the second inequality used that $\mathcal{H}_S(S)$ is spread, and the last inequality holds for $C \geq 2$. This together with $\mathcal{H}_S(S)$ being $\frac{\tau r}{n}$ -spread implies \mathcal{H}'_S is $q := \frac{2\tau r}{n}$ -spread. Similarly if one defines $\mathcal{H}'_T \subseteq \mathcal{H}_T(T)$ by deleting the vertices of $S \setminus T$ we get that this is q -spread.

Randomly partition the vertices of $V(\mathcal{H}) \setminus (S \cup T)$ into two sets V_1, V_2 of size at least $\frac{1}{2}(n - 2k) \geq \frac{1}{4}n$, where this holds if C is sufficiently large. By hypothesis this is at least $\frac{1}{4}C\tau r \log r = \frac{1}{8}Cq \log rn$. Thus by Theorem 9.1, if C is sufficiently large, then with positive probability both V_1, V_2 contain edges e', f' of $\mathcal{H}'_S, \mathcal{H}'_T$ respectively, which by definition means e', f' contains no vertices of T, S . This means $e = e' \cup S, f = f' \cup T$ are edges of \mathcal{H} with $e \cap f = S \cap T$, proving the result. \square

A simple application of this result gives the t -intersecting version of the Erdős-Ko-Rado theorem (albeit with suboptimal bounds on n).

Theorem 11.4. *Let \mathcal{H} be an n -vertex r -graph such that $|e \cap f| \geq t$ for all $e, f \in \mathcal{H}$. If n is sufficiently large in terms of r , then $|\mathcal{H}| \leq \binom{n-t}{r-t}$ with equality holding if and only if \mathcal{H} consists of every edge containing some fixed set T of size t .*

Proof. Apply Theorem 11.2 with $k = t$ and $\tau = \frac{n}{Cr \max\{\log r, k\}}$ with C as in Lemma 11.3, and let $\mathcal{H}', \mathcal{S}$ be the resulting families. Note that by Theorem 11.2,

$$|\mathcal{H}'| \leq \tau^{-k-1} \binom{n}{r} = O(n^{r-t-1}).$$

By Lemma 11.3, our hypothesis on \mathcal{H} , and the fact that $|S| \leq k = t$ for all $S \in \mathcal{S}$, we see that \mathcal{S} is either empty or consists of a single set T of size t . In the former case $|\mathcal{H}| = |\mathcal{H}'| = O(n^{r-t-1}) < \binom{n-t}{r-t}$ and there is nothing to prove, so we may assume such a T exists. This implies that every element of $\mathcal{H} \setminus \mathcal{H}'$ contains T .

First consider the case that \mathcal{H} contains some e which does not contain T . By our observation above, this means that every element of $\mathcal{H} \setminus \mathcal{H}'$ contains both T and some additional element of e . This implies

$$|\mathcal{H} \setminus \mathcal{H}'| + |\mathcal{H}'| \leq r \binom{n-t-1}{r-t-1} + |\mathcal{H}'| = O(n^{r-t-1}) < \binom{n-t}{r-t}.$$

Thus we can assume every element of \mathcal{H} contains T , which means $|\mathcal{H}| \leq \binom{n-t}{r-t}$. \square

The above argument actually gives the following stability result: for all r, t there exists a constant $c' = c'(r, t)$ such that if \mathcal{H} is t -intersecting with $|\mathcal{H}| > c' \binom{n-t}{r-t}$, then there exists a set of size t which is contained in every edge of \mathcal{H} .

Another application is a bound for how large an intersecting hypergraph \mathcal{H} can be if it's "far" from the extremal example, i.e. a star. There are many ways to make the notion of "far" precise. One way is to demand that every vertex of \mathcal{H} to be contained in the same number of edges, i.e. to demand that \mathcal{H} be regular.

Theorem 11.5 ([77]). *There exists an absolute constant $C > 0$ such that if $n \geq Cr \max\{\log r, k\}$, then any n -vertex intersecting r -graph \mathcal{H} has $|\mathcal{H}| \leq 2^{-k} \binom{n}{r}$.*

This result is roughly optimized when $k \approx n/r$, giving an upper bound of roughly $2^{-n/r} \binom{n}{r}$, which is typically much stronger than the bound $\binom{n-1}{r-1}$ given by Erdős-Ko-Rado for (not necessarily regular) intersecting families. Note that in particular this bound implies that regular intersecting hypergraphs cannot exist if $n \gg r$ **Which is maybe obvious by elementary means.**

Proof. Apply Theorem 11.2 with $\tau = 2$, which we can do if C is sufficiently large. Let $\mathcal{H}', \mathcal{S}$ be the corresponding families that we get. Note that by Theorem 11.2 we have

$$|\mathcal{H} \setminus \mathcal{H}'| \geq \frac{1}{2}|\mathcal{H}|.$$

We claim that every edge $e \in \mathcal{H} \setminus \mathcal{H}'$ intersects every $S \in \mathcal{S}$. Indeed, by the first property of Theorem 11.2, there exists some $S' \subseteq e$ with $S' \in \mathcal{S}$, and by Lemma 11.3, this S' (and hence e) intersects S .

Pick any $S \in \mathcal{S}$. By the previous claim, some vertex $x \in S$ must satisfy

$$d(x) \geq \frac{1}{|S|}|\mathcal{H} \setminus \mathcal{H}'| \geq \frac{1}{2k}|\mathcal{H}|,$$

where this last step used $|S| \leq k$ for any $S \in \mathcal{S}$ and $|\mathcal{H} \setminus \mathcal{H}'| \geq \frac{1}{2}|\mathcal{H}|$. However, since \mathcal{H} is regular, by the handshaking lemma we must have

$$d(x) = \frac{r}{n}|\mathcal{H}|.$$

This contradicts the previous bound if $n > 2rk$, giving the result. \square

11.1 Further Results

One can push the ideas of this section significantly further. Here we sketch out some of these ideas, and we refer the reader to [77] for the details.

One important direction is that one can consider different “ambient families.” That is, up to this point we were considering r -uniform hypergraphs, i.e. $\mathcal{H} \subseteq \mathcal{A} := \binom{[n]}{r}$. Alternatively, one can identify subsets of $[n^2]$ by their 0-1 characteristic vectors, which can in turn be written as n -dimensional 0-1 matrices. In particular, by letting \mathcal{A} denote the set of such vectors corresponding to permutation matrices, we can now consider “intersection” problems for sets of permutations $\mathcal{H} \subseteq \mathcal{A}$. Here one can again go through similar steps to develop a notion of homogenous families (with respect to our new choice of \mathcal{A}) in order to get results about sets of permutations \mathcal{H} which are intersecting (i.e. such that any two permutations $\pi, \sigma \in \mathcal{H}$ have $\pi(i) = \sigma(i)$ for some i , i.e. if their corresponding 0-1 matrices have a common entry equal to 1).

Through Lemma 11.3, we showed that if \mathcal{H} is t -intersecting and τ is small enough, then we can guarantee that \mathcal{S} is t -intersecting. Through a more refined argument, one can show that if τ is very small, then in fact this same conclusion holds if we only impose the much weaker condition that $|e \cap f| \neq t - 1$ for $e, f \in \mathcal{H}$.

Part IV

Entropy

12 Introduction

Throughout this part we let \log denote logarithms base 2 unless stated otherwise, and we define $x \log x = 0$ whenever $x = 0$.

Let X be a discrete random variable and $p_x = \Pr[X = x]$. The *binary entropy* of this random variable is defined as

$$H[X] = - \sum_{x \in \text{supp}(X)} p_x \log(p_x),$$

where the sum ranges over all x in the support of X (i.e. those x with $p_x > 0$).

The definition for $H[X]$ is quite strange if one has never seen it before. Roughly speaking, $H[X]$ can be thought of as measuring how much “information” the random variable X carries. For example, one can easily check that $H[X] = 0$ if and only if X is deterministic, corresponding to the fact that knowing the outcome of a deterministic process gives no new information.

12.1 The Main Properties

Before getting into applications, let us start by recording the most common properties of entropy that will be used throughout this part, and for this it will be useful to establish some notation.

Given a random variable X and an event E , we define $X|E$ to be the random variable X conditioned on the event E . Given a pair of random variables X, Y we define

$$H[X|Y] := \mathbb{E}_{y \sim Y} H[X|Y = y] = - \sum_y \Pr[Y = y] \sum_x \Pr[X = x|Y = y] \log(\Pr[X = x|Y = y]).$$

The expression $H[X|Y]$ is commonly referred to as *conditional entropy*. For convenience, we will often denote vectors of random variables (X_1, \dots, X_n) as simply X_1, \dots, X_n , e.g. by writing $H[X_1, \dots, X_n]$ instead of $H[(X_1, \dots, X_n)]$. For an integer i we let $X_{<i} := (X_1, \dots, X_{i-1})$. Slightly more generally, if X is a random vector indexed by a set S with a total ordering $<$, then we let $X_{<s}$ denote the elements of X indexed by $t < s$.

We now state our list of properties about the entropy function. The reader is not expected to memorize this right away, though it might be a good idea to what extent these properties agree with the intuition¹ that $H[X]$ measures the information of X .

Claim 12.1. *The following properties hold for any random variable X .*

¹For example, Subadditivity says that the total information in the vector (X, Y) is at most the sum of the information of X and Y , Dropping Conditioning says that knowing less at the start can only lead to more information gained, and Data Processing says that one can’t do anything to a random variable Y in order to give it more information than it already has.

- (Non-negativity) We have $H[X] \geq 0$ with equality if and only if X is deterministic.
- (Maximality Principle) We have

$$H[X] \leq \log |\text{supp}(X)|,$$

with equality if and only if X is uniformly distributed on $\text{supp}(X)$.

- (Chain Rule) For two random variables X, Y we have

$$H[X, Y] = H[X] + H[Y|X].$$

More generally, given random variables X_1, \dots, X_n we have

$$H[X_1, \dots, X_n] = \sum_i H[X_i | X_1, \dots, X_{i-1}].$$

- (Subadditivity) For random variables X_1, \dots, X_n we have

$$H[X_1, \dots, X_n] \leq \sum_i H[X_i].$$

- (Dropping Conditioning) For random variables X, Y, Z we have

$$H[X|Y] \leq H[X],$$

with equality if and only if X is independent of Y . Similarly

$$H[X|Y, Z] \leq H[X|Y],$$

with equality if and only if X conditioned on Y has the same distribution as X conditioned on both Y and Z .

- (Data Processing Inequality) If X, Y, Z are random variables such that Z is a function of Y , then

$$H[X|Z] \leq H[X|Y].$$

Not only does the function $H[X] = -\sum_x p_x \log(p_x)$ satisfy these properties, it is in fact essentially the unique function satisfying (somewhat weaker versions of) these properties. Because of this, H will be the “right” function for us to use to encode how much “information” X has. For more on this see e.g. [60, Chapter 7].

We will not prove the claim since it’s a slight detour from our main goal of applying entropy to combinatorics problems. Most proofs can be found in standard texts on entropy, e.g. the survey by Galvin [54] (which contains even more properties, especially around conditional entropy). The only exceptions might be proofs for the “only if” portion of Dropping Conditioning (which we will never use); as well as the Data Processing Inequality (whose name is not entirely standard), but this follows from observing that $H[X|Y, Z] = H[X|Y]$ by definition and that $H[X|Y, Z] \geq H[X|Z]$ by Dropping Conditioning.

There is plenty of redundancy in the list of properties above. For example Subadditivity follows from the Chain Rule and Dropping Conditioning, but this property is used so frequently that it will be useful to list it as a separate property. For our purposes, the most useful property from this list will be the if and only if portion of the Maximality Principle. More precisely, we have the following consequence of it (our name for which is non-standard).

Corollary 12.2 (Fundamental Property of Entropy). *If \mathcal{X} is a set and $X \in \mathcal{X}$ is chosen uniformly at random, then*

$$|\mathcal{X}| = 2^{H[X]}.$$

This corollary shows that bounding the size of a set is equivalent to bounding the entropy of a corresponding random variable, and sometimes this latter perspective can be easier to work with, especially when it comes to proving upper bounds.

12.2 Basic Application 1: Binomial Coefficients

We begin with a typical usage for entropy by upper bounding (sums of) binomial coefficients. For this we define for $p \in [0, 1]$ the binary entropy function

$$H(p) := -p \log_2(p) - (1 - p) \log_2(1 - p),$$

which is simply the entropy of a Bernoulli variable with probability of success p .

Proposition 12.3. *For all $k \leq n/2$, we have*

$$\sum_{i=0}^k \binom{n}{i} \leq 2^{H(k/n) \cdot n}.$$

Proof. Let \mathcal{X} denote the set of binary strings of length n with at most k 1's. Note that

$$|\mathcal{X}| = \sum_{i=0}^k \binom{n}{i},$$

so by the Fundamental Property of Entropy, proving our desired bound $\log |\mathcal{X}| \leq H(k/n) \cdot n$ is equivalent to showing $H[X] \leq H(k/n) \cdot n$ where $X \in \mathcal{X}$ is chosen uniformly at random. For this we use the following.

Claim 12.4. *Each of the random variables X_i is a Bernoulli random variable with probability of success $p \leq k/n \leq 1/2$.*

Proof. By the symmetry of the problem, we see that each X_i is Bernoulli with the same probability of success p . By construction we deterministically have $\sum X_i \leq k$ for all $X \in \mathcal{X}$, so it must be that $p \leq k/n$, and this is at most $1/2$ by hypothesis on k . \square

The claim above implies that for all i ,

$$H[X_i] = H(p) \leq H(k/n),$$

with this last step using the (easy to prove) fact that $H(p)$ is increasing for $p \leq 1/2$. This together with Subadditivity gives

$$H[X] \leq \sum H[X_i] = H(k/n) \cdot n,$$

proving the result. \square

We note that this bound is essentially tight in the sense that if k is linear in n , then $\binom{n}{k} = 2^{(1+o(1))H(k/n) \cdot n}$.

The framework in our proof above is typical for the entropy method: we (1) started with a uniform random object X and then (2) used entropy to upper bound the size of $\text{supp}(X)$. Although this is the most common framework for using entropy, it is also possible to (1') start with a non-uniform random object X and then (2') use entropy to lower bound the size of $\text{supp}(X)$. We study one such example in the next subsection.

12.3 Basic Application 2: Walks in Graphs

One of the most famous open problems in extremal combinatorics is Sidorenko's conjecture stated below, where we recall that $\text{hom}(F, G)$ denote the number of homomorphisms from a graph F to a graph G .

Conjecture 12.5 (Sidorenko's Conjecture). *If F is a bipartite graph, then every n -vertex graph G satisfies*

$$\frac{\text{hom}(F, G)}{n^{|V(F)|}} \geq \left(\frac{2e(G)}{n^2} \right)^{e(F)}.$$

This bound is asymptotically best possible for $G = G_{n,p}$, so the conjecture morally says that every graph G contains as many (homomorphic) copies of each bipartite graph as one would expect in the random graph of the same density. Note that the bound does not hold for F non-bipartite (via taking $G = K_2$).

There are very few examples of graphs F for which Sidorenko's Conjecture is known. Here we outline a nice entropy proof which shows that Sidorenko's Conjecture holds for all F which are paths. In this setting, we observe that homomorphisms from paths of length k to G are equivalent to walks of length k in G , i.e. sequences of vertices (x_1, \dots, x_{k+1}) such that $x_i \sim x_{i+1}$ for all $1 \leq i \leq k$. As such, the following result (commonly known as the Blakey-Roy Theorem (though they weren't quite the first to prove it)) is equivalent to Sidorenko's Conjecture for paths.

Theorem 12.6 (Blakey-Roy [15]). *If G is an n -vertex graph with $m \geq 1$ edges, then the number of walks of length k in G is at least*

$$2m(2m/n)^{k-2}.$$

Note that this bound is tight whenever G is regular.

Proof. Let $X = (X_1, \dots, X_{k+1})$ be a random walk of length k in G chosen in the following *non-uniform* way: choose the pair (X_1, X_2) uniformly at random amongst all pairs such that $X_1 \sim X_2$, and given X_{i-1} for $i \geq 3$, we choose X_i uniformly at random amongst the neighbors of X_{i-1} . Observe that X is indeed always a walk of length k (with us implicitly using that (X_1, X_2) exists due to $m \geq 1$). We can express its entropy as

$$\begin{aligned}
H[X] &= \sum_{i=1}^{k+1} H[X_i | X_{<i}] \\
&= H[X_1] + H[X_1 | X_2] + \sum_{i=3}^{k+1} H[X_i | X_{i-1}] \\
&= H[X_1, X_2] + \sum_{i=3}^{k+1} H[X_i, X_{i-1}] - H[X_{i-1}], \tag{13}
\end{aligned}$$

where here the first and last equality used the Chain Rule, and the second used the equality case of Dropping Conditioning (since X_i depends only on X_{i-1}).

We claim (crucially) that (X_{i-1}, X_i) is uniformly random amongst all pairs such that $X_{i-1} \sim X_i$. This is true for $i = 2$ by construction, so assume we have proven it true up to some value $i \geq 3$. In this case, for any $(y, z) \in V(G)^2$ such that $y \sim z$, we have (by conditioning on every possible value that $X_{i-2} \sim X_{i-1}$ can take on)

$$\begin{aligned}
\Pr[(X_{i-1}, X_i) = (y, z)] &= \sum_{x \in N(y)} \Pr[(X_{i-2}, X_{i-1}) = (x, y)] \cdot \Pr[X_i = z | (X_{i-2}, X_{i-1}) = (x, y)] \\
&= \sum_{x \in N(y)} \frac{1}{2m} \cdot \frac{1}{\deg(y)} = \frac{1}{2m},
\end{aligned}$$

with this last equality using the hypothesis that (X_{i-2}, X_{i-1}) is distributed uniformly at random and that X_i is a uniform random neighbor of X_{i-1} . This establishes the claim.

With this claim, we have by the Maximization Principle that $H[X_i, X_{i-1}] = \log(2m)$ for all i , and also that $H[X_{i-1}] \leq \log(n)$ for all i . Using this with (13) gives

$$H[X] \geq (k-1) \log(2m) - (k-2) \log(n).$$

Let W_k denote the set of walks of length k in G . Since X is a random element from W_k , we have $H[X] \leq \log |W_k|$, which combined with the lower bound for $H[X]$ above gives the desired result. \square

To emphasize, proof above would theoretically have worked if we considered a uniform random element $\tilde{X} \in W_k$ instead of the non-uniform X described above, in the sense that the Maximization Principle gives

$$H[\tilde{X}] \geq H[X] \geq (k-1) \log(2m) - (k-2) \log(n).$$

However, it is not clear how one would prove the lower bound of $(k-1) \log(2m) - (k-2) \log(n)$ for $H[\tilde{X}]$ directly without going through X first. More generally, it can be useful when working

with lower bounds from entropy to work with a distribution for X which is “natural” to the problem rather than one which is uniform. We also emphasize that to obtain tight examples, it is crucial that our choice of X is actually distributed uniformly whenever we are working with an extremal example (as otherwise the Maximization Principle shows that we can not hope to obtain a tight bound).

The exact same proof as above can easily be extended to prove Sidorenko’s Conjecture for all trees at the cost of more complicated notation. A different and somewhat more involved entropy argument can be used to show Sidorenko’s conjecture holds whenever F has a vertex which is adjacent to every vertex in the other partition set. This result was originally proven by Conlon, Fox, and Sudakov [28] using Dependent Random Choice **And I’m not sure who gave the entropy proof for this, though it can be found in notes of Yufei Zhao.**

12.4 Further Applications

While one can prove a number of results using only the entropy properties listed above (see for example Theorem 15.2), we can greatly increase the power of this method by adding in a few extra tools which we explore in the forthcoming two sections. Much more can be said regarding the entropy method than we do in this text, and we refer the reader to the survey by Galvin [54] for much more on this topic.

13 Shearer's Lemma

In this section we consider a strengthening of the Subadditivity property of entropy known as Shearer's lemma, which plays a central role in many applications of the entropy method (so much so that it is often listed as one of the basic properties of entropy). For this, given a vector $X = (X_1, \dots, X_n)$ and a set $F = \{i_1, \dots, i_k\} \subseteq [n]$, we define $X_F = (X_{i_1}, \dots, X_{i_k})$.

Lemma 13.1 (Shearer's Lemma). *Let $X = (X_1, \dots, X_n)$ and let $\mathcal{F} \subseteq 2^{[n]}$ be a set system such that every $i \in [n]$ is contained in at least d sets of \mathcal{F} , i.e. such that \mathcal{F} has minimum degree at least d . Then*

$$H[X] \leq d^{-1} \sum_{F \in \mathcal{F}} H[X_F].$$

Note that if \mathcal{F} consists of the singletons $\{i\}$, then Shearer's lemma with $d = 1$ exactly recovers subadditivity.

The statement of Lemma 13.1 is a little strange at first glance. Roughly speaking, the intuition is that because each i appears at least d times in \mathcal{F} , the sum on the righthand side in principle has as much information as d independent copies of X , and hence dividing by d should in principle be an upper bound for the amount of information of X . This intuition can easily be made precise if the X_i are mutually independent, as in this case we have

$$d \cdot H[X] = \sum_i d \cdot H[X_i] \leq \sum_i \deg_{\mathcal{F}}(i) \cdot H[X_i] = \sum_{F \in \mathcal{F}} H[X_F].$$

A similar line of reasoning gives the result in general.

Proof. Our proof will go through the same logic as in the independent case, except that we will replace both of our equalities with an application of the Chain Rule. To this end, given i, F with $i \in F \in \mathcal{F}$, we define $(i, F) = \{j \in F : j < i\}$. With this we have

$$\begin{aligned} d \cdot H[X] &= \sum_i d \cdot H[X_i | X_{<i}] \\ &\leq \sum_i \deg_{\mathcal{F}}(i) \cdot H[X_i | X_{<i}] = \sum_i \sum_{F \in \mathcal{F}} H[X_i | X_{<i}] \\ &\leq \sum_i \sum_{F \in \mathcal{F}} H[X_i | X_{(i,F)}] = \sum_{F \in \mathcal{F}} H[X_F], \end{aligned}$$

where this last inequality used Dropping Conditioning. □

We begin with a basic application of Shearer's lemma to a problem from geometry. Given a vector $x \in \mathbb{Z}^n$ and $i \in [n]$, we define the projection $\pi_i(x) = (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$, and given $S \subseteq \mathbb{Z}^n$ we define $\pi_i(S) = \{\pi_i(x) : x \in S\}$. A natural extremal question is: given the sizes of $\pi_i(S)$ for all i , how large can S itself be?

A natural candidate for a construction here is to take S to be a cube whose i th side length is $|\pi_i(S)|^{-(n-2)/(n-1)} \prod_j |\pi_j(S)|^{1/(n-1)}$. Indeed, it is easy to check that each $\pi_i(S)$ has the prescribed size, and with this we have $|S| = \prod |\pi_i(S)|^{1/(n-1)}$. It turns out this is best possible.

Theorem 13.2 (Discrete Loomis-Whitney Theorem). *If $S \subseteq \mathbb{Z}^n$, then*

$$|S| \leq \prod |\pi_i(S)|^{1/(n-1)}.$$

This result is essentially equivalent to the Loomis-Whitney Theorem [79], which says that if K is a measurable body in \mathbb{R}^n , then $\text{vol}_n(K) \leq \prod \text{vol}_{n-1}(\pi_i(K))^{1/(n-1)}$, where here π_i is the corresponding projection map and vol_n is the n -dimensional volume measure. Indeed, one can show that it suffices to prove Loomis-Whitney when K is the union of axis-aligned unit cubes via an approximation argument, and in this case one can replace the center of each unit cube of K by its center, which exactly translates to the present problem.

Proof. As per usual, we start by picking X to be a uniformly random element of S . Observe that, in the notation of Shearer's lemma, $\pi_i(X) = X_{[n] \setminus \{i\}}$. Motivated by this, we define \mathcal{F} to consist of all sets of the form $[n] \setminus \{i\}$. Since this set system has minimum degree $n - 1$, Shearer's lemma gives

$$\log_2 |S| = H[X] \leq \frac{1}{n-1} \sum H[X_{[n] \setminus \{i\}}] = \frac{1}{n-1} \sum H[\pi_i(X)] \leq \frac{1}{n-1} \sum \log_2 |\pi_i(S)|,$$

with this last step using $\pi_i(X) \in \pi_i(S)$ (since $X \in S$) together with the Maximization Principle. Exponentiating both sides gives the result. \square

We next look at a more involved application of Shearer's Lemma involving homomorphisms between graphs. To this end, we let $\text{hom}(G, H)$ denote the number of homomorphisms ϕ from G to H . For example, $\text{hom}(G, K_r)$ is the number of proper r -colorings of G . We will also want to allow our target graph H to contain loops. For example, if H is a K_2 with a loop on one vertex, then one can check that $\text{hom}(G, H)$ counts the number of independent sets of G .

The fundamental question we want to ask is: given a target graph H and some family of graphs \mathcal{G} , which $G \in \mathcal{G}$ maximizes $\text{hom}(G, H)$? One natural family to consider is the set of n vertex d -regular graphs. While this problem is still open for general H , there is a very clean solution when we restrict to bipartite G .

Theorem 13.3 ([69, 110]). *If G is a bipartite n -vertex d -regular graph, then for any graph H (possibly with loops), we have*

$$\text{hom}(G, H) \leq \text{hom}(K_{d,d}, H)^{n/2d}.$$

This bound is sharp by considering G to be the disjoint union of $n/2d$ copies of $K_{d,d}$. This result was first proven in the case when H is a K_2 with a loop (i.e. when counting independent sets in G) by Kahn [69], with his argument later generalized considerably by Galvin and Tetali [56] to the present theorem.

Proof. It will be convenient to express our homomorphisms as vectors. To this end, given a homomorphism $\phi : V(G) \rightarrow V(H)$, we define the vector $X^\phi \in V(G)^{V(H)}$ by having $X_u^\phi = \phi(u)$ for each $u \in V(G)$. Let ϕ be a uniform random homomorphism from G to H , and let $X = X^\phi$.

Let $U \cup V$ denote a bipartition of G , noting that $|U| = |V| = n/2$ since G is regular. By the Maximization Principle and the chain rule, we have

$$\log(\text{hom}(G, H)) = H[X] = H[X_U] + H[X_V|X_U].$$

Because G is d -regular, every $u \in U$ is contained in exactly d of the sets $\mathcal{F} := \{N(v) : u \in U\}$, so by Shearer's lemma we have

$$H[X_U] \leq d^{-1} \sum_{v \in V} H[X_{N(v)}].$$

By Subadditivity and dropping Conditioning we have

$$H[X_V|X_U] \leq \sum_{v \in V} H[X_v|X_U] \leq \sum_{v \in V} H[X_v|X_{N(v)}],$$

so in total we find

$$\log(\text{hom}(G, H)) = H[X] \leq d^{-1} \sum_{v \in V} H[X_{N(v)}] + d \cdot H[X_v|X_{N(v)}], \quad (14)$$

and it now suffices to upper bound each term in the sum.

Fix some $v \in V$. For each vector x of length d indexed by $V(H)$, let $A(x)$ denote the set of vertices of $V(H)$ which are adjacent to every vertex of x_1, \dots, x_d . Equivalently, $A(x)$ is the set of “available” values that X_v can take given that $X_{N(v)} = x$ (here we use that X corresponds to a homomorphism, i.e. X_v must be adjacent to every vertex of $X_{N(v)}$). We have $H[X_v|X_{N(v)} = x] \leq \log |A(x)|$ by the Maximization Principle, which together with the definition of (conditional) entropy gives

$$\begin{aligned} H[X_{N(v)}] + d \cdot H[X_v|X_{N(v)}] &= \sum_x \Pr[X_{N(v)} = x] (-\log(\Pr[X_{N(v)} = x]) + d \cdot H[X_v|X_{N(v)} = x]) \\ &\leq \sum_x \Pr[X_{N(v)} = x] \left(\log \left(\frac{|A(x)|^d}{\Pr[X_{N(v)} = x]} \right) \right) \\ &\leq \sum_x \log(|A(x)|^d) = \log \left(\prod_x |A(x)|^d \right), \end{aligned}$$

where the second inequality used Jensen's inequality together with the convexity of \log .

Crucially, we observe that $\prod_x |A(x)|^d$ where x ranges all d -length vectors indexed of H is exactly equal to $\text{hom}(K_{d,d}, H)$. Indeed, each homomorphism from $K_{d,d}$ can be uniquely identified by first choosing a vector x based on how the vertices on the bottom half of $K_{d,d}$ map to H , and after this each of the d remaining vertices of $K_{d,d}$ can map to any vertex in $A(x)$ while maintaining the homomorphism property. Combining this with the inequality above gives

$$H[X_{N(v)}] + d \cdot H[X_v|X_{N(v)}] \leq \log(\text{hom}(K_{d,d}, H)),$$

and plugging this into (14) and using $|V| = n/2$ gives

$$\log(\text{hom}(G, H)) \leq d^{-1} \sum_{v \in V} \log(\text{hom}(K_{d,d}, H)) = \log(\text{hom}(K_{d,d}, H))^{n/2d}.$$

□

The result above gives optimal bounds for $\text{hom}(G, H)$ when G is bipartite, and an elegant idea of Zhao's allows one to immediately lift this result to all G for certain H . In particular, we get the following.

Theorem 13.4 (Zhao [110]). *Let $i(H)$ denote the number of independent sets of a graph H . If G is an n -vertex d -regular graph, then $i(G) \leq i(K_{d,d})^{n/2d}$.*

Proof Sketch. Consider the “bipartite double cover” $G \times K_2$, i.e. the graph with vertex set $V(G) \times \{1, 2\}$ where $(u, s) \sim (v, t)$ if and only if $u \sim_G v$ and $s \neq t$. A simple but clever injectivity argument shows $i(G \times K_2) \geq i(G)^2$. Because $G \times K_2$ is a d -regular $(2n)$ -vertex bipartite graph, we have by Theorem 13.3 (applied when H is K_2 with a loop) that

$$i(G)^2 \leq i(G \times K_2) \leq i(K_{d,d})^{n/d},$$

giving the result. □

This same bipartite double cover trick can be used to extend Theorem 13.3 to all d -regular graphs for a few other choices of H [111], though there exist H for which such an extension is impossible (e.g. H being two isolated vertices with loops fails for $G = K_3$). Finally, we note a substantial extension of Theorem 13.3 due to Sah, Sawhney, Stoner, and Zhao [98], who showed that any graph G (not necessarily regular) which is triangle-free and has no isolated vertices satisfies for all graphs H that

$$\text{hom}(G, H) \leq \prod_{uv \in E(G)} \text{hom}(K_{d_u, d_v}, H)^{\frac{1}{d_u d_v}}.$$

There proof does not use entropy at all, but instead a very elaborate set of analytic inequalities.

14 Random Chain Rules

This section concerns a strengthening of the chain rule. Historically, this strengthening was first used to give an entropy proof of the following result.

Theorem 14.1 (Brégman’s Theorem [18]). *If G is a bipartite graph with bipartition $U \cup V$ such that $|U| = |V| = n$, then the number of perfect matchings of G is at most*

$$\prod_{u \in U} (\deg(u)!)^{1/\deg(u)}.$$

Observe that this bound is tight by considering disjoint unions of complete bipartite graphs. This was originally proven by Brégman [18], and later Radhakrishnan [91] gave an elegant entropy-based proof that we present below.

Before stating the key lemma needed to prove Theorem 14.1, let’s first try and prove this result naively from first principles and see where things go wrong.

As usual, we start with a uniform random perfect matching M of G . We then translate M into a vector X indexed by U by having $X_u \in V$ be the unique neighbor of u in M . We fix some arbitrary ordering $<$ of the vertices of U , and then apply the chain rule to obtain

$$H[X] = \sum_u H[X_u | X_{<u}].$$

At this point, the naive entropy argument calls for upper bounding $H[X_u | X_{<u}]$ by log of the number of possible values X_u can take given the values of $X_{<u}$. To this end, we define $A_{<u} \subseteq N(u)$ to be the set of neighbors of u which do not appear in $X_{<u}$ (i.e. this is the set of available neighbors of u for M given the information in $X_{<u}$). The Maximality Principle then gives

$$\sum_u H[X_u | X_{<u}] \leq \sum_u \log(|A_{<u}|).$$

Unfortunately, for any given u and ordering $<$, we can’t say anything about $|A_{<u}|$ other than $|A_{<u}| \leq \deg(u)$. Applying this worst-case bound for all u gives a trivial upper bound of $\prod_u \deg(u)$ in the end.

While it is true that worst case we can have $|A_{<u}| = \deg(u)$ for any given u , intuitively we should “typically” have $|A_{<u}| \approx \frac{1}{2} \deg(u)$, since for a “random” M we would expect around half of u ’s neighbors to be matched in M to vertices appearing before u in $<$ and half to be matched to vertices after u . Again, this intuition may not hold for a given M and $<$, but this intuition can be made precise if we consider a random ordering $<$ instead of a fixed one. To this end, one can consider the following random variant of the chain rule.

Lemma 14.2 (Random Chain Rule). *Let X be a vector indexed by a set S and let $<$ be a random ordering of S . Then*

$$H[X] = \sum_{s \in S} \mathbb{E}_{<} [H[X_s | X_{<s}]].$$

Indeed, the proof of this follows from the fact that equality holds for any fixed $<$ (by the usual chain rule), and hence equality also holds when one takes expectations. With this we can quickly adapt our previous failed attempt to give Theorem 14.1.

Proof of Theorem 14.1. Let $<$ denote a uniform random ordering of U . Keeping all of the notation from the argument above, we have by the random chain rule that

$$H[X] = \sum_u \mathbb{E}_{<}[H[X_u | X_{<u}]] \leq \sum_u \mathbb{E}_{<}[\log(|A_{<u}|)]. \quad (15)$$

We claim that $|A_{<u}|$ is distributed uniformly at random amongst $[\deg(u)]$. Indeed, fix any perfect matching M (so now all the randomness lies in the much simpler random object $<$) and let v denote the neighbor of u in M . We then observe that $|A_{<u}| = i$ if and only if $i - 1$ vertices of $N(u) \setminus \{v\}$ have their neighbors in M appear after u under $<$. Since $<$ gives a uniform random ordering on $N(u)$ regardless of our choice of M , we conclude that $|A_{<u}|$ is indeed equally likely to be any value in $[\deg(u)]$. Again, we emphasize that this result holds regardless of the fixed value of M , and hence $|A_{<u}|$ continues to be uniformly distributed even if we do not condition on M .

With this claim, we can write (15) above as

$$H[X] \leq \sum_u \frac{1}{\deg(u)} \sum_{i=1}^{\deg(u)} \log(i) = \sum_u \frac{\log(\deg(u)!)}{\deg(u)}.$$

Exponentiating both sides gives the result. \square

Unpublished work of Kahn and Lovász generalizes Brégman's theorem to non-bipartite graphs by proving that every graph has at most $\prod_{u \in V(G)} d(u)!^{1/2d(u)}$ perfect matchings. An entropy proof of this result was given by Cutler and Radcliffe [32], with this proof being complicated by the fact that $|A_{<u}|$ is no longer uniformly distributed if u has edges inside $N(u)$. A short proof of Alon and Friedland [3] deduces this result of Kahn and Lovász directly from Brégman's theorem.

Our proof above used the random chain rule to count the number of 1-factors (i.e. perfect matchings) in a graph. We next use a somewhat more complex version of this argument to count 1-factorizations in K_n , which we recall are ordered partitions of $E(G)$ into perfect matchings. For example, K_4 has 6 different 1-factorizations, namely $(\{12, 34\}, \{13, 24\}, \{14, 23\})$ and all of its permutations.

I don't know if there's a relevant citation here.

Theorem 14.3. *The number of 1-factorizations of K_n when n is even is at most $((1 + o(1))n/e^2)^{\binom{n}{2}}$.*

Observe that this improves upon the trivial upper bound $(n-1)^{\binom{n}{2}}$ (which is just the number of ways to partition the edge set of K_n into $n-1$ edge-disjoint graphs).

Proof. For this proof, it will be slightly more convenient to work with a base e notion of entropy rather than the usual base 2 notion. To this end, if X is a random variable with $p_x = \Pr[X = x]$, then we define

$$H_e[X] = - \sum_x p_x \log_e(P_X).$$

Note that $H_e[X] = \log_e(2)H[X]$, and in particular, essentially all of the properties for $H[X]$ continue to hold for $H_e[X]$.

Let M denote a uniformly random 1-factorization of K_n . We will think of M as assigning to each edge uv of K_n a color in $[n-1]$ such that the edges in color i form a perfect matching (equivalently, M is a proper edge coloring of K_n with $n-1$ colors). Let X be the vector indexed by $E(K_n)$ where X_{uv} equals the color assigned to uv by M .

As before, we will consider a uniformly random ordering $<$ on $E(K_n)$, but for technical reasons we will want to form this ordering in a slightly more complex way. To this end, assign to each edge uv a random weight w_{uv} chosen independently and uniformly from $[0, 1]$, then let $<$ be the ordering of $E(K_n)$ which has $uv < xy$ iff $w_{uv} < w_{xy}$. Again we emphasize that $<$ has the same distribution as if we just chose it to be uniformly at random, but it will be convenient for us to have these extra w_{uv} parameters to work with. An application of the random chain rule then gives

$$H_e[X] = \sum_{uv} \mathbb{E}_<[H_e[X_{uv}|X_{<uv}]] \leq \sum_{uv} \mathbb{E}_<[\log_e(|A_{<uv}|)], \quad (16)$$

where here A_{uv} denotes the set of colors that are “available” for uv given $X_{<uv}$; i.e. A_{uv} consists of the colors i that do not lie on any edge xy which intersects uv and which has $xy < uv$ under the coloring M .

It remains to estimate $\mathbb{E}_<[\log_e(|A_{<uv}|)]$, and for this, it will suffice to condition on the 1-factorization M and prove an upper bound that is independent of M . From now on we fix M , noting that $A_{<uv}$ will always contain the color c which M assigns to uv . Observe that a color $i \in [n-1] \setminus \{c\}$ will be in $A_{<uv}$ if and only if uv appears in the ordering before the two edges incident to uv which are colored i by M (note that exactly two such edges exist since M is a 1-factorization). Conditional on the value w_{uv} , the probability that this happens for any given i is $(1 - w_{uv})^2$. Thus in total, we have

$$\mathbb{E}_<[\log_e(|A_{<uv}|)|w_{uv}, M] = \log_e(1 + (n-2)(1 - w_{uv})^2).$$

As w_{uv} was distributed uniformly at random in $[0, 1]$, we find

$$\mathbb{E}_<[\log_e(|A_{<uv}|)|M] = \int_0^1 \log_e(1 + (n-2)(1-x)^2) dx.$$

Summing this over all edges uv together with (16) gives

$$H_e[X] \leq \binom{n}{2} \int_0^1 \log_e(1 + (n-2)(1-x)^2) dx = \log_e(n) - 2 + o(1),$$

where this last equality follows from some fiddly integral analysis¹ (with the intuition being that the integrand is close to $\log_e(n(1-x)^2) = \log_e(n) + 2\log_e(1-x)$, and this ends up integrating to the desired value). Exponentiating both sides by e gives that the total number of 1-factorizations is at most $((1 + o(1))n/e^2)^{\binom{n}{2}}$ as desired.

□

¹Slightly more precisely, one can argue that the integral evaluated from $1 - n^{-1}$ to 1 is at most $O(n^{-1} \log n) = o(1)$, and outside of this range the difference between the integrand $\log_e(1 + (n-2)(1-x)^2)$ and $\log_e(n(1-x)^2)$ is $o(1)$.

As an aside, if in (16) we used the Chain Rule instead of the Random Chain Rule, then we would pessimistically use $|A_{<uv}| \leq n - 1$ for all uv , and this would give the trivial upper bound $(n - 1)^{\binom{n}{2}}$.

15 The Union-Closed Sets Conjecture

Up to this point we've used entropy to bound the size of some set, and this is certainly the most common way to utilize entropy for combinatorial problems. However, there are more exotic ways that entropy can be used to solve problems. We illustrate one such example here involving a major breakthrough by Gilmer for the Union-Closed Sets Conjecture. For this, we say that a set system \mathcal{F} is *union-closed* if for all $A, B \in \mathcal{F}$, we also have $A \cup B \in \mathcal{F}$. The following (frustratingly) simple extremal question related to union-closed families is typically attributed to Frankl from 1979 [45].

Conjecture 15.1 (Union-Closed Sets Conjecture). *If $\mathcal{F} \neq \{\emptyset\}$ is union-closed, then there exists some x in at least half of the elements of \mathcal{F} , i.e. such that $\deg(x) \geq \frac{1}{2}|\mathcal{F}|$.*

We omit discussing the history of this conjecture prior to 2022 (most of which can be found in the survey by Bruhn and Schaudt [19]). The only thing we note is that after many years of work, the best bound known was that there exists an x with $\deg(x) = \Omega(\frac{|\mathcal{F}|}{\log_2(|\mathcal{F}|)})$ due to Knill [76]. This bound stood for nearly 30 years until the following breakthrough result.

Theorem 15.2 (Gilmer [57]). *There exists a constant $c > 0$ such that if $\mathcal{F} \neq \{\emptyset\}$ is a union closed set system, then there exists an element x such that $\deg(x) \geq c|\mathcal{F}|$.*

Gilmer originally proved this with $c = .01$, though his methods were quickly optimized to give $c = \frac{3-\sqrt{5}}{2} \approx .381$ by a variety of authors, and our particular approach for achieving this constant will be closest to that of Chase and Lovett [24].

The key insight of Gilmer's approach is to look at the contrapositive of the Union-Closed Sets Conjecture, i.e. by showing that if a set system has $\deg(x) < c|\mathcal{F}|$ for all x , then \mathcal{F} can not be union-closed. In particular, we will reach this union-closed conclusion through the following entropy result.

Theorem 15.3. *There exists a constant $c > 0$ such that the following holds. If \mathcal{F} is a set system with $|\mathcal{F}| > 1$ and the property that $\deg(x) < c|\mathcal{F}|$ for all x , and if $A, B \in \mathcal{F}$ are chosen independently and uniformly at random, then*

$$H[A \cup B] > H[A].$$

Let us assume this result for the moment and show how it implies Gilmer's theorem.

Proof of Theorem 15.2. Let $\mathcal{F} \neq \{\emptyset\}$ be union-closed. The result is trivial if \mathcal{F} has only 1 element, so assume this is not the case.

Let $A, B \in \mathcal{F}$ be chosen independently and uniformly at random. Because \mathcal{F} is union-closed, the random variable $A \cup B$ always lies in \mathcal{F} . As such, the Maximality Principle implies.

$$H[A \cup B] \leq \log |\mathcal{F}| = H[A],$$

with this last step using that A is chosen uniformly at random. This contradicts the conclusion of Theorem 15.3, so it must be the case that $\deg(x) \geq c|\mathcal{F}|$ for some element x . \square

The proof above illustrates that the Union-Closed Sets Conjecture would follow if Theorem 15.3 held with $c = .5$. However, it turns out that $c = \frac{3-\sqrt{5}}{2}$ is the best constant one can prove here.

We now move onto our proof of Theorem 15.3. To start, we will attempt to build some intuition by working out how one might go about trying to solve this problem, though the eager reader is welcome to jump ahead to Lemma 15.4 for the formal details.

One way you might think of proving Theorem 15.3 is through induction on the ground set of \mathcal{F} . To this end, we can think of A, B as characteristic vectors in $\{0, 1\}^n$. In this case, the chain rule implies that the conclusion of Theorem 15.3 is equivalent to having

$$\sum_{i \leq n} H[(A \cup B)_i | (A \cup B)_{<i}] > \sum_{i \leq n} H[A_i | A_{<i}].$$

Given this formulation, one might hope to prove this inequality term by term, and indeed this is what we will do.

In particular, when $i = 1$ we have that A_1, B_1 are just Bernoulli random variables with some common probability of failure¹ p . In this case, $(A \cup B)_1$ is a Bernoulli random variable with probability of failure p^2 . As such, we need to show that if p is large (i.e. if the element 1 fails to be in a large proportion of edges of \mathcal{F}), then

$$H(p^2) > H(p),$$

where here $H(x) := -x \log_2(x) - (1-x) \log_2(1-x)$ is the entropy function for a Bernoulli random variable with probability x . Intuitively, this inequality will hold if and only if p^2 is closer to $1/2$ than p is to $1/2$, so the cutoff point should be when these two distances from $1/2$ are the same. That is, we need to solve for $.5 - p^2 = p - .5$, or equivalently to compute the roots of $p^2 + p - 1 = 0$. One can quickly check that the positive root of this is $p = \phi := \frac{\sqrt{5}-1}{2}$.

In conclusion, this first step of our induction (heuristically) seems to hold provided $p > \phi$ and fails otherwise. For our full proof, we will in fact need the following generalization of the claim that $H(p^2) > H(p)$ when $p > \phi$.

Lemma 15.4. *For all $p, q \in [0, 1]$, we have*

$$H(pq) \geq \frac{1}{2\phi}(pH(q) + qH(p)),$$

where again $H(x) := -x \log_2(x) - (1-x) \log_2(1-x)$ and $\phi := \frac{\sqrt{5}-1}{2}$.

The proof of this purely analytical lemma is somewhat annoying to prove, so we'll omit it from the text here; see [17, 24] for the details. With this lemma and our approach outlined above, we'll quickly be able to derive the following.

Lemma 15.5 ([24]). *If $A, B \in \{0, 1\}^n$ are independent random variables with $\Pr[A_i = 0], \Pr[B_i = 0] \geq p$ for all i , then*

$$H[A \cup B] \geq \frac{p}{2\phi}(H[A] + H[B]).$$

¹We let p denote the probability of failure here (rather than the convention of having p denote the probability of success) in order to make some of the later algebra cleaner.

Proof. By the Chain Rule and Data Processing Inequality, we have

$$H[A \cup B] = \sum_{i \leq n} H[(A \cup B)_i | (A \cup B)_{<i}] \geq \sum_{i \leq n} H[(A \cup B)_i | A_{<i}, B_{<i}]. \quad (17)$$

Fix some $1 \leq i \leq n$. For each $x \in \{0, 1\}^i$, define $p(x) = \Pr[A_i = 0 | A_{<i} = x]$ and $q(y) = \Pr[B_i = 0 | B_{<i} = y]$. By Lemma 15.4 and the independence of A, B , we have for all x, y that

$$H[(A \cup B)_i | A_{<i} = x, B_{<i} = y] = H(p(x)q(y)) \geq \frac{1}{2\phi} (p(x)H(q(y)) + q(y)H(p(x))).$$

Multiplying this inequality by $\Pr[A_{<i} = x, B_{<i} = y] = \Pr[A_{<i} = x] \cdot \Pr[B_{<i} = y]$ and summing over all x, y gives

$$\begin{aligned} H[(A \cup B)_i | A_{<i}, B_{<i}] &\geq \frac{1}{2\phi} \left(\mathbb{E}_{A_{<i}, B_{<i}} [p(x)H(q(y))] + \mathbb{E}_{A_{<i}, B_{<i}} [q(y)H(p(x))] \right) \\ &= \frac{1}{2\phi} (\mathbb{E}_{A_{<i}} [p(x)] \cdot \mathbb{E}_{B_{<i}} [H(q(y))] + \mathbb{E}_{B_{<i}} [q(y)] \cdot \mathbb{E}_{A_{<i}} [H(p(x))]) \\ &= \frac{1}{2\phi} (\Pr[A_i = 0] \cdot H[B_i | B_{<i}] + \Pr[B_i = 0] \cdot H[A_i | A_{<i}]) \\ &\geq \frac{p}{2\phi} (H[B_i | B_{<i}] + H[A_i | A_{<i}]), \end{aligned}$$

where the second equality used the definition of conditional entropy and the definition of $q(y)$, and the last inequality used the hypothesis $\Pr[A_i = 0], \Pr[B_i = 0] \geq p$. Combining this with (17) gives

$$H[A \cup B] \geq \frac{p}{2\phi} \sum_{i \leq n} H[A_i | A_{<i}] + H[B_i | B_{<i}] = \frac{p}{2\phi} (H[A] + H[B]),$$

with this last equality using the Chain Rule. \square

We can now complete the proof of Theorem 15.3 with $c = 1 - \phi = \frac{3-\sqrt{5}}{2} \approx .381$.

Proof of Theorem 15.3. Let $c = 1 - \phi$ and identify $A, B, A \cup B$ by their characteristic vectors in $\{0, 1\}^n$. Note that our hypothesis $|\mathcal{F}| > 1$ implies $H(A), H(B) > 0$. Because $\deg(x) < c|\mathcal{F}|$ for all x , we have $\Pr[A_i = 0], \Pr[B_i = 0] > 1 - c = \phi$ for all i . Hence the previous lemma (together with $H(A), H(B) > 0$) implies

$$H[A \cup B] > \frac{1}{2} (H[A] + H[B]) = H[A],$$

proving the result. \square

Before moving on, let us take a few moments to comment on some strengthenings of Theorem 15.2. As noted earlier, it turns out that $c = \frac{3-\sqrt{5}}{2}$ is the best c one can take so that Theorem 15.3 remains true. Moreover, this value of c (rather than $c = 1/2$) turns out to be the optimal value for an analogous “approximate” version of the union closed conjecture, i.e. a variant where we only require that most of the unions $A \cup B$ lie in \mathcal{F} ; see [24] for more on this.

Given the above, some sort of genuinely new approach is needed to get beyond this $c = \frac{3-\sqrt{5}}{2}$ barrier for the original Union-Closed Sets Conjecture, and such an approach was found by Sawin [99]. In essence, his idea is to look at three random sets $A, B, C \in \mathcal{F}$ such that B is independent of A and C but (crucially) A, C are correlated in some way. Again, \mathcal{F} being union closed implies that we must have $H[A \cup B], H[A \cup C] \leq H[A]$, and hence that

$$\alpha H[A \cup B] + (1 - \alpha) H[A \cup C] \leq H[A]$$

for all α . Sawin then argues that this inequality fails to hold for some α if every element is in at most a $\frac{3-\sqrt{5}}{2} + \epsilon$ proportion of the elements of \mathcal{F} , giving the desired improvement. Cambie [23] obtained the best bounds one can get via this approach of Sawin, showing that one can take $c \approx .38235$ (cf the previous value of $c = \frac{3-\sqrt{5}}{2} \approx .38197$).

Part V

Hypergraph Containers

This part is heavily based off of lecture notes by Balogh [9]. Throughout this section we let $\mathcal{I}(H)$ denote the set of independent sets of a hypergraph H and $\mathcal{I}_m(H)$ the set of independent sets of size m . We adopt the notation $\binom{n}{\leq k}$ to denote the number of subsets of $[n]$ of size at most k . Many of the bounds in this part will be rough approximations to the truth in order to emphasize the intuition of the results and techniques rather than the nitty gritty detail that is actually required.

16 Introduction

Many problems in extremal combinatorics can be stated in terms of independent sets of hypergraphs. For example, one can define \mathcal{H}_n^{AP} to be the 3-graph on $[n]$ where every triple $S \subseteq [n]$ is a hyperedge if and only if S is a 3-term arithmetic progression. Thus Roth's theorem is equivalent to saying $\alpha(\mathcal{H}_n^{AP}) = o(n)$. Similarly one can define \mathcal{H}_n^Δ to be the 3-graph whose vertex set is $E(K_n)$ and whose hyperedges are triples of edges in K_n which form a triangle. Independent sets of \mathcal{H}_n^Δ are triangle-free subgraphs of K_n , so Mantel's theorem says $\alpha(\mathcal{H}_n^\Delta) = \lfloor n^2/4 \rfloor$.

This part is dedicated to a powerful method of upper bounding the size of $\mathcal{I}(H)$. Observe that for any hypergraph H we have

$$2^{\alpha(H)} \leq |\mathcal{I}(H)| \leq \binom{n}{\alpha(H)} 2^{\alpha(H)} \leq (2n)^{\alpha(H)}.$$

In particular, the upper bound follows because every independent set is a subset of a set of size $\alpha(H)$. More generally, we say that a collection \mathcal{C} of subsets $C \subseteq V(H)$ is a set of *containers* for H if every independent set $I \in \mathcal{I}(H)$ is a subset of some $C \in \mathcal{C}$. If such a set of containers exists, then

$$|\mathcal{I}(H)| \leq \sum_{C \in \mathcal{C}} 2^{|C|} \leq |\mathcal{C}| 2^{\max_{C \in \mathcal{C}} |C|}. \quad (18)$$

Thus we will get an effective upper bound on $|\mathcal{I}(H)|$ whenever we can find a small collection of containers, each of which are relatively small. Sometimes we will be interested in finding the number of independent sets of H of size m . The same reasoning as above gives

$$\binom{\alpha(H)}{m} \leq |\mathcal{I}_m(H)| \leq |\mathcal{C}| \binom{\max_{C \in \mathcal{C}} |C|}{m}. \quad (19)$$

The method of hypergraph containers gives a systematic way of obtaining such a collection of containers whenever H satisfies some fairly mild conditions. The main condition we need is that the codegrees of H to be relatively small, and in practice this often corresponds to having some notion of supersaturation.

17 Graph Containers

While the general method of containers involves bounding independent sets of *hypergraphs*, one can get pretty far by only considering independent sets of *graphs*. To this end we prove the following graph container lemma, which will be the main workhorse for the rest of this section. Recall that a collection \mathcal{C} of subsets $C \subseteq V(G)$ is a set of *containers* for G if every independent set $I \in \mathcal{I}(G)$ is a subset of some $C \in \mathcal{C}$.

Lemma 17.1. *Let G be an n -vertex graph and $t > 0$ a positive number. There exists a collection \mathcal{C} of containers such that*

$$(a) \quad |\mathcal{C}| \leq \binom{n}{\leq n/t}.$$

$$(b) \quad \Delta(G[C]) < t - 1 \text{ for all } C \in \mathcal{C}.$$

In other words, there exists a small set of containers \mathcal{C} such that each $C \in \mathcal{C}$ is “small” in the sense that it induces a graph with small maximum degree.

Proof. Our proof will be algorithmic: we construct a (deterministic) algorithm which takes as input a set $I \subseteq V(G)$ and which outputs a pair $(S(I), A(I))$ such that $S(I) \subseteq I \subseteq S(I) \cup A(I)$, and we will ultimately use $\{S(I) \cup A(I) : I \in \mathcal{I}(G)\}$ as our set of containers. We now describe the algorithm.

Fix an arbitrary ordering of $V(G)$. As input we take in an independent set $I \subseteq V(G)$. We initially set $S = \emptyset$ and $A = V(G)$ (the former corresponds to a set of “selected” vertices which are in I , and the latter to the set of “available” vertices which could possibly be in I given the current stage of the algorithm). The algorithm proceeds as follows:

Step 1 If $\Delta(G[A]) < t - 1$, output $(S(I), A(I))$. Otherwise proceed to Step 2.

Step 2 Let v be the vertex of maximum degree in $G[A]$, with ties being broken according to the fixed ordering of $V(G)$. If $v \notin I$, then set $A = A - v$ and repeat Step 1. Otherwise proceed to Step 3.

Step 3 Set $A = A - v - N_{G[A]}(v)$, $S = S \cup \{v\}$. Proceed to Step 1.

Let’s reiterate what’s going on here. It’s not difficult to show inductively that we always have $I \subseteq S \cup A$, so $S \cup A$ serves as a container set for I , and we would like to trim this set down as much as possible. We do this by selecting a vertex $v \in A \cap I$ and adding it to S . If v has large degree in $G[A]$, then v being in the independent set I means that its many neighbors are not, so we get to remove all of these vertices from A while maintaining $I \subseteq S \cup A$. In particular, since we keep going so long as $G[A]$ has large maximum degree, we know at each step of this process that we’re removing many vertices from A .

Define

$$\mathcal{C} = \{S(I) \cup A(I) : I \in \mathcal{I}(G)\},$$

which is a set of containers since $I \subseteq S(I) \cup A(I)$ at every step of the algorithm. Since we terminate the algorithm precisely when $\Delta(G[A(I)]) = \Delta(G[S(I) \cup A(I)]) < t - 1$ (the equality holds since $S(I)$ has no neighbors in $S(I) \cup A(I)$), (b) holds. It thus remains to verify (a). To do this, we note the following which is easy to verify.

Claim 17.2. *Let I_1, I_2 be two independent sets and let $(S_1, A_1), (S_2, A_2)$ be their outputs from the algorithm. If $S_1 = S_2$, then $A_1 = A_2$.*

This claim implies that given $S(I)$, the container $S(I) \cup A(I)$ is uniquely determined¹. In particular, if we always have $|S(I)| \leq n/t$, then the number of containers will be at most $\binom{n}{\leq n/t}$. And indeed, each round of the algorithm has $\Delta(G[A]) \geq t - 1$, so every time a vertex is added to S at least $1 + (t - 1) = t$ vertices are removed from A . In particular, at most n/t vertices can be added to S , giving the result. \square

Actually, a closer inspection of the proof gives the following.

Lemma 17.3. *Let G be a graph on n vertices and $t \in \mathbb{R}$. There is a collection \mathcal{C} of containers and functions*

$$f : \mathcal{I}(G) \rightarrow \binom{V(G)}{\leq n/t}, \quad g : \binom{V(G)}{\leq n/t} \rightarrow \mathcal{C}$$

such that the following hold.

- (a) *The function g is a surjection. In particular, $|\mathcal{C}| \leq \binom{n}{\leq n/t}$.*
- (b) *We have $\Delta(G[C]) < t - 1$ for all $C \in \mathcal{C}$.*
- (c) *For every $I \in \mathcal{I}(G)$ we have*

$$f(I) \subseteq I \subseteq g(f(I)).$$

Proof. Consider the exact same algorithm as before. Define $f(I) = S(I)$ and $g(S) = C(S)$ (if $S \neq S(I)$ for any I , then assign g arbitrarily). It's not hard to check that this works. \square

The extra source of power of this lemma is that for each $I \in \mathcal{I}$ we are given some set $S = f(I)$ contained in I . In many examples this extra information is needed to get tight upper bounds when counting independent sets, though for pedagogical purposes we will often work with the simpler Lemma 17.1 to get close to tight results.

In the coming subsections we'll show how to use Lemma 17.1 to solve several combinatorial problems. All of the proofs will be very similar to each other, though they'll become increasingly sophisticated as we go along.

Before going on, let us briefly note that there are many variants of Lemma 17.1 that one can prove using a similar approach. These variants of Lemma 17.1 are both a blessing and a curse since they give many options for how to solve a given problem (and it isn't always clear which is best).

¹Because of this, S is often called a "certificate" or "fingerprint" of I .

17.1 Regular Graphs

Our first application of Lemma 17.1 will be to count the number of independent sets in a d -regular graph. As a point of reference, it is not difficult to show that if G consists of $n/2d$ disjoint copies of $K_{d,d}$, then

$$|\mathcal{I}(G)| = (2^{d+1} - 1)^{n/2d} = 2^{n/2 + n/2d + o(n)}.$$

Thus for d -regular graphs, we can't possibly hope to prove an upper bound on $|\mathcal{I}(G)|$ stronger than roughly $2^{n/2}$ when d is large. We can prove that this is close to best possible using containers.

Theorem 17.4. *Let G be a d -regular n -vertex graph with $\log n \ll d \ll n/2$. Then*

$$|\mathcal{I}(G)| \leq 2^{n/2 + o(n)}.$$

In fact, it turns out that $|\mathcal{I}(G)| \leq (2^{d+1} - 1)^{n/2d}$ for all d -regular n -vertex graphs. This was proven for bipartite graphs by Kahn [69] using entropy, and the problem was solved in full by Zhao [110]. As far as I'm aware, the proof of Theorem 17.4 presented here is due to Balogh.

As a first step to proving Theorem 17.4, we will apply Lemma 17.1 to our graph G to get a collection of containers \mathcal{C} . We would like to conclude the result by the observation from (18):

$$|\mathcal{I}(G)| \leq \sum_{C \in \mathcal{C}} 2^{|C|} \leq |\mathcal{C}| 2^{\max_{C \in \mathcal{C}} |C|},$$

but there's an issue with this. Namely, Lemma 17.1 only tells us that each $C \in \mathcal{C}$ induces a graph in G with small maximum degree. For a general graph this tells us nothing about $|C|$, but fortunately in d -regular graphs, $G[C]$ having small maximum degree is only possible if C is small. The following states a precise version of the contrapositive of the previous sentence.

Lemma 17.5. *For any $\epsilon > 0$, if G is a d -regular graph and $C \subseteq V(G)$ with $|C| = n/2 + \epsilon n$, then $\Delta(G[C]) \geq 2\epsilon d$.*

This lemma is a form of supersaturation: a d -regular graph can have a subset of size $n/2$ with $G[C]$ empty (e.g. if G is bipartite), but if C is just a bit larger than this, then it must have relatively high maximum degree. As we will see, supersaturation results are almost always a necessary ingredient for applying the method of containers.

Proof. Because the maximum degree is always at least the average degree, we have

$$\Delta(G[C]) \geq 2e(G[C])/|C| \geq 2e(G[C])/n$$

, so it will suffice to show that $e(G[C])$ is large. To do this, we let $\overline{C} = V(G) \setminus C$ and note that

$$d|C| = \sum_{v \in C} d(v) = 2e(G[C]) + e(C, \overline{C}) \leq 2e(G[C]) + d|\overline{C}|.$$

Because $|C| = n/2 + \epsilon n$ and $|\overline{C}| = n/2 - \epsilon n$, in total this implies

$$2e(G[C]) \geq 2\epsilon dn.$$

Combining this with the observation at the start gives the result. \square

Corollary 17.6. *For all t , if G is an n -vertex d -regular graph, then there exists a set of containers \mathcal{C} with $|\mathcal{C}| \leq \binom{n}{\leq n/t}$ and $|C| \leq \frac{1}{2}n + \frac{t}{d}n$ for all $C \in \mathcal{C}$.*

Proof. Let \mathcal{C} be a set of containers as guaranteed by Lemma 17.1. Because $\Delta(G[C]) < t-1 \leq t$, Lemma 17.5 implies that $|C| \leq \frac{1}{2}n + \frac{t}{d}n$. \square

With this we can prove Theorem 17.4.

Proof of Theorem 17.4. At this point all we need to do is use (18) after applying Corollary 17.6 with a carefully chosen value of t . Note that

$$|\mathcal{C}| \approx \binom{n}{n/t} \approx 2^{n \log(t)/t},$$

and we already know $2^{\max |C|} \approx 2^{\frac{1}{2}n + \frac{t}{d}n}$. Thus to minimize $|\mathcal{C}| \cdot 2^{\max |C|}$ we should choose t so that $\frac{t}{d} \approx \log(t)/t$, and in particular $t = \sqrt{d \log n}$ is a reasonable choice. One can verify with a more formal argument that this does indeed give the desired result after applying (18). \square

We note that the statement of Corollary 17.6 and the optimization of t in the proof of Theorem 17.4 is in some sense independent¹ of the problem of determining $|\mathcal{I}(G)|$ for G a d -regular graph. That is, these results are effective for other problems which involve counting independent sets of d -regular graphs.

For example, recall that a q -coloring of a graph G is a map $\chi : G \rightarrow [q]$ such that $\chi(u) \neq \chi(v)$ whenever $uv \in E(G)$. Equivalently, a q -coloring is a partition of $V(G)$ into independent sets I_1, \dots, I_q . With this latter formulation, we can use containers to get an effective bound on the number of q -colorings of G , which we'll denote by $X_q(G)$.

Again, let's consider a test case to figure out how strong of a bound we could possibly hope to prove. Let G be $n/2d$ disjoint copies of $K_{d,d}$. We know that G has close to as many independent sets as it could possibly have, so it seems plausible that it would have many q -colorings as well. In particular, one can prove that $X_q(G) \approx (q/2)^n$, and once again we can prove that this is essentially best possible.

Theorem 17.7 ([55]). *Let G be an n -vertex d -regular graph and q an integer such that $q^2 \log n \ll d$. Then*

$$X_q(G) \leq (q/2 + o(1))^n.$$

We note that a stronger result was proven by Galvin [55] with a somewhat more involved proof.

Proof. By the same reasoning as in Theorem 17.4, there exists a set of containers \mathcal{C} for G such that $|\mathcal{C}| \approx 2^{\sqrt{\frac{\log n}{d}}n}$ and $|C| \approx \frac{1}{2}n$ for each $C \in \mathcal{C}$. Consider all vectors of the form (C_1, \dots, C_q) with $C_i \in \mathcal{C}$, noting that the number of such vectors is at most $|\mathcal{C}|^q = 2^{o(n)}$.

Observe that every q -coloring can be identified by a vector (I_1, \dots, I_q) where each I_j is an independent set and $\bigcup I_j = V(G)$. Each of these vectors is “contained” in some “container

¹Ha.

vector" (C_1, \dots, C_q) with $C_j \in \mathcal{C}$ in the sense that $I_j \subseteq C_j$ for all j . Thus it's enough to count how many q -colorings each container vector contains.

A naive upper bound for the number of q -colorings contained in (C_1, \dots, C_q) is roughly $2^{qn/2}$ since this is the number of ways to choose an independent set from each C_i . This bound is too weak, so we have to utilize the extra information that the I_j partition $V(G)$.

To this end, assume $V(G) = \{v_1, \dots, v_n\}$. Given (C_1, \dots, C_q) , let a_i be the number of containers C_j with $v_i \in C_j$. It's not difficult to see that the number of q -colorings contained in this vector is then at most $\prod a_i$, and by the AMGM inequality this is at most $(\sum a_i/n)^n = (\sum |C_j|/n)^n$.

Each of the q containers has size at most roughly $n/2 + \sqrt{\frac{\log n}{d}}n$, so this gives the desired result. \square

17.2 A Randomized Sperner's Theorem

Throughout this subsection we fix an integer n and define $N := 2^n$ and $m = \binom{n}{\lfloor n/2 \rfloor}$. An *antichain* of $[n]$ is a subset $S \subseteq 2^{[n]}$ such that $A \not\subseteq B$ for any distinct $A, B \in S$. For example, $\binom{[n]}{k}$ is an antichain for all k . A famous result of Sperner's says that an antichain of $[n]$ has size at most m .

Our first goal is to count the number of antichains of $[n]$. To do this, we form a graph where independent sets correspond to antichains. Let G_N denote the graph whose vertex set is $2^{[n]}$ and where A, B are adjacent to each other if either $A \subseteq B$ or $B \subseteq A$. Analogous to Lemma 17.5, we need a supersaturation lemma for G_N which says that any collection of vertices that is much larger than m induces many edges. In particular, the following suffices.

Lemma 17.8 ([74, 11]). *If $C \subseteq 2^{[n]}$ has $|C| > (1 + \epsilon)m$ with $0 < \epsilon \leq 1/3$, then $e(G_N[C]) \geq \epsilon mn/2$.*

We won't prove this, but we will briefly comment on some intuition for the result. Intuitively, if you want to build a set of size $(1 + \epsilon)m$ which induces few edges, then a good place to start is with the middle layer $\binom{[n]}{\lfloor n/2 \rfloor}$ since this is a maximum independent set. From there one could greedily choose ϵm sets which have as few neighbors as possible in this middle layer, and in particular choosing them all from $\binom{[n]}{\lfloor n/2 \rfloor + 1}$ gives a total of $(\lfloor n/2 \rfloor + 1) \cdot \epsilon m \geq \epsilon mn/2$ edges. Kleitman [74] proved that this is indeed the best construction, and the exact numerical computation¹ was done by Balogh, Mycroft, and Treglown [11].

With our supersaturation lemma in hand, we can easily prove the following result of Kleitman.

Theorem 17.9 ([75]). *The number of antichains of $[n]$ is $2^{m+o(m)}$.*

Proof. We obtain a set of containers \mathcal{C} for G_N by applying lemma 17.1 to G_N with a parameter t to be determined later. Let ϵ be such that $(1 + \epsilon)m = \max_{C \in \mathcal{C}} |C|$ and let C be a container

¹The numbers in Lemma 17.8 are slightly different from those that appear in [11], but it's not difficult to refine their proof to give this result.

achieving this bound. By Lemma 17.8 we have¹

$$\Delta(G_N[C]) \geq 2e(G_N[C])/|C| \geq \epsilon mn/(1 + \epsilon)m \approx \epsilon n.$$

By assumption this quantity is at most t , or equivalently we roughly have $\max_{|C| \in \mathcal{C}} |C| \leq (1 + t/n)m$. By (18) we have an upper bound of roughly

$$\binom{N}{N/t} 2^{(1+t/n)m} \approx 2^{\frac{N}{t} \log(t) + (1+t/n)m}.$$

This quantity is optimized when $N \log(t)/t \approx tm/n$. We have $m \approx N/\sqrt{n}$, so in total we want $t \approx n^{3/4}/\sqrt{\log n}$, and one can verify that this choice of t gives the desired bound. \square

We next prove a random version of Sperner's theorem. The setup is as follows. Choose a subset $R_p \subseteq 2^{[n]}$ by including each set in R_p independently and with probability p . How large is the size of a largest antichain in R_p (in expectation)? In terms of our graph G_N , this is equivalent to computing $\mathbb{E}[\alpha(G_N[R_p])]$.

Let's consider some simple cases first. If $p = 1$, then $G_N[R_p] = G_N$ and we know its independence number is m . Somewhat more generally, we always have the lower bound $\mathbb{E}[\alpha(G_N[R_p])] \geq pm$ since this is the expected size of the set $\binom{[n]}{\lfloor n/2 \rfloor} \cap R_p$. However (as will often be the case), the behavior of $\mathbb{E}[\alpha(G_N[R_p])]$ changes considerably when p is very small.

For example, if $p^2 3^n \ll p 2^n$, then asymptotically we have $\mathbb{E}[\alpha(G_N[R_p])] \sim |R_p|$ by a simple deletion argument. Even above the deletion threshold it is possible to improve on the trivial lower bound. Indeed, construct an independent set I by keeping each vertex in $\binom{[n]}{\lfloor n/2 \rfloor} \cap R_p$ together with all the vertices in $\binom{[n]}{\lfloor n/2 \rfloor - 1} \cap R_p$ which are not contained in any of the vertices of $\binom{[n]}{\lfloor n/2 \rfloor} \cap R_p$. The expected number of vertices we get from this first part is $p \binom{n}{\lfloor n/2 \rfloor}$, and from the second is² $p(1 - p)^{n+1-\lfloor n/2 \rfloor} \binom{n}{\lfloor n/2 \rfloor - 1}$. In particular, if $p = c/n$ for a fixed constant c , then this asymptotically gives $(1 + \epsilon)pm$ for some $\epsilon > 0$.

It turns out that for larger p we do have $\mathbb{E}[\alpha(G_N[R_p])] \sim pm$. More precisely we have the following due to Balogh, Mycroft, and Treglown [11].

Theorem 17.10 ([11]). *For any $\epsilon > 0$, there exists a constant c so that if $p > c/n$, then a.a.s. $\alpha(G_N[R_p]) \leq (1 + \epsilon)pm$.*

Roughly speaking the approach we would like to use is as follows. Observe that $\alpha(G_N[R_p]) \geq k$ if and only if $G_N[R_p]$ contains an independent set of size k . The expected number of such sets in $G_N[R_p]$ is exactly $p^k \mathcal{I}_k(G_N)$, and if this quantity is small then we can conclude the result by Markov's inequality. Thus to solve this problem (and in general to solve extremal problems in random sets), we need to get effective upper bounds on $\mathcal{I}_k(G_N)$.

Unfortunately a naive application of Lemmas 17.1 and 17.8 together with (19) turns out to be too weak. The bottleneck here is the supersaturation result from Lemma 17.8. While the

¹Implicitly this assumes $\epsilon \leq 1/3$. One can get around this by taking $C' \subseteq C$ a set of size exactly $4m/3$, but ultimately this computation is just to obtain intuition for what value t should be.

²To keep a vertex it has to be in R_p and all its neighbors in $\binom{[n]}{\lfloor n/2 \rfloor - 1}$ have to be out.

stated bound is essentially tight for $|C| = (1 + \epsilon)m$ with $0 < \epsilon \leq 1/3$ and n sufficiently large, the bound is not tight when e.g. $|C| = (2 + \epsilon)m$. Intuitively in this case the extremal example should come from taking two middle layers together with ϵm of the layer right above these two. In particular, each of the ϵm vertices will have degree about n^2 , so we expect around ϵmn^2 edges in $G_N[C]$. And indeed, this is the case.

Lemma 17.11 ([74, 11]). *If $C \subseteq 2^{[n]}$ has $|C| > (2 + \epsilon)m$ with $0 < \epsilon \leq 1/3$, then $e(G_N[C]) \geq \epsilon mn^2/9$.*

With this we can prove the main result.

Proof of Theorem 17.10. The key idea is to apply the container lemma twice using the two different levels of supersaturation from Lemmas 17.8 and 17.11. In particular, let \mathcal{C}_1 be a set of containers coming from Lemma 17.1 using G_N and some t_1 , and for each $C_1 \in \mathcal{C}_1$, let $\mathcal{C}_2(C_1)$ be a set of containers coming from Lemma 17.3 using $G_N[C_1]$ and some t_2 .

We first want to choose t_1 so that each $C_1 \in \mathcal{C}_1$ has size roughly $\alpha(G_N) = m$. Observe that if $|C_1| > 3m$, then by Lemma 17.11, every $C'_1 \subseteq C_1$ of size $3m$ has

$$\Delta(G_N[C_1]) \geq \Delta(G_N[C'_1]) \geq 2e(G_N[C'_1])/|C'_1| \geq n^2/81.$$

Thus if we take $t_1 = n^{1.99}$ in Lemma 17.1, we find for n sufficiently large that $|C_1| \leq 3m$ for all $C_1 \in \mathcal{C}_1$.

We now want to choose t_2 so that each $C_2 \in \mathcal{C}_2(C_1)$ has size very close to m . Let $G' = G_N[C_1]$. If $|C_2| > (1 + \gamma)m$ with $\gamma \leq 1/3$ we find that

$$\Delta(G'[C_2]) \leq 2e(G'[C_2])/|C_2| \leq \gamma n/3.$$

With some foresight we take $t_2 = \epsilon n/12$ to guarantee that $|C_2| \leq (1 + \epsilon/4)m$ for all $C_2 \in \mathcal{C}_2(C_1)$ by Lemma 17.8.

Recall that we want to show with high probability no independent set I of size $(1 + \epsilon)pm$ lies in the random set R_p . To this end, we note that we can identify each I with a pair (C_1, S_2) where

- $C_1 \in \mathcal{C}_1$ contains I ,
- S_2 is the set $f(I)$ from Lemma 17.3, i.e. $|S_2| \leq |C_1|/t_2$, $S_2 \subseteq I$, and S_2 determines a set $C_2 \supseteq I$ in $\mathcal{C}_2(C_1)$.

Given this pair, if $I \subseteq R_p$ has size at least $(1 + \epsilon)m$, then (1) $S_2 \subseteq R_p$ since $S_2 \subseteq I$, and (2)

$$|R_p \cap (C_2 \setminus S_2)| \geq (1 + \epsilon)pm - |S_2|,$$

since $C_2 \setminus S_2$ contains $I \setminus S_2$. Observe that for $p = c/n$ with $c \gg \epsilon^{-2}$ we have

$$|S_2| \leq \frac{|C_1|}{t_1} \approx \frac{m}{\epsilon n} \leq \frac{\epsilon pm}{2}.$$

With this in mind, we define $A(S_2)$ to be the event that $S_2 \subseteq R_p$ and $B(S_2)$ to be the event that $|R_p \cap (C_2 \setminus S_2)| \geq (1 + \epsilon/2)pm$. Observe that $A(S_2)$ and $B(S_2)$ are independent events, so by a union bound over all pairs (C_1, S_2) , we see that the probability that $\alpha(G_N[R_p]) \geq (1 + \epsilon)pm$ is at most

$$\sum_{C_1 \in \mathcal{C}_1} \sum_{S_2: |S_2| \leq |C_1|/t_1} \Pr[A(S_2)] \cdot \Pr[B(S_2)]. \quad (20)$$

We have $\Pr[A(S_2)] = p^{|S_2|}$. By a Chernoff bound it is not difficult to show that $\Pr[B(S_2)] \leq e^{-c'\epsilon^2 pm}$ for some $c' > 0$. Thus if we fix some C_1 in the first term of the sum we get

$$\sum_{s \leq |C_1|/t_2} \binom{|C_1|}{s} p^s \cdot e^{-c'\epsilon^2 pm} \leq \sum_{s \leq |C_1|/t_2} (ep|C_1|/s)^s \cdot e^{-c'\epsilon^2 pm} \approx (p \cdot \epsilon n / 12)^{12|C_1|/\epsilon n} \cdot e^{-c'\epsilon^2 pm} \leq (\epsilon c)^{36m/\epsilon n} \cdot e^{-c'\epsilon^2 cm/n},$$

where the approximation only looked at the term with $s = |C_1|/t_2$. Note that for $c \gg \epsilon^{-4}$ the second term dominates, so this bound is roughly

$$e^{-c'\epsilon^2 cm/n}. \quad (21)$$

Note that this is the critical place where we used the two applications of the container lemma: if we only applied the container lemma once to $V(G_N)$ instead of using C_1 , then the first term here would be roughly $e^{N/\epsilon n}$ instead of $e^{m/\epsilon n}$, which would dominate the expression.

Returning to (20), we sum the bound of (21) for each element in C_1 , which multiplies (21) by

$$\binom{N}{\leq N/t_1} \approx e^{N \log(t_1)/t_1} \approx e^{mn^{-1.49} \log(n)}.$$

This is much smaller than $e^{-c'\epsilon^2 cm/n}$, so the probability in (20) tends to 0 as n tends towards infinity as desired. \square

We note that one can get almost as strong a result if one only uses Lemma 17.1, i.e. if one doesn't use the more refined Lemma 17.3. Indeed, the main consequence of using this refined lemma was the extra term $\Pr[A(S_2)] = p^s$ appearing in (20). If one omits this term, then the same proof will go through provided $p = c \log n/n$. This is a common phenomenon in containers: if you don't use the fact you have certificates $S \subseteq I$, then you'll end up with a bound which is worse by a log factor.

17.3 Counting Sidon Sets

This is a neat example where you use roughly $\log n$ iterated supersaturation lemmas to get the right result. I may write this at some point, but in any case it will be nearly identical to Balogh's notes [9].

18 A Proof of an r -Uniform Container Lemma

In the previous section, we saw how Lemma 17.1 allowed us to effectively count the number of independent sets in “sufficiently nice” graphs. In this section we present a proof of a hypergraph container lemma which applies to “sufficiently nice” hypergraphs, but two things should be noted.

The first is that there are many different variants of hypergraph container lemmas, though most of them are quite similar and broadly speaking apply only to hypergraphs with small codegrees.

The second is that, quite frankly, one doesn’t need to know the proof of the container lemma to use it or its variants. As such, the reader may just want to glance at the definition below, and then skip over to latter sections to see some nice applications before jumping back over here whenever they want to see the full proof.

18.1 An Informal Discussion

The hypergraph container lemma we prove comes from [?] (though we deviate somewhat from their notation). We’ll formally state this as Theorem 18.5 below, but roughly our goal will be to prove the following. Recall that $\Delta_\ell(H)$ denotes the maximum ℓ -degree of H , i.e. the maximum number of edges containing a given set of ℓ vertices.

Proposition 18.1 (Informal). *If H is an r -graph such that for all $1 \leq \ell \leq r$ we have*

$$\Delta_\ell(H) \ll q^{\ell-1} \frac{e(H)}{v(H)},$$

then there exists $\delta > 0$, $\mathcal{S} \subseteq \binom{V(H)}{\leq q v(H)}$, and functions $f : \mathcal{S} \rightarrow \binom{V(H)}{\leq (1-\delta)v(H)}$ and $g : \mathcal{I}(H) \rightarrow \mathcal{S}$ such that

$$g(I) \subseteq I \subseteq f(g(I)) \cup g(I).$$

In other words, if H has small codegrees, then one can find a set of small certificates \mathcal{S} which are each associated with a container $f(S)$ which is of size at most $(1 - \delta)v(H)$.

The proof of Proposition 18.1 will in essence be a proof by induction on the uniformity r , and the inductive step of the proof uses an algorithm which is similar to the one used in Lemma 17.1.

Definition 1 (Informal). The *Scythe Algorithm* takes as input a pair (H_{k+1}, I) with H_{k+1} a $(k + 1)$ -uniform hypergraph and $I \subseteq V(H_{k+1})$ an independent set. It then outputs a triple (H_k, A_k, S_k) with H_k a k -uniform hypergraph such that $I \subseteq V(H_k)$ is an independent set with $I \subseteq A_k \cup S_k$, and S_k is a small set which uniquely determines H_k and A_k given H_{k+1} . Moreover, if H_{k+1} is “nice”, then either H_k will be “nice” or A_k will be small.

Given such an algorithm, we can start with any “nice” r -uniform hypergraph H_r and independent set I . We then repeatedly apply this algorithm until we get some H_k which is not “nice”, at which point $\bigcup_{i \geq k} S_i$ is a small certificate which determines a small container $A_k \cup \bigcup_{i \geq k} S_i$ for I .

While it's not a priori clear what the “nice” conditions should be, they should in particular guarantee that H_{k+1} has few independent sets, as otherwise there's no hope of this method being effective. In particular, a reasonable set of conditions is to enforce that H_{k+1} has many edges and relatively low codegrees, and this will ultimately be the conditions that we use.

Now that we know what we want our algorithm to do, how should it actually work in practice? Perhaps the most naive approach is to do what we did in Lemma 17.1, where we iteratively select the vertex u of $I \cap H_{k+1}$ which has the largest degree and then adds this to S_k . Once we identify such a u , we know that I does not contain any k -set of the form $e \setminus \{u\}$ for any $e \ni u$ which is an edge in H_{k+1} , so it is natural to make all of these k -sets edges of H_k . If we do this repeatedly, then A_k will be relatively small and I will be an independent set of H_k . Moreover, if we enforce from the start that we'll run this procedure at most s times, then we will have $|S_k| \leq s$, giving a small certificate.

Unfortunately we have to be more careful than this. Namely, we need to ensure that H_k is “nice”, and in particular that it has small codegrees. As it currently stands this might not work out, e.g. there may be some ℓ -set T which is in many edges containing vertices of S_k . To get around this, we define $D_\ell(H_k, \Delta)$ to be the set of “dangerous” ℓ -sets of $V(H_k)$ which have degree at least $\Delta/2$, where we think of Δ as being the maximum ℓ -degree we want H_k to have (which is analogous to the t parameter used in the algorithm of Lemma 17.1).

We now adjust our naive algorithm by making it so that whenever a set T gets added to $D_\ell(H_k, \Delta)$, we delete from H_{k+1} all of the edges that contain T . This ensures that T never passes over the enforced codegree threshold. With this it turns out that our algorithm will succeed.

18.2 A Formal Algorithm

Motivated by our discussion in the previous section, we make the following definitions. For any hypergraph H' , we define the *max-degree order* on $V(H')$ as follows. Fix an arbitrary ordering of $V(H')$. For each integer j , recursively define u_j to be the maximum-degree vertex in the hypergraph $H'[V(H') \setminus \{u_1, \dots, u_{j-1}\}]$ with ties broken based on the ordering of $V(H')$. The max-degree order is then the ordering u_1, u_2, \dots , and for all j we define $W_{H'}(u_j) = \{u_1, \dots, u_j\}$.

For any k -uniform hypergraph H' , integer $\ell \leq k$, and real number Δ , we define

$$D_\ell(H', \Delta) = \left\{ T \in \binom{V(H')}{\ell} : \deg_{H'}(T) \geq \frac{1}{2}\Delta \right\}.$$

Definition 2. The Scythe Algorithm is defined as follows. It takes as input a $(k+1)$ -uniform hypergraph H_{k+1} with $k \geq 1$, an independent set $I \subseteq V(H_{k+1})$, and parameters $s, \Delta_1^k, \dots, \Delta_k^k$.

At the start of the algorithm, we set $H_{k+1}^{(0)} = H_{k+1}$, $S_k^{(0)} = \emptyset$, and we let $H_k^{(0)}$ be the empty hypergraph on $V(H_{k+1})$. For $j = 0, \dots, s-1$, the algorithm proceeds as follows:

Step 1: If $I \cap V(H_{k+1}^{(j)}) = \emptyset$, then set $H_k = H_k^{(0)}$, $A_k = \emptyset$, $S_k = S_k^{(j)}$. If this happens, stop the algorithm and output (H_k, A_k, S_k) .

Step 2: Let u_j be the vertex of $I \cap V(H_{k+1}^{(j)})$ which is first according to the max-degree ordering of $H_{k+1}^{(j)}$. Set $S_k^{(j+1)} = S_k^{(j)} \cup \{u_j\}$.

Step 3: Let $H_k^{(j+1)}$ be the hypergraph on $V(H)$ defined by

$$H_k^{(j+1)} \cup \{e \setminus \{u_j\} : e \in H_{k+1}^{(j)}, u_j \in e\}.$$

Step 4: Let $H_{k+1}^{(j+1)}$ be the hypergraph on $V(H_{k+1}^{(j)}) \setminus W_{H_{k+1}^{(j)}}(u_j)$ with

$$H_{k+1}^{(j+1)} = \{e \in H_{k+1}^{(j)} : e \cap W_{H_{k+1}^{(j)}}(u_j) = \emptyset \text{ and } T \not\subseteq e \text{ for all } T \in \bigcup_{\ell=1}^k D_\ell(H_k^{(j+1)}, \Delta_\ell^k)\}.$$

After running through the above procedure, set $H_k = H_k^{(s)}$, $A_k = V(H_{k+1}^{(s)})$, and $S_k = S_k^{(s)}$. Output (H_k, A_k, S_k) . ■

We emphasize that this algorithm allows the hypergraphs H_{k+1} and H_k to have repeated edges. We now analyze this algorithm through a series of lemmas. We omit many of the proofs since most are either straightforward or analogous to what was done in Lemma 17.1.

Lemma 18.2. *Assume one runs the Scythe Algorithm with parameters $s, \Delta_1^k, \dots, \Delta_k^k$ on inputs $(H_{k+1}, I), (H_{k+1}, I')$ and that the algorithm outputs $(H_k, A_k, S_k), (H'_k, A'_k, S'_k)$, respectively. If $S_k \subseteq I'$ and $S'_k \subseteq I$, then $(H_k, A_k, S_k) = (H'_k, A'_k, S'_k)$.*

For the rest of this subsection we will assume that we have run the Scythe Algorithm with parameters $s, \Delta_1^k, \dots, \Delta_k^k$ on (H_{k+1}, I) which outputs some (H_k, A_k, S_k) . We observe the following basic properties.

Lemma 18.3. *The following hold:*

- I is an independent set of H_k ,
- $S_k \subseteq I \subseteq A_k \cup S_k$,
- Both H_k and A_k are determined by H_{k+1} and S_k ,
- $|S_k| \leq s$,
- For all $\ell \leq k$ we have $\Delta_\ell(H_k) \leq \frac{1}{2}\Delta_\ell^k + \Delta_{\ell+1}(H_{k+1})$.

We now turn to the main lemma of this subsection, which roughly says that if H_{k+1} is “nice,” then either H_k will also be “nice” or A_k will be small.

Lemma 18.4. *Either*

$$e(H_k) \geq \min \left\{ \frac{s}{v(H)}, \frac{\Delta_1^k}{\Delta_1(H_{k+1})}, \dots, \frac{\Delta_k^k}{\Delta_k(H_{k+1})} \right\} \cdot \frac{e(H_{k+1})}{(k+1)2^{k+2}},$$

or

$$|A_k| \leq v(H_{k+1}) - \frac{e(H_{k+1})}{4 \cdot \Delta_1(H_{k+1})}.$$

Note that this lemma will be most “effective” when each ratio of the minimum is roughly the same. And indeed, we will end up choosing our parameters so that these ratios are all at least $s/v(H)$.

Proof. If the algorithm ever stops at Step 1, then $|A_k| = 0$ and there is nothing to prove, so we can assume that Steps 2 through 4 are completed a total of s times. We observe that

$$|A_k| = v(H_{k+1}^{(s)}) = v(H_{k+1}) - \sum_{j=0}^{s-1} |W_j(u_j)|,$$

where for ease of notation we let $W_j = W_{H_{k+1}^{(j)}}$. Thus we can assume

$$\sum_{j=0}^{s-1} |W_j(u_j)| < \frac{e(H_{k+1})}{4\Delta_1(H_{k+1})}. \quad (22)$$

By construction, for all j we have

$$e(H_k^{(j+1)}) - e(H_k^{(j)}) = \deg_{H_{k+1}^{(j)}}(u_j).$$

Because u_j is the largest element of $I \cap V(H_{k+1}^{(j)})$ in the max-degree order, the degree of u_j is at least as large as the average degree of the subgraph of $H_{k+1}^{(j)}$ after deleting $W_j(u_j) \setminus \{u_j\}$, and again by definition of the max-degree order, this is at least as large as the average degree of $H_{k+1}^{(j)}$ (which has at most $v(H)$ vertices). In total then we find

$$e(H_k) = \sum_{j=0}^{s-1} e(H_k^{(j+1)}) - e(H_k^{(j)}) \geq \sum_{j=0}^{s-1} \frac{(k+1)e(H_{k+1}^{(j+1)})}{v(H)}.$$

If we have $(k+1)e(H_{k+1}^{(j+1)}) \geq e(H_{k+1})$ for all j , then the above sum is at least $(s/v(H)) \cdot e(H_{k+1})$, giving the desired result. Thus we can assume this fails for some j , and this implies

$$e(H_{k+1}^{(s)}) \leq e(H_{k+1}^{(j+1)}) < \frac{e(H_{k+1})}{k+1}. \quad (23)$$

This means that many edges of H_{k+1} were deleted in Step 4 of the algorithm. We claim that this implies that one of the sets $D_\ell(H_k, \Delta_\ell^k)$ is large. Indeed, observe that

$$e(H_{k+1}^{(j)}) - e(H_{k+1}^{(j+1)}) \leq |W_j(u_j)| \cdot \Delta_1(H_k) + \sum_{\ell} |D_\ell(H_k^{(j+1)}, \Delta_\ell^k) \setminus D_\ell(H_k^{(j)}, \Delta_\ell^k)| \cdot \Delta_\ell(H_{k+1}),$$

since edges are either deleted by deleting vertices in $W_j(u_j)$ or by deleting ℓ -sets which are in D_ℓ for $H_k^{(j+1)}$ but not $H_k^{(j)}$. Summing this over all j gives

$$\begin{aligned} e(H_{k+1}) - e(H_{k+1}^{(s)}) &\leq \sum_j |W_j(u_j)| \cdot \Delta_1(H_{k+1}) + \sum_{\ell} |D_\ell(H_k, \Delta_\ell^k)| \cdot \Delta_\ell(H_{k+1}) \\ &< \frac{k \cdot e(H_{k+1})}{2(k+1)} + \sum_{\ell} |D_\ell(H_k, \Delta_\ell^k)| \cdot \Delta_\ell(H_{k+1}), \end{aligned}$$

where this last step used (22) and $k \geq 1$. Using (23) shows that

$$\frac{k \cdot e(H_{k+1})}{2(k+1)} \geq \sum_{\ell} |D_{\ell}(H_k, \Delta_{\ell}^k)| \cdot \Delta_{\ell}(H_{k+1}),$$

so for some ℓ we must have

$$|D_{\ell}(H_k, \Delta_{\ell}^k)| \geq \frac{e(H_{k+1})}{2(k+1)\Delta_{\ell}(H_{k+1})}.$$

With ℓ as above, the handshaking lemma and definition of D_{ℓ} implies

$$e(H_k) = \binom{k}{\ell}^{-1} \sum_{T \in \binom{V(H_k)}{\ell}} \deg_{H_k}(T) \geq \binom{k}{\ell}^{-1} \cdot |D_{\ell}(H_k, \Delta_{\ell}^k)| \cdot \frac{1}{2} \Delta_{\ell}^k \geq \frac{e(H_{k+1}) \cdot \Delta_{\ell}^k}{(k+1)2^{k+2}\Delta_{\ell}(H_{k+1})},$$

giving the desired result. \square

18.3 Proof of The Formal Result

We are now ready to prove the following formal statement.

Theorem 18.5 ([10]). *For every integer $r \geq 2$ and $c \geq 1$, there exists $\delta > 0$ such that the following holds. Let $q \in (0, 1)$ and suppose H is an r -uniform hypergraph such that for every $1 \leq \ell \leq r$ we have*

$$\Delta_{\ell}(H) \leq cq^{\ell-1} \cdot \frac{e(H)}{v(H)}.$$

Then there exists $\mathcal{S} \subseteq \binom{V(H)}{\leq (r-1)q \cdot v(H)}$ and functions $f : \mathcal{S} \rightarrow \binom{V(H)}{\leq (1-\delta)v(H)}$ and $g : \mathcal{I}(H) \rightarrow \mathcal{S}$ such that for every $I \in \mathcal{I}(H)$ we have

$$g(I) \subseteq I \subseteq f(g(I)) \cup g(I).$$

Moreover, $S \cap f(S) = \emptyset$ for all $S \in \mathcal{S}$, and if $I, I' \in \mathcal{I}(H)$ satisfy $g(I) \subseteq I'$, $g(I') \subseteq I$, then $g(I) = g(I')$.

Proof. For all $\ell \leq r$ let $\Delta_{\ell}^r := \Delta_{\ell}(H)$, and inductively define

$$\Delta_{\ell}^k := \max\{2 \cdot \Delta_{\ell+1}^{k+1}, q \cdot \Delta_{\ell}^{k+1}\}.$$

The following is straightforward to prove given the hypothesis of the theorem.

Claim 18.6. *For all $k < r$ we have $\Delta_1^{k+1} \leq C2^r q^{r-k-1} \frac{e(H)}{v(H)}$.*

For each $I \in \mathcal{I}(H)$, iteratively run through the Scythe Algorithm with the parameters above and $s = q \cdot v(H)$, starting with $H_r = H$. Let $(H_{r-1}, A_{r-1}, S_{r-1}), \dots, (H_1, A_1, S_1)$ denote the outputs of this algorithm.

It is straightforward to show that $\Delta_{\ell}(H_k) \leq \Delta_{\ell}^k$ for all ℓ, k by using induction and Lemma 18.3. For $k < r$ we define $c_k = (Cr2^{r+1})^{k-r}$. Let K be the smallest integer such that $|A_K| \leq$

$(1 - c_K)v(H)$, and if no such $K \geq 1$ exists we set $K = 0$. It is straightforward to prove that $e(H_k) \geq c_k q^{k-r} e(H)$ for all $k > K$ by using induction, Lemma 18.4, and Claim 18.6.

Let $\delta := c_1$. Before we define our functions, for technical reasons it will be convenient to first define a function $f^* : \mathcal{I}(H) \rightarrow \binom{V(H)}{\leq (1-\delta)v(H)}$ before defining f . Pick some I and let the hypergraphs H_k and integer K be as defined above. Observe that if $K \geq 1$, then $|A_K| \leq (1 - \delta)v(H)$, so in this case we will set

$$g(I) = \bigcup_{k \geq K} S_k, \quad f^*(I) = A_K.$$

If $K = 0$, then we set

$$g(I) = \bigcup_{k \geq 1} S_k, \quad f^*(I) = \{v \in V(H_1) : \{v\} \notin H_1\}.$$

Note that in this case $|f^*(I)| = v(H) - e(H_1)$, which is at most $(1 - \delta)v(H)$ by our observations above. Lastly, we define $\mathcal{S} = \{g(I) : I \in \mathcal{I}(H)\}$ and $f(S) = f^*(I)$ for any $I \in g^{-1}(S)$. The fact that f is well defined is implied by the following claim, which itself follows from Lemma 18.3.

Claim 18.7. *If $I, I' \in \mathcal{I}(H)$ with $g(I) \subseteq I'$ and $g(I') \subseteq I$, then $g(I) = g(I')$ and $f^*(I) = f^*(I')$.*

The fact that these definitions give the desired result, except possibly the condition $S \cap f(S) = \emptyset$, can be checked by using the properties from Lemma 18.3. This last condition can be established by taking $f'(S) := f(S) \setminus S$ if needed. \square

The following weaker version of Theorem 18.5 is often good enough for most applications and is conceptually simpler.

Corollary 18.8. *For every integer $r \geq 2$ and $c \geq 1$, there exists $\delta > 0$ such that the following holds. Let $q \in (0, 1)$ and suppose H is an r -uniform hypergraph such that for every $1 \leq \ell \leq r$ we have*

$$\Delta_\ell(H) \leq cq^{\ell-1} \cdot \frac{e(H)}{v(H)}.$$

Then there exists a collection of sets \mathcal{C} such that every independent set of H is a subset of some $C \in \mathcal{C}$, and moreover, $|C| \leq (1 - \delta)v(H)$ for all $C \in \mathcal{C}$ and $|\mathcal{C}| \leq \binom{v(H)}{\leq (r-1)q \cdot v(H)}$.

Proof. In the notation of Theorem 18.5, we let $\mathcal{C} = \{f(g(I)) \cup g(I) : I \in \mathcal{I}(H)\}$. \square

On its own, Corollary 18.8 (and even Theorem 18.5) isn't terribly useful since the containers it generates are rather large, and in practice one needs to reapply this lemma to each $C \in \mathcal{C}$ which is large, and to keep repeating this argument until the contains are sufficiently small. Because δ depends only on r and C , this only needs to be done a constant number of times. However, to reapply the lemma, each large $C \in \mathcal{C}$ must satisfy essentially the same hypothesis as H . While a generic hypergraph will fail to have this property, many nice hypergraphs will.

19 Hypergraph Containers and Triangle-Free Graphs

Let us restate our weak container theorem Corollary 18.8 for the special case of 3-uniform hypergraphs.

Theorem 19.1 ([10]). *For every $c \geq 1$, there exists $\delta > 0$ such that the following holds. Let $q \in (0, 1)$ and suppose H is a 3-uniform hypergraph such that*

$$\Delta_1(H) \leq c \frac{e(H)}{v(H)},$$

$$\Delta_2(H) \leq cq \frac{e(H)}{v(H)},$$

$$\Delta_3(H) \leq cq^2 \frac{e(H)}{v(H)}.$$

Then there exists a collection of sets \mathcal{C} such that every independent set of H is a subset of some $C \in \mathcal{C}$, and moreover, $|C| \leq (1 - \delta)v(H)$ for all $C \in \mathcal{C}$ and $|\mathcal{C}| \leq \binom{v(H)}{\leq 2q \cdot v(H)}$.

Note that for simple hypergraphs we have $\Delta_3(H) = 1$, so this last bound is equivalent to lower bounding the average degree by $c^{-1}q^{-2}$. One important consequence of Theorem 19.1 is the following.

Theorem 19.2. *For all $n, \epsilon > 0$, there exists a collection of n -vertex graphs \mathcal{C} such that*

(a) *Every triangle-free graph $G \subseteq K_n$ is a subgraph of some $C \in \mathcal{C}$,*

(b) *Every $C \in \mathcal{C}$ has less than ϵn^3 triangles, and*

(c) *We have $|\mathcal{C}| = n^{O_\epsilon(n^{3/2})}$.*

That is, there exists a small set of nearly triangle-free graphs which contains every triangle-free graph.

Proof. Start with $\mathcal{C} = \{K_n\}$, and note that \mathcal{C} trivially satisfies (a). Iteratively proceed as follows. If every $C \in \mathcal{C}$ has less than ϵn^3 triangles then output the current collection \mathcal{C} . Otherwise, let $C \in \mathcal{C}$ be such that it contains at least ϵn^3 triangles. Form a 3-graph H with vertex set $E(C)$ where three edges of C form a hyperedge in H if they form a triangle. Note that $e(H) \geq \epsilon n^3$ and $v(H) = e(C) \leq n^2$. Every edge is contained in at most n triangles, so $\Delta_1(H) \leq n \leq \epsilon^{-1} \frac{e(H)}{v(H)}$. We also have $\Delta_2(H) = \Delta_3(H) = 1 \leq \epsilon^{-1} (n^{-1/2})^2 \frac{e(H)}{v(H)}$. With this we see that we can apply Theorem 19.1 with $q = n^{-1/2}$ and $c = \epsilon^{-1}$. This gives a collection of containers \mathcal{C}' for C , i.e. subgraphs $C' \subseteq C$ such that every triangle-free subgraph of C is contained in some $C' \in \mathcal{C}'$. Remove C from \mathcal{C} and add every $C' \in \mathcal{C}'$ to \mathcal{C} . Repeat this process.

Let \mathcal{C} be the final collection that this algorithm produces. It is straightforward to show that (a) holds inductively, and (b) holds by construction. To show that the final collection is small, first

note that each time we apply the container lemma, the number of new graphs we create is at most $\binom{v(H)}{\leq 2n^{-1/2}v(H)} = n^{O(n^{3/2})}$. Second, observe that each time we apply the container lemma to C , the graphs in \mathcal{C}' have at most $(1-\delta)e(C)$ edges, where δ depends only on ϵ . Because we only iterate on C which have at least ϵn^2 edges (since they need at least ϵn^3 triangles), we iteratively apply the lemma at most some bounded number of times $b = b(\epsilon)$ to reach any element in the final collection \mathcal{C} . Thus the total number of containers we create is $\left(n^{O(n^{3/2})}\right)^b = n^{O_\epsilon(n^{3/2})}$ as desired. \square

For Theorem 19.2 to be useful, we need to get a handle on graphs with at most ϵn^3 triangles. As is typical with containers, this will come from a supersaturation lemma.

Lemma 19.3. *For every $\delta > 0$ there exists an $\epsilon > 0$ such that if G is an n -vertex graph with $e(G) \geq (\frac{1}{2} + \delta)\binom{n}{2}$, then G contains at least ϵn^3 triangles.*

I'm not crazy about the ordering of δ, ϵ but I admit the final thing should be about ϵ ...well actually in a lot of the applications it kind of makes more sense to do it the other way. Maybe do the K_r version in general depending on what I need later on.

Proof. **TODO** \square

With this we can prove the following counting result.

Theorem 19.4. *The number of n -vertex triangle-free graphs is equal to*

$$2^{(1+o(1))n^2/4}.$$

Proof. The lower bound comes from considering all of the subgraphs of $K_{n/2, n/2}$. For the upper bound, fix some $\delta > 0$ and let ϵ be as in Lemma 19.3. Let \mathcal{C} be the containers guaranteed by Theorem 19.2 with parameter ϵ . Because every triangle-free graph is a subgraph of some $C \in \mathcal{C}$, the number of triangle-free graphs is at most

$$\sum_{C \in \mathcal{C}} 2^{|C|} \leq n^{O(n^{3/2})} \cdot 2^{\max_{C \in \mathcal{C}} e(C)},$$

Since each $C \in \mathcal{C}$ has less than ϵn^3 and at triangles, Lemma 19.3 implies $e(C) \leq (\frac{1}{2} + \delta)\binom{n}{2}$ for all $C \in \mathcal{C}$. In total we get an upper bound of

$$2^{(\frac{1}{2} + \delta)\binom{n}{2} + O(n^{3/2} \log n)},$$

and letting δ tend towards 0 gives the result. \square

While containers most directly allow one to solve problems that are equivalent to counting the number of independent sets of a hypergraph, there are other related problems which they're effective for. For example, in the next subsection we show how containers can be used to prove probabilistic analogs of classical extremal results. In the section after this we show how one can use containers to count special kinds of independent sets, namely maximal independent sets.

19.1 Mantel's Theorem in Random Graphs

Given two graphs G, F , we let $\text{ex}(G, F)$ denote the largest F -free subgraph of G . For example, $\text{ex}(K_n, F) = \text{ex}(n, F)$. The following result can be viewed as a random version of Mantel's theorem.

Theorem 19.5. *Define $\text{ex}(G_{n,p}, K_3)$ to be the largest triangle-free subgraph of $G_{n,p}$. We have $\text{ex}(G_{n,p}, K_3) = (1 + o(1))pn^2/4$ whp provided $p \gg n^{-1/2} \log n$.*

Proof. The lower bound follows by considering $G_{n,p} \cap K_{n/2, n/2}$, which is always triangle-free and which has $(1 + o(1))pn^2/4$ edges whp. For the upper bound, fix $\delta > 0$, and let $\epsilon > 0$ be as in Lemma 19.3. Let \mathcal{C} be the set of containers given by Theorem 19.2 with parameter ϵ , and as before we have $e(C) \leq (1/2 + \delta)\binom{n}{2}$ for all $C \in \mathcal{C}$. Because every triangle-free graph is contained in some $C \in \mathcal{C}$, in order to have $\text{ex}(G_{n,p}, K_3) \geq (1 + 4\delta)pn^2/4$, there must exist some $C \in \mathcal{C}$ such that $|G_{n,p} \cap C| \geq (1 + 4\delta)pn^2/4$. Let E_C be the event that this bound holds. Observe that $|G_{n,p} \cap C|$ is a binomial random variable with probability p and at most $(1 + 2\delta)n^2/4$ trials. By the Chernoff bound, we find $\Pr[E_C] \leq e^{-O_\delta(pn^2)}$. In total then, we have

$$\Pr[\text{ex}(G_{n,p}, K_3) \geq (1 + 4\delta)pn^2/4] \leq \Pr\left[\bigcup_{C \in \mathcal{C}} E_C\right] \leq n^{O_\delta(n^{3/2})} \cdot e^{-O_\delta(pn^2)} \rightarrow 0,$$

with this last step holding by hypothesis on p . We conclude the result by taking δ arbitrarily close to 0. \square

We note that for $p \ll n^{-1/2}$, a simple deletion argument shows that for $p \ll n^{-1/2}$ there exist triangle-free subgraphs with $(1 + o(1))p\binom{n}{2}$ edges, and this is certainly best possible since $G_{n,p}$ has at most this many edges asymptotically. Thus the bound for p in Theorem 19.5 is almost optimal. In fact, we can obtain the optimal bound in this theorem by using the strong container theorem Theorem 18.5, which in the case of 3-graphs can be written as follows.

Theorem 19.6. *For every $c \geq 1$, there exists $\delta > 0$ such that the following holds. Let $q \in (0, 1)$ and suppose H is a 3-uniform hypergraph such that*

$$\begin{aligned}\Delta_1(H) &\leq c \frac{e(H)}{v(H)}, \\ \Delta_2(H) &\leq cq \frac{e(H)}{v(H)}, \\ \Delta_3(H) &\leq cq^2 \frac{e(H)}{v(H)}.\end{aligned}$$

Then there exists $\mathcal{S} \subseteq \binom{V(H)}{\leq 2q \cdot v(H)}$ and functions $f : \mathcal{S} \rightarrow \binom{V(H)}{\leq (1-\delta)v(H)}$ and $g : \mathcal{I}(H) \rightarrow \mathcal{S}$ such that for every $I \in \mathcal{I}(H)$ we have

$$g(I) \subseteq I \subseteq f(g(I)) \cup g(I).$$

Moreover, $S \cap f(S) = \emptyset$ for all $S \in \mathcal{S}$, and if $I, I' \in \mathcal{I}(H)$ satisfy $g(I) \subseteq I'$, $g(I') \subseteq I$, then $g(I) = g(I')$.

This allows us to construct the following “strong” set of containers for triangle-free graphs.

Theorem 19.7. *Let $\mathcal{G}_n, \mathcal{T}_n$ denote the set of all n -vertex graphs and all n -vertex triangle-free graphs, respectively. For all $n, \epsilon > 0$, there exists a set of graphs \mathcal{S} with at most $O_\epsilon(n^{3/2})$ edges, as well as functions $f : \mathcal{S} \rightarrow \mathcal{G}_n$ and $g : \mathcal{T}_n \rightarrow \mathcal{S}$ such that for every $G \in \mathcal{T}_n$, we have*

$$g(G) \subseteq G \subseteq f(g(G)) \cup g(G),$$

and such that $f(S)$ has less than ϵn^3 triangles for all $S \in \mathcal{S}$.

Proof. We start with \mathcal{S} consisting only of the empty graph and define $g(G) = \emptyset$ and $f(\emptyset) = K_n$. Iteratively assume we have constructed some \mathcal{S}, f, g satisfying all of the conditions except possibly that each $S \in \mathcal{S}$ has at most $O_\epsilon(n^{3/2})$ edges and that $f(S)$ has less than ϵn^3 triangles (which holds for our initial step). If $f(S)$ has less than ϵn^3 triangles for all $S \in \mathcal{S}$ then we end the procedure. Otherwise, let S be such that $C = f(S)$ has at least ϵn^3 triangles. By repeating our computations from the proof of Theorem 19.2, we see that we can apply Theorem 19.6 to the 3-graph H encoding triangles of C , and we let \mathcal{S}_C, f_C, g_C be the output of this theorem.

Claim 19.8. *Let $\mathcal{S}' := (\mathcal{S} \setminus \{S\}) \cup \{S_C \cup S : S_C \in \mathcal{S}_C\}$, define $g'(G) = g(G)$ if $g(G) \neq S$ and $g'(G) = g_C(G - S)$ otherwise, and define $f'(S') = f(S')$ if $S' \in \mathcal{S} \setminus \{S\}$ and $f'(S') = f_C(S' - S)$ otherwise. These maps are well defined and satisfy the conditions of the theorem except possibly that each $S \in \mathcal{S}'$ has at most $O_\epsilon(n^{3/2})$ edges and that $f'(S)$ has less than ϵn^3 triangles.*

Proof. First observe that because $C \cap S = \emptyset$, each element of \mathcal{S}_C (which is a subgraph of C) is disjoint from S . This implies that all of the elements $S_C \cup S$ for $S_C \in \mathcal{S}_C$ are distinct. Moreover, none of these elements are equal to any element of $\mathcal{S} \setminus \{S\}$. Indeed, if $S_C \cup S = S' \in \mathcal{S}$, then \mathcal{S} would contain two elements with $S \subsetneq S'$. The last condition of Theorem 19.6 then implies that we must have $S = S'$. This all implies that g', f' are well defined maps, and it is not difficult to check that they inherit all of the other desired properties. \square

With this we can keep applying Theorem 19.6 until we get \mathcal{S}, f, g which satisfies all of the conditions except possibly that $e(S)$ is small. As in the proof of Theorem 19.2, one can check that each $S \in \mathcal{S}$ is obtained by applying Theorem 19.6 at most $O_\epsilon(1)$ times, and each time its applied at most $O(n^{3/2})$ edges get added to S . With this we can conclude the result. \square

We note that there exists a somewhat stronger version of Theorem 19.6 (and more generally Theorem 18.5) which allows one to prove the previous result with less work. However, the theorem statement is somewhat more complicated conceptually (involving things called (\mathcal{F}, ϵ) -dense families), so for this exposition we have opted to use the simpler version. In any case, with this enhanced version of Theorem 19.2, we can improve upon our threshold for the random Mantel theorem by dropping a logarithmic term.

Theorem 19.9. *Define $\text{ex}(G_{n,p}, K_3)$ to be the largest triangle-free subgraph of $G_{n,p}$. We have $\text{ex}(G_{n,p}, K_3) = (1 + o(1))pn^2/4$ whp provided $p \gg n^{-1/2}$.*

Proof. The lower bound follows by considering $G_{n,p} \cap K_{n/2, n/2}$, which is always triangle-free and which has $(1 + o(1))pn^2/4$ edges whp. For the upper bound, fix $\delta > 0$, and let $\epsilon > 0$

be as in Lemma 19.3. Let \mathcal{S}, f, g be as in Theorem 19.7. Note that each $f(S)$ has at most $(1/4 + 2\delta)n^2$ edges by Lemma 19.3. For each $S \in \mathcal{S}$, let E_S be the event that $S \subseteq G_{n,p}$ and that $|f(S) \cap G_{n,p}| \geq (1 + 4\delta)pn^2/4$. Note that in order to have $\text{ex}(G_{n,p}, K_3) \geq (1 + 4\delta)pn^2 + O_\epsilon(n^{3/2})$, some E_S event must occur, and moreover that $\Pr[E_S] = p^{|S|} \cdot e^{-O_\delta(pn^2)}$. With this we have

$$\Pr[\text{ex}(G_{n,p}, K_3) \geq (1 + 4\delta)pn^2/4 + O_\epsilon(n^{3/2})] \leq \Pr\left[\bigcup_{S \in \mathcal{S}} E_S\right] \leq \sum_{s=0}^{O_\epsilon(n^{3/2})} \sum_{S \in \mathcal{S}: |S|=s} p^s e^{-O_\delta(pn^2)}.$$

As the number of $S \in \mathcal{S}$ with $|S| = s$ is trivially at most $\binom{n^2}{s} \leq (en^2/s)^s$, we find that the above is at most

$$\sum_{s=0}^{O_\epsilon(n^{3/2})} (epn^2/s)^s e^{-O_\delta(pn^2)}.$$

One can check that the function $(epn^2/s)^s$ is increasing for $s \leq pn^2$. Since we know $s \leq C_\epsilon n^{3/2}$ for some suitable C_ϵ , we get that the sum above is at most

$$C_\epsilon n^{3/2} \cdot (eC_\epsilon^{-1}pn^{1/2})^{C_\epsilon n^{3/2}} e^{-O_\delta(pn^2)},$$

and this tends to 0 provided $pn^{1/2} \rightarrow \infty$ (since $pn^2 \gg n^{3/2} \log(pn^{1/2})$), proving the result. \square

Note that in this proof, the main extra power we gained by utilizing Theorem 19.7 is that S must be contained in our subgraph. This makes it so that the $S \in \mathcal{S}$ with many edges “cost more”, allowing us to gain.

We note that in general, it is very common that by using the weak container lemma, one ends up getting tight bounds up to a logarithmic factor, and this extra factor can usually be remedied by utilizing the strong container lemma in some straightforward (if slightly more tedious) way.

19.2 Maximal Triangle-Free Graphs

In this subsection we use containers to count maximal independent sets, i.e. those that are maximal with respect to set inclusion. To do this, we again apply the container lemma to find a small collection of containers \mathcal{C} for a hypergraph H . We then argue that each $C \in \mathcal{C}$ contains few maximal independent sets, which gives the result.

In order for this approach to be effective, we need a supersaturation result saying that if $H[C]$ has few edges, then C contains few maximal independent sets, and these results are typically a bit more complicated to prove compared to the non-maximal setting, and often these proofs invoke facts about the number of maximal independent sets in special kinds of graphs.

One case where we can pull off the scheme outlined above is in counting maximal triangle-free graphs. By Theorem 19.4, we know that there are $2^{n^2/4 + o(n^2)}$ triangle-free graphs on n vertices, and moreover, it is a well known result of Erdős, Kleitman, and Rothschild [41] that almost all of these graphs are bipartite. However, a bipartite triangle-free graph is a maximal triangle-free graph if and only if it is a complete bipartite graph, and there are less than 2^n such graphs. A different set of constructions gives the following.

Proposition 19.10. *There are at least $2^{n^2/8}$ maximal triangle-free graphs on n -vertices.*

Proof. Write the vertices of $[n]$ as $a_1, \dots, a_{n/4}, b_1, \dots, b_{n/4}, c_1, \dots, c_{n/2}$. Add every edge of the form $a_i b_i$. For every $1 \leq i \leq n/4$ and $1 \leq j \leq n/2$, add exactly one of the edges $a_i c_j$ or $b_i c_j$. In total this gives $2^{(n/4)(n/2)} = 2^{n^2/8}$ different graphs $\{G_1, G_2, \dots\}$, and it's not hard to see that each of these are triangle-free. Let G'_k be any maximal triangle-free graph containing G_k (so $G'_k = G_k$ if G_k is maximal). One can check that each of the G'_k graphs are distinct from each other, giving the desired result. \square

Our goal is to show that this result is best possible. This was originally proven by Balogh and Petříčková [12], and our proof follows their same approach.

Similar to the proof of Theorem 19.4 where we counted the number of triangle-free graphs, we'll begin by constructing a set of using Theorem 19.2 to obtain a set of containers \mathcal{C} where each element is a graph which contains few triangles. From there it remains to show that each of these $C \in \mathcal{C}$ contains at most roughly $2^{n^2/8}$ maximal triangle-free graphs.

Essentially the only thing we know about each $C \in \mathcal{C}$ is that they contain few triangles, and fortunately quite a lot can be said about such graphs: the triangle removal lemma says that we can delete a small number of edges from C to get a triangle-free graph, and supersaturation says that $e(C)$ is not much larger than $\frac{1}{4}n^2$. With these ideas in mind we prove the following.

Lemma 19.11. *Let $\epsilon, \gamma > 0$ be constants. Let $\delta(\epsilon)$ be as in **REF**, $\delta(\gamma)$ be as in **REF**, and $\delta = \min\{\delta(\epsilon), \delta(\gamma)\}$. If G is an n -vertex graph with at most δn^3 triangles, then the number of maximal triangle-free subgraphs of G is at most*

$$2^{n^2/8 + \gamma n^2/2 + \epsilon n^2}.$$

Proof. By **Triangle-removal**, there exists a set $F \subseteq E(G)$ of size at most ϵn^2 such that $G - F$ is triangle-free. For $F' \subseteq F$, let $\mathcal{M}(F')$ denote the set of maximal triangle-free subgraphs $G' \subseteq G$ with $G' \cap F = F'$. Observe that the $\mathcal{M}(F')$ sets partition the maximal triangle-free subgraphs into $2^{\epsilon n^2}$ sets, so it suffices to show that for any $F' \subseteq F$,

$$|\mathcal{M}(F')| \leq 2^{n^2/8 + \gamma n^2/2}.$$

The result is trivial if F' contains a triangle (since there are 0 graphs which contain F' and which are triangle-free), so from now on we'll assume F' is triangle-free. Define an auxiliary graph T via

$$V(T) = G - (F - F') - \{e : \exists f, g \in F' \text{ and } e, f, g \text{ form a triangle}\},$$

and where two edges $f, g \in V(T)$ are adjacent in T if there exists an edge $e \in F'$ which forms a triangle with f, g . We make these definitions so that the following holds.

Claim 19.12. *Every $G' \in \mathcal{M}(F')$ is a maximal independent set of T .*

Proof. Fix some $G' \in \mathcal{M}(F')$. We first claim that

$$G' \subseteq V(T) = G - (F - F') - \{e : \exists f, g \in F' \text{ and } e, f, g \text{ form a triangle}\}.$$

Indeed, we must have $G' \subseteq G - (F - F')$ in order to have $G' \cap F = F'$, and given this, G' cannot contain any edge e for which there exist $f, g \in F'$ forming a triangle with e since G' is triangle-free. This proves the claim.

We next claim that G' is an independent set of T . Indeed, if $f, g \in G'$ were adjacent in T , then there exists some edge $e \in F' \subseteq G'$ forming a triangle with f, g , contradicting G' being triangle-free.

It remains to show that G' is a maximal independent set. To this end, consider any $f \in V(T) \setminus G'$. Because G' is a maximal triangle-free graph, there must exist edges $e, g \in G'$ which form a triangle with f . Observe that at least one of e, g must be in F' , since by assumption of $G - F$ being triangle-free, every triangle must contain at least one edge of F (and the only such edges in $V(T)$ are in F'). Thus we may assume $e \in F'$. We also note that g does not form a triangle with two edges of F' , as this would contradict $g \in G'$ and G' being triangle-free. Thus we have $f, g \in V(T)$, and $e \in F'$ implies that these two edges are adjacent in T . This implies that G' is indeed a maximal independent of T , completing the proof. \square

It remains to show that T has few maximal independent sets. For this we make the following key observation.

Claim 19.13. *The graph T is triangle-free.*

Proof. Assume for contradiction that there existed edges f_1, f_2, f_3 forming a triangle in T . By definition of T , this is only possible if these edges are intersecting, and hence these edges either form a triangle or a star. If they form a triangle, then by definition of T we in fact have $f_i \in F'$ for all i , and hence F' forms a triangle, a contradiction to our assumption on F' . If these edges form a star, say with leaves x, y, z , then again by definition of T we must have $xy, yz, xz \in F'$, a contradiction to our assumption of F' being triangle-free. We conclude the result. \square

Thus we've reduced the problem to upper bounding the number of maximal independent sets in a triangle-free graph T . A well known result of Moon and Moser [87] says that an N -vertex triangle-free graph has at most $2^{N/2}$ maximal independent sets, so the number of maximal independent sets of T is at most

$$2^{|V(T)|/2} \leq 2^{e(G)/2} \leq 2^{n^2/8 + \gamma n^2/2},$$

where the last step used $\delta \leq \delta(\gamma)$. This implies the result. \square

From here one can prove that there are at most $2^{n^2/8 + o(n^2)}$ maximal triangle-free graphs on n -vertices analogous to how we proved Theorem 19.4 I should check this more carefully.

Before moving on, it is natural to ask how many maximal K_4 -free graphs there are. Unfortunately the present argument completely fails to generalize to this setting.

Essentially, the issue is that T can be viewed as the link graph of a 3-uniform hypergraph. More precisely, let H be the 3-graph with $V(H) = G - (F - F')$ and whose hyperedges are triangles. Then the link hypergraph $H[F']$ contains the graph T together with a 1-edge on every edge e such that there exist $f, g \in F'$ forming a triangle with e . For the K_4 problem, one could again consider the link set of some 6-graph, but the structure you end up getting is some very

non-uniform hypergraph which avoids some strange set of subgraphs, and as such the analysis becomes somewhat unwieldy.

20 **TODO** Counting F -free graphs

Ferber-McKinley-Samotij.

Part VI

Linear Algebra Methods

Very roughly speaking, the *linear algebra method* in combinatorics works as follows:

1. Associate a “linear algebraic object” M to your problem (e.g. a matrix, list of vectors).
2. Determine algebraic information about M (e.g. its rank, eigenvalues, eigenvectors),
3. Use this algebraic information to say something about your original problem.

The linear algebra method applies to a broad range of problems. We only scratch the surface here, and we refer the reader to books by Babai and Frankl [8] and by Matoušek [83] for a more thorough treatment of this versatile method.

21 Modular Intersections

We begin with a very classical application of the linear algebra method: Oddtown.

Consider the following (somewhat whimsical) setup. The city of Oddtown has a variety of clubs, each of which follows the following odd set of rules: each club must have an odd number of people, and every two distinct clubs must have an even number of people in common.

The main question now becomes: if Oddtown has n people, what's the maximum number of clubs it can have? Equivalently, if $\mathcal{F} = \{F_1, F_2, \dots, F_m\} \subseteq 2^{[n]}$ is a set system such that $|F_i|$ is odd for all i and such that $|F_i \cap F_j|$ is even for all $i \neq j$, then what is the maximum size of \mathcal{F} ?

A very simple construction is to take $F_i = \{i\}$ for all i , which trivially satisfies the stated conditions. However, it's far from the only construction. For example, if n is even one can also take each F_i to be either $\{i\}$ or $[n] \setminus \{i\}$, and there are many, many more constructions achieving a bound of n (in fact, there's close to 2^{n^2} non-isomorphic constructions due to Szegedy [8, Exercise 1.1.14]).

Given all of these constructions, it seems plausible that (1) the true answer is indeed n , and (2) proving this might be difficult (since we have to come up with an argument that somehow deals with all of these constructions in a unified way). Fortunately, the linear algebra method manages to give a unified approach for all of these constructions in an extremely elegant way. More generally, we note that it's a good rule of thumb is that if there are many distinct looking constructions, then the linear algebra method might come in handy.

Theorem 21.1 (Oddtown). *Let $\mathcal{F} \subseteq 2^{[n]}$ be a set system such that $|F|$ is odd for all $F \in \mathcal{F}$ and such that $|F \cap F'|$ is even for all $F \neq F' \in \mathcal{F}$. Then $|\mathcal{F}| \leq n$.*

Proof. Given a set $F \subseteq [n]$, define its characteristic vector $\chi_F \in \mathbb{F}_2^n$ by having $(\chi_F)_i = 1$ if $i \in F$ and $(\chi_F)_i = 0$ otherwise. Note crucially that for any F, F' , the dot product satisfies

$$\langle \chi_F, \chi_{F'} \rangle = |F \cap F'| \pmod{2}.$$

We claim that $\{\chi_F : F \in \mathcal{F}\}$ is a set of linearly independent vectors. Indeed, say we had

$$\sum_{F \in \mathcal{F}} \lambda_F \chi_F = 0.$$

Take any $F' \in \mathcal{F}$ and apply the dot product on both sides to get

$$\sum_{F \in \mathcal{F}} \lambda_F \langle \chi_F, \chi_{F'} \rangle = 0.$$

By the observation above and the hypothesis of the theorem, we see $\langle \chi_F, \chi_{F'} \rangle = 0$ if $F \neq F'$ and that $\langle \chi_{F'}, \chi_{F'} \rangle = 1$. Thus the above says $\lambda_{F'} = 0$, and as $F' \in \mathcal{F}$ was arbitrary, we conclude that these vectors are indeed linearly independent.

Since we have $|\mathcal{F}|$ linearly independent vectors in \mathbb{F}_2^n , we must have $|\mathcal{F}| \leq n$, giving the result. \square

Since that was so clean, let's give another (essentially equivalent) proof of this result using slightly different language, which in some situations might be simpler to think about.

Proof. Write $\mathcal{F} = \{F_1, \dots, F_m\}$ and let M be the $m \times n$ matrix over \mathbb{F}_2 which has $M_{i,j} = 1$ if $j \in F_i$ and $M_{i,j} = 0$ otherwise. Let $L = MM^T$. We claim that L is the $m \times m$ identity matrix. Indeed, one can verify that $L_{i,j} \equiv |F_i \cap F_j| \pmod{2}$, so the hypothesis of the theorem gives this claim.

Using the general fact

$$\text{rank}(AB) \leq \text{rank}(A) \leq \#\text{columns of } A,$$

which is valid for any two matrices A, B for which AB makes sense; we see that we can apply this with $A = B^T = M$ to conclude that

$$m = \text{rank}(I_m) \leq \text{rank}(M) \leq n,$$

proving the result. □

While the proof of Oddtown is extremely nice, one might complain that the problem itself is rather ad hoc. Here we give a far reaching generalization of the Oddtown problem which, in addition to being nice on its own, has a number of important applications.

To this end, we adopt the following (non-standard) notation. For an integer p and a set $L \subseteq \{0, 1, \dots, p-1\}$, we say that a family \mathcal{F} is (p, L) -*modular intersecting* if $|F| \pmod{p} \notin L$ for all $F \in \mathcal{F}$ and if $|F \cap F'| \pmod{p} \in L$ for all $F \neq F' \in \mathcal{F}$.

For example, a family is $(2, \{0\})$ -modular intersecting if and only if it follows the rules of Oddtown. By following a somewhat similar strategy as in Oddtown, we can prove the following.

Proposition 21.2. *For any prime p and $L \subseteq \{0, 1, \dots, p-1\}$, if $\mathcal{F} \subseteq 2^{[n]}$ is (p, L) -modular intersecting then*

$$|\mathcal{F}| \leq \sum_{i=0}^{|L|} \binom{n+i-1}{i}.$$

Proof. Similar to before, we define the characteristic vector $\chi_F \in \mathbb{F}_p^n$ by having $(\chi_F)_i = 1$ if $i \in F$ and $(\chi_F)_i = 0$ otherwise, and we again observe that $\langle \chi_F, \chi_{F'} \rangle = |F \cap F'| \pmod{p}$. Unfortunately we can't conclude that these vectors are linearly independent like we could in the Oddtown case, but we can do this if we generalize our vectors somewhat.

To this end, for each $F \in \mathcal{F}$, define the polynomial $p_F : \mathbb{F}_p^n \rightarrow \mathbb{F}_p$ by

$$p_F(\vec{x}) = \prod_{\ell \in L} (\langle \chi_F, \vec{x} \rangle - \ell).$$

Note that by hypothesis, we have $p_F(\chi_{F'}) = 0$ if $F \neq F' \in \mathcal{F}$ (since $\langle \chi_F, \chi_{F'} \rangle \equiv |F \cap F'| \equiv \ell \pmod{p}$ for some $\ell \in L$) and that $p_F(\chi_F) \neq 0$ (since $|F| \not\equiv \ell \pmod{p}$ for any $\ell \in L$).

We claim that the polynomials $\{p_F : F \in \mathcal{F}\}$ are linearly independent. Indeed, say we had

$$\sum_{F \in \mathcal{F}} \lambda_F p_F = 0.$$

By plugging in $\chi_{F'}$ for some $F' \in \mathcal{F}$ into both sides of this equality, the observation above implies $c_{F'} \lambda_{F'} = 0$ for some $c_{F'} \neq 0$, and hence $\lambda_{F'} = 0$, proving the claim.

Observe that each polynomial p_F has degree at most $|L|$. It is a basic exercise to show that the dimension of the space of polynomials in n variables with degree at most $|L|$ is equal to $\sum_{i=0}^{|L|} \binom{n+i-1}{i}$. Since we constructed a set of $|\mathcal{F}|$ linearly independent polynomials that lie in a space of dimension $\sum_{i=0}^{|L|} \binom{n+i-1}{i}$, we must have $|\mathcal{F}| \leq \sum_{i=0}^{|L|} \binom{n+i-1}{i}$, proving the result. **Maybe add notation like $V_{n,\leq d}$ for this vector space.** \square

The bound of Proposition 21.2 is relatively close to tight. Indeed, if $L = \{0, 1, \dots, s-1\}$, then $\mathcal{F} = \binom{[n]}{s}$ is (p, L) -modular intersecting and has size $\binom{n}{s} \approx \sum_{i=0}^s \binom{n+i-1}{i}$. However, one can do better by refining our argument (though we emphasize that the present weaker bound of Proposition 21.2 is all one typically needs for most applications).

Towards refining this proof/bound, one might first turn to the simplest case of $p = 2$, $L = \{0\}$. As this is just Oddtown, we know the answer here should be n , but the bound we get is $n + 1$. By carefully analyzing the proof, one realizes that the one place our argument isn't sharp is when we argue about the dimension of the space spanned by the p_F polynomials. Indeed, we naively said that they lived in the space of degree at most 1 polynomials, but in fact they are all in the span of the monomials $\{x_i\}$, which gives the optimal bound.

We are thus left with the problem of trying to find a smaller subspace for us to work in while achieving the same conclusion, and there are a couple of semi-standard ways of doing this. One way is to try and find polynomials which are more “efficient”, i.e. of lower degree, for which our conclusion still holds. And indeed, if you look back through the proof of Proposition 21.2, one sees that although we defined our polynomials to have codomain \mathbb{F}_p^n , we only fed them 0-1 vectors to show linear independence. As such, it would suffice to look at polynomials \bar{p}_F which agree with the p_F polynomials on 0-1 vectors, and this can be done simply by replacing each x_i^α in the expansion of p_F by x_i (i.e. by replacing p_F by a multilinear function). For example, if $p_F(\vec{x}) = 3x_1x_2^2 + 2x_1^3x_3^2$ then we would want to look at $\bar{p}_F(\vec{x}) = 3x_1x_2 + 2x_1x_3$. By using this we can prove the following.

Theorem 21.3. *For any prime p and $L \subseteq \{0, 1, \dots, p-1\}$, if $\mathcal{F} \subseteq 2^{[n]}$ is (p, L) -modular intersecting then*

$$|\mathcal{F}| \leq \sum_{i=0}^{|L|} \binom{n}{i}.$$

We note that this theorem was originally proven by Deza, Frankl, and Singhi [35], with the following simpler proof due to Alon, Babai, and Suzuki [2].

Proof. Define p_F as before, and define the multilinear polynomial \bar{p}_F by replacing each x_i^α in the expansion of p_F by x_i . Note that $p_F(\chi_{F'}) = \bar{p}_F(\chi_{F'})$ for all F' , so by following the same proof as before we conclude that the \bar{p}_F polynomials are linearly independent. Moreover, these linearly independent polynomials all lie in the space of multilinear polynomials in n variables with degree at most $|L|$, which is a space of dimension $\sum_{i=0}^{|L|} \binom{n}{i}$. We thus must have $|\mathcal{F}| \leq \sum_{i=0}^{|L|} \binom{n}{i}$, proving the result. \square

The reader might still complain that, although we have improved our bound, it still fails to be tight for Oddtown. As far as we are aware, it is still an open problem as to whether the present bound can be replaced by $|\mathcal{F}| \leq \binom{n}{|L|}$ for all L , which would be tight whenever $L = \{0, 1, \dots, s-1\}$. However, such a general result is known to hold when \mathcal{F} is uniform.

Theorem 21.4 (Frankl-Wilson [48]). *For any prime p and $L \subseteq \{0, 1, \dots, p-1\}$, if $\mathcal{F} \subseteq 2^{[n]}$ is (p, L) -modular intersecting and k -uniform, then*

$$|\mathcal{F}| \leq \binom{n}{|L|}.$$

We omit their proof, which utilizes the powerful tool of incidence matrices, and instead refer the reader to [8, Chapter 7] for more on this. Instead, we will prove a weaker version of this result which holds in the non-modular setting.

For this, given a set of integers L , we say that a set system \mathcal{F} is L -intersecting if $|F \cap F'| \in L$ for all $F \neq F' \in \mathcal{F}$.

Theorem 21.5 (Ray-Chaudhuri-Wilson [94]). *If L is a set of integers and $\mathcal{F} \subseteq 2^{[n]}$ is L -intersecting and k -uniform, then*

$$|\mathcal{F}| \leq \binom{n}{|L|}.$$

Proof. Note that we may assume $L \subseteq \{0, 1, \dots, k-1\}$ since these are the only intersections that can actually occur, and in particular we may assume $|L| \leq k$.

Similar to before, we define polynomials $p_F : \mathbb{R}^n \rightarrow \mathbb{R}$ by $p_F(\vec{x}) = \prod_{\ell \in L} (\langle \chi_F, \vec{x} \rangle - \ell)$ and we define \bar{p}_F to be their multilinearizations. As before, we find that these \bar{p}_F polynomials are linearly independent, giving a bound of $\sum_{i=0}^{|L|} \binom{n}{i}$ since these polynomials lie in the vector space V of multilinear polynomials of degree at most $|L|$.

Observe that the bound above fails to be sharp if and only if these \bar{p}_F do not span all of V . In particular, to prove the result, it suffices to find a set of $\sum_{i=0}^{|L|-1} \binom{n}{i}$ polynomials of V which are linearly independent with the \bar{p}_F vectors.

Say we write these supposed polynomials as q_I for $I \subseteq [n]$ with $|I| \leq |L| - 1$, what would be a good choice for these? Well, to mimic the previous proof we would like to have $q_I(\chi_F) = 0$ for all $F \in \mathcal{F}$. Since all we know about the individual elements of \mathcal{F} is that they each of size k , perhaps a reasonable condition is to demand

$$q_I(\vec{x}) = q'_I(\vec{x}) \left(\sum_{j=1}^n x_j - k \right),$$

where q'_I is some appropriate polynomial of degree at most $|L| - 1$. Indeed, if we do this and if we have

$$\sum_F \lambda_F \bar{p}_F + \sum_I \lambda_I q_I = 0,$$

then by plugging in $\chi_{F'}$ into both sides of this equality we see that $\lambda_F = 0$ for all $F \in \mathcal{F}$. As such, all we have to do is choose the q'_I polynomials such that the q_I vectors themselves are

linearly independent. One possibility is to set

$$q_I(\vec{x}) = \prod_{i \in I} x_i \cdot \left(\sum_{j=1}^n x_j - k \right).$$

Note that with this, for any set $J \subseteq [n]$ with $|J| \leq |L| - 1 < k$, we have $q_I(\chi_J) = 0$ if $I \not\subseteq J$, and we have $q_I(\chi_J) \neq 0$ if $I \subseteq J$ (here we implicitly use $|J| < k$ to ensure $\sum_{j=1}^n (\chi_J)_j - k \neq 0$).

With this in mind, say we had

$$\sum_I \lambda_I q_I = 0,$$

and let J be a smallest set such that $\lambda_J \neq 0$. By plugging χ_J into both sides of this expression, the observation above implies that we get $c_J \lambda_J = 0$ for some $c_J \neq 0$, and hence $\lambda_J = 0$, a contradiction. We conclude that the q_I polynomials are linearly independent, and hence all of the polynomials $\{\bar{p}_F\}_F \cup \{q_I\}_I$ are linearly independent vectors. Unfortunately the q_I polynomials aren't in V , i.e. aren't multilinear, but this can be easily remedied by considering $\{\bar{p}_F\}_F \cup \{\bar{q}_I\}_I$, and again we conclude that these vectors (in V) are linearly independent, giving the desired bound on $|\mathcal{F}|$. \square

We note that the exact proof as written almost goes through if instead of demanding $|F| = k$ for $F \in \mathcal{F}$, we only demand $|F| \equiv k \pmod{p}$. A careful reading shows that this will go through if we assume $k \notin [|L| - 1] \pmod{p}$ (which holds if e.g. $|L| \leq k$), but without this assumption we can't assume $q_I(\chi_J) \neq 0$ whenever $I \subseteq J$. Nevertheless, it does turn out that this bound does continue to hold under the weaker hypothesis $|F| \equiv k \pmod{p}$, but of the proof of this ends up being somewhat more involved and we refer the interested reader to [8, Theorem 5.37].

22 Pseudo-Adjacency Matrices

Given a graph G , we define its adjacency matrix A to be the matrix whose rows and columns are indexed by $V(G)$ and where $A_{i,j} = 1$ if $i \sim j$ and $A_{i,j} = 0$ otherwise. While A is a natural way to encode a graph G , it is not at all obvious that its spectral properties should tell you anything about G , but this remarkably does turn out to be the case! For example, we'll see below that $\sigma(A)$ can give effective bounds on its maximum degree, as well as its independence number.

Many of these results which hold for A (and which perhaps are most naturally proven by thinking about A) can be tweaked to hold for a wider class of “pseudo-adjacency matrices”, which exhibit behavior similar to A but which are cooked up to deal with the specific problem at hand. We exhibit this phenomenon in the next two subsection. In each subsection, we begin by highlighting important lemmas relating the spectrum of A to G , after which we generalize these lemmas apply to a slightly broader class of pseudo-adjacency matrices M , and we then carefully choose such an M to solve a cool problem.

Before moving on, we note that much more broadly, the area of *spectral graph theory* concerns studying matrices M associated to graphs G and how the spectral properties of M relate to combinatorial properties of G . We will not have time to dive very deeply into this fascinating area, and we refer the interested reader to [the appendix](#) for more on this.

22.1 Huang's Theorem

In this subsection we give a simple proof of Huang's, which solved what used to be a 30 year old problem known as the sensitivity conjecture. Our proof relies on two basic results from linear algebra, both of which are used heavily throughout spectral graph theory. First we have the Rayleigh quotient, which gives an analytic way to compute the eigenvalues of a symmetric matrix. Here $\lambda_1(M)$ denotes the largest eigenvalue of M .

Lemma 22.1. *Let M be a real symmetric matrix. Then*

$$\lambda_1(M) = \max_{x \neq 0} \frac{x^* M x}{x^* x},$$

and any x achieving equality is an eigenvector corresponding to $\lambda_1(M)$.

As a small application [Actually do we even use the Raleigh quotient here?](#), we use the Rayleigh quotient to establish a connection between eigenvalues and combinatorial information of graphs.

Lemma 22.2. *Let G be a graph and A its adjacency matrix. Then*

$$\lambda_1(A) \leq \Delta,$$

where Δ is the maximum degree of G .

Proof. let x be an eigenvector of A corresponding to $\lambda_1(A)$ and let $v \in V(G)$ be such that $|x_v|$ is maximized. Then by our definitions, we have

$$|\lambda_1(A)x_v| = |(Ax)_v| = \left| \sum_u A_{v,u}x_u \right| \leq \sum_{u \sim v} |x_u| \leq \deg(v)|x_v| \leq \Delta|x_v|.$$

This shows $|\lambda_1(A)| \leq \Delta$, proving the result. \square

Examining this proof, we see that we hardly used any of the properties of A in our argument. In particular, word for word the same argument gives the following.

Lemma 22.3. *Let G be a graph and M a symmetric matrix such that $M_{i,j} = \pm 1$ if $ij \in E(G)$ and $M_{i,j} = 0$ otherwise. Then*

$$\lambda_1(M) \leq \Delta,$$

where $\lambda_1(M)$ is the largest eigenvalue of M and Δ is the maximum degree of G .

Next we have the Cauchy interlacing theorem, which allows us to bound the eigenvalues of submatrices of B in terms of the eigenvalues of B .

Theorem 22.4 (Cauchy interlacing theorem). *Let B be a real symmetric $n \times n$ matrix and C an $m \times m$ principal submatrix of B with $m \leq n$. If B has eigenvalues $\lambda_1 \geq \dots \geq \lambda_n$ and C has eigenvalues $\mu_1 \geq \dots \geq \mu_m$, then for all i*

$$\lambda_i \geq \mu_i \geq \lambda_{i+n-m}.$$

Remark 22.5. Personally, I always forget the exact statement of Cauchy's interlacing theorem, so here's a bit of "mnemonic" to help remember it. By the Rayleigh quotient we always have

$$\mu_1 = \max_{x \neq 0} \frac{x^* C x}{x^* x} \leq \max_{x \neq 0} \frac{x^* B x}{x^* x} = \lambda_1,$$

so that gives the first inequality. We also have $\mu_1 \geq \lambda_1$ if $m = n$, and each time we decrease m this bound has to get weaker, so we end up getting $\mu_1 \geq \lambda_{1+n-m}$ in general. We then get the rest of the inequalities by translating $\lambda_1 \geq \mu_1 \geq \lambda_{1+n-m}$ by $i - 1$.

With these standard spectral graph theory lemmas established, we can now state the main theorem of this subsection.

Theorem 22.6 (Huang [65]). *Let Q_n be the hypercube graph on 2^n vertices. If $V \subseteq V(Q_n)$ is a subset of size $2^{n-1} + 1$, then the induced subgraph $Q_n[V]$ has maximum degree at least \sqrt{n} .*

This result is sharp in several ways. First, it is easy to find subsets of size 2^{n-1} such that $Q_n[V]$ is the empty graph, so in order to get any non-trivial lower bound on the maximum degree one needs V to have size at least $2^{n-1} + 1$. Second, Chung et. al. [25] proved that there exist choices of V such that $Q_n[V]$ has maximum degree $\lceil \sqrt{n} \rceil$, so this bound is essentially best possible.

It was shown by Gotsman and Linial [59] that proving a result of this form is equivalent to showing that two notions of "sensitivity" for Boolean functions are equivalent, which led to a

great deal of interest in resolving it. Nevertheless, it remained unanswered for 30 years until Huang came up with the following remarkable proof.

The key idea is to define the $2^n \times 2^n$ matrix B_n recursively by

$$B_0 = [0], \quad B_n = \begin{bmatrix} B_{n-1} & I \\ I & -B_{n-1} \end{bmatrix},$$

where here I denotes the identity matrix of dimension 2^{n-1} . Observe that if the negative sign in the definition of B_n wasn't there, then this would just define the adjacency matrix of Q_n . Thus this is a sort of “twisted adjacency matrix” which has -1 's in some of the positions where there are usually 1 's. This choice of signings turns out to spread out the spectrum of B_n in a nice way.

Lemma 22.7. *The spectrum $\sigma(B_n)$ consists of $\pm\sqrt{n}$ each occurring with multiplicity 2^{n-1} .*

Proof. It is straightforward to prove by induction that $B_n^2 = nI$, which implies that every eigenvalue λ of B_n satisfies $\lambda^2 = n$. Thus $\sigma(B_n)$ consists of $\pm\sqrt{n}$, and each must appear with equal multiplicity because $\text{Tr}(B_n) = 0$. \square

Shockingly, we have everything we need for our proof.

Proof of Theorem 22.6. Let $B = B_n$ be as described above. Let $V \subseteq V(Q_n)$ be any subset of size $2^{n-1} + 1$ and let C be the submatrix of B indexed by the rows and columns corresponding to B . Let $G = Q_n[V]$. Observe that C satisfies the conditions for M of Lemma 22.3 since B is a (symmetrically) signed version of the adjacency matrix. By Lemma 22.3, the Cauchy interlacing theorem, and the previous lemma, we conclude that

$$\Delta(G) \geq \lambda_1(C) \geq \lambda_{2^{n-1}}(B) = \sqrt{n},$$

proving the result. \square

22.2 Hoffman's Bound and Erdős-Ko-Rado

One of the most important results in spectral graph theory is Hoffman's bound, which gives an upper bound on the independence number of a regular graph in terms of its eigenvalues.

Lemma 22.8 (Hoffman). *Let G be a non-empty n vertex d -regular graph and A its adjacency matrix. Then*

$$\frac{\alpha(G)}{n} \leq \frac{-\lambda_{\min}}{d - \lambda_{\min}},$$

where λ_{\min} is the smallest eigenvalue of A .

Proof. Let I be an independent set of size $\alpha = \alpha(G)$, and let x be the characteristic vector with $x_i = 1$ if $i \in I$ and $x_i = 0$ otherwise. Observe that because I is an independent set, we have

$$x^T A x = 0.$$

Let v_1, \dots, v_n be an orthonormal eigenbasis for A with eigenvalues $\lambda_1, \dots, \lambda_n$. Since G is regular, the all 1's vector $\mathbf{1}$ is an eigenvector with eigenvalue equal to d , so we can assume $v_1 = \mathbf{1}/\sqrt{n}$ and $\lambda_1 = d$. Writing $x = \sum c_i v_i$ for some real numbers c_i , we see that

$$\alpha = x^T x = \sum c_i^2,$$

and

$$\alpha/\sqrt{n} = \langle x, v_1 \rangle = c_1.$$

Putting all of this together, we find

$$\begin{aligned} 0 = x^T A x &= x^T \sum c_i \lambda_i v_i = \sum c_i^2 \lambda_i = (\alpha^2/n)d + \sum_{i \neq 1} c_i^2 \lambda_i \\ &\geq (\alpha^2/n)d + \sum_{i \neq 1} c_i^2 \lambda_{\min} = (\alpha^2/n)d + (\alpha - \alpha^2/n)\lambda_{\min}. \end{aligned}$$

Dividing both sides by α and rearranging gives

$$\alpha(\lambda_{\min} - d)/n \geq \lambda_{\min}.$$

Dividing both sides by $\lambda_{\min} - d$ (which is negative because $\lambda_{\min} < 0$ since G is non-empty) gives the result. \square

This result gives yet another proof of Erdős-Ko-Rado.

Sketch of Proof of Erdős-Ko-Rado. Let G be the graph with vertex set $\binom{[n]}{r}$ where two sets are adjacent if and only if they are disjoint. Observe that independent sets of G are exactly intersecting families, so Erdős-Ko-Rado is equivalent to saying that

$$\alpha(G) = \binom{n-1}{r-1}$$

provided $n \geq 2r$. In this case, one can verify that G has eigenvalues $(-1)^j \binom{n-r-j}{r-j}$ for $0 \leq j \leq r$ (each appearing with multiplicity $\binom{n}{j} - \binom{n}{j-1}$), and hence

$$\lambda_{\min} = -\binom{n-r-1}{r-1} = -\frac{r}{n-r} \binom{n-r}{r}.$$

Since G has $\binom{n}{r} = \frac{n}{r} \binom{n-1}{r-1}$ vertices and is regular with degree $\binom{n-r}{r}$, plugging things into Lemma 22.8 gives $\alpha(G) \leq \binom{n-1}{r-1}$ as desired. \square

Perhaps the biggest constraint in Lemma 22.8 is the need for G to be regular, which we used in this proof to guarantee that the all 1's vector is an eigenvector, allowing us to extract $\alpha = \langle x, \mathbf{1} \rangle$. We can get around this issue if we change how we measure independent sets. To this end, given a graph G together with a vector w indexed by $V(G)$ and a set of vertices I , define $|I|_w = \sum_{i \in I} w_i^2$, and define $\alpha_w(G) = \max_I |I|_w$ where I ranges over all independent sets of G .

With this we can generalize Lemma 22.8 by both allowing G not to be regular, and by allowing A to only be a “pseudo-adjacency matrix.” It will also be important for our main application to allow G to have loops, where we emphasize that no independent set of G can contain a vertex which has a loop.

Lemma 22.9. *Let G be a graph with loops and M a matrix such that $M_{i,j} = 0$ whenever $i \not\sim j$ and such that M has a basis of eigenvectors. If λ_{\min} is the smallest eigenvalue of M , and if w is a unit eigenvector of M with eigenvalue $\lambda > \lambda_{\min}$, then*

$$\alpha_w(G) \leq \frac{-\lambda_{\min}}{\lambda_{\min} - \lambda}.$$

The proof of Lemma 22.9 is nearly identical to that of Lemma 22.8, and we leave the details as an exercise to the reader.

In order to use Lemma 22.9, we need a problem where we care about independent sets which are weighted in non-standard ways. Motivated by our application of Lemma 22.8, we will do this for a weighted version of the Erdős-Ko-Rado problem.

Specifically, given an n -vertex set system \mathcal{F} , we define its p -biased measure by

$$\mu_p(\mathcal{F}) = \sum_{F \in \mathcal{F}} (1-p)^{n-|F|} p^{|F|}.$$

In this setting, our main question is: what is the largest size (in the p -biased sense) of an intersecting set system? Here we emphasize that we consider the non-uniform families, as otherwise this is just equivalent to Erdős-Ko-Rado. The answer to this question is more or less what one would expect (after observing that $\mu_p(\mathcal{F}) = p$ when \mathcal{F} consists of all sets containing a given element).

Theorem 22.10. *If \mathcal{F} is an n -vertex intersecting family and $p \in [0, 1/2]$, then $\mu_p(\mathcal{F}) \leq p$.*

One can show that this result is false for $p > 1/2$. It also turns out that for $p < 1/2$ the unique construction with $\mu_p(\mathcal{F}) = p$ is the star, but we'll refrain from going into this.

One can also get uniqueness but I don't know how much extra work this is. Also note that this result is false for larger p

Just like with the original EKR theorem, there exist many proofs of Theorem 22.10 (in fact, Friedgut has a talk going through 5 and a half proofs of this theorem [REF](#)). The spectral proof we present here is due to Friedgut [50], and this proof ends up having the advantage that a closer analysis of the proof gives stability results.

Mimicking our proof of EKR using Lemma 22.8 described above, we define the graph G_n on $2^{[n]}$ where two sets are adjacent if and only if they are disjoint. Note that G has a loop at the vertex \emptyset . Again, independent sets of G_n are intersecting families, and proving Theorem 22.10 is equivalent to showing $\alpha_w(G_n) \leq p$ where w is the vector defined by $w_S = (1-p)^{n-|S|} p^{|S|}$ (which is a unit vector).

To complete the proof using Lemma 22.9, it remains to cook up a pseudo-adjacency matrix M_n which in particular has w as an eigenvector. To develop some intuition, we focus on the case $n = 1$. Here G_1 is just an edge where one vertex has a loop, so the M_1 we are looking must have the form $M_1 = \begin{bmatrix} * & * \\ * & 0 \end{bmatrix}$. To limit our search space a little, we will try and find such a matrix which is symmetric (though this isn't required to use Lemma 22.9), at which point we

can scale M appropriately so that it has the form

$$M_1 = \begin{bmatrix} c & 1 \\ 1 & 0 \end{bmatrix},$$

for some value c to be determined. Again, we must have that $w = (\sqrt{1-p}, \sqrt{p})$ is an eigenvector for this matrix, which means there exists some λ with $\lambda w_1 = cw_1 + w_2$ and with $\lambda w_2 = w_1$. Plugging things in, we see that $\lambda = \sqrt{(1-p)/p}$, and hence that

$$M_1 = \begin{bmatrix} \sqrt{\frac{1-p}{p}} - \sqrt{\frac{p}{1-p}} & 1 \\ 1 & 0 \end{bmatrix}$$

We also observe that the other eigenvector of this matrix is $w' = (\sqrt{p}, -\sqrt{1-p})$, which has eigenvalue $\lambda' = -\sqrt{p/(1-p)}$. Plugging this into Lemma 22.9, we find

$$\alpha_p(G) \leq \frac{\sqrt{p/(1-p)}}{\sqrt{(1-p)/p} + \sqrt{p/(1-p)}} = \frac{p}{1-p+p} = p,$$

proving the result when $n = 1$.

It remains to generalize M_1 to work for larger n . While it isn't hard to do this directly, we'll do this in one go with the following linear algebra fact about tensor products.

Lemma 22.11. *Given real symmetric matrices B_1, B_2 with rows and columns indexed by V_1, V_2 respectively, define the tensor product matrix $B_1 \otimes B_2$ by having its rows and columns indexed by $V_1 \times V_2$ with $(B_1 \otimes B_2)_{(i_1, i_2), (j_1, j_2)} = (B_1)_{i_1, j_1} (B_2)_{i_2, j_2}$.*

If $u_1, \dots, u_{|V_1|}$ and $v_1, \dots, v_{|V_2|}$ are orthonormal sets of eigenvectors for M_1, M_2 with u_i having eigenvalue μ_i and v_i having eigenvalue λ_i , then the vectors $u_i \otimes v_j$ defined by $(u_i \otimes v_j)_{(k_1, k_2)} = (u_i)_{k_1} (v_j)_{k_2}$ are an orthonormal set of eigenvectors for $M_1 \otimes M_2$ with eigenvalue $\mu_i \lambda_j$.

This is pretty easy to prove once you unwind the definitions. With this in mind, we inductively define $M_n = M_{n-1} \otimes M_1$ (i.e. M_n is the n th tensor power of M_1). Equivalently, this is the matrix defined by $(M_n)_{S,T} = 0$ whenever $S \cap T = \emptyset$ and which has $(M_n)_{S,T} = (\sqrt{\frac{1-p}{p}} - \sqrt{\frac{p}{1-p}})^{n-|S \cup T|}$ otherwise. The point is that by the lemma above, the vector w with $w_S = (1-p)^{(n-|S|)/2} p^{|S|/2}$ is an eigenvector corresponding to the eigenvalue $((1-p)/p)^{n/2}$. Further, if $p \leq 1/2$, then we have

$$\lambda_{\min} = \min_i (-1)^i (p/(1-p))^{i/2} ((1-p)/p)^{(n-i)/2} = -\sqrt{p/(1-p)} \cdot ((1-p)/p)^{(n-1)/2}.$$

Plugging this into Lemma 22.9 gives

$$\alpha_w(G_n) \leq \frac{\sqrt{p/(1-p)} \cdot ((1-p)/p)^{(n-1)/2}}{((1-p)/p)^{n/2} + \sqrt{p/(1-p)} \cdot ((1-p)/p)^{(n-1)/2}} = p,$$

proving Theorem 22.10.

Comment on uniqueness and stability; maybe go through this for $t = 1$.

With some work, one can extend the approach of this subsection to work for t -intersecting families. Unfortunately the most naive approach of solving the problem for $n = 1$ and then tensorizing doesn't work (since the $n = 1$ case is too small to capture the situation for $t > 1$). The way Friedgut gets around this in [50] is by doing linear algebra over the ring $\mathbb{R}[X]/(X^t = 0)$ instead of just \mathbb{R} . In this setting, he roughly considers matrices where the S, T entry is equal to $c_{S,T}X^{|S \cap T|}$ for some real number $c_{S,T}$, which makes it so that these entries are 0 whenever $|S \cap T| \geq t$. A lot of the steps become more intricate in this setup, but eventually everything does end up going through.

23 The Polynomial Method I: Slice Rank

Roughly speaking, the *polynomial method* is the method of using polynomials to solve problems, though there is no real agreed upon definition of what does and does not constitute an instance of this method.

One might argue that the modular intersection section used this method, in that we obtained our results by cooking up a system of (low degree) polynomials which “encoded” our problem (i.e. by choosing them so that the roots of the polynomials corresponded to intersections being of a given size). Over the next two sections, we will look at two other systematic ways one can take (low degree) polynomials which “encode” your problem and translate them into bounds for an extremal problem: slice rank and combinatorial Nullstellensatz. We note that these two methods are independent of each other and can be read in any order.

23.1 Slice Rank

In the previous section we considered proofs using linear algebra. Here we go a step further and use *multilinear* algebra.

To this end, given a set S and an integer k , we let S^k denote the set of all k -tuples of elements of S . Given a field \mathbb{F} , we will say that any function of the form $f : S^k \rightarrow \mathbb{F}$ is a k -*tensor*. Note that when $k = 2$, we can express f as a matrix whose rows and columns are indexed by S , and as such we can think of k -tensors as “higher order” matrices. Our main goal here is to use “ranks” of tensors to obtain bounds for combinatorial problems, analogous to what we did in our second proof of Oddtown.

There are various non-equivalent ways one can generalize the notion of rank from matrices to tensors. One way is through *slice rank*. For this, we say that a k -tensor f is a *slice* if

$$f(x_1, \dots, x_k) = g(x_i)h(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_k),$$

where i is some integer, g is a 1-tensor, and h is a $(k - 1)$ -tensor. Note that when $k = 2$, slices are exactly rank 1 matrices, i.e. the outer product of two vectors g and h . We define the slice rank $sr(f)$ of a k -tensor f to be the smallest integer r such that f can be written as the sum of r slices. Again note that in the case $k = 2$ this exactly corresponds to the usual notion of rank.

Although not every property of matrix rank carries over to the setting of slice rank, one important property that does is the fact that the slice rank of a “diagonal tensor” equals the number of non-zero entries it has. To this end, we say that a tensor is *proper diagonal* if $f(\vec{x}) \neq 0$ if and only if $x_i = x_j$ for all i, j .

Lemma 23.1. *If $f : S^k \rightarrow \mathbb{F}$ is a proper diagonal k -tensor, then $sr(f) = |S|$.*

Proof Sketch. We first show $sr(f) \leq |S|$. For each $a \in S$, let $g_a(x)$ be the 1-tensor with $g_a(a) = 1$ and $g_a(b) = 0$ otherwise. We observe

$$f(\vec{x}) = \sum_{a \in S} g_a(x_1)f(a, x_2, \dots, x_k),$$

proving that f has slice rank at most $|S|$ as desired.

The proof of the lower bound $sr(f) \geq |S|$ is a somewhat fiddly induction argument that takes about two pages to do; we refer the reader to a nice writeup by Martinez [82, Lemma 2.2.9] for the full details. \square

Our general approach using Lemma 23.1 will be as follows:

- (1) Start with some set S which has some desired properties.
- (2) Using the properties of S , we construct a proper diagonal k -tensor $f : S^k \rightarrow \mathbb{F}$ which is “simple” (e.g. a polynomial of low degree).
- (3) Using the fact that f is “simple”, we find some explicit way to write f as the sum of r slices.
- (4) In total, steps (2) and (3) imply

$$|S| = sr(f) \leq r,$$

giving us an effective bound on sets S with our desired properties.

We note that this sort of approach was first used by Croot, Lev, and Pach [31]. This method was then adapted by Ellenberg and Gijswijt [39] before being systemetized by Tao [106].

We will use the general approach outlined above to give very short proofs of two problems which were previously thought to be incredibly difficult. Throughout this section, we suggest before the reader goes through each proof that they first sit down and try to construct an f as in the framework outlined above in order to solve the problem.

23.2 The Capset Problem

Given an abelian group G , we say that a triple of elements $(x, y, z) \in G^3$ forms a 3-term arithmetic progression (or 3AP for short) if $y - x = z - y$, and we say this is a non-trivial 3AP if we do not have $x = y = z$. A fundamental question in additive combinatorics is to determine how large a subset $S \subseteq G$ can be if it contains no non-trivial 3AP. For example, Roth’s theorem famously says that when $G = \mathbb{Z}$ such a set must have density 0.

Another natural case to consider is \mathbb{F}_p^n for prime p . The case $p = 2$ is trivial (if $S \subseteq \mathbb{F}_2^n$ contains two distinct elements x, y , then (x, y, x) forms a non-trivial 3AP). As such, the first interesting case is to look at \mathbb{F}_3^n , and in this setting sets $S \subseteq \mathbb{F}_3^n$ without non-trivial 3AP’s are referred to as cap sets. Using a difficult Fourier analytic argument, Bateman and Katz [13] showed that capsets can have size at most $O(3^n/n^{1+\epsilon})$ and Edel [38] gave the best known lower bound of about 2.2^n . At one point Tao mentioned that this capset problem was perhaps his favorite open problem and that he thought the answer was probably $(3 - o(1))^n$. It was thus a major shock when Ellenberg and Gijswijt [39] gave a remarkably short proof showing an upper bound of $(3 - \epsilon)^n$.

Theorem 23.2 (Ellenberg-Gijswijt [39]). *For all primes p , there exists a real number $\epsilon > 0$ such that any set $S \subseteq \mathbb{F}_p^n$ which contains no non-trivial 3AP has $|S| \leq (p - \epsilon)^n$.*

Again, the reader is encouraged to try and come up with a tensor f which might work to give the result before going through the details of this proof.

Proof. For slight ease of notation we will only deal with the case $p = 3$, the proof for general p being almost identical. As outlined above, our goal will be to use S to construct a proper diagonal tensor f of low degree. Given the inputs of the problem, perhaps the most natural kind of tensor to consider is one of the form $f : S^3 \rightarrow \mathbb{F}_3$, where again we want to ensure that if $x, y, z \in S$ then $f(x, y, z) \neq 0$ if and only if $x = y = z$. Critically, because of our hypothesis that S contains no non-trivial 3AP, we see that we want $f(x, y, z) \neq 0$ if and only if x, y, z is a 3AP, and as such our goal is essentially equivalent to constructing an f which is the indicator function for 3AP's!

By definition, (x, y, z) is a 3AP if and only if $y - x = z - y$, and rearranging gives $x - 2y + z = 0$. As such, we're essentially left with the problem of constructing an indicator function for $x - 2y + z \neq 0$, or equivalently that $x_i - 2y_i + z_i \neq 0$ for all i . And this is somewhat easy: simply take

$$f(x, y, z) = \prod_{i=1}^n (1 - (x_i - 2y_i + z_i)^2),$$

which one can readily check is an indicator function for 3AP's in S , and hence is a proper diagonal tensor.

It remains to estimate the slice rank of f . That is, we want to show that f can be written as the sum of a small number of functions of the form e.g. $g(x)h(y, z)$. To this end, we make the observation that by the definition of f given above, each of its monomials has degree at most $2n$, and as such each of its monomials has one of its x -degree, y -degree, or z -degree is at most $2n/3$. The idea now is to group each of these monomials with the same low degree variable and then bound the slice rank of each of these groups by using the low degree variable.

To be somewhat more precise, given a vector $\alpha \in \{0, 1, 2\}^n$, define $x^\alpha = \prod x_i^{\alpha_i}$, and similarly define y^β and z^γ . Let $|\alpha| := \sum \alpha_i$, and let \mathcal{M} denote the set of all triples (α, β, γ) with $\alpha, \beta, \gamma \in \{0, 1, 2\}^n$ and $|\alpha| + |\beta| + |\gamma| \leq 2n$. Observe that each monomial of f is of the form $x^\alpha y^\beta z^\gamma$ for some $(\alpha, \beta, \gamma) \in \mathcal{M}$.

Let $\mathcal{M}_x \subseteq \mathcal{M}$ be the set of triples with $|\alpha| \leq 2n/3$, let $\mathcal{M}_y \subseteq \mathcal{M} \setminus \mathcal{M}_x$ be those with $|\beta| \leq 2n/3$, and let $\mathcal{M}_z \subseteq \mathcal{M} \setminus (\mathcal{M}_x \cup \mathcal{M}_y)$ be those with $|\gamma| \leq 2n/3$. As noted above, $\mathcal{M}_x \cup \mathcal{M}_y \cup \mathcal{M}_z$ partition \mathcal{M} . Given α with $|\alpha| \leq 2n/3$, define the polynomial

$$h_\alpha(y, z) = \sum_{\beta, \gamma: (\alpha, \beta, \gamma) \in \mathcal{M}_x} c_{\alpha, \beta, \gamma} y^\beta z^\gamma,$$

where $c_{\alpha, \beta, \gamma}$ is the (possibly 0) coefficient of $x^\alpha y^\beta z^\gamma$ in f . Similarly define $h_\beta(x, z)$ and $h_\gamma(x, y)$. By definition and the fact that $\mathcal{M}_x \cup \mathcal{M}_y \cup \mathcal{M}_z$ partition \mathcal{M} , we have

$$f(x, y, z) = \sum_{\alpha: |\alpha| \leq 2n/3} x^\alpha h_\alpha(y, z) + \sum_{\beta: |\beta| \leq 2n/3} y^\beta h_\beta(x, z) + \sum_{\gamma: |\gamma| \leq 2n/3} z^\gamma h_\gamma(x, y).$$

With this, we conclude that the slice rank of f is at most 3 times the number of terms in each sum, which is equal to the number of vectors $\alpha \in \{0, 1, 2\}^n$ with $|\alpha| \leq 2n/3$. It is not convince oneself that the number of such vectors is at most $(3 - \epsilon)^n$ for some ϵ (essentially because a random $\alpha \in \{0, 1, 2\}^n$ behaves like the sum of two binomial random variables with total expectation n), proving the result. \square

We note in particular in the case $p = 3$ one can go through the analysis and get an upper bound of roughly $|S| = O(2.756^n)$ for sets $S \subseteq \mathbb{F}_3^n$ not containing a capset. In fact, it turns out that this upper bound continues to hold for a slightly more general problem: we say that three sets $X, Y, Z \subseteq \mathbb{F}_3^n$ contain no rainbow 3AP if the only 3AP's $(x, y, z) \in X \times Y \times Z$ are trivial. With a bit more work one can show that the above proof yields that if X, Y, Z contains no rainbow 3AP then $|X| + |Y| + |Z| = O(2.756^n)$, and moreover there exist constructions that essentially match this upper bound. Thus any improvement to the bound for capsets has to somehow utilize that we only allow triples from S^3 and not from three distinct sets. We refer to the interested reader to the nice survey article by Grochow [61] for more on this topic.

23.3 Nonuniform Sunflowers

Recall that a k -sunflower is a collection of sets h_1, \dots, h_k such that $h_i \cap h_j$ is equal to the same set for all $i \neq j$. In the setting of uniform hypergraphs, the famous Erdős-Rado sunflower conjecture says that there exists a constant $C = C(k)$ such that every r -uniform hypergraph with at least C^r hyperedges contains a k -sunflower.

Here we consider an analogous conjecture in the non-uniform setting due to Erdős and Szemerédi [42]: there exists a constant $\epsilon = \epsilon(k) > 0$ such that for all $n \geq \epsilon^{-1}$, every $\mathcal{H} \subseteq 2^{[n]}$ with at least $(2 - \epsilon)^n$ hyperedges contains a k -sunflower. This conjecture turns out to be weaker than the Erdős-Rado sunflower conjecture [42].

The slice rank method allows us to give an easy proof of this non-uniform sunflower conjecture when $k = 3$.

Theorem 23.3 (Naslund-Sawin [90]). *If \mathcal{H} is an n -vertex hypergraph without any 3-sunflower, then*

$$|\mathcal{H}| \leq (n + 1) \sum_{t \leq n/3} \binom{n}{t} \leq 1.89^n.$$

Proof. Let S be the set of characteristic vectors corresponding to the hyperedges of \mathcal{H} . As a first attempt for this problem, one would probably try to work with the set S directly to construct a tensor f , but this causes some technical issues pop up.

To get around these issues, we let $S_r \subseteq S$ be those characteristic vectors with r 1's. The insight here is that $\max_r |S_r| \leq |S| \leq (n + 1) \cdot \max_r |S_r|$, so up to some negligible factor it suffices to bound the size of each S_r set. This will allow us to get around the previously mentioned issues.

With this, our goal is to construct a 3-tensor f , now with domain S_r^3 , such that $f(x, y, z) \neq 0$ if and only if $x = y = z$. The crucial insight here is that if we do not have $x = y = z$, then there exists some i such that $x_i + y_i + z_i = 2$. Indeed, if this were not the case and x, y, z were

all pairwise distinct from each other, then this would imply that every i is either contained in 0, 1, or 3 of the sets x, y, z , which would imply these distinct hyperedges form a 3-sunflower. If, say $x = y \neq z$, then the non-existence of such an i would imply $x \subseteq z$, a contradiction to $x \neq z$ both being in S_r , i.e. both having the same number of elements.

With this observation in mind, we see that our desired f is essentially the indicator function for [no coordinate i having $x_i + y_i + z_i = 2$]. Thus we can take $f : S_r^3 \rightarrow \mathbb{F}_3$ by defining

$$f(x, y, z) = \prod_{i=1}^n (x_i + y_i + z_i - 2),$$

and one can easily check that f is a proper diagonal tensor given that $S_r \subseteq S$ contains no 3-sunflowers, so it remains to bound the slice rank of f . Similar to before, we observe that each monomial of f has either x , y or z degree at most $n/3$, so by a similar trick as before we conclude

$$\max_r |S_r| \leq sr(f) \leq \sum_{t \leq n/3} \binom{n}{t} \leq 2^{nH(1/3)},$$

where $H(p) = -p \log_2(p) - (1-p) \log_2(1-p)$ is the binary entropy function. Evaluating this (and multiplying by $n+1$) gives the desired upper bound. \square

24 The Polynomial Method II: Combinatorial Nullstellensatz

Roughly speaking, the main idea of the previous section was to construct low degree polynomials f that were capable of encoding any set S with a given property; and from there we used slice rank to show that any such set S must be small in terms of $\deg(f)$.

In this subsection we take a subtly different approach: we construct polynomials f in terms of some given set S (generally with degree $|S|$), and then use the properties of S to argue that f must have relatively small degrees.

To emphasize, in the previous subsection we constructed f which was independent of the specific S we were given with some property, but here we construct f which will be defined based off of whatever S is given to us. This is somewhat more reminiscent of the approach we took with modular intersections.

In any case, the main tool we will use for this new approach is the following result of Alon's, which is inspired by Hilbert's Nullstellensatz theorem from algebraic geometry. Roughly speaking, it says that if f has small degree, then f can not vanish on a large cartesian product.

Theorem 24.1 (Alon's Combinatorial Nullstellensatz [1]). *Let \mathbb{F} be a field and $f \in \mathbb{F}[x_1, \dots, x_n]$ a polynomial. If t_1, \dots, t_n are non-negative integers such that $\deg(f) = \sum t_i > 0$ and such that the monomial $\prod x_i^{t_i}$ has a non-zero coefficient in f , then for any sets $S_i \subseteq \mathbb{F}$ satisfying $|S_i| > t_i$ for all i , there exist elements $s_i \in S_i$ such that $f(s_1, \dots, s_n) \neq 0$.*

We will somewhat informally write the conclusion of this statement as $f(S_1, \dots, S_n) \neq 0$ (and we will also write $f(S_1, \dots, S_n) = 0$ whenever this conclusion fails to hold). We note that this result is best possible, as can be seen by taking the function $\prod_{s \in S_1} (x_1 - s)g(x_2, \dots, x_n)$ for any set $S_1 \subseteq \mathbb{F}$ and function g .

Proof. We give a short proof due to Michałek [86] by induction on $\deg(f)$, the case $\deg(f) = 1$ being straightforward. Let f, t_1, \dots, t_n , and S_1, \dots, S_n be as in the hypothesis, and assume we have proven the result for all polynomials of degree smaller than $\deg(f) > 1$. Without loss of generality, we can assume $t_1 > 1$, and we let $s \in S_1$ be an arbitrary element. Using polynomial division, we can write

$$f(x_1, \dots, x_n) = (x_1 - s)g(x_1, \dots, x_n) + h(x_1, \dots, x_n)$$

such that $\deg(g) = \deg(f) - 1$ and such that h does not depend on x_1 . If we assume for contradiction that $f(S_1 \times \dots \times S_n) = 0$, then in particular we have

$$0 = f(\{s\} \times S_2 \times \dots \times S_n) = 0 + h(\{s\} \times S_2 \times \dots \times S_n).$$

Because h does not depend on x_1 , this implies $h(S_1 \times \dots \times S_n) = 0$. Again using that h does not depend on x_1 and the hypothesis of the theorem, g must have a monomial $x_1^{t_1-1} \prod_{i \neq 1} x_i^{t_i}$ with a non-zero coefficient, so by induction we can find some $(s_1, \dots, s_n) \in (S_1 \setminus \{s\}) \times S_2 \times \dots \times S_n$ which g does not vanish on. Note that $h(s_1, \dots, s_n) = 0$ by our previous observation, so in total we conclude that $f(s_1, \dots, s_n) = (s_1 - s)g(s_1, \dots, s_n) \neq 0$, a contradiction. \square

We will now showcase a slew of examples which use the combinatorial Nullstellensatz, with many more examples being found in Alon's original paper [1]. We note that in most cases, proofs of these results were known well before the introduction of the combinatorial Nullstellensatz, but this tool allows for elegant and unified solutions to all of them. Again, the reader is encouraged to try and figure out which polynomial to use before reading through the proof.

24.1 Cauchy-Davenport

We begin with a basic result from additive combinatorics. Given two subsets A, B of an abelian group G , we define $A + B = \{a + b : a \in A, b \in B\}$. If $G = \mathbb{Z}$, then it is not difficult to show that $|A + B| \geq |A| + |B| - 1$, which is best possible. However, if G is a finite group, then this bound no longer holds for the simple reason that one could have $|A| + |B| - 1 > |G|$. The Cauchy-Davenport theorem says that this is the only obstruction for \mathbb{F}_p .

Proposition 24.2 (Cauchy-Davenport). *If $A, B \subseteq \mathbb{F}_p$ for some prime number p , then $|A + B| \geq \min\{|A| + |B| - 1, p\}$.*

Proof. Let's think for a moment about what parameters we might choose when utilizing Theorem 24.1. Seemingly we should take $\mathbb{F} = \mathbb{F}_p$. Since we ultimately want to conclude something about the size of $A + B$, it seems reasonable that we want to construct our polynomial f to depend on $A + B$, and in particular to have degree equal to $|A + B|$. The only relevant sets we have that could play the role of the S_i are $A, B, A + B$ and \mathbb{F}_p . Since we're already planning to use $A + B$ to construct our polynomial, it perhaps makes the most sense to try and use $S_1 = A$ and $S_2 = B$.

Roughly then, we want to construct a polynomial $f(x, y)$ of degree equal to $|A + B|$, which means that if $|A + B| = \deg(f) \leq (|A| - 1) + (|B| - 1)$, then by the combinatorial Nullstellensatz we will have $f(A, B) \neq 0$. We want to then derive a contradiction from this and from this; that is, we want to define f in such a way that we obviously have $f(A, B) = 0$.

With the discussion above in mind, we want to define an f such that $f(A, B) = 0$ and such that f has degree $|A + B|$. An obvious candidate for this is to take

$$f(x, y) = \prod_{c \in A+B} (x + y - c),$$

which has all of the properties described above.

To complete the proof, we assume for contradiction that $|A + B| \leq \min\{|A| + |B| - 2, p - 1\}$. To apply the Nullstellensatz with $S_1 = A$, $S_2 = B$, we need to show that there exists a monomial $x^{t_1}y^{t_2}$ with $t_1 + t_2 = |A + B|$, $t_1 < |A|$, and $t_2 < |B|$. To this end, consider $t_1 = |A| - 1$ and $t_2 = |A + B| - |A| + 1 \leq |B| - 1$, with this inequality coming from our assumption on $A + B$. It is not difficult to see that the coefficient of $x^{t_1}y^{t_2}$ in f is equal to $\binom{|A+B|}{|A|-1}$. Crucially, because $|A + B| < p$, this binomial coefficient is not a multiple of p and hence is non-zero. We can thus apply the Nullstellensatz with $S_1 = A$, $S_2 = B$ to conclude that $f(A, B) \neq 0$, a contradiction to how we defined f . \square

24.2 Covering the Hypercube

How many hyperplanes in \mathbb{R}^n does it take to cover all of the points of the hypercube $\{0, 1\}^n$? It's not difficult to see that the minimum you need is 2. To make things more interesting, we ask the following variant for affine hyperplanes: how many hyperplanes do you need to cover all of $\{0, 1\}^n \setminus (0, \dots, 0)$ such that none of the hyperplanes contain $(0, \dots, 0)$? It isn't too difficult to work out that n hyperplanes suffice, but proving this is best possible isn't as easy. However, this becomes doable with the Nullstellensatz.

Proposition 24.3 (Alon-Füredi [4]). *If h_1, \dots, h_m are a set of hyperplanes in \mathbb{R}^n which do not contain the origin and which contain every other point of $\{0, 1\}^n$, then $m \geq n$.*

Proof. Again let us discuss what sort of parameters we might choose in applying the Nullstellensatz. The most natural choice for field is probably \mathbb{R} . Again our polynomial f should probably have degree m , and a natural way to do this is to take f to be a product over terms indexed by the h_j . Motivated by this, for each j let $a_j \in \mathbb{R}^n$ and $b_j \in \mathbb{R}$ be such that h_j contains all of the points $x \in \mathbb{R}^n$ satisfying $\langle x, a_j \rangle = b_j$, and we note for later that $b_j \neq 0$ for all j since the origin is not contained in any h_j . Then a natural choice of $f : \mathbb{R}^n \rightarrow \mathbb{R}$ to consider is

$$f(x) = \prod_{i=1}^m (b_j - \langle x, a_j \rangle),$$

since by hypothesis we know that our choice of f vanishes on all of $\{0, 1\}^n$ except the origin.

At this point, if we assume for contradiction that $m \leq n - 1$, then we know f has degree at most $n - 1$, and hence there's a chance we can find a monomial of the form $\prod_{i \neq k} x_i$ for some k with a non-zero coefficient. In this case we could apply the Nullstellensatz with $S_i = \{0, 1\}$ for $i \neq k$ and $S_k = \{1\}$ to find a point which isn't the origin that f fails to vanish on, a contradiction to how we defined f .

While the above approach is theoretically possible, it's not at all clear how you could correctly choose the value of k given the very limited information of the problem setup. Thus instead of doing the above, we'll modify our polynomial f so that it vanishes at every point of $\{0, 1\}^n$ (including the origin). Since f only fails to vanish at the origin, this can easily be done by taking

$$g(x) = \prod_{j=1}^m (b_j - \langle x, a_j \rangle) - \prod_{j=1}^m b_j \cdot \prod_{i=1}^n (1 - x_i),$$

and it is straightforward to check that g now vanishes on all of $\{0, 1\}^n$. Moreover, because $f(x)$ had degree $m \leq n - 1$ and because $\prod b_j \neq 0$ (as noted at the start of the proof); we have that $g(x)$ has degree equal to n with its only monomial of this degree equal to $\prod x_i$. Thus we can apply the Nullstellensatz with $S_i = \{0, 1\}$ for all i to conclude that $g(\{0, 1\}^n) \neq 0$, a contradiction to how we constructed g . We conclude that we must have $m \geq n$ as desired. \square

We note that this proof features a common trick with the Nullstellensatz: often you will have some “main term” in your polynomial which encodes the bulk of your problem, and from there you add in some additional terms to forbid certain “bad” witnesses s_1, \dots, s_n .

24.3 Regular Subgraphs mod p

Our last example concerns finding subgraphs of graphs which are “regular mod p ”. The proof will showcase a useful trick one can utilize together with the Nullstellensatz: over the finite field \mathbb{F}_p , the function $1 - x^{p-1}$ is the indicator function for x being equal to 0 (since $x^{p-1} = 1$ whenever $x \neq 0$).

Proposition 24.4. *Let p be a prime. If G is an n -vertex graph with more than $(p-1)n$ edges, then G has a non-empty subgraph H where every degree is a multiple of p .*

Proof. Again let's consider the parameters. We should take $\mathbb{F} = \mathbb{F}_p$. One might first consider taking $f : \mathbb{F}_p^n \rightarrow \mathbb{F}_p$ with our variables indexed by vertices of G , but in fact the more appropriate domain is $\mathbb{F}_p^{e(G)}$ where each variable x_e corresponds to an edge $e \in G$. This is because we ultimately want to construct a subgraph of G , which is most naturally thought of as a subset of the edges. Since we want our edges to either be in or out of this subgraph, it seems most natural to take $S_e = \{0, 1\}$, where we'll think of $x_e = 1$ meaning $e \in H$. As in the proof of Alon-Füredi, we'll need to ensure $f(0) = 0$ so that H will be non-empty.

Given this setup, we need to cook up an f which encodes the degrees of our vertices being a multiple of p in H . The degree of a vertex v in H is exactly equal to $\sum_{e \ni v} x_e$, and as noted before the proof, we can turn this into an indicator function for being a multiple of p by taking $1 - (\sum_{e \ni v} x_e)^{p-1}$. We again want to make sure we vanish at $x = 0$, so in total the function we come to is

$$f(x) = \prod_v \left(1 - \left(\sum_{e \ni v} x_e \right)^{p-1} \right) - \prod_e (1 - x_e^{p-1}).$$

Note that the degree of the left term is $(p-1)n$ while the degree on the right is $e(G) > (p-1)n$. Thus f is a polynomial of degree $e(G)$ with $\prod_e x_e^{p-1}$ a monomial achieving this. Thus we can apply the Nullstellensatz with $S_e = \{0, 1\}$ for all e to conclude there's some $x \in \{0, 1\}^n$ on which f does not vanish. By construction we know $x \neq 0$, and hence the graph $H = \{e : x_e \neq 0\}$ is a non-empty subgraph of G , which by construction has every vertex a multiple of p (since if some vertex failed to have this, then we would have $f(x) = 0$). \square

Part VII

Other Methods

25 The Delta-system Method

Recall that a k -sunflower (also called a delta-system) is a hypergraph \mathcal{S} with edges e_1, \dots, e_k such that there exists a set K called the *kernel* which has $e_i \cap e_j = K$ for all $i \neq j$. Roughly speaking, the Delta-system method is any proof using the following observation, which is usually credited to Deza, Erdős, and Frankl [34].

Lemma 25.1. *If \mathcal{H} is an r -graph which contains an $(r+1)$ -sunflower with kernel K , then for every edge $e \in \mathcal{H}$, there exists an edge $f \in \mathcal{H}$ with $e \cap f \subseteq K$.*

Proof. Let e_1, \dots, e_{r+1} be the edges of the sunflower. Since each of the sets $e_1 \setminus K, \dots, e_{r+1} \setminus K$ are non-empty disjoint sets, one of these $e_i \setminus K$ sets must be disjoint from e . Taking $f = e_i$ gives the result. \square

An effective tool to use in conjunction with this observation is Füredi's intersection semilattice lemma (which itself is proven using the Delta-system method). The full statement is a little intimidating, so we'll start by just stating a consequence of it.

Lemma 25.2 (Weak intersection semilattice lemma). *For all r, s , there exists $c = c(r, s) > 0$ such that for every r -graph \mathcal{H} , there exists a subgraph $\mathcal{H}' \subseteq \mathcal{H}$ with $|\mathcal{H}'| \geq c|\mathcal{H}|$ such that for all $e, f \in \mathcal{H}'$, $e \cap f$ is the kernel of an s -sunflower.*

That is, we can approximate \mathcal{H} by a hypergraph \mathcal{H}' such that any two edges of \mathcal{H}' are petals of a large sunflower. This quickly gives the following strengthening of the Erdős-Rado sunflower lemma due to Mubayi and Zhao [89].

Corollary 25.3 ([89]). *For all r, s there exists a constant $C = C(r, s)$ such that if \mathcal{H} is an n -vertex r -graph with $|\mathcal{H}| \geq Cn^{r-t-1}$, then \mathcal{H} contains an s -sunflower which has core of size at most t .*

Proof. We may assume n is sufficiently large in terms of r , as otherwise one can trivially find a sufficiently large C . Let $C = 2c^{-1}$ with c the constant from the previous lemma. Then $|\mathcal{H}'| \geq 2n^{r-t-1} > \binom{n-t-1}{r-t-1}$. By the Erdős-Ko-Rado theorem for $(t+1)$ -intersecting hypergraphs (see Theorem 11.4), \mathcal{H}' must contain two edges e, f which intersect in less than $t+1$ vertices. By assumption $e \cap f$ is the core of a sunflower with at least s petals, proving the result. \square

We'll now state the full intersection semilattice lemma. For this, if \mathcal{H} is an r -partite r -graph with partition $\bigcup V_i$ and if S is a set of vertices, then we define $\text{proj}(S) := \{i : S \cap V_i \neq \emptyset\}$. That is, $\text{proj}(S)$ records which coordinates its vertices are in. Given a hypergraph \mathcal{H} and a set of vertices S , define $d^*(S)$ to be the largest integer d such that there exist edges $e_1, \dots, e_d \in \mathcal{H}$ with $e_i \cap e_j = S$ for all $i \neq j$. In other words, $d^*(K)$ is the size of the largest sunflower which contains K as its kernel.

It's unfortunate that here \mathcal{H}' is the part we can approximate while it's the opposite for spread approximations. Probably change one of these, most likely the spread approximation one.

Lemma 25.4 (Intersection semilattice lemma). *For all r, s , there exists $c = c(r, s) > 0$ such that for every r -graph \mathcal{H} , there exists an r -partite subgraph $\mathcal{H}' \subseteq \mathcal{H}$ with $|\mathcal{H}'| \geq c|\mathcal{H}|$ and a hypergraph $\mathcal{J} \subseteq 2^{[r]}$ not containing the edge of size r such that:*

- (1) \mathcal{J} is intersection closed, i.e. $I, J \in \mathcal{J}$ implies $I \cap J \in \mathcal{J}$.
- (2) For every $e \in \mathcal{H}'$, $\{\text{proj}(e \cap f) : f \in \mathcal{H}' \setminus \{e\}\} = \mathcal{J}$.
- (3) $d_{\mathcal{H}'}^*(e \cap f) \geq s$ for all $e, f \in \mathcal{H}'$.

Note that if we ignore (1) and (2) we get back Lemma 25.2. Roughly speaking, this result says that we can approximate a large chunk of \mathcal{H} , namely \mathcal{H}' , by a small hypergraph \mathcal{J} such that for any edge $e \in \mathcal{H}'$, the hypergraph \mathcal{J} tells you exactly how other edges can intersect e , and moreover, (3) guarantees that each possible intersection occurs at least s times. We say that an \mathcal{H}' as in the conclusion of this lemma is (s, \mathcal{J}) -homogeneous.

We postpone proving this result for the moment and instead look at some consequences. For $\mathcal{J} \subseteq 2^{[r]} \setminus [r]$, define the rank

$$\text{rank}(\mathcal{J}) = \min\{|T| : T \subseteq [r], T \not\subseteq I \forall I \in \mathcal{J}\},$$

i.e. this is the smallest integer t such that there exists a t -set not contained in an edge of \mathcal{J} . For example, $\text{rank}(\mathcal{J}) > 1$ if and only if every vertex of $[r]$ is contained in an edge of \mathcal{J} .

Lemma 25.5. *If \mathcal{H}' is an n -vertex (s, \mathcal{J}) -homogeneous r -graph, then $|\mathcal{H}'| \leq \binom{n}{\text{rank}(\mathcal{J})}$.*

Note that s does not appear in this bound. Before looking at the proof, the reader may want to try proving this result for themselves when $\text{rank}(\mathcal{J}) = 1$ in order to get a sense for the definitions.

Proof. Let $T \subseteq [r]$ be a set such that $|T| = \text{rank}(\mathcal{J})$ and such that $T \not\subseteq J$ for all $J \in \mathcal{J}$. Given an edge $e \in \mathcal{H}'$, let $\phi(e) = e \cap \bigcup_{i \in T} V_i$. We claim that ϕ is injective. Indeed, if $\phi(e) = \phi(f)$, then $T \subseteq \text{proj}(e \cap f) \in \mathcal{J}$, a contradiction to our assumption on T . Since ϕ maps edges of \mathcal{H}' injectively to sets of size $\text{rank}(\mathcal{J})$, we conclude the result. \square

We can use this result to give yet another proof of the Erdős-Ko-Rado theorem for t -intersecting hypergraphs, whose statement we recall below.

Theorem 25.6. *Let \mathcal{H} be an n -vertex r -graph such that $|e \cap f| \geq t$ for all $e, f \in \mathcal{H}$. If n is sufficiently large in terms of r , then $|\mathcal{H}| \leq \binom{n-t}{r-t}$ with equality holding if and only if \mathcal{H} consists of every edge containing some fixed set T of size t .*

Proof. Apply Lemma 25.4 with $s = r + 1$ and let $\mathcal{H}', \mathcal{J}$ be the resulting hypergraphs with $\bigcup V_i$ the r -partition of \mathcal{H}' . First note that if $\text{rank}(\mathcal{J}) < r - t$, then by Lemma 25.5 we have

$$|\mathcal{H}| \leq c^{-1}|\mathcal{H}'| \leq c^{-1} \binom{n}{r-t-1} < \binom{n-t}{r-t},$$

where this last step holds for n sufficiently large in terms of $c = c(r, r+1)$. Thus if $|\mathcal{H}| \geq \binom{n-t}{r-t}$, we must have $\text{rank}(\mathcal{J}) \geq r-t$.

Let S be with $|S| = \text{rank}(\mathcal{J})$ such that no edge of \mathcal{J} contains S , and let $T = [r] \setminus S$. By the above we may assume $|S| \geq r-t$, and hence $|T| \leq t$. By definition of $\text{rank}(\mathcal{J}) = |S|$, for every $i \in S$, there exists an edge $J_i \in \mathcal{J}$ such that $S \setminus \{i\} \subseteq J_i$. Note that $i \notin J_i$ by assumption of S not being contained in any edge of \mathcal{J} , which means $J := \bigcap_{i \in S} J_i \subseteq T$. Because \mathcal{J} is intersection closed, we have $J \in \mathcal{J}$.

We claim that $|J| \geq t$. Indeed, by definition of $J \in \mathcal{J}$, there exist two edges of $\mathcal{H}' \subseteq \mathcal{H}$ whose intersection is exactly J , and by the t -intersecting property we must have $|J| \geq t$. Because $|T| \leq t$ and $J \subseteq T$, we conclude that $J = T \in \mathcal{J}$.

Now let $e \in \mathcal{H}'$ and $K = e \cap \bigcup_{i \in T} V_i$, noting that $|K| = t$. Then Lemma 25.4 guarantees that there is a sunflower with at least $r+1$ petals and K as its kernel. This implies that for every edge $f \in \mathcal{H}$, there exists an edge $e' \in \mathcal{H}'$ which contains K and which is disjoint from $f \setminus K$. Thus to have $|e' \cap f| \geq t$, we must have $K \subseteq f$. In other words, every edge of \mathcal{H} must contain the t -set K . This implies the result. \square

The above argument actually gives the following stability result: for all r, t there exists a constant $c' = c'(r, t)$ such that if \mathcal{H} is t -intersecting with $|\mathcal{H}| > c' \binom{n-t}{r-t}$, then there exists a set of size t which is contained in every edge of \mathcal{H} .

Remark 25.7. Stronger versions of Theorem 11.4 are known. For example, [REF](#) determined the maximum size of a t -intersecting family for *any* value n . In another direction, Frankl and Füredi [46] showed that the conclusion of the theorem holds if we only impose the hypothesis $|e \cap f| \neq t$ for $e \neq f$ provided $r \geq 2t+2$, with their proof using a somewhat more involved version of the Delta-system method.

A similar argument works for more general kinds of intersection problems. Given a set $L \subseteq \{0, 1, \dots, r-1\}$, we say that a hypergraph \mathcal{H} is an (n, r, L) -system if it's an n -vertex r -graph such that $|e \cap f| \in L$ for all $e, f \in \mathcal{H}$ distinct. For example, $(n, r, \{t, t+1, \dots, r\})$ -systems are t -intersecting hypergraphs. Little is known about how large (n, r, L) -systems can be for general L , but one can get effective bounds in terms of ranks. To this end, for any $L \subseteq \{0, 1, \dots, r-1\}$, define

$$\text{rank}(r, L) = \max_{\mathcal{J}} \text{rank}(\mathcal{J}),$$

where the maximum ranges over all $\mathcal{J} \subseteq 2^{[r]}$ containing no edges of size r which are intersection-closed with $|J| \in L$ for all $J \in \mathcal{J}$.

Theorem 25.8. *If \mathcal{H} is an (n, r, L) -system, then*

$$|\mathcal{H}| = O(n^{\text{rank}(r, L)}).$$

Proof. Let $\mathcal{H}', \mathcal{J}$ be as in Lemma 25.4. Observe that \mathcal{J} is intersection closed and that $|J| \in L$ for all $J \in \mathcal{J}$ (since otherwise two edges of $\mathcal{H}' \subseteq \mathcal{H}$ would fail to have $|e \cap e'| \in L$). Thus letting c be the constant from Lemma 25.4, we have

$$|\mathcal{H}| \leq c^{-1} e(\mathcal{H}') \leq c^{-1} \binom{n}{\text{rank}(\mathcal{J})} \leq c^{-1} \binom{n}{\text{rank}(r, L)},$$

where this second inequality used Lemma 25.5 and the last inequality used the definition of $\text{rank}(r, L)$. We conclude the result. \square

It's conjectured by Frankl that for all r, L there exist (n, r, L) -systems of size $\omega(n^{\text{rank}(r, L)-1})$. This is unknown in general, but see e.g. [47, Theorem 16.6] for a construction of size $\Omega(n^{1+1/(r-1)})$ whenever $\text{rank}(r, L) \geq 2$.

One of the best general bounds on the size of (n, r, L) -systems is the Deza-Erdős-Frankl theorem, which states that such a system \mathcal{H} satisfies

$$|\mathcal{H}| \leq \prod_{\ell \in L} \frac{n - \ell}{r - \ell} = O(n^{|L|}),$$

provided n is sufficiently large. One can prove this result using a variant of the Delta-system method, though we omit doing so here. Instead, we give an easy proof of the asymptotic result by utilizing the following.

Lemma 25.9. *Every $L \subseteq \{0, 1, \dots, r-1\}$ satisfies $\text{rank}(r, L) \leq |L|$.*

Proof. We prove the result by induction on $|L|$, the case $|L| = 0$ being trivial. Assume we have proven the result for all L with $|L| < k$. We first consider the case $|L| = k$ and $0 \in L$. Let \mathcal{J} be an intersection closed hypergraph on $2^{[r]} \setminus [r]$ with $|J| \in L$ for all $J \in \mathcal{J}$ and $\text{rank}(\mathcal{J}) = \text{rank}(r, L)$. For any vertex $x \in [r]$, let $\mathcal{J}_x = \{J - x : x \in J \in \mathcal{J}\}$ be the link hypergraph. If $L' = \{\ell - 1 : \ell \in L, \ell \neq 0\}$, then we see that \mathcal{J}_x is intersection closed with $|J| \in L'$ for all $J \in \mathcal{J}_x$. Because $|L'| = |L| - 1$, our inductive hypothesis implies that $\text{rank}(r-1, L') \leq |L| - 1$, i.e. there exists some set T of size $|L| - 1$ in $[r] \setminus \{x\}$ which is not contained in any edge of \mathcal{J}_x , which implies $T \cup \{x\}$ is a set of size $|L|$ not contained in any edge of \mathcal{J} . We conclude $\text{rank}(r, L) = \text{rank}(\mathcal{J}) \leq |L|$.

Now assume $0 \notin L$, and again let \mathcal{J} be an intersection closed hypergraph on $2^{[r]} \setminus [r]$ with $|J| \in L$ for all $J \in \mathcal{J}$ with $\text{rank}(\mathcal{J}) = \text{rank}(r, L)$. Let $I = \bigcap_{J \in \mathcal{J}} J$. Note that by definition I is contained in every edge of \mathcal{J} , and by the intersection closed property we have $I \in \mathcal{J}$, and hence $|I| \in L$. Define $L' = \{\ell - |I| : \ell \in L\}$, and note that the link hypergraph $\mathcal{J}_I = \{J \setminus I : J \in \mathcal{J}\}$ has all of its edge sizes lying in L' . Since $|L'| \leq |L|$ and $0 \in L'$, the previous case implies $\text{rank}(\mathcal{J}_I) \leq |L|$, which implies there exists some set $J \subseteq [r] \setminus I$ of size $|L|$ not contained in an edge of \mathcal{J}_I , and hence this set continues to not be contained in an edge of \mathcal{J} (since every edge of \mathcal{J} is the union of an edge of \mathcal{J}_I with I). We conclude $\text{rank}(r, L) = \text{rank}(\mathcal{J}) \leq |L|$, proving the result. \square

This together with the previous theorem immediately gives the following.

Corollary 25.10. *If \mathcal{H} is an (n, r, L) -system, then*

$$e(\mathcal{H}) = O(n^{|L|}).$$

It is not difficult to show that this result is tight if $L = \{0, 1, \dots, t-1\}$.

Before moving on, we note that while all of our applications here came from extremal set theory, the intersection semilattice lemma has application to other areas of extremal combinatorics as well. See for example [88], where this lemma is used to bound the Turán number of a class of linear hypergraphs called “expansions.”

25.1 Proof of the Semilattice Intersection Lemma

We first recall that for every r -graph \mathcal{H} there exists a subgraph $\mathcal{H}' \subseteq \mathcal{H}$ which is r -partite and which keeps a constant proportion of its edges. Thus it suffices to prove that if \mathcal{H} has r -partition $\bigcup V_i$, then one can find a subgraph $\mathcal{H}' \subseteq \mathcal{H}$ and $\mathcal{J} \subseteq 2^{[r]} \setminus [r]$ such that $|\mathcal{H}'| \geq c(r, s)|\mathcal{H}|$ and

- (1) \mathcal{J} is intersection closed, i.e. $I, J \in \mathcal{J}$ implies $I \cap J \in \mathcal{J}$.
- (2) For every $e \in \mathcal{H}'$, $\{\text{proj}(e \cap f) : f \in \mathcal{H}' \setminus \{e\}\} = \mathcal{J}$.
- (3) $d_{\mathcal{H}'}^*(e \cap f) \geq s$ for all $e, f \in \mathcal{H}'$.

Claim 25.11. *If there exists \mathcal{H}' , \mathcal{J} satisfying (2) and (3) with $s \geq r+1$, then they automatically satisfy (1).*

Proof. Let $J_1, J_2 \in \mathcal{J}$. This means that for any edge $e \in \mathcal{H}'$, there exist edges e_1, e_2 such that e, e_i intersect exactly in the coordinates of J_i and that this intersection is the kernel of a sunflower in \mathcal{H}' on at least $r+1$ petals. In particular, there must exist an edge $f \in \mathcal{H}'$ which contains $e \cap e_1$ and which is otherwise disjoint from e_2 (namely, f is one of the edges of the sunflower with core $e \cap e_1$). With this $\text{proj}(f \cap e_2) = J_1 \cap J_2$, so necessarily $J_1 \cap J_2 \in \mathcal{J}$. \square

With this claim in mind, we only have to find \mathcal{H}' , \mathcal{J} satisfying (2) and (3) (this is immediate if $s \geq r+1$, and for all other values of s we can take $c(r, s) = c(r, r+1)$ and apply the $s = r+1$ result). For the rest of the proof, given a hypergraph \mathcal{H} , we define

$$\mathcal{I}(\mathcal{H}) = \{\text{proj}(e \cap f) : e, f \in \mathcal{H}, e \neq f\}.$$

Note that if (2) and (3) are satisfied for some \mathcal{J} , then it must be that $\mathcal{J} = \mathcal{I}(\mathcal{H})$.

Claim 25.12. *For any r -partite r -graph \mathcal{H} , one can decompose \mathcal{H} as $\mathcal{H} = \mathcal{H}_0 \cup \bigcup_{I \in \mathcal{I}(\mathcal{H})} \mathcal{H}_I$ such that \mathcal{H}_0 satisfies (2) and (3) for some \mathcal{J} , and such that $I \neq \text{proj}(K)$ for any set K which is the kernel of a sunflower on at least s petals in \mathcal{H}_I .*

Proof. Initially start with $\mathcal{H}_0 = \mathcal{H}$ and $\mathcal{H}_I = \emptyset$ for all $I \in \mathcal{I}$. Consider the following procedure. If at any point \mathcal{H}_0 satisfies (2) and (3) for some set \mathcal{J} , then we stop and output the current sets. Otherwise, it is not difficult to see that there must exist some edge $e \in \mathcal{H}_0$ and $K \subseteq e$ such that $\text{proj}(K) \in \{\text{proj}(f \cap g) : f, g \in \mathcal{H}_0, f \neq g\}$ but K is not the kernel of a sunflower with at least s petals in \mathcal{H}_0 (otherwise the conditions would be satisfied with $\mathcal{J} = \{\text{proj}(f \cap g) : f, g \in \mathcal{H}_0, f \neq g\}$). Delete e from \mathcal{H}_0 and add it to $\mathcal{H}_{\text{proj}(K)}$.

We claim that this procedure gives the desired result. Indeed, \mathcal{H}_0 satisfies (2) and (3) by construction. Assume for contradiction that there existed some I with \mathcal{H}_I containing a sunflower on at least s petals with kernel K satisfying $\text{proj}(K) = I$. Let e be the first edge of this sunflower that was added to \mathcal{H}_I during the procedure. This implies that every edge of the sunflower was in \mathcal{H}_0 right before e was removed, i.e. that \mathcal{H}_0 contains a sunflower with at least s petals and kernel K . This contradicts us removing e from \mathcal{H}_0 at this step, so we include no such sunflower exists in any \mathcal{H}_I . \square

Claim 25.13. *Let \mathcal{H} be an r -partite r -graph and I a set such that $I \neq \text{proj}(K)$ for any K which is the kernel of a sunflower with at least s petals. Then one can decompose \mathcal{H} as $\mathcal{H} = \mathcal{H}^1 \cup \dots \cup \mathcal{H}^{r(s-1)}$ such that $I \notin \mathcal{I}(\mathcal{H}^j)$ for all j .*

Proof. Note that for each edge $e \in \mathcal{H}$, there exists a unique set $K \subseteq V$ with $\text{proj}(K) = I$. For each set of this form, let $\mathcal{H}(K)$ be its link hypergraph, i.e. $\mathcal{H}(K) = \{e \setminus K : K \subseteq e \in \mathcal{H}\}$. By hypothesis, none of the $\mathcal{H}(K)$ sets contain a matching of size s (since this translates to a sunflower of size s in \mathcal{H} with kernel K). It is not difficult to see that one can decompose each $(r - |K|)$ -graph $\mathcal{H}(K)$ into at most $(r - |K|)(s - 1) \leq r(s - 1)$ intersecting hypergraphs $\mathcal{H}^1(K), \dots, \mathcal{H}^{r(s-1)}(K)$ (e.g. by taking a largest matching M in $\mathcal{H}(K)$ and then assigning edges to $\mathcal{H}^i(K)$ if they contain the i th vertex which is contained in an edge of M). Let $\mathcal{H}^i[K] = \{e' \cup K : e' \in \mathcal{H}^i(K)\}$ and let $\mathcal{H}^i = \bigcup_K \mathcal{H}^i(K)$. Note that this decomposes \mathcal{H} , and that $K \neq e \cap f$ for any K with $\text{proj}(K) = I$ and $e, f \in \mathcal{H}^i$ (as this would imply $e \setminus K, f \setminus K \in \mathcal{H}^i(K)$, and hence e, f contain an additional vertex since $\mathcal{H}^i(K)$ is intersecting). \square

By repeatedly applying the above two times a bounded number of times, one can decompose \mathcal{H} as $\bigcup \mathcal{H}_i$ where each \mathcal{H}_i satisfies (2) and (3). Taking the largest of these hypergraphs gives the desired result.

25.2 Other Hypergraph Approximations

The results of this section are very similar in spirit to those of Section 11. To close this section, we briefly compare and contrast these results.

By using the Delta-system method (and more precisely the semi-lattice intersection lemma), we are able to conclude that many of the links of each $e \in \mathcal{H}'$ contains large sunflowers. In contrast, the spreadness of Theorem 11.2 not only gives us large sunflowers, but the stronger fact that a random partitioning of our vertex set is likely to give a large sunflower (which is a more robust condition). Moreover, the “error term” of Theorem 11.2 is typically much smaller than that of Lemma 25.4 (which only approximates a constant proportion of \mathcal{H} , which is in some sense necessary since it can only perfectly approximate r -partite r -graphs). On the other hand, the approximating hypergraph of Theorem 11.2 is more “complex”, in the sense that it is not just a hypergraph on $[r]$ but all of $V(\mathcal{H})$, and moreover here you lose the homogeneity of Lemma 25.4 which implies that every two edges have the same intersection pattern. In conclusion, the two methods are overall incomparable to each other, with each finding different uses in different situations.

We also note that there is another famous hypergraph approximation called the Junta method, which was developed by Dinur and Friedgut [36]. This roughly says that if \mathcal{H} is an intersecting hypergraph, then there exists a hypergraph \mathcal{J} on a small set of vertices J of \mathcal{H} such that almost every edge $e \in \mathcal{H}$ has $e \cap J \in \mathcal{J}$. We omit going into this in detail and refer the reader to [36].

Finally, we emphasize a fundamental weakness in all of these approaches, which is that they only work when n is quite large. In many cases we’re okay with this, but for some results like the t -intersecting Erdős-Ko-Rado theorem, it is of interest in nailing down the exact dependency on n . In cases such as these a more careful argument is needed.

26 Expansion and α -maximality

This almost surely needs to be rewritten to incorporate improvements to the bounds, as well as the claimed simplification mentioned at the end of “Towards the Erdős-Gallai Cycle Decomposition conjecture”

One of the nice features of random graphs is that they have good expansion properties; e.g. any set of vertices $B \subseteq V(G_{n,p})$ is likely to have about $pn|B|$ edges leaving B provided B is not too small. It is too much to ask that a graph has such strong expansion properties in general, but it is often the case that one can find subgraphs of arbitrary graphs which have reasonable expansion properties. There are many techniques in the field that achieve this end. The focus on this chapter will be an approach introduced by Tomon [108] which has the advantage of being both simple to state and powerful in applications.

Definition 3. Given a real number α , we say that a graph G is α -maximal if $e(G)/v(G)^{1+\alpha} = \max_{H \subseteq G} e(H)/v(H)^{1+\alpha}$. Equivalently, this says that if $e(G) = \gamma \cdot v(G)^{1+\alpha}$, then $e(H) \leq \gamma \cdot v(H)^{1+\alpha}$ for all $H \subseteq G$.

Observe that every graph has an α -maximal subgraph.

The motivation for this definition is that often in extremal graph theory, one wants to prove that graphs with $e(G) \geq \gamma v(G)^{1+\alpha}$ contain some desired structure. If such a result were true, then in particular any α -maximal subgraph of G must contain this structure, so being α -maximal is essentially the hardest case that one can consider. Moreover, it turns out that by reducing to α -maximal graphs, one gains a lot of nice expansion properties. Here and throughout this section we let $N(B)$ be the set of vertices $y \notin B$ which are adjacent to a vertex in B , and we let $d(G)$ denote the average degree of G .

Proposition 26.1. *Let G be an n -vertex α -maximal graph with $\alpha \in (0, 1]$ and $d(G) = \gamma n^\alpha$, and let $B \subseteq V(G)$ be such that $|B| \leq n/2$.*

- (i) *If G is non-empty, then $\gamma \geq \frac{1}{2}$.*
- (ii) *The minimum degree of G is at least $\frac{1}{2}d(G) = \frac{1}{2}\gamma n^\alpha$.*
- (iii) *We have $e(B, N(B)) \geq \frac{1}{4}\gamma n^\alpha |B|(1 + \alpha - (2|B|/n)^\alpha)$.*
- (iv) *We have $|N(B)| > |B|((1 + \frac{1}{2}\alpha)(\frac{n}{2|B|})^{\alpha/(1+\alpha)} - 1)$.*

The main benefit of (i) is that the bound is an absolute constant independent of α . Condition (ii) is obviously convenient to have. Note that in $G_{n,p}$ with $p = \gamma n^\alpha/n$, we have $\mathbb{E}[e(B, N(B))] \approx \gamma n^\alpha |B|$ as long as $|B| \leq n/2$, so the level of expansion in (iii) is about as much as we could hope for. The bound for (iv) is roughly $(n^\alpha |B|)^{1/(1+\alpha)}$, which is best possible when $|B| \approx n$ **Though beyond this I don't have much intuition for why this is a reasonable condition to shoot for.**

Proof. For (i), taking $H \subseteq G$ to be a single edge implies $d(H)/v(H)^\alpha = 2^{-\alpha} \geq \frac{1}{2}$, so the same bound holds for G .

For (ii), let v be a vertex of minimum degree δ and let $H = G - v$. We have $d(H)/v(H)^\alpha \leq d(G)/v(G)^\alpha$ by definition of α -maximality, which is equivalent to

$$\frac{d(G)n - 2\delta}{(n-1)^{1+\alpha}} \leq \frac{d(G)}{n^\alpha}.$$

This implies

$$\frac{1}{2}d(G)(n - \frac{(n-1)^{1+\alpha}}{n^\alpha}) \leq \delta,$$

showing $\delta \geq \frac{1}{2}d(G)$.

For (iii) and (iv), let $C = V(G) \setminus B$. With this we have

$$\begin{aligned} e(B, N(B)) &= e(G[B \cup C]) - e(G[B]) - e(G[C]) = \frac{1}{2}\gamma(|B| + |C|)^{1+\alpha} - e(G[B]) - e(G[C]) \\ &\geq \frac{1}{2}\gamma|C|^{1+\alpha}((1 + |B|/|C|)^{1+\alpha} - \frac{1}{2}\gamma|B|^{1+\alpha} - \frac{1}{2}\gamma|C|^{1+\alpha}). \end{aligned}$$

Using $(1 + |B|/|C|)^{1+\alpha} \geq 1 + (1 + \alpha)|B|/|C|$ and that $|C| \geq (n/2)^\alpha \geq \frac{1}{2}n^\alpha$, we find that this is at least

$$\frac{1}{2}\gamma(1 + \alpha)|B||C|^\alpha - \frac{1}{2}\gamma|B|^{1+\alpha} \geq \frac{1}{2}\gamma|B|(\frac{1}{2}(1 + \alpha)n^\alpha - |B|^\alpha),$$

giving (iii).

Similarly for (iv) we observe

$$e(G[B \cup N(B)]) \geq e(G[B \cup C]) - e(G[C]) \geq \frac{1}{2}\gamma(|B| + |C|)^{1+\alpha} - \frac{1}{2}\gamma|C|^{1+\alpha} \geq \frac{1}{2}\gamma(1 + \alpha)|B||C|^\alpha.$$

However, by α -maximality we have $e(G[B \cup N(B)]) \leq \frac{1}{2}\gamma(|B| + |N(B)|)^{1+\alpha}$. Combining these inequalities gives

$$|N(B)| \geq ((1 + \alpha)|B||C|^\alpha)^{1/(1+\alpha)} - |B|,$$

giving the result. \square

Our main application of α -maximal graphs will be to something called rainbow Turán numbers, which were first introduced by Keevash, Mubayi, Sudakov, and Verstraëte [72]. We say that a colored graph F is *rainbow* if all of the colors of its edges are distinct. Given a set of graphs \mathcal{F} , we define $\text{ex}^*(n, \mathcal{F})$ to be the maximum number of edges a *properly colored* n -vertex graph G can have without containing a rainbow copy of any $F \in \mathcal{F}$.

Note that $\text{ex}(n, \mathcal{F}) \leq \text{ex}^*(n, \mathcal{F})$ for all \mathcal{F} (since we can take any extremal \mathcal{F} -free graph and give each edge a distinct color), and in general these two quantities can be somewhat far from each other. Indeed, let \mathcal{C} denote the set of all cycles, which means $\text{ex}(n, \mathcal{C}) = n - 1$. On the other hand, we have $\text{ex}^*(n, \mathcal{C}) \geq n \log_2 n$ when n is a power of 2. This is because one can take G to be an n -vertex hypercube where an edge uv is colored i if u, v differ in the i th bit. It is not difficult to see that this is a proper coloring which contains no rainbow cycles.

Even though the problem of determining $\text{ex}(n, \mathcal{C})$ is easy, determining $\text{ex}^*(n, \mathcal{C})$ is an open and seemingly difficult problem. The first non-trivial upper bounds on $\text{ex}^*(n, \mathcal{C})$ were established by Das, Lee, and Sudakov [33], and later O. Janzer [66] managed to prove $\text{ex}^*(n, \mathcal{C}) = O((\log n)^4 n)$. Currently the best known upper bound is the following result due to Tomon [108].

Theorem 26.2 ([108]). *We have $\text{ex}^*(n, \mathcal{C}) = (\log n)^{2+o(1)}n$.*

The main lemma we need to prove this is the following. Here a Q -rainbow path refers to a rainbow path which only uses colors in the set Q .

Lemma 26.3. *Given $p_c \in (0, 1)$, there exists a constant C such that the following holds. Let $\lambda > C(\log \log n)^{10}$, and let G be an n -vertex α -maximal graph with proper coloring $c : E(G) \rightarrow R$ and $d(G) > C\lambda^2\alpha^{-2}n^\alpha$. If $Q \subseteq R$ is chosen by including each color independently and with probability p_c , then for every $v \in V(G)$, with probability at least $1 - O(\alpha^{-1}e^{-\Omega(\lambda^{1/2})})$ at least $n/3$ vertices of G can be reached by a Q -rainbow path.*

Before proving this lemma, let us first show how this implies the main result.

Proof of Theorem 26.2. Let G be an n -vertex graph with $e(G) \geq 2(\log n)^{2+\epsilon}n$ and $c : E(G) \rightarrow R$ a proper coloring, and let $\alpha = 1/\log_2(G)$ and $\lambda = (\log n)^{\epsilon/10}$. Let H be a subgraph of G maximizing $d(H)/v(H)^\alpha$ and $m = v(H)$. Note that H is α -maximal and $d(H) \geq d(G) \cdot (v(H)/v(G))^\alpha \geq \frac{1}{2}d(G)$ due to our choice of α , and this quantity is at least $C\lambda^2\alpha^{-2}m^\alpha$ for n sufficiently large.

Pick some $v \in V(H)$. Partition R into four parts Q_1, Q_2, Q_3, Q_4 be independently and uniformly at random assigning each color to one of these sets, and let B_i be the set of vertices that can be reached by v with a Q_i -path. By Lemma 26.3 with $p_c = 1/4$, we see that with probability at least $4/5$ we have $|B_i| \geq n/3$, so there exists some partition Q_1, \dots, Q_4 such that $|B_i| \geq n/3$ holds for all i .

Note that $B_i \cap B_j \neq \emptyset$ for some $i \neq j$, and let $w \in B_i \cap B_j$. By definition this means there exist rainbow paths P_i, P_j from v to w using colors in Q_i, Q_j . Thus the union of these two paths is a rainbow graph which contains a cycle, proving the result. \square

It remains to prove Lemma 26.3. Given a graph G and a proper coloring $c : E(G) \rightarrow R$, define $N_{Q,\phi}(v)$ with $\phi : V(G) \rightarrow 2^{V(G) \cup R}$ to be the set of vertices w with $vw \in E(G)$, $c(vw) \in Q \setminus \phi(v)$, and $w \notin \phi(v)$. That is, this is the neighborhood if we restrict to colors in Q and forbid some set of neighbors/colors for v to use. We define $N_{Q,\phi}(B) = \bigcup_{v \in B} N_{Q,\phi}(v) \setminus B$. To prove Lemma 26.3, we show that α -maximal graphs have vertex expansion about as strong as in Proposition 26.1 even when forbidding some colors/vertices.

Lemma 26.4. *Let $p_c, \alpha \in (0, 1]$, let n be a positive integer and $\lambda > 10^{10}$. Let G be an n -vertex graph, $c : E(G) \rightarrow R$ a proper edge coloring, and $B \subseteq V(G)$ such that the following hold:*

- G is α -maximal
- $d := d(G) \geq \lambda(p_c \cdot \alpha)^{-1}$,
- $\phi : V(G) \rightarrow 2^{V(G) \cup R}$ is such that $|\phi(v)| \leq d\alpha/32$ for all $v \in V(G)$, and
- $2\lambda^2 p_c^{-1} < |B| < n/2$.

Let $Q \subseteq R$ be obtained by including each color independently and with probability p_c . Then with probability at least $1 - e^{-\Omega(\lambda^{1/2})}$ we have

$$|N_{Q,\phi}(B)| \geq \frac{1}{4}|B| \min \left\{ \frac{d \cdot p_c \cdot \alpha}{64\lambda^{1/2}}, \left(\frac{n}{2|B|} \right)^{\alpha/(1+\alpha)} - 1 \right\}$$

Proof. Let $d = \gamma n^\alpha$, and let H be the bipartite (uncolored) graph on $B \cup N(B)$ such that $x \in B$ and $y \in N(B)$ are adjacent if $xy \in E(G)$, $y \notin \phi(x)$ and $c(xy) \notin \phi(x)$. Let $H_Q \subseteq H$ be the (random) subgraph which only includes edges xy with $c(xy) \in Q$. Thus our problem is equivalent to showing $|N_{H_Q}(B)|$ is large with high probability.

Since we're aiming something comparable to that of Proposition 26.1 (namely, this is basically what we get when the right term in the lemma achieves the minimum), one might try to just naively replicate that proof. This almost works, but to get things to occur with high probability we need the vertices of $N_H(B)$ to have large degrees. To this end, let $S \subseteq N_H(B)$ be the vertices w such that $|N_H(w) \cap B| \geq \lambda^{1/2} p_c^{-1} =: \Delta$, and let $T = N_G(B) \setminus S$.

Claim 26.5. *If $e_G(B, T) \leq d\alpha|B|/16$, then the result follows.*

Note that the claim involves edges of the original graph G , not H .

Proof. Let $C = V(G) \setminus B$, noting that $|C|^\alpha \geq (\frac{1}{2}n)^\alpha \geq \frac{1}{2}n^\alpha$, and since $d \geq \frac{1}{2}\gamma n^\alpha$ by Proposition 26.1, we conclude

$$e_G(B, T) \leq \frac{1}{8}\alpha\gamma|B||C|^\alpha.$$

Note that

$$E(G) = E(G[B \cup S]) \cup E(G[C]) \cup E(G[B, T]),$$

where $E(G[B', T])$ denotes the set of edges of G with one end in B' and the other in T . To see this, we note that vertices of B can only be adjacent to vertices of $B \cup N_G(B) = B \cup S \cup T$. With this we have

$$\begin{aligned} e_G(B \cup S) &\geq e(G) - e(G[C]) - e_G(B, T) \\ &\geq \frac{1}{2}\gamma(|B| + |C|)^{1+\alpha} - \frac{1}{2}\gamma|C|^{1+\alpha} - \frac{1}{8}\alpha\gamma|B||C|^\alpha, \end{aligned}$$

where this inequality used $e(G) = \frac{1}{2}\gamma n^{1+\alpha} = \frac{1}{2}\gamma(|B'| + |C|)^{1+\alpha}$, α -maximality, and the inequality noted above. Note that

$$(|B| + |C|)^{1+\alpha} = |C|^{1+\alpha}(1 + |B|/|C|)^{1+\alpha} \geq |C|^{1+\alpha} + (1 + \alpha)|B||C|^\alpha.$$

Using this gives

$$e_G(B \cup S) \geq \frac{1}{2}\gamma(1 + \alpha)|B||C|^\alpha - \frac{1}{8}\alpha\gamma|B||C|^\alpha \geq \frac{1}{2}\gamma(1 + \frac{1}{2}\alpha)|B||C|^\alpha.$$

By α -maximality we have $e_G(B \cup S) \leq \frac{1}{2}\gamma(|B| + |S|)^{1+\alpha}$, so in total this implies

$$|S| \geq \left((1 + \frac{1}{2}\alpha)|B||C|^\alpha \right)^{1/(1+\alpha)} - |B|.$$

For $w \in S$, let X_w be the indicator random variable for the event $w \notin N_{H_Q}(B)$. Then

$$\Pr[X_w = 1] = (1 - p_c)^{d_{H_Q}(w)} \leq (1 - p_c)^\Delta \leq e^{-\lambda^{1/2}}.$$

Thus if $X = \sum_{w \in S} X_w$ then $\mathbb{E}[X] \leq |S|e^{-\lambda^{1/2}}$, and by Markov's inequality this means $\Pr[X \geq |S|/2] \leq 2e^{-\lambda^{1/2}}$. This gives the result. \square

From now on we assume $e_G(B, T) > d\alpha|B|/16$. For each $w \in T$, let Y_w be the indicator random variable for the event $w \in N_{H_Q}(B)$. Then by the inequality $(1 - a)^b \geq 1 - \frac{1}{2}ab$ for $ab < \frac{1}{2}$,

$$\mathbb{E}[Y_w] = 1 - (1 - p_c)^{d_H(w)} \geq \frac{1}{2} \min\{1, p_c \cdot d_H(w)\}.$$

We partition T into two sets based on which term prevails in this minimum. Namely, let $T_1 = \{w \in T : d_H(w) \leq p_c^{-1}\}$ and $T_2 = T \setminus T_1$. If $Y = \sum_{w \in T} Y_w$, then

$$\mathbb{E}[Y] \geq \sum_{w \in T_1} \frac{1}{2} p_c \cdot d_H(w) + \frac{1}{2} |T_2| = \frac{1}{2} p_c e_H(B, T_1) + \frac{1}{2} |T_2| \geq \frac{1}{2} p_c e_H(B, T_1) + \frac{1}{2} \lambda^{-1/2} p_c e_H(B, T_2),$$

where this last step used that each vertex of $T_2 \subseteq T$ has degree at most $\Delta = \lambda^{1/2} p_c^{-1}$ in H . Thus

$$\mathbb{E}[Y] \geq \frac{1}{2} \lambda^{-1/2} p_c e_H(B, T).$$

By hypothesis,

$$e_H(B, T) \geq e_G(B, T) - \sum_{v \in B} |\phi(v)| \geq \frac{1}{32} d\alpha|B|.$$

This together with the hypothesis $d \geq \lambda^{1/2} (p_c \cdot \alpha)^{-1}$ implies $\mathbb{E}[Y] \geq \frac{1}{64} \lambda^{1/2} |B|$.

Note that Y is a function of which colors survive in Q . Each color appears at most $|B|$ times since G is properly colored, so changing Q by a single element changes Y by at most $|B|$, i.e. Y is $|B|$ -Lipschitz. By the multiplicative Azuma inequality (Lemma 6.6), we have $\Pr[Y \leq \frac{1}{2} \mathbb{E}[Y]] \leq e^{-\Omega(\lambda)}$. Since $Y = |N_{H_Q}(B)|$, we conclude the result. \square

Proof of Lemma 26.3. Similar to our proofs involving spread hypergraphs, we will iteratively generate random sets Q_i a total of $\ell = 100\alpha^{-1} \log \log(n)$ times and take $Q = \bigcup Q_i$, iteratively arguing that each Q_i is likely to have good properties.

Let q_c be the unique solution to $p_c = 1 - (1 - q_c)^\ell$; the main take away being that $q_c = \Omega(p_c/\ell)$. For $1 \leq i \leq \ell$, let Q_i be obtained by including each color of R independently and with probability q_c (and independent of any other Q_j set), noting that $\bigcup_{i=1}^\ell Q_i$ has the same distribution as Q .

Let B_i be the set of vertices x that can be reached from v by some $(Q_1 \cup \dots \cup Q_i)$ -rainbow path P_x of length at most i . Let $\phi_i : V(G) \rightarrow 2^{V(G) \cup R}$ be the function which maps to the vertices and colors of P_x if $x \in B_i$, and otherwise $\phi_i(x) = \emptyset$. Note that $|\phi_i(x)| \leq 2i \leq 2\ell$. We wish to show that $|B_i|$ is rapidly increasing with high probability.

First note that B_1 is just the set of neighbors of v with $c(vx) \in Q_1$. By Proposition 26.1 we have $d(v) \geq \frac{1}{2}d$, so $\mathbb{E}[|B_1|] \geq \frac{1}{2}dq_c$. Thus by the Chernoff bound we will have $|B_1| \geq \frac{1}{4}dq_c > 2\lambda^2q_c^{-1}$ with high probability, so from now on we assume this is the case..

Note that $N_{Q_i, \phi_i}(B_i \cap U_i) \subseteq B_{i+1}$. Since $d \geq \lambda^2(\alpha \cdot q_c)^{-1}n^\alpha$ by hypothesis, and since $|B_i| \geq |B_1| > 2\lambda^2q_c^{-1}$, as long as $|B_i| < n/3$ we can apply Lemma 26.4 to get

$$N_{Q_i, \phi_i}(B_i) \geq \frac{1}{4}|B_i| \min \left\{ \frac{d \cdot q_c \cdot \alpha}{64\lambda^{1/2}}, \left(\frac{n}{2|B_i|} \right)^{\alpha/(1+\alpha)} - 1 \right\}$$

with probability at least $1 - e^{-\Omega(\lambda^{1/2})}$. Note that the leftside of the minimum is always at least n^α , so the minimum is always achieved by the righthand side. Using this and $\alpha \leq 1$ gives

$$|N_{Q_i, \phi_i}(B_i)| \geq \frac{1}{4}|B_i|((n/2|B_i|)^{\alpha/2} - 1)$$

with probability at least $1 - e^{-\Omega(\lambda^{1/2})}$. Thus with probability at least $1 - \ell e^{-\Omega(\lambda^{1/2})} = 1 - O(\alpha^{-1}e^{-\Omega(\lambda^{1/2})})$ this holds for all i (note that the $\log \log(n)$ gets absorbed by $e^{-\Omega(\lambda^{1/2})}$ since $\lambda \geq (\log \log n)^2$). We claim that this implies $|B_{\ell-1}| \geq n/3$. And indeed, using that $|B_{i+1}| \geq |B_i| + |N_{Q_i, \phi_i}(B_i)|$ (since B_i and $N(B_i) \supseteq N_{Q_i, \phi_i}(B_i)$ by definition), one can prove by induction that $|B_i| \geq (n/2)^{1-(1-\alpha/16)^i}$ provided $|B_{i-1}| \leq n/3$. This gives the result. \square

Tomon [108] proved several other nice results using a result which extends Lemma 26.3 in two ways. The first way is by enforcing short paths from v provided we don't require v to reach as many vertices (and it is easy to adapt our current proof to achieve this end). The other extension is that it allows one to sample a random set of vertices $U \subseteq V(G)$ in addition to a random set of colors, and which guarantees short paths from v to a large set of vertices. To state such a result, we say that a path is a (U, Q) -rainbow path if it is a rainbow path whose internal vertices all lie in U and whose colors all lie in U .

Lemma 26.6. *There exists a sufficiently large constant C such that the following holds. Let $p, p_c, \alpha \in (0, 1]$, n a positive integer, $\tau \in [1/\log_3 n, \frac{1}{2})$, and $\lambda > C(\log \log n)^{10}$. Let G be an n -vertex α -maximal graph with proper edge coloring $c : E(G) \rightarrow R$ with average degree $d = d(G)$ satisfying either $d > C\lambda^2(\alpha^2 \cdot p_c^2)^{-1}n^\alpha$ if $p = 1$, and otherwise $d > C\lambda^2(\alpha^3 \cdot p \cdot p_c^2)^{-1}n^\alpha$.*

Let $U \subseteq V(G)$ be obtained by including each vertex independently and with probability p , and similarly define $Q \subseteq R$. For each $v \in V(G)$, with probability at least $1 - O(\alpha^{-1}e^{-\Omega(\lambda^{1/2})})$, at least $n^{1-\tau}$ vertices of G can be reached from v by a (U, Q) -path of length at most $O(\alpha^{-1} \log(1/\tau))$.

To prove this, one needs to extend Lemma 26.4 to say that the same conclusion holds for $N_{Q, \phi}(U)$ with $U \subseteq B$ obtained by including each vertex independently and with probability p . This isn't too hard to prove if the vertices of B all have reasonable maximum degree, and one extra case deals with the situation where this doesn't happen.

With Lemma 26.6 it is possible to prove results about rainbow Turán numbers of subdivisions of K_t . To this end, let \mathcal{K}_t denote the set of subdivisions of K_t (i.e. the graphs which can be obtained by subdividing each edge of K_t some number of times). Mader [81] showed $\text{ex}(n, \mathcal{K}_t) = O_t(n)$, and again the hypercube shows $\text{ex}^*(n, \mathcal{K}_t) = \Omega(n \log n)$ for $t \geq 3$. Jiang, Letzter, Methuku, and Yepremyan [67] showed $\text{ex}^*(n, \mathcal{K}_t) = O((\log n)^{60}n)$. These bounds were improved significantly by Tomon [108].

Theorem 26.7 ([108]). *For all fixed t we have $\text{ex}^*(n, \mathcal{K}_t) \leq (\log n)^{6+o(1)}n$.*

Proof Sketch. Let G be an n -vertex graph with $e(G) \geq 2(\log n)^{6+\epsilon}n$ and $c : E(G) \rightarrow R$ a proper coloring, and let $\alpha = 1/\log_2(G)$, $s = (\log n)^{1+\epsilon/10}$, $p = p_c = 1/s$, and $\lambda = (\log n)^{\epsilon/10}$. For some slight ease of notation we assume G is α -maximal (though it's easy for the rest of the proof to go through by reducing to an α -maximal subgraph). Note that by our choice of parameters, $d(G) \geq C\lambda(\alpha^3 p \cdot p_c^2)^{-1}n^\alpha$.

Define an auxiliary graph H where two vertices v, w are adjacent if there exist at least $s/6$ internally disjoint paths from v to w such that no color is used more than once in the union of the paths.

We claim that if H has minimum degree at least $n/6$, then G contains a rainbow K_t -subdivision. Indeed, by Theorem 5.4, H (easily) contains a 1-subdivision of K_t . One can then greedily replace each edge with a rainbow path which doesn't use any vertices or colors that have already been used.

To show that H has this minimum degree, we apply Lemma 26.6 with the stated parameters and $\tau = 1/\log_3(n)$ to any vertex v . By a similar argument to before, this implies there exist partitions U_1, \dots, U_s and Q_1, \dots, Q_s such that the sets B_i of vertices we can reach from v with a (U_i, Q_i) -rainbow path all have size at least $n/3$. This implies that there exist at least $n/6$ vertices w in at least s of the B_i sets, proving that $d_H(v) \geq n/6$. \square

Part VIII

Bibliography

References

- [1] N. Alon. Combinatorial nullstellensatz. *Combinatorics, Probability and Computing*, 8(1-2):7–29, 1999.
- [2] N. Alon, L. Babai, and H. Suzuki. Multilinear polynomials and frankl-ray-chaudhuri-wilson type intersection theorems. *Journal of Combinatorial Theory, Series A*, 58(2):165–180, 1991.
- [3] N. Alon and S. Friedland. The maximum number of perfect matchings in graphs with a given degree sequence. *arXiv preprint arXiv:0803.2578*, 2008.
- [4] N. Alon and Z. Füredi. Covering the cube by affine hyperplanes. *European journal of combinatorics*, 14(2):79–83, 1993.
- [5] N. Alon, M. Krivelevich, and B. Sudakov. Turán numbers of bipartite graphs and related ramsey-type questions. *Combinatorics, Probability and Computing*, 12(5-6):477–494, 2003.
- [6] N. Alon and J. H. Spencer. *The probabilistic method*. John Wiley & Sons, 2016.
- [7] R. Alweiss, S. Lovett, K. Wu, and J. Zhang. Improved bounds for the sunflower lemma. In *Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing*, pages 624–630, 2020.
- [8] L. Babai and P. Frankl. *Linear algebra methods in combinatorics*. University of Chicago, 1988.
- [9] J. Balogh. [Container Method notes](#).
- [10] J. Balogh, R. Morris, and W. Samotij. Independent sets in hypergraphs. *Journal of the American Mathematical Society*, 28(3):669–709, 2015.
- [11] J. Balogh, R. Mycroft, and A. Treglown. A random version of sperner’s theorem. *Journal of Combinatorial Theory, Series A*, 128:104–110, 2014.
- [12] J. Balogh and Š. Petříčková. The number of the maximal triangle-free graphs. *Bulletin of the London Mathematical society*, 46(5):1003–1006, 2014.
- [13] M. Bateman and N. Katz. New bounds on cap sets. *Journal of the American Mathematical Society*, 25(2):585–613, 2012.
- [14] T. Bell, S. Chueluecha, and L. Warnke. Note on sunflowers. *Discrete Mathematics*, 344(7):112367, 2021.

- [15] G. R. Blakley and P. Roy. A hölder type inequality for symmetric matrices with non-negative entries. *Proceedings of the American Mathematical Society*, 16(6):1244–1245, 1965.
- [16] B. Bollobás. *Modern graph theory*, volume 184. Springer Science & Business Media, 2013.
- [17] R. B. Boppana. A useful inequality for the binary entropy function. *arXiv preprint arXiv:2301.09664*, 2023.
- [18] L. M. Brègman. Some properties of nonnegative matrices and their permanents. In *Doklady Akademii Nauk*, volume 211, pages 27–30. Russian Academy of Sciences, 1973.
- [19] H. Bruhn and O. Schaudt. The journey of the union-closed sets conjecture. *Graphs and Combinatorics*, 31:2043–2074, 2015.
- [20] B. Bukh. Random algebraic construction of extremal graphs. *arXiv preprint arXiv:1409.3856*, 2014.
- [21] B. Bukh. Extremal graphs without exponentially-small bicliques. *arXiv preprint arXiv:2107.04167*, 2021.
- [22] B. Bukh and D. Conlon. Rational exponents in extremal graph theory. *Journal of the European Mathematical Society*, 20(7):1747–1757, 2018.
- [23] S. Cambie. Better bounds for the union-closed sets conjecture using the entropy approach. *arXiv preprint arXiv:2212.12500*, 2022.
- [24] Z. Chase and S. Lovett. Approximate union closed conjecture. *arXiv preprint arXiv:2211.11689*, 2022.
- [25] F. R. Chung, Z. Füredi, R. L. Graham, and P. Seymour. On induced subgraphs of the cube. *Journal of Combinatorial Theory, Series A*, 49(1):180–187, 1988.
- [26] D. Conlon. Graphs with few paths of prescribed length between any two vertices. *Bulletin of the London Mathematical Society*, 51(6):1015–1021, 2019.
- [27] D. Conlon and A. Ferber. Lower bounds for multicolor ramsey numbers. *Advances in Mathematics*, 378:107528, 2021.
- [28] D. Conlon, J. Fox, and B. Sudakov. An approximate version of sidorenko’s conjecture. *Geometric and Functional Analysis*, 20:1354–1366, 2010.
- [29] D. Conlon, C. Pohoata, and D. Zakharov. Random multilinear maps and the erdos box problem. *arXiv preprint arXiv:2011.09024*, 2020.
- [30] D. W. Cranston and D. B. West. An introduction to the discharging method via graph coloring. *Discrete Mathematics*, 340(4):766–793, 2017.
- [31] E. Croot, V. F. Lev, and P. P. Pach. Progression-free sets in are exponentially small. *Annals of Mathematics*, pages 331–337, 2017.

- [32] J. Cutler and A. Radcliffe. An entropy proof of the kahn-lovász theorem. *the electronic journal of combinatorics*, 18(1):P10, 2011.
- [33] S. Das, C. Lee, and B. Sudakov. Rainbow turán problem for even cycles. *European Journal of Combinatorics*, 34(5):905–915, 2013.
- [34] M. Deza, P. Erdős, and P. Frankl. Intersection properties of systems of finite sets. *Proceedings of the London Mathematical Society*, 3(2):369–384, 1978.
- [35] M. Deza, P. Frankl, and N. M. Singhi. On functions of strength t . *Combinatorica*, 3(3-4):331–339, 1983.
- [36] I. Dinur and E. Friedgut. Intersecting families are essentially contained in juntas. *Combinatorics, Probability and Computing*, 18(1-2):107–122, 2009.
- [37] D. P. Dubhashi and A. Panconesi. *Concentration of measure for the analysis of randomized algorithms*. Cambridge University Press, 2009.
- [38] Y. Edel. Extensions of generalized product caps. *Designs, Codes and Cryptography*, 31:5–14, 2004.
- [39] J. S. Ellenberg and D. Gijswijt. On large subsets of with no three-term arithmetic progression. *Annals of Mathematics*, pages 339–343, 2017.
- [40] P. Erdős. Some remarks on the theory of graphs. *Bulletin of the American Mathematical Society*, 53(4):292–294, 1947.
- [41] P. Erdos, D. J. Kleitman, and B. L. Rothschild. Asymptotic enumeration of kn -free graphs. *Colloquio Internazionale sulle Teorie Combinatorie (Rome, 1973)*, 2(17):19–27, 1976.
- [42] P. Erdős and E. Szemerédi. Combinatorial properties of systems of sets. *Journal of Combinatorial Theory, Series A*, 24(3):308–313, 1978.
- [43] R. J. Faudree and M. Simonovits. On a class of degenerate extremal graph problems. *Combinatorica*, 3:83–93, 1983.
- [44] J. Fox and B. Sudakov. Dependent random choice. *Random Structures & Algorithms*, 38(1-2):68–99, 2011.
- [45] P. Frankl. Extremal set systems. In *Handbook of combinatorics (vol. 2)*, pages 1293–1329. 1996.
- [46] P. Frankl and Z. Füredi. Forbidding just one intersection. *Journal of Combinatorial Theory, Series A*, 39(2):160–176, 1985.
- [47] P. Frankl and N. Tokushige. *Extremal problems for finite sets*, volume 86. American Mathematical Soc., 2018.
- [48] P. Frankl and R. M. Wilson. Intersection theorems with geometric consequences. *Combinatorica*, 1:357–368, 1981.

- [49] K. Frankston, J. Kahn, B. Narayanan, and J. Park. Thresholds versus fractional expectation-thresholds. *arXiv preprint arXiv:1910.13433*, 2019.
- [50] E. Friedgut. On the measure of intersecting families, uniqueness and stability. *Combinatorica*, 28(5):503–528, 2008.
- [51] A. Frieze and M. Karoński. *Introduction to random graphs*. Cambridge University Press, 2016.
- [52] Z. Füredi. On a turán type problem of erdős. *Combinatorica*, 11(1):75–79, 1991.
- [53] Z. Füredi and M. Simonovits. The history of degenerate (bipartite) extremal graph problems. In *Erdős Centennial*, pages 169–264. Springer, 2013.
- [54] D. Galvin. Three tutorial lectures on entropy and counting. *arXiv preprint arXiv:1406.7872*, 2014.
- [55] D. Galvin. Counting colorings of a regular graph. *Graphs and Combinatorics*, 31(3):629–638, 2015.
- [56] D. Galvin and P. Tetali. On weighted graph homomorphisms. *DIMACS Series in Discrete Mathematics and Theoretical Computer Science*, 63:97–104, 2004.
- [57] J. Gilmer. A constant lower bound for the union-closed sets conjecture. *arXiv preprint arXiv:2211.09055*, 2022.
- [58] S. Glock, D. Kühn, A. Lo, and D. Osthus. The existence of designs via iterative absorption: hypergraph f -designs for arbitrary f . *arXiv preprint arXiv:1611.06827*, 2016.
- [59] C. Gotsman and N. Linial. The equivalence of two problems on the cube. *Journal of Combinatorial Theory, Series A*, 61(1):142–146, 1992.
- [60] W. Gowers. Topics in combinatorics. *Notes taken by Paul Russell*, 2004.
- [61] J. Grochow. New applications of the polynomial method: the cap set conjecture and beyond. *Bulletin of the American Mathematical Society*, 56(1):29–64, 2019.
- [62] D. S. Gunderson, V. Rödl, and A. Sidorenko. Extremal problems for sets forming boolean algebras and complete partite hypergraphs. *Journal of Combinatorial Theory, Series A*, 88(2):342–367, 1999.
- [63] A. Heckel. Random triangles in random graphs. *Random Structures & Algorithms*, 2018.
- [64] A. Heckel and O. Riordan. How does the chromatic number of a random graph vary? *arXiv preprint arXiv:2103.14014*, 2021.
- [65] H. Huang. Induced subgraphs of hypercubes and a proof of the sensitivity conjecture. *Annals of Mathematics*, 190(3):949–955, 2019.
- [66] O. Janzer. Rainbow turán number of even cycles, repeated patterns and blow-ups of cycles. *arXiv preprint arXiv:2006.01062*, 2020.

- [67] T. Jiang, S. Letzter, A. Methuku, and L. Yepremyan. Rainbow clique subdivisions and blow-ups. *arXiv preprint arXiv:2108.08814*, 2021.
- [68] A. Johansson, J. Kahn, and V. Vu. Factors in random graphs. *Random Structures & Algorithms*, 33(1):1–28, 2008.
- [69] J. Kahn. An entropy approach to the hard-core model on bipartite graphs. *Combinatorics, Probability and Computing*, 10(3):219–237, 2001.
- [70] J. Kahn, B. Narayanan, and J. Park. The threshold for the square of a hamilton cycle. *arXiv preprint arXiv:2010.08592*, 2020.
- [71] P. Keevash. The existence of designs. *arXiv preprint arXiv:1401.3665*, 2014.
- [72] P. Keevash, D. Mubayi, B. Sudakov, and J. Verstraëte. Rainbow turán problems. *Combinatorics, Probability and Computing*, 16(1):109–126, 2007.
- [73] J. H. Kim. The ramsey number $r(3, t)$ has order of magnitude $t^2/\log t$. *Random Structures & Algorithms*, 7(3):173–207, 1995.
- [74] D. Kleitman. A conjecture of erdős-katona on commensurable pairs among subsets of an n -set. In *Theory of Graphs, Proc. Colloq. held at Tihany, Hungary*, pages 187–207, 1966.
- [75] D. Kleitman. On dedekind’s problem: the number of monotone boolean functions. *Proceedings of the American Mathematical Society*, 21(3):677–682, 1969.
- [76] E. Knill. Graph generated union-closed families of sets. *arXiv preprint math/9409215*, 229, 1994.
- [77] A. Kupavskii and D. Zakharov. Spread approximations for forbidden intersections problems. *arXiv preprint arXiv:2203.13379*, 2022.
- [78] W. Kuszmaul and Q. Qi. The multiplicative version of azuma’s inequality, with an application to contention analysis. *arXiv preprint arXiv:2102.05077*, 2021.
- [79] L. H. Loomis and H. Whitney. An inequality related to the isoperimetric inequality. 1949.
- [80] L. Lovász. *Combinatorial problems and exercises*, volume 361. American Mathematical Soc., 2007.
- [81] W. Mader. Homomorphieeigenschaften und mittlere kantendichte von graphen. *Mathematische Annalen*, 174(4):265–268, 1967.
- [82] T. C. Martinez. The slice rank polynomial method. 2021.
- [83] J. Matoušek. *Thirty-three miniatures: Mathematical and Algorithmic applications of Linear Algebra*. American Mathematical Society Providence, RI, 2010.
- [84] J. Matoušek. *Lectures on discrete geometry*, volume 212. Springer Science & Business Media, 2013.

- [85] C. McDiarmid. General first-passage percolation. *Advances in Applied Probability*, 15(1):149–161, 1983.
- [86] M. Michałek. A short proof of combinatorial nullstellensatz. *The American Mathematical Monthly*, 117(9):821–823, 2010.
- [87] J. W. Moon and L. Moser. On cliques in graphs. *Israel journal of Mathematics*, 3:23–28, 1965.
- [88] D. Mubayi and J. Verstraëte. A survey of turán problems for expansions. *Recent trends in combinatorics*, pages 117–143, 2016.
- [89] D. Mubayi and Y. Zhao. Forbidding complete hypergraphs as traces. *Graphs and Combinatorics*, 23(6):667–679, 2007.
- [90] E. Naslund and W. Sawin. Upper bounds for sunflower-free sets. In *Forum of Mathematics, Sigma*, volume 5, page e15. Cambridge University Press, 2017.
- [91] J. Radhakrishnan. An entropy proof of bregman’s theorem. *Journal of combinatorial theory, Series A*, 77(1):161–164, 1997.
- [92] A. Rao. Coding for sunflowers. *arXiv preprint arXiv:1909.04774*, 2019.
- [93] A. Rao. Sunflowers: from soil to oil. *Bulletin of the American Mathematical Society*, 60(1):29–38, 2023.
- [94] D. K. Ray-Chaudhuri and R. M. Wilson. On t-designs. 1975.
- [95] O. Riordan. Random cliques in random graphs. *arXiv preprint arXiv:1802.01948*, 2018.
- [96] V. Rödl. On a packing and covering problem. *European Journal of Combinatorics*, 6(1):69–78, 1985.
- [97] D. Romik. *The surprising mathematics of longest increasing subsequences*. Number 4. Cambridge University Press, 2015.
- [98] A. Sah, M. Sawhney, D. Stoner, and Y. Zhao. The number of independent sets in an irregular graph. *Journal of Combinatorial Theory, Series B*, 138:172–195, 2019.
- [99] W. Sawin. An improved lower bound for multicolor ramsey numbers and a problem of erdős. *Journal of Combinatorial Theory, Series A*, 188:105579, 2022.
- [100] J. B. Shearer. On a problem of spencer. *Combinatorica*, 5(3):241–245, 1985.
- [101] A. Sheffer. Polynomial methods and incidence theory.
- [102] M. Simonovits and E. Szemerédi. Embedding graphs into larger graphs: results, methods, and problems. In *Building Bridges II*, pages 445–592. Springer, 2019.
- [103] J. Spencer. Ramsey’s theorem—a new lower bound. *Journal of Combinatorial Theory, Series A*, 18(1):108–115, 1975.

- [104] J. Spencer. *Asymptopia*, volume 71. American Mathematical Soc., 2014.
- [105] S. Spiro. A smoother notion of spread hypergraphs. *arXiv preprint arXiv:2106.11882*, 2021.
- [106] T. Tao. A symmetric formulation of the croot-lev-pach-ellenberg-gijswijt capset bound. *blog post*, 2016.
- [107] T. Tao and V. H. Vu. *Additive combinatorics*, volume 105. Cambridge University Press, 2006.
- [108] I. Tomon. Robust (rainbow) subdivisions and simplicial cycles. *arXiv preprint arXiv:2201.12309*, 2022.
- [109] Y. Wigderson. An improved lower bound on multicolor ramsey numbers. *Proceedings of the American Mathematical Society*, 149(6):2371–2374, 2021.
- [110] Y. Zhao. The number of independent sets in a regular graph. *Combinatorics, Probability and Computing*, 19(2):315–320, 2010.
- [111] Y. Zhao. The bipartite swapping trick on graph homomorphisms. *SIAM Journal on Discrete Mathematics*, 25(2):660–680, 2011.
- [112] Y. Zhao. *Graph theory and additive combinatorics: exploring structure and randomness*. Cambridge University Press, 2023.

Part IX

Appendix: Spectral Graph Theory

TODO