

Polynomial Relations Between Matrices of Graphs

Sam Spiro, Rutgers University

Joint with Bryce Frederickson, Paul Horn, and Sabrina Lato



Spectral Graph Theory

Spectral Graph Theory

G

Spectral Graph Theory

$$G \rightarrow M_G$$

Spectral Graph Theory

$$G \rightarrow M_G \rightarrow \{\lambda_1, \dots, \lambda_n\}$$

Spectral Graph Theory

$$G \rightarrow M_G \rightarrow \{\lambda_1, \dots, \lambda_n\} \rightarrow \text{Properties of } G$$

Spectral Graph Theory

Define the adjacency matrix A by

$$A_{ij} = \begin{cases} 1 & ij \in E(G) \\ 0 & ij \notin E(G) \end{cases}$$

Spectral Graph Theory

Define the adjacency matrix A by

$$A_{ij} = \begin{cases} 1 & ij \in E(G) \\ 0 & ij \notin E(G) \end{cases}$$

Theorem

If $\{\lambda_1, \dots, \lambda_n\}$ are the eigenvalues of A , then the total number of closed walks (walks from a vertex to itself) of length k in G is

$$\lambda_1^k + \dots + \lambda_n^k.$$

Spectral Graph Theory

Let $D = \text{diag}(d_1, \dots, d_n)$ be the diagonal matrix of degrees of G .

Spectral Graph Theory

Let $D = \text{diag}(d_1, \dots, d_n)$ be the diagonal matrix of degrees of G . We define the Laplacian matrix L by

$$L = D - A.$$

Spectral Graph Theory

Let $D = \text{diag}(d_1, \dots, d_n)$ be the diagonal matrix of degrees of G . We define the Laplacian matrix L by

$$L = D - A.$$

Example

If G is the path on 3 vertices, then

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, \quad D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad L = \begin{bmatrix} 1 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 1 \end{bmatrix}.$$

Polynomial Relations

If G is a d -regular graph, then we have the nice relation

$$A = dI - L,$$

and from this one can easily translate between eigenvalues of A and L .

Polynomial Relations

If G is a d -regular graph, then we have the nice relation

$$A = dI - L,$$

and from this one can easily translate between eigenvalues of A and L .

If G is biregular (i.e. bipartite where every vertex on the same side has the same degree d_i), then

$$A^2 = (d_1I - L)(d_2I - L)$$

Polynomial Relations

If G is a d -regular graph, then we have the nice relation

$$A = dI - L,$$

and from this one can easily translate between eigenvalues of A and L .

If G is biregular (i.e. bipartite where every vertex on the same side has the same degree d_i), then

$$A^2 = (d_1I - L)(d_2I - L),$$

and using this one can also translate between the eigenvalues of A and L .

Polynomial Relations

Question

Do there exist “triangular graphs”, i.e. those with

$$A^3 = f(L)$$

for some polynomial f ?

Polynomial Relations

Question

Do there exist “triregular graphs”, i.e. those with

$$A^3 = f(L)$$

for some polynomial f ?

Theorem (S. 2018)

No.

Polynomial Relations

Theorem (S. 2018)

Let G be a connected graph. If there exists a positive integer r and polynomial f such that

$$A^r = f(L),$$

then G is either regular or biregular.

Polynomial Relations

Theorem (S. 2018)

Let G be a connected graph. If there exists a positive integer r and polynomial f such that

$$A^r = f(L),$$

then G is either regular or biregular.

Table: Graphs Satisfying $X^r = f(Y)$

X/Y	A	L	Q	\mathcal{L}
A		Reg, Bireg	Reg, Bireg	Reg, Bireg
L	Reg		Reg	Reg
Q	Reg	Reg		Reg
\mathcal{L}	Reg, Bireg	Reg	Reg	

Polynomial Relations

Question (Vague)

When do there exist “nice” polynomials f, g such that $f(A) = g(L)$?

Polynomial Relations

Question (Vague)

When do there exist “nice” polynomials f, g such that $f(A) = g(L)$?

This trivially holds if $f = g = 0$.

Polynomial Relations

Question (Vague)

When do there exist “nice” polynomials f, g such that $f(A) = g(L)$?

This trivially holds if $f = g = 0$. To avoid issues like this, we'll say that a relationship is *proper* if $f(A) \neq cI$ for some $c \in \mathbb{R}$.

Polynomial Relations

Question (Vague)

When do there exist “nice” polynomials f, g such that $f(A) = g(L)$?

This trivially holds if $f = g = 0$. To avoid issues like this, we'll say that a relationship is *proper* if $f(A) \neq cI$ for some $c \in \mathbb{R}$.

For this project, we decided to focus on (proper) relations $f(A) = g(L)$ when at least one of f, g has low degrees.

Low Degree Relations

Low Degree Relations

It's easy to show that having $f(A) = g(L)$ with $\deg f = 1$ or $\deg g = 1$ implies G is regular.

Low Degree Relations


It's easy to show that having $f(A) = g(L)$ with $\deg f = 1$ or $\deg g = 1$ implies G is regular.

Theorem (FHLS 2023+)

If G is connected and $f(A) = g(L)$ is proper with $\deg f, \deg g \leq 2$, then G is either regular or biregular.

Low Degree Relations

We define the *join* $G \vee H$ by taking $G \cup H$ and adding all edges between G and H



Low Degree Relations

Theorem (FHLS 2023+)

Let G be a k -regular m -vertex graph and H a d -regular n -vertex graph such that $G \vee H$ is not regular. Let A_G, A_H be the adjacency matrices of G, H and let A, L be the adjacency matrix and Laplacian matrix of $G \vee H$.

There exist polynomials f, g such that $f(A) = g(L)$ with $\deg g \leq 2$ if and only if there exists no $\mu \neq k, d$ which is an eigenvalue of both A_G and A_H and no eigenvalue of A_G or A_H is equal to $\frac{k+d-\sqrt{(k-d)^2+4mn}}{2}$.

Low Degree Relations

Theorem (FHLS 2023+)

Let G be a k -regular m -vertex graph and H a d -regular n -vertex graph such that $G \vee H$ is not regular. Let A_G, A_H be the adjacency matrices of G, H and let A, L be the adjacency matrix and Laplacian matrix of $G \vee H$.

There exist polynomials f, g such that $f(A) = g(L)$ with $\deg g \leq 2$ if and only if there exists no $\mu \neq k, d$ which is an eigenvalue of both A_G and A_H and no eigenvalue of A_G or A_H is equal to $\frac{k+d-\sqrt{(k-d)^2+4mn}}{2}$.

Informally, this says the join of two regular graphs G, H has $f(A) = g(L)$ with $\deg(g) \leq 2$ if and only if G, H share no eigenvalues and neither of their eigenvalues equal $\frac{k+d-\sqrt{(k-d)^2+4mn}}{2}$.

Proofs

Theorem (FHLS 2023+)

If G is connected and $f(A) = g(L)$ is proper with $\deg f, \deg g \leq 2$, then G is either regular or biregular.

Proofs

Theorem (FHLS 2023+)

If G is connected and $f(A) = g(L)$ is proper with $\deg f, \deg g \leq 2$, then G is either regular or biregular.

$$f(x) = ax^2 + bx + c, \quad g(x) = \alpha x^2 + \beta x + \gamma.$$

Proofs

Theorem (FHLS 2023+)

If G is connected and $f(A) = g(L)$ is proper with $\deg f, \deg g \leq 2$, then G is either regular or biregular.

$$f(x) = ax^2 + bx + c, \quad g(x) = \alpha x^2 + \beta x + \gamma.$$

If $a = 0$ or $\alpha = 0$ then it is easy to show G is regular.

Proofs

Theorem (FHLS 2023+)

If G is connected and $f(A) = g(L)$ is proper with $\deg f, \deg g \leq 2$, then G is either regular or biregular.

$$f(x) = ax^2 + bx + c, \quad g(x) = \alpha x^2 + \beta x + \gamma.$$

If $a = 0$ or $\alpha = 0$ then it is easy to show G is regular. Assuming $G \neq K_n$, there exist vertices u, v at distance 2 in G .

Proofs

Theorem (FHLS 2023+)

If G is connected and $f(A) = g(L)$ is proper with $\deg f, \deg g \leq 2$, then G is either regular or biregular.

$$f(x) = ax^2 + bx + c, \quad g(x) = \alpha x^2 + \beta x + \gamma.$$

If $a = 0$ or $\alpha = 0$ then it is easy to show G is regular. Assuming $G \neq K_n$, there exist vertices u, v at distance 2 in G .

$$f(A)_{u,v} = a \cdot A_{u,v}^2 + b \cdot A_{u,v} + c \cdot I_{u,v}$$

Proofs

Theorem (FHLS 2023+)

If G is connected and $f(A) = g(L)$ is proper with $\deg f, \deg g \leq 2$, then G is either regular or biregular.

$$f(x) = ax^2 + bx + c, \quad g(x) = \alpha x^2 + \beta x + \gamma.$$

If $a = 0$ or $\alpha = 0$ then it is easy to show G is regular. Assuming $G \neq K_n$, there exist vertices u, v at distance 2 in G .

$$f(A)_{u,v} = a \cdot A_{u,v}^2 + b \cdot A_{u,v} + c \cdot I_{u,v} = a \cdot d(u, v) \neq 0.$$

Proofs

Theorem (FHLS 2023+)

If G is connected and $f(A) = g(L)$ is proper with $\deg f, \deg g \leq 2$, then G is either regular or biregular.

$$f(x) = ax^2 + bx + c, \quad g(x) = \alpha x^2 + \beta x + \gamma.$$

If $a = 0$ or $\alpha = 0$ then it is easy to show G is regular. Assuming $G \neq K_n$, there exist vertices u, v at distance 2 in G .

$$f(A)_{u,v} = a \cdot A_{u,v}^2 + b \cdot A_{u,v} + c \cdot I_{u,v} = a \cdot d(u, v) \neq 0.$$

$$g(L)_{u,v} = \alpha d(u, v).$$

Proofs

Theorem (FHLS 2023+)

If G is connected and $f(A) = g(L)$ is proper with $\deg f, \deg g \leq 2$, then G is either regular or biregular.

$$f(x) = ax^2 + bx + c, \quad g(x) = \alpha x^2 + \beta x + \gamma.$$

If $a = 0$ or $\alpha = 0$ then it is easy to show G is regular. Assuming $G \neq K_n$, there exist vertices u, v at distance 2 in G .

$$f(A)_{u,v} = a \cdot A_{u,v}^2 + b \cdot A_{u,v} + c \cdot I_{u,v} = a \cdot d(u, v) \neq 0.$$

$$g(L)_{u,v} = \alpha d(u, v).$$

Having $f(A)_{u,v} = g(L)_{u,v}$ means

$$ad(u, v) = \alpha d(u, v)$$

Proofs

Theorem (FHLS 2023+)

If G is connected and $f(A) = g(L)$ is proper with $\deg f, \deg g \leq 2$, then G is either regular or biregular.

$$f(x) = ax^2 + bx + c, \quad g(x) = \alpha x^2 + \beta x + \gamma.$$

If $a = 0$ or $\alpha = 0$ then it is easy to show G is regular. Assuming $G \neq K_n$, there exist vertices u, v at distance 2 in G .

$$f(A)_{u,v} = a \cdot A_{u,v}^2 + b \cdot A_{u,v} + c \cdot I_{u,v} = a \cdot d(u, v) \neq 0.$$

$$g(L)_{u,v} = \alpha d(u, v).$$

Having $f(A)_{u,v} = g(L)_{u,v}$ means

$$ad(u, v) = \alpha d(u, v) \implies a = \alpha.$$

Proofs

$$f(x) = x^2 + bx + c, \quad g(x) = x^2 + \beta x + \gamma.$$

Proofs

$$f(x) = x^2 + bx + c, \quad g(x) = x^2 + \beta x + \gamma.$$

Let u, v be two adjacent vertices of G .

Proofs

$$f(x) = x^2 + bx + c, \quad g(x) = x^2 + \beta x + \gamma.$$

Let u, v be two adjacent vertices of G .

$$f(A)_{u,v} = d(u, v) + b$$

Proofs

$$f(x) = x^2 + bx + c, \quad g(x) = x^2 + \beta x + \gamma.$$

Let u, v be two adjacent vertices of G .

$$f(A)_{u,v} = d(u, v) + b, \quad g(L)_{u,v} = d(u, v) - d(u) - d(v) - \beta$$

Proofs

$$f(x) = x^2 + bx + c, \quad g(x) = x^2 + \beta x + \gamma.$$

Let u, v be two adjacent vertices of G .

$$f(A)_{u,v} = d(u, v) + b, \quad g(L)_{u,v} = d(u, v) - d(u) - d(v) - \beta$$

$$\implies d(u) + d(v) = -b - \beta \quad \forall uv \in E(G).$$

Proofs

$$f(x) = x^2 + bx + c, \quad g(x) = x^2 + \beta x + \gamma.$$

Let u, v be two adjacent vertices of G .

$$f(A)_{u,v} = d(u, v) + b, \quad g(L)_{u,v} = d(u, v) - d(u) - d(v) - \beta$$

$$\implies d(u) + d(v) = -b - \beta \quad \forall uv \in E(G).$$

We thus have $d(u) + d(v)$ equal to a common value for all $uv \in E(G)$.

Proofs

$$f(x) = x^2 + bx + c, \quad g(x) = x^2 + \beta x + \gamma.$$

Let u, v be two adjacent vertices of G .

$$f(A)_{u,v} = d(u, v) + b, \quad g(L)_{u,v} = d(u, v) - d(u) - d(v) - \beta$$

$$\implies d(u) + d(v) = -b - \beta \quad \forall uv \in E(G).$$

We thus have $d(u) + d(v)$ equal to a common value for all $uv \in E(G)$.
This common value must be $\delta + \Delta$

Proofs

$$f(x) = x^2 + bx + c, \quad g(x) = x^2 + \beta x + \gamma.$$

Let u, v be two adjacent vertices of G .

$$f(A)_{u,v} = d(u, v) + b, \quad g(L)_{u,v} = d(u, v) - d(u) - d(v) - \beta$$

$$\implies d(u) + d(v) = -b - \beta \quad \forall uv \in E(G).$$

We thus have $d(u) + d(v)$ equal to a common value for all $uv \in E(G)$.

This common value must be $\delta + \Delta$, which means every vertex of minimum degree is only adjacent to vertices of maximum degree and vice versa.

Proofs

$$f(x) = x^2 + bx + c, \quad g(x) = x^2 + \beta x + \gamma.$$

Let u, v be two adjacent vertices of G .

$$f(A)_{u,v} = d(u, v) + b, \quad g(L)_{u,v} = d(u, v) - d(u) - d(v) - \beta$$

$$\implies d(u) + d(v) = -b - \beta \quad \forall uv \in E(G).$$

We thus have $d(u) + d(v)$ equal to a common value for all $uv \in E(G)$.

This common value must be $\delta + \Delta$, which means every vertex of minimum degree is only adjacent to vertices of maximum degree and vice versa.

This means G is either regular or biregular. □

Proofs

Proposition

For $G \vee H$, under reasonable conditions there exist a proper relation $f(A) = g(L)$ with $\deg f = 2$, $\deg g = 3$.

Proofs

Proposition

For $G \vee H$, under reasonable conditions there exist a proper relation $f(A) = g(L)$ with $\deg f = 2$, $\deg g = 3$.

Under “reasonable conditions”, the vector space spanned by

$$\{A^i\}_{i=0}^{\infty} \cup \{L^i\}_{i=0}^{\infty}$$

has dimension at most 5.

Proofs

Proposition

For $G \vee H$, under reasonable conditions there exist a proper relation $f(A) = g(L)$ with $\deg f = 2$, $\deg g = 3$.

Under “reasonable conditions”, the vector space spanned by

$$\{A^i\}_{i=0}^{\infty} \cup \{L^i\}_{i=0}^{\infty}$$

has dimension at most 5. Thus there exists a non-trivial linear combination of

$$\{I, A, A^2, L, L^2, L^3\}$$

equal to 0.

Proofs

Proposition

For $G \vee H$, under reasonable conditions there exist a proper relation $f(A) = g(L)$ with $\deg f = 2$, $\deg g = 3$.

Under “reasonable conditions”, the vector space spanned by

$$\{A^i\}_{i=0}^{\infty} \cup \{L^i\}_{i=0}^{\infty}$$

has dimension at most 5. Thus there exists a non-trivial linear combination of

$$\{I, A, A^2, L, L^2, L^3\}$$

equal to 0. This is exactly a proper relation of the desired degrees. □

Open Problems

Open Problems

Conjecture

If $f(A) = g(L)$ is proper with $\deg f \leq 2$, $g \leq 3$, then G is either regular, biregular, or the join of two regular graphs.

Open Problems

Conjecture

If $f(A) = g(L)$ is proper with $\deg f \leq 2$, $g \leq 3$, then G is either regular, biregular, or the join of two regular graphs.

Question

If $f(A) = g(L)$ proper with $\deg f = 2$, does G have at most 2 degrees?

Open Problems

Conjecture

If $f(A) = g(L)$ is proper with $\deg f \leq 2$, $g \leq 3$, then G is either regular, biregular, or the join of two regular graphs.

Question

If $f(A) = g(L)$ proper with $\deg f = 2$, does G have at most 2 degrees?
More generally, does G have at most $\deg f$ degrees?

Open Problems

Problem

Can you come up with a guess as to which graphs have $f(A) = g(L)$ with $\deg f = \deg g = 3$? What about $\{\deg f, \deg g\} = \{2, 4\}$?

Open Problems

Problem

Can you come up with a guess as to which graphs have $f(A) = g(L)$ with $\deg f = \deg g = 3$? What about $\{\deg f, \deg g\} = \{2, 4\}$?

Conjecture

If $f(A) = g(\mathcal{L})$ with $\deg f, \deg g \leq 2$, then G is either regular or biregular.