

Generalized Turán Problems for Trees and More

Sam Spiro, Georgia State University

Based on Joint Work with Sean English



Turán Problems

The Turán number $\text{ex}(n, F)$ is defined to be the maximum number of edges that an F -free graph on n vertices can have.

Turán Problems

The Turán number $\text{ex}(n, F)$ is defined to be the maximum number of edges that an F -free graph on n vertices can have.

Theorem

$$\text{ex}(n, K_2) = 0.$$

Turán Problems

Theorem (Turán 1941)

$$\text{ex}(n, K_r) = \left\lfloor \binom{r-1}{2} \frac{n^2}{(r-1)^2} \right\rfloor$$

Turán Problems

Theorem (Turán 1941)

$$\text{ex}(n, K_r) = \left\lfloor \binom{r-1}{2} \frac{n^2}{(r-1)^2} \right\rfloor$$

Theorem (Kővari-Sós-Turán 1954)

$$\text{ex}(n, K_{s,t}) = O(n^{2-1/s}).$$

Turán Problems

Theorem (Turán 1941)

$$\text{ex}(n, K_r) = \left\lfloor \binom{r-1}{2} \frac{n^2}{(r-1)^2} \right\rfloor$$

Theorem (Kővari-Sós-Turán 1954)

$$\text{ex}(n, K_{s,t}) = O(n^{2-1/s}).$$

Theorem (Folklore)

If T is a tree with $e(T) \geq 2$, then

$$\text{ex}(n, T) = \Theta(n).$$

Turán Problems

Given a family of graphs \mathcal{F} , we say G is \mathcal{F} -free if G is F -free for all $F \in \mathcal{F}$. We define $\text{ex}(n, \mathcal{F})$ to be the maximum number of edges in an n -vertex \mathcal{F} -free graph.

Turán Problems

Given a family of graphs \mathcal{F} , we say G is \mathcal{F} -free if G is F -free for all $F \in \mathcal{F}$. We define $\text{ex}(n, \mathcal{F})$ to be the maximum number of edges in an n -vertex \mathcal{F} -free graph.

Conjecture (Compactness Conjecture)

If \mathcal{F} is a finite family of graphs which does not include both a star and a matching, then $\text{ex}(n, \mathcal{F}) = \Theta(\text{ex}(n, F))$ for some $F \in \mathcal{F}$.

Generalized Turán Problems

Generalized Turán Problems

Definition (Alon-Shikhelman 2016)

Given a graph H and a family of graphs \mathcal{F} , we define the generalized Turán number $\text{ex}(n, H, \mathcal{F})$ to be the maximum number of copies of H in an n -vertex \mathcal{F} -free graph.

Generalized Turán Problems

Definition (Alon-Shikhelman 2016)

Given a graph H and a family of graphs \mathcal{F} , we define the generalized Turán number $\text{ex}(n, H, \mathcal{F})$ to be the maximum number of copies of H in an n -vertex \mathcal{F} -free graph.

Proposition

$$\text{ex}(n, K_2, \mathcal{F}) = \text{ex}(n, \mathcal{F}).$$

Generalized Turán Problems

Definition (Alon-Shikhelman 2016)

Given a graph H and a family of graphs \mathcal{F} , we define the generalized Turán number $\text{ex}(n, H, \mathcal{F})$ to be the maximum number of copies of H in an n -vertex \mathcal{F} -free graph.

Proposition

$$\text{ex}(n, K_2, \mathcal{F}) = \text{ex}(n, \mathcal{F}).$$

See the recent survey by Gerbner and Palmer for way more history than I'm going to give here.

Generalized Turán Problems

Theorem (Gerbner-Palmer 2019)

For all $r \geq 2$ and families \mathcal{F} we have

$$\text{ex}(n, K_r, \mathcal{F}) \leq \text{ex}(n, \mathcal{F})^{r/2}.$$

Generalized Turán Problems

Theorem (Gerbner-Palmer 2019)

For all $r \geq 2$ and families \mathcal{F} we have

$$\text{ex}(n, K_r, \mathcal{F}) \leq \text{ex}(n, \mathcal{F})^{r/2}.$$

Theorem (Gerbner 2023)

Determines $\text{ex}(n, K_{2,t}, F)$ asymptotically for any non-bipartite graph F .

Generalized Turán Problems

Theorem (Gerbner-Palmer 2019)

For all $r \geq 2$ and families \mathcal{F} we have

$$\text{ex}(n, K_r, \mathcal{F}) \leq \text{ex}(n, \mathcal{F})^{r/2}.$$

Theorem (Gerbner 2023)

Determines $\text{ex}(n, K_{2,t}, F)$ asymptotically for any non-bipartite graph F .

Theorem (Füredi-Kündgen 2006)

If $\text{ex}(n, \mathcal{F}) = \Theta(n^{2-\beta})$ and \mathcal{F} does not contain a star, then

$$\text{ex}(n, K_{1,t}, \mathcal{F}) = \tilde{\Theta}(\max\{n^t, n^{t+1-t\beta}\}).$$

Generalized Turán Problems

Question

Can we prove bounds on $\text{ex}(n, T, \mathcal{F})$ for arbitrary trees T (for some possibly fixed \mathcal{F})?

Generalized Turán Problems

Question

Can we prove bounds on $\text{ex}(n, T, \mathcal{F})$ for arbitrary trees T (for some possibly fixed \mathcal{F})?

Theorem (Gerbner 2023)

Determines the order of magnitude of $\text{ex}(n, T, K_{2,t})$ for every tree T .

Generalized Turán Problems

Question

Can we prove bounds on $\text{ex}(n, T, \mathcal{F})$ for arbitrary trees T (for some possibly fixed \mathcal{F})?

Theorem (Gerbner 2023)

Determines the order of magnitude of $\text{ex}(n, T, K_{2,t})$ for every tree T .

Theorem (Alon-Shikhelman 2016)

If T is a tree and F is a tree, then $\text{ex}(n, T, F) = \Theta(n^k)$ for some integer k .

Generalized Turán Problems

Question

Can we prove bounds on $\text{ex}(n, T, \mathcal{F})$ for arbitrary trees T (for some possibly fixed \mathcal{F})?

Theorem (Gerbner 2023)

Determines the order of magnitude of $\text{ex}(n, T, K_{2,t})$ for every tree T .

Theorem (Alon-Shikhelman 2016)

If T is a tree and F is a tree, then $\text{ex}(n, T, F) = \Theta(n^k)$ for some integer k .

Theorem (Letzter 2019)

If H is any graph and F is a tree, then $\text{ex}(n, H, F) = \Theta(n^k)$ for some integer k .

A General Theorem for Trees

Theorem (English-S. 2025+)

For any tree T , integer $k \geq 1$, and family of graphs \mathcal{F} , either

$$\text{ex}(n, T, \mathcal{F}) = \Omega(n^k),$$

or

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

Moreover, we can determine which of these two cases happen for a given T, k, \mathcal{F} .

A General Theorem for Trees

Theorem (English-S. 2025+)

Either $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$ or $\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1})$.

A General Theorem for Trees

Theorem (English-S. 2025+)

Either $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$ or $\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1})$.

Corollary

If T is a tree and \mathcal{F} contains a forest, then $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$ for some integer k .

A General Theorem for Trees

Theorem (English-S. 2025+)

Either $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$ or $\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1})$.

Corollary

If T is a tree and \mathcal{F} contains a forest, then $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$ for some integer k .

Proof.

Let k be the smallest integer such that $\text{ex}(n, T, \mathcal{F}) = O(n^k)$.

A General Theorem for Trees

Theorem (English-S. 2025+)

Either $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$ or $\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1})$.

Corollary

If T is a tree and \mathcal{F} contains a forest, then $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$ for some integer k .

Proof.

Let k be the smallest integer such that $\text{ex}(n, T, \mathcal{F}) = O(n^k)$. If $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$ then we're done

A General Theorem for Trees

Theorem (English-S. 2025+)

Either $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$ or $\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1})$.

Corollary

If T is a tree and \mathcal{F} contains a forest, then $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$ for some integer k .

Proof.

Let k be the smallest integer such that $\text{ex}(n, T, \mathcal{F}) = O(n^k)$. If $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$ then we're done, otherwise our theorem implies that

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1})$$

A General Theorem for Trees

Theorem (English-S. 2025+)

Either $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$ or $\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1})$.

Corollary

If T is a tree and \mathcal{F} contains a forest, then $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$ for some integer k .

Proof.

Let k be the smallest integer such that $\text{ex}(n, T, \mathcal{F}) = O(n^k)$. If $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$ then we're done, otherwise our theorem implies that

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}) = O(n^{k-1}).$$



A General Theorem for Trees

Corollary

If $T \neq K_2$ is a tree with $\ell \geq 2$ leaves, then any family \mathcal{F} with $\text{ex}(n, T, \mathcal{F}) = O(n^\ell)$ has $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$ for some integer k .

In particular, if $\text{ex}(n, T, \mathcal{F}) = o(n^\ell)$ then $\text{ex}(n, T, \mathcal{F}) = O(n^{\ell-1})$.

A General Theorem for Trees

Corollary

If $T \neq K_2$ is a tree with $\ell \geq 2$ leaves, then any family \mathcal{F} with $\text{ex}(n, T, \mathcal{F}) = O(n^\ell)$ has $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$ for some integer k .

In particular, if $\text{ex}(n, T, \mathcal{F}) = o(n^\ell)$ then $\text{ex}(n, T, \mathcal{F}) = O(n^{\ell-1})$.

Proof.

If \mathcal{F} contains a forest then we are done by the previous result.

A General Theorem for Trees

Corollary

If $T \neq K_2$ is a tree with $\ell \geq 2$ leaves, then any family \mathcal{F} with $\text{ex}(n, T, \mathcal{F}) = O(n^\ell)$ has $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$ for some integer k .

In particular, if $\text{ex}(n, T, \mathcal{F}) = o(n^\ell)$ then $\text{ex}(n, T, \mathcal{F}) = O(n^{\ell-1})$.

Proof.

If \mathcal{F} contains a forest then we are done by the previous result. Otherwise, the graph G obtained by duplicating each leaf of T a total of n/ℓ times is \mathcal{F} -free

A General Theorem for Trees

Corollary

If $T \neq K_2$ is a tree with $\ell \geq 2$ leaves, then any family \mathcal{F} with $\text{ex}(n, T, \mathcal{F}) = O(n^\ell)$ has $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$ for some integer k .

In particular, if $\text{ex}(n, T, \mathcal{F}) = o(n^\ell)$ then $\text{ex}(n, T, \mathcal{F}) = O(n^{\ell-1})$.

Proof.

If \mathcal{F} contains a forest then we are done by the previous result. Otherwise, the graph G obtained by duplicating each leaf of T a total of n/ℓ times is \mathcal{F} -free and contains $\Omega(n^\ell)$ copies of T . □

A General Theorem for Trees

Theorem (English-S. 2025++)

If T is a tree with $\ell \geq 2$ leaves, then every family \mathcal{F} with $\text{ex}(n, T, \mathcal{F}) = o(n^{\ell+1})$ satisfies

$$\text{ex}(n, T, \mathcal{F}) = O\left(n^{\ell + \frac{\ell^2 - \ell}{e(T) - 1}}\right).$$

A General Theorem for Trees

Theorem (English-S. 2025++)

If T is a tree with $\ell \geq 2$ leaves, then every family \mathcal{F} with $\text{ex}(n, T, \mathcal{F}) = o(n^{\ell+1})$ satisfies

$$\text{ex}(n, T, \mathcal{F}) = O\left(n^{\ell + \frac{\ell^2 - \ell}{e(T) - 1}}\right).$$

Moreover, this bound is best possible for essentially all values of $\ell, e(T)$.

A General Theorem for Trees

Theorem (English-S. 2025++)

If T is a tree with $\ell \geq 2$ leaves, then every family \mathcal{F} with $\text{ex}(n, T, \mathcal{F}) = o(n^{\ell+1})$ satisfies

$$\text{ex}(n, T, \mathcal{F}) = O\left(n^{\ell + \frac{\ell^2 - \ell}{e(T) - 1}}\right).$$

Moreover, this bound is best possible for essentially all values of $\ell, e(T)$.

Theorem (English-S. 20??++)

For the path graph P_t , every graph F with $\text{ex}(n, P_t, F) = O(n^{\alpha(P_t)})$ has $\text{ex}(n, P_t, F) = \Theta(n^k)$ for some integer k .

Stability for Generalized Turán Problems

Stability for Generalized Turán Problems

Proposition (English-Halfpap-Krueger 2024)

For stars $K_{1,t}$, every family of graphs \mathcal{F} either satisfies $\text{ex}(n, K_{1,t}, \mathcal{F}) = \Omega(n^t)$ or $\text{ex}(n, K_{1,t}, \mathcal{F}) = O(n)$.

Stability for Generalized Turán Problems

Proposition (English-Halfpap-Krueger 2024)

For stars $K_{1,t}$, every family of graphs \mathcal{F} either satisfies $\text{ex}(n, K_{1,t}, \mathcal{F}) = \Omega(n^t)$ or $\text{ex}(n, K_{1,t}, \mathcal{F}) = O(n)$.

Proof.

If \mathcal{F} does not contain a subgraph of a star, then $G = K_{1,n-1}$ is \mathcal{F} -free and shows $\text{ex}(n, K_{1,t}, \mathcal{F}) = \Omega(n^t)$.

Stability for Generalized Turán Problems

Proposition (English-Halfpap-Krueger 2024)

For stars $K_{1,t}$, every family of graphs \mathcal{F} either satisfies $\text{ex}(n, K_{1,t}, \mathcal{F}) = \Omega(n^t)$ or $\text{ex}(n, K_{1,t}, \mathcal{F}) = O(n)$.

Proof.

If \mathcal{F} does not contain a subgraph of a star, then $G = K_{1,n-1}$ is \mathcal{F} -free and shows $\text{ex}(n, K_{1,t}, \mathcal{F}) = \Omega(n^t)$. Otherwise, \mathcal{F} must contain some subgraph of some star $K_{1,r}$, which means $\text{ex}(n, K_{1,t}, \mathcal{F}) = O_r(n)$. \square

Stability for Generalized Turán Problems

Proposition (English-Halfpap-Krueger 2024)

For stars $K_{1,t}$, every family of graphs \mathcal{F} either satisfies $\text{ex}(n, K_{1,t}, \mathcal{F}) = \Omega(n^t)$ or $\text{ex}(n, K_{1,t}, \mathcal{F}) = O(n)$.

Proof.

If \mathcal{F} does not contain a subgraph of a star, then $G = K_{1,n-1}$ is \mathcal{F} -free and shows $\text{ex}(n, K_{1,t}, \mathcal{F}) = \Omega(n^t)$. Otherwise, \mathcal{F} must contain some subgraph of some star $K_{1,r}$, which means $\text{ex}(n, K_{1,t}, \mathcal{F}) = O_r(n)$. \square

We want to generalize this idea by saying that for *any* graph H , there exists some “simple” family \mathcal{F}_H such that the behavior of $\text{ex}(n, H, \mathcal{F})$ depends on how \mathcal{F} “interacts” with \mathcal{F}_H .

Stability for Generalized Turán Problems


Definition

Given a graph H , a subset $R \subseteq V(H)$, and an integer q , we define the sunflower-power H_R^q to be the graph obtained by taking q copies of H which all agree on R and which are otherwise disjoint.

Stability for Generalized Turán Problems

Definition


Given a graph H , a subset $R \subseteq V(H)$, and an integer q , we define the sunflower-power H_R^q to be the graph obtained by taking q copies of H which all agree on R and which are otherwise disjoint.




Stability for Generalized Turán Problems

Definition

Given a graph H , a subset $R \subseteq V(H)$, and an integer q , we define the sunflower-power H_R^q to be the graph obtained by taking q copies of H which all agree on R and which are otherwise disjoint.




$$(P_5)_{\{x_3, x_4\}}^3$$




Stability for Generalized Turán Problems

Definition

Given a graph H , a subset $R \subseteq V(H)$, and an integer q , we define the sunflower-power H_R^q to be the graph obtained by taking q copies of H which all agree on R and which are otherwise disjoint.



$(P_5)_{\{x_3, x_4\}}^3$




We define

$$\mathcal{F}_{H,k}^q = \{H_R^q : H - R \text{ has at least } k \text{ connected components}\}.$$


Stability for Generalized Turán Problems

Definition

Given a graph H , a subset $R \subseteq V(H)$, and an integer q , we define the sunflower-power H_R^q to be the graph obtained by taking q copies of H which all agree on R and which are otherwise disjoint.



$(P_5)_{\{x_3, x_4\}}^3$



We define

$$\mathcal{F}_{H,k}^q = \{H_R^q : H - R \text{ has at least } k \text{ connected components}\}.$$

Claim

Every graph in $\mathcal{F}_{H,k}^q$ has at least q^k copies of H .

Stability for Generalized Turán Problems

Proposition (Key Observation)

For every H, k, \mathcal{F} , either $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$ or there exists some q such that every \mathcal{F} -free graph is $\mathcal{F}_{H,k}^q$ -free.

Stability for Generalized Turán Problems

Proposition (Key Observation)

For every H, k, \mathcal{F} , either $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$ or there exists some q such that every \mathcal{F} -free graph is $\mathcal{F}_{H,k}^q$ -free.

Proof.

If some $H_R^q \in \mathcal{F}_{H,k}^q$ is \mathcal{F} -free for all q , then the previous claim with $q \approx n/v(H)$ shows $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$.

Stability for Generalized Turán Problems

Proposition (Key Observation)

For every H, k, \mathcal{F} , either $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$ or there exists some q such that every \mathcal{F} -free graph is $\mathcal{F}_{H,k}^q$ -free.

Proof.

If some $H_R^q \in \mathcal{F}_{H,k}^q$ is \mathcal{F} -free for all q , then the previous claim with $q \approx n/v(H)$ shows $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$. Otherwise there exists some q so that every $H_R^q \in \mathcal{F}_{H,k}^q$ contains an element of \mathcal{F} as a subgraph. \square

Stability for Generalized Turán Problems

Proposition (Key Observation)

For every H, k, \mathcal{F} , either $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$ or there exists some q such that every \mathcal{F} -free graph is $\mathcal{F}_{H,k}^q$ -free.

Proof.

If some $H_R^q \in \mathcal{F}_{H,k}^q$ is \mathcal{F} -free for all q , then the previous claim with $q \approx n/v(H)$ shows $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$. Otherwise there exists some q so that every $H_R^q \in \mathcal{F}_{H,k}^q$ contains an element of \mathcal{F} as a subgraph. \square

Corollary (General Stability for Generalized Turán)

If $\text{ex}(n, H, \mathcal{F}_{H,k}^q) = O_q(n^\beta)$, then every family \mathcal{F} either satisfies $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$ or $\text{ex}(n, H, \mathcal{F}) = O(n^\beta)$.

Stability for Generalized Turán Problems

Proposition

For every graph H and family \mathcal{F} , either $\text{ex}(n, H, \mathcal{F}) = \Omega(n)$ or $\text{ex}(n, H, \mathcal{F}) = O(1)$.

Stability for Generalized Turán Problems

Proposition

For every graph H and family \mathcal{F} , either $\text{ex}(n, H, \mathcal{F}) = \Omega(n)$ or $\text{ex}(n, H, \mathcal{F}) = O(1)$.


By the previous corollary, it suffices to prove $\text{ex}(n, H, \mathcal{F}_{H,1}^q) = O_q(1)$.

Stability for Generalized Turán Problems

Proposition

For every graph H and family \mathcal{F} , either $\text{ex}(n, H, \mathcal{F}) = \Omega(n)$ or $\text{ex}(n, H, \mathcal{F}) = O(1)$.

By the previous corollary, it suffices to prove $\text{ex}(n, H, \mathcal{F}_{H,1}^q) = O_q(1)$.




Stability for Generalized Turán Problems

Proposition

For every graph H and family \mathcal{F} , either $\text{ex}(n, H, \mathcal{F}) = \Omega(n)$ or $\text{ex}(n, H, \mathcal{F}) = O(1)$.

By the previous corollary, it suffices to prove $\text{ex}(n, H, \mathcal{F}_{H,1}^q) = O_q(1)$.



This is essentially equivalent to the Erdős-Rado Sunflower lemma. □

Stability for Generalized Turán Problems

Theorem (English-S. 2025+)

For any tree T , integer $k \geq 1$, and family of graphs \mathcal{F} , either

$$\text{ex}(n, T, \mathcal{F}) = \Omega(n^k),$$

or

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

Stability for Generalized Turán Problems

Theorem (English-S. 2025+)

For any tree T , integer $k \geq 1$, and family of graphs \mathcal{F} , either

$$\text{ex}(n, T, \mathcal{F}) = \Omega(n^k),$$

or

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

Proposition (Key Observation)

For every H, k, \mathcal{F} , either $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$ or there exists some q such that every \mathcal{F} -free graph is $\mathcal{F}_{H,k}^q$ -free.

Stability for Generalized Turán Problems

Theorem (English-S. 2025+)

For any tree T , integer $k \geq 1$, and family of graphs \mathcal{F} , either

$$\text{ex}(n, T, \mathcal{F}) = \Omega(n^k),$$

or

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

Proposition (Key Observation)

For every H, k, \mathcal{F} , either $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$ or there exists some q such that every \mathcal{F} -free graph is $\mathcal{F}_{H,k}^q$ -free.

Theorem (English-S. 2025+)

If T is a tree and if G is $\mathcal{F}_{T,k}^q$ -free, then the number of copies of T in G is at most $O(e(G)^{k-1})$.

Proof Sketch

Proof Vibes

Proof Vibes

Theorem (English-S. 2025+)

If T is a tree and if G is $\mathcal{F}_{T,k}^q$ -free, then the number of copies of T in G is at most $O(e(G)^{k-1})$.

Proof Vibes

Theorem (English-S. 2025+)

If T is a tree and if G is $\mathcal{F}_{T,k}^q$ -free, then the number of copies of T in G is at most $O(e(G)^{k-1})$.


Strategy: identify each copy of T in G by a set of $k - 1$ edges E such that each E identifies at most $O(1)$ copies of T .

Proof Vibes

Theorem (English-S. 2025+)

If T is a tree and if G is $\mathcal{F}_{T,k}^q$ -free, then the number of copies of T in G is at most $O(e(G)^{k-1})$.

Strategy: identify each copy of T in G by a set of $k-1$ edges E such that each E identifies at most $O(1)$ copies of T .




Proof Vibes

Theorem (English-S. 2025+)

If T is a tree and if G is $\mathcal{F}_{T,k}^q$ -free, then the number of copies of T in G is at most $O(e(G)^{k-1})$.

Strategy: identify each copy of T in G by a set of $k - 1$ edges E such that each E identifies at most $O(1)$ copies of T .




For this example, to specify a given copy of P_5 , we **must** identify its last edge.

Proof Vibes

Theorem (English-S. 2025+)

If T is a tree and if G is $\mathcal{F}_{T,k}^q$ -free, then the number of copies of T in G is at most $O(e(G)^{k-1})$.

Strategy: identify each copy of T in G by a set of $k - 1$ edges E such that each E identifies at most $O(1)$ copies of T .



For this example, to specify a given copy of P_5 , we **must** identify its last edge. More generally, whatever set of edges E we choose to identify a given copy K in a graph, the edges of E **must** intersect every subtree $K' \subseteq K$ which has “many extensions.”

Proof Vibes

Question

Given a set of subtrees \mathcal{T} of a tree T , when can we guarantee that there exist a set of $k - 1$ edges E which intersect all of the edge sets of these subtrees?

Proof Vibes


Question

Given a set of subtrees \mathcal{T} of a tree T , when can we guarantee that there exist a set of $k - 1$ edges E which intersect all of the edge sets of these subtrees?


Theorem (Helly Property for Trees)

If \mathcal{T} is a set of subtrees of a tree T such that the vertex sets of the subtrees pairwise intersect, then there exists a vertex $v \in V(T)$ which intersects the vertex set of every subtree of \mathcal{T} .

Proof Vibes



Proof Vibes



Definition

We call a subtree $T' \subseteq T$ leaf-cuttable if $e(T) \geq 1$ and if every edge in $E(T) \setminus E(T')$ which intersects T' intersects a leaf of T' .

Proof Vibes

Theorem (English-S. 2025+)

If \mathcal{T} is a set of leaf-cuttable subtrees of a tree T such that the edge sets of the subtrees pairwise intersect, then there exists an edge $e \in E(T)$ which intersects the edge set of every subtree of \mathcal{T} .

Proof Vibes

Theorem (English-S. 2025+)

If \mathcal{T} is a set of leaf-cuttable subtrees of a tree T such that the edge sets of the subtrees pairwise intersect, then there exists an edge $e \in E(T)$ which intersects the edge set of every subtree of \mathcal{T} .

Theorem (English-S. 2025+)

If \mathcal{T} is a set of leaf-cuttable subtrees of a tree T such that every k subtrees from \mathcal{T} contains two subtrees with intersecting edge sets, then there exists a set $E \subseteq E(T)$ of at most $k - 1$ edges which intersects the edge set of every subtree of \mathcal{T} .

Proof Vibes

Theorem (English-S. 2025+)

If \mathcal{T} is a set of leaf-cuttable subtrees of a tree T such that the edge sets of the subtrees pairwise intersect, then there exists an edge $e \in E(T)$ which intersects the edge set of every subtree of \mathcal{T} .

Theorem (English-S. 2025+)

If \mathcal{T} is a set of leaf-cuttable subtrees of a tree T such that every k subtrees from \mathcal{T} contains two subtrees with intersecting edge sets, then there exists a set $E \subseteq E(T)$ of at most $k - 1$ edges which intersects the edge set of every subtree of \mathcal{T} .

This result implies the vertex-Helly result (and also König's Theorem for trees).

Proof Vibes

We are now at the “starting line” of the proof, the rest of which requires us to do a few more things:

Proof Vibes

We are now at the “starting line” of the proof, the rest of which requires us to do a few more things:

- Verify that the “highly extendable” subtrees \mathcal{T}_K are leaf-cuttable (easy),

Proof Vibes

We are now at the “starting line” of the proof, the rest of which requires us to do a few more things:

- Verify that the “highly extendable” subtrees \mathcal{T}_K are leaf-cuttable (easy),
- Verify that any k subtrees from \mathcal{T}_K have a pair with intersecting edge sets (true because $\mathcal{F}_{T,k}^q$ -free, but not so easy to prove),

Proof Vibes

We are now at the “starting line” of the proof, the rest of which requires us to do a few more things:

- Verify that the “highly extendable” subtrees \mathcal{T}_K are leaf-cuttable (easy),
- Verify that any k subtrees from \mathcal{T}_K have a pair with intersecting edge sets (true because $\mathcal{F}_{T,k}^q$ -free, but not so easy to prove),
- Verify that finding this set of $k - 1$ edges for each \mathcal{T}_K is enough to give the desired bound (annoying but doable).

Going Further

Theorem (English-S. 2025+)

For any tree T , integer $k \geq 1$, and family of graphs \mathcal{F} , either

$$\text{ex}(n, T, \mathcal{F}) = \Omega(n^k),$$

or every \mathcal{F} -free graph is $\mathcal{F}_{T,k}^q$ -free and

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

Going Further

Theorem (English-S. 2025+)

For any tree T , integer $k \geq 1$, and family of graphs \mathcal{F} , either

$$\text{ex}(n, T, \mathcal{F}) = \Omega(n^k),$$

or every \mathcal{F} -free graph is $\mathcal{F}_{T,k}^q$ -free and

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

Question

Does this hold with T replaced by some more general type of graph?

Going Further

Theorem (English-S. 2025+)

For any tree T , integer $k \geq 1$, and family of graphs \mathcal{F} , either

$$\text{ex}(n, T, \mathcal{F}) = \Omega(n^k),$$

or every \mathcal{F} -free graph is $\mathcal{F}_{T,k}^q$ -free and

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

Question

Does this hold with T replaced by some more general type of graph?

If this result holds at $k = \alpha(H) + 1$, then

$$\text{ex}(n, H, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{\alpha(H)})$$

Going Further

Theorem (English-S. 2025+)

For any tree T , integer $k \geq 1$, and family of graphs \mathcal{F} , either

$$\text{ex}(n, T, \mathcal{F}) = \Omega(n^k),$$

or every \mathcal{F} -free graph is $\mathcal{F}_{T,k}^q$ -free and

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

Question

Does this hold with T replaced by some more general type of graph?

If this result holds at $k = \alpha(H) + 1$, then

$$\text{ex}(n, H, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{\alpha(H)}),$$

and this holds for bipartite graphs without isolated edges by König's Theorem.

Going Further

Question

Is it the case that for every bipartite graph H without isolated vertices, integer k , and family of graphs \mathcal{F} that either

$$\text{ex}(n, H, \mathcal{F}) = \Omega(n^k),$$

or

$$\text{ex}(n, H, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

Going Further

Question

Is it the case that for every bipartite graph H without isolated vertices, integer k , and family of graphs \mathcal{F} that either

$$\text{ex}(n, H, \mathcal{F}) = \Omega(n^k),$$

or

$$\text{ex}(n, H, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

Fact

This is false for C_4 and $k = 2$:(

Going Further

We've seen a lot of ways that $\text{ex}(n, H, \mathcal{F})$ *can't* behave like; what about the ways it *can* behave?

Going Further

We've seen a lot of ways that $\text{ex}(n, H, \mathcal{F})$ *can't* behave like; what about the ways it *can* behave?

Definition

We say that a rational number r is *realizable* for a graph H if there exists a (finite) family of graphs \mathcal{F} such that $\text{ex}(n, H, \mathcal{F}) = \Theta(n^r)$.

Going Further

We've seen a lot of ways that $\text{ex}(n, H, \mathcal{F})$ *can't* behave like; what about the ways it *can* behave?

Definition

We say that a rational number r is *realizable* for a graph H if there exists a (finite) family of graphs \mathcal{F} such that $\text{ex}(n, H, \mathcal{F}) = \Theta(n^r)$.

Theorem (Bukh-Conlon 2018)

Every rational in $[1, 2]$ is realizable for $H = K_2$.

Going Further

Theorem (English-Halfpap-Krueger 2024)

Every rational in $[1, t]$ is realizable for $H = K_t$.

Going Further

Theorem (English-Halfpap-Krueger 2024)

Every rational in $[1, t]$ is realizable for $H = K_t$.

Theorem (English-Halfpap-Krueger 2024)

Every rational in $[t, t + 1]$ is realizable for $H = K_{1,t}$ and no rational in $(1, t)$ is.

Going Further

Theorem (English-S. 2025+)

For every graph H of maximum degree Δ , every rational in the range

$$\left[v(H) - \frac{e(H)}{2\Delta^2}, v(H) \right]$$

is realizable.

Going Further

Theorem (English-S. 2025+)

For every graph H of maximum degree Δ , every rational in the range

$$\left[v(H) - \frac{e(H)}{2\Delta^2}, v(H) \right]$$

is realizable.

Theorem (English-S. 2025+)

For every tree T of maximum degree Δ , every rational in the range

$$\left[v(T) - \frac{e(T)}{\Delta}, v(T) \right]$$

is realizable.

This is best possible for stars.

Open Problems

Question

Given a graph H , what rationals r are realizable?

Open Problems

Question

Given a graph H , what rationals r are realizable?

From our general stability theorem, we know that if $\text{ex}(n, H, \mathcal{F}_{H,k}^q) = o(n^\beta)$, then no rational in (β, k) is realizable.

Open Problems

Question

Given a graph H , what rationals r are realizable?

From our general stability theorem, we know that if $\text{ex}(n, H, \mathcal{F}_{H,k}^q) = o(n^\beta)$, then no rational in (β, k) is realizable.

Question

Is this the only way an exponent can fail to be rational?

Open Problems

Question

Given a graph H , what rationals r are realizable?

From our general stability theorem, we know that if $\text{ex}(n, H, \mathcal{F}_{H,k}^q) = o(n^\beta)$, then no rational in (β, k) is realizable.

Question

Is this the only way an exponent can fail to be rational?

We believe we can verify that this is true for all graphs on at most 4 vertices through various ad hoc techniques.

Open Problems

Question

Is it the case that every rational in $[\alpha(H), v(H)]$ is realizable?

This is true and best possible for cliques and stars.

Open Problems

Question

Is it the case that every rational in $[\alpha(H), v(H)]$ is realizable?

This is true and best possible for cliques and stars.

Conjecture

For every graph H , every rational in

$$\left[v(H) - \frac{e(H)}{\Delta}, v(H) \right]$$

is realizable.

Open Problems

Problem

Can you solve the realizable problem for simple families of graphs, e.g. paths?


Open Problems

Problem

Can you solve the realizable problem for simple families of graphs, e.g. paths?

Conjecture

We have $\text{ex}(n, P_7, F) = O(n^4)$ where F is two C_4 's sharing a vertex.



This is easy if we avoid just C_4 : we can specify the last two edges of P_7 and x_4 in $(n^{3/2})^2 \cdot n = n^4$ ways, and this uniquely determines the P_7 .

Open Problems

Say that a rational r is strongly realizable if there exists a single graph F such that $\text{ex}(n, H, F) = \Theta(n^r)$.

Open Problems

Say that a rational r is strongly realizable if there exists a single graph F such that $\text{ex}(n, H, F) = \Theta(n^r)$.

Question

Is it the case that for all H , every rational above $\alpha(H)$ is strongly realizable?

This is unknown even for $H = K_2$.

Open Problems

Say that a rational r is strongly realizable if there exists a single graph F such that $\text{ex}(n, H, F) = \Theta(n^r)$.

Question

Is it the case that for all H , every rational above $\alpha(H)$ is strongly realizable?

This is unknown even for $H = K_2$.

Question

Is it the case that for all H , the number of strongly realizable exponents below $\alpha(H)$ is $O_H(1)$?

Open Problems

Say that a rational r is strongly realizable if there exists a single graph F such that $\text{ex}(n, H, F) = \Theta(n^r)$.

Question

Is it the case that for all H , every rational above $\alpha(H)$ is strongly realizable?

This is unknown even for $H = K_2$.

Question

Is it the case that for all H , the number of strongly realizable exponents below $\alpha(H)$ is $O_H(1)$?

The idea here is that any F of this form must be a subgraph of every member of $\mathcal{F}_{H, \alpha(H)}^q \neq \emptyset$ for some q . For any fixed q this is a finite number of possibilities, and I don't think arbitrarily large q should give new behaviors (analogous to $F \subseteq K_{2,t}$ implying $\text{ex}(n, F) = \Theta(n^r)$ for some $r \in \{0, 1, 3/2\}$).

Open Problems

Our new Helly theorem for edge sets of trees turns out to imply the usual Helly theorem for vertex sets.

Question

Does the usual vertex Helly Theorem imply the edge Helly Theorem?

Open Problems

Our new Helly theorem for edge sets of trees turns out to imply the usual Helly theorem for vertex sets.

Question

Does the usual vertex Helly Theorem imply the edge Helly Theorem? In particular, can you derive König's Theorem for trees?

Open Problems

Our new Helly theorem for edge sets of trees turns out to imply the usual Helly theorem for vertex sets.

Question

Does the usual vertex Helly Theorem imply the edge Helly Theorem? In particular, can you derive König's Theorem for trees?

I'm willing to offer up to \$1 for a proof* or disproof of either of these questions.

Open Problems

Our new Helly theorem for edge sets of trees turns out to imply the usual Helly theorem for vertex sets.

Question

Does the usual vertex Helly Theorem imply the edge Helly Theorem? In particular, can you derive König's Theorem for trees?

I'm willing to offer up to \$1 for a proof* or disproof of either of these questions.

Question

Can one prove that if T_1, \dots are “reasonable” subtrees of a tree T which pairwise intersect in at least one path of length 2, then there exists a path of length 2 common to each T_i ?

Open Problems

Question

Can one say anything about generalized Turán numbers of trees T satisfying

$$n^{\ell+1} \ll \text{ex}(n, T, \mathcal{F}) \ll n^{\ell+2}?$$

For this it would suffice to prove upper bounds on $\text{ex}(n, \mathcal{F}_{T, \ell+2}^q)$.

Open Problems: Supersaturation

Open Problems: Supersaturation

Proposition

If G contains $kn^{2-1/s}$ edges with $k = \omega(1)$, then it contains at least $\Omega(k^{st}n^s)$ copies of $K_{s,t}$.

Note that this is best possible by considering a random graph with $kn^{2-1/s}$ edges.

Open Problems: Supersaturation

Proposition

If G contains $kn^{2-1/s}$ edges with $k = \omega(1)$, then it contains at least $\Omega(k^{st}n^s)$ copies of $K_{s,t}$.

Note that this is best possible by considering a random graph with $kn^{2-1/s}$ edges.

Question

Does the same phenomenon hold if G contains many copies of K_r ?

Open Problems: Supersaturation

Conjecture (Dubroff-Gunby-Narayanan-S.)

If $2 \leq r \leq s \leq t$ and if G contains at least $kn^{r - \binom{r}{2}/s}$ copies of K_r , then it contains at least $k^{st/\binom{r}{2}} n^{s - o(1)}$ copies of $K_{s,t}$.

Open Problems: Supersaturation

Conjecture (Dubroff-Gunby-Narayanan-S.)

If $2 \leq r \leq s \leq t$ and if G contains at least $kn^{r - \binom{r}{2}/s}$ copies of K_r , then it contains at least $k^{st/\binom{r}{2}} n^{s - o(1)}$ copies of $K_{s,t}$.

Things seem to behave differently if $r > s$ and k is small.

Open Problems: Supersaturation

Conjecture (Dubroff-Gunby-Narayanan-S.)

If $2 \leq r \leq s \leq t$ and if G contains at least $kn^{r - \binom{r}{2}/s}$ copies of K_r , then it contains at least $k^{st/\binom{r}{2}} n^{s-o(1)}$ copies of $K_{s,t}$.

Things seem to behave differently if $r > s$ and k is small.

Conjecture (Dubroff-Gunby-Narayanan-S.)

For all $1 \leq k \leq n^{1/2t}$, there exist n -vertex graphs with $kn^{3/2}$ triangles and at most $k^t n^{3/2+o(1)}$ copies of $K_{2,t}$.