

# Generalized Turán Problems for Trees and More

Sam Spiro, Georgia State University

Based on Joint Work with Sean English



Feedback can be given at [binatorics.com](http://binatorics.com)

Done

## Leaderboard

Current Rankings

My Points

**Sam Spiro**

15400 pts



You made it on the board! Tell people how awesome you are.

[Share](#)**June Lee**

14700 pts

**Jacob Platnick**

12100 pts



4

**Irena Swanson**

11000 pts



5

**Shannon Auckly**

9800 pts



# Turán Problems

The Turán number  $\text{ex}(n, F)$  is defined to be the maximum number of edges that an  $F$ -free graph on  $n$  vertices can have.

# Turán Problems

The Turán number  $\text{ex}(n, F)$  is defined to be the maximum number of edges that an  $F$ -free graph on  $n$  vertices can have.

## Theorem

$$\text{ex}(n, K_2) = 0.$$

# Turán Problems

## Theorem (Turán 1941)

$$\text{ex}(n, K_r) = \left\lfloor \binom{r-1}{2} \frac{n^2}{(r-1)^2} \right\rfloor$$

# Turán Problems

## Theorem (Turán 1941)

$$\text{ex}(n, K_r) = \left\lfloor \binom{r-1}{2} \frac{n^2}{(r-1)^2} \right\rfloor$$

## Theorem (Kővari-Sós-Turán 1954)

$$\text{ex}(n, K_{s,t}) = O(n^{2-1/s}).$$

# Turán Problems

## Theorem (Turán 1941)

$$\text{ex}(n, K_r) = \left\lfloor \binom{r-1}{2} \frac{n^2}{(r-1)^2} \right\rfloor$$

## Theorem (Kővari-Sós-Turán 1954)

$$\text{ex}(n, K_{s,t}) = O(n^{2-1/s}).$$

## Theorem (Folklore)

*If  $T$  is a tree with  $e(T) \geq 2$ , then*

$$\text{ex}(n, T) = \Theta(n).$$

# Turán Problems

Given a family of graphs  $\mathcal{F}$ , we say  $G$  is  $\mathcal{F}$ -free if  $G$  is  $F$ -free for all  $F \in \mathcal{F}$ . We define  $\text{ex}(n, \mathcal{F})$  to be the maximum number of edges in an  $n$ -vertex  $\mathcal{F}$ -free graph.



# Turán Problems

Given a family of graphs  $\mathcal{F}$ , we say  $G$  is  $\mathcal{F}$ -free if  $G$  is  $F$ -free for all  $F \in \mathcal{F}$ . We define  $\text{ex}(n, \mathcal{F})$  to be the maximum number of edges in an  $n$ -vertex  $\mathcal{F}$ -free graph.

## Conjecture (Compactness Conjecture)

*If  $\mathcal{F}$  is a finite family of graphs, then  $\text{ex}(n, \mathcal{F}) = \Theta(\text{ex}(n, F))$  for some  $F \in \mathcal{F}$ .*

# Generalized Turán Problems

# Generalized Turán Problems

## Definition (Alon-Shikhelman 2016)

Given a graph  $H$  and a family of graphs  $\mathcal{F}$ , we define the generalized Turán number  $\text{ex}(n, H, \mathcal{F})$  to be the maximum number of copies of  $H$  in an  $n$ -vertex  $\mathcal{F}$ -free graph.

# Generalized Turán Problems

## Definition (Alon-Shikhelman 2016)

Given a graph  $H$  and a family of graphs  $\mathcal{F}$ , we define the generalized Turán number  $\text{ex}(n, H, \mathcal{F})$  to be the maximum number of copies of  $H$  in an  $n$ -vertex  $\mathcal{F}$ -free graph.

## Proposition

$$\text{ex}(n, K_2, \mathcal{F}) = \text{ex}(n, \mathcal{F}).$$

# Generalized Turán Problems

## Definition (Alon-Shikhelman 2016)

Given a graph  $H$  and a family of graphs  $\mathcal{F}$ , we define the generalized Turán number  $\text{ex}(n, H, \mathcal{F})$  to be the maximum number of copies of  $H$  in an  $n$ -vertex  $\mathcal{F}$ -free graph.

## Proposition

$$\text{ex}(n, K_2, \mathcal{F}) = \text{ex}(n, \mathcal{F}).$$

See the recent survey by Gerbner and Palmer for way more history than I'm going to give here.

# Generalized Turán Problems

## Theorem (Gerbner-Palmer 2019)

*For all  $r \geq 2$  and families  $\mathcal{F}$  we have*

$$\text{ex}(n, K_r, \mathcal{F}) \leq \text{ex}(n, \mathcal{F})^{r/2}.$$

# Generalized Turán Problems

## Theorem (Gerbner-Palmer 2019)

*For all  $r \geq 2$  and families  $\mathcal{F}$  we have*

$$\text{ex}(n, K_r, \mathcal{F}) \leq \text{ex}(n, \mathcal{F})^{r/2}.$$

## Theorem (Gerbner 2023)

*Determines  $\text{ex}(n, K_{2,t}, F)$  asymptotically for any non-bipartite graph  $F$ .*

# Generalized Turán Problems

## Theorem (Gerbner-Palmer 2019)

*For all  $r \geq 2$  and families  $\mathcal{F}$  we have*

$$\text{ex}(n, K_r, \mathcal{F}) \leq \text{ex}(n, \mathcal{F})^{r/2}.$$

## Theorem (Gerbner 2023)

*Determines  $\text{ex}(n, K_{2,t}, F)$  asymptotically for any non-bipartite graph  $F$ .*

## Theorem (Füredi-Kündgen 2006)

*If  $\text{ex}(n, \mathcal{F}) = \Theta(n^{2-\beta})$  and  $\mathcal{F}$  does not contain a star, then*

$$\text{ex}(n, K_{1,t}, \mathcal{F}) = \tilde{\Theta}(\max\{n^t, n^{t+1-t\beta}\}).$$



# Generalized Turán Problems

## Question

Can we prove bounds on  $\text{ex}(n, T, \mathcal{F})$  for arbitrary trees  $T$  (for some possibly fixed  $\mathcal{F}$ )?

# Generalized Turán Problems

## Question

Can we prove bounds on  $\text{ex}(n, T, \mathcal{F})$  for arbitrary trees  $T$  (for some possibly fixed  $\mathcal{F}$ )?

## Theorem (Gerbner 2023)

*Determines the order of magnitude of  $\text{ex}(n, T, K_{2,t})$  for every tree  $T$ .*

# Generalized Turán Problems

## Question

Can we prove bounds on  $\text{ex}(n, T, \mathcal{F})$  for arbitrary trees  $T$  (for some possibly fixed  $\mathcal{F}$ )?

## Theorem (Gerbner 2023)

*Determines the order of magnitude of  $\text{ex}(n, T, K_{2,t})$  for every tree  $T$ .*

## Theorem (Alon-Shikhelman 2016)

*If  $T$  is a tree and  $F$  is a tree, then  $\text{ex}(n, T, F) = \Theta(n^k)$  for some integer  $k$ .*

# Generalized Turán Problems

## Question

Can we prove bounds on  $\text{ex}(n, T, \mathcal{F})$  for arbitrary trees  $T$  (for some possibly fixed  $\mathcal{F}$ )?

## Theorem (Gerbner 2023)

*Determines the order of magnitude of  $\text{ex}(n, T, K_{2,t})$  for every tree  $T$ .*

## Theorem (Alon-Shikhelman 2016)

*If  $T$  is a tree and  $F$  is a tree, then  $\text{ex}(n, T, F) = \Theta(n^k)$  for some integer  $k$ .*

## Theorem (Letzter 2019)

*If  $H$  is any graph and  $F$  is a tree, then  $\text{ex}(n, H, F) = \Theta(n^k)$  for some integer  $k$ .*

# A General Theorem for Trees

## Theorem (English-S. 2025+)

*For any tree  $T$ , integer  $k \geq 1$ , and family of graphs  $\mathcal{F}$ , either*

$$\text{ex}(n, T, \mathcal{F}) = \Omega(n^k),$$

*or*

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

*Moreover, we can determine which of these two cases happen for a given  $T, k, \mathcal{F}$ .*

# A General Theorem for Trees

Theorem (English-S. 2025+)

*Either  $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$  or  $\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1})$ .*

# A General Theorem for Trees

## Theorem (English-S. 2025+)

*Either  $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$  or  $\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1})$ .*

## Corollary

*If  $T$  is a tree and  $\mathcal{F}$  contains a forest, then  $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$  for some integer  $k$ .*

# A General Theorem for Trees

## Theorem (English-S. 2025+)

*Either  $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$  or  $\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1})$ .*

## Corollary

*If  $T$  is a tree and  $\mathcal{F}$  contains a forest, then  $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$  for some integer  $k$ .*

## Proof.

Let  $k$  be the smallest integer such that  $\text{ex}(n, T, \mathcal{F}) = O(n^k)$ .



# A General Theorem for Trees

## Theorem (English-S. 2025+)

*Either  $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$  or  $\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1})$ .*

## Corollary

*If  $T$  is a tree and  $\mathcal{F}$  contains a forest, then  $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$  for some integer  $k$ .*

## Proof.

Let  $k$  be the smallest integer such that  $\text{ex}(n, T, \mathcal{F}) = O(n^k)$ . If  $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$  then we're done

# A General Theorem for Trees

## Theorem (English-S. 2025+)

*Either  $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$  or  $\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1})$ .*

## Corollary

*If  $T$  is a tree and  $\mathcal{F}$  contains a forest, then  $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$  for some integer  $k$ .*

## Proof.

Let  $k$  be the smallest integer such that  $\text{ex}(n, T, \mathcal{F}) = O(n^k)$ . If  $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$  then we're done, otherwise our theorem implies that

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1})$$

# A General Theorem for Trees

## Theorem (English-S. 2025+)

*Either  $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$  or  $\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1})$ .*

## Corollary

*If  $T$  is a tree and  $\mathcal{F}$  contains a forest, then  $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$  for some integer  $k$ .*

## Proof.

Let  $k$  be the smallest integer such that  $\text{ex}(n, T, \mathcal{F}) = O(n^k)$ . If  $\text{ex}(n, T, \mathcal{F}) = \Omega(n^k)$  then we're done, otherwise our theorem implies that

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}) = O(n^{k-1}).$$



# A General Theorem for Trees

## Corollary

*If  $T \neq K_2$  is a tree with  $\ell \geq 2$  leaves, then any family  $\mathcal{F}$  with  $\text{ex}(n, T, \mathcal{F}) = O(n^\ell)$  has  $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$  for some integer  $k$ .*

In particular, if  $\text{ex}(n, T, \mathcal{F}) = o(n^\ell)$  then  $\text{ex}(n, T, \mathcal{F}) = O(n^{\ell-1})$ .

# A General Theorem for Trees

## Corollary

*If  $T \neq K_2$  is a tree with  $\ell \geq 2$  leaves, then any family  $\mathcal{F}$  with  $\text{ex}(n, T, \mathcal{F}) = O(n^\ell)$  has  $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$  for some integer  $k$ .*

In particular, if  $\text{ex}(n, T, \mathcal{F}) = o(n^\ell)$  then  $\text{ex}(n, T, \mathcal{F}) = O(n^{\ell-1})$ .

## Proof.

If  $\mathcal{F}$  contains a forest then we are done by the previous result.

# A General Theorem for Trees

## Corollary

*If  $T \neq K_2$  is a tree with  $\ell \geq 2$  leaves, then any family  $\mathcal{F}$  with  $\text{ex}(n, T, \mathcal{F}) = O(n^\ell)$  has  $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$  for some integer  $k$ .*

In particular, if  $\text{ex}(n, T, \mathcal{F}) = o(n^\ell)$  then  $\text{ex}(n, T, \mathcal{F}) = O(n^{\ell-1})$ .

## Proof.

If  $\mathcal{F}$  contains a forest then we are done by the previous result. Otherwise, the graph  $G$  obtained by duplicating each leaf of  $T$  a total of  $n/\ell$  times is  $\mathcal{F}$ -free

# A General Theorem for Trees

## Corollary

*If  $T \neq K_2$  is a tree with  $\ell \geq 2$  leaves, then any family  $\mathcal{F}$  with  $\text{ex}(n, T, \mathcal{F}) = O(n^\ell)$  has  $\text{ex}(n, T, \mathcal{F}) = \Theta(n^k)$  for some integer  $k$ .*

In particular, if  $\text{ex}(n, T, \mathcal{F}) = o(n^\ell)$  then  $\text{ex}(n, T, \mathcal{F}) = O(n^{\ell-1})$ .

## Proof.

If  $\mathcal{F}$  contains a forest then we are done by the previous result. Otherwise, the graph  $G$  obtained by duplicating each leaf of  $T$  a total of  $n/\ell$  times is  $\mathcal{F}$ -free and contains  $\Omega(n^\ell)$  copies of  $T$ . □

# A General Theorem for Trees

## Theorem (English-S. 2025++)

*If  $T$  is a tree with  $\ell \geq 2$  leaves, then every family  $\mathcal{F}$  with  $\text{ex}(n, T, \mathcal{F}) = o(n^{\ell+1})$  satisfies*

$$\text{ex}(n, T, \mathcal{F}) = O\left(n^{\ell + \frac{\ell^2 - \ell}{e(T) - 1}}\right).$$



# A General Theorem for Trees

## Theorem (English-S. 2025++)

*If  $T$  is a tree with  $\ell \geq 2$  leaves, then every family  $\mathcal{F}$  with  $\text{ex}(n, T, \mathcal{F}) = o(n^{\ell+1})$  satisfies*

$$\text{ex}(n, T, \mathcal{F}) = O\left(n^{\ell + \frac{\ell^2 - \ell}{e(T) - 1}}\right).$$

*Moreover, this bound is best possible for essentially all values of  $\ell, e(T)$ .*

# A General Theorem for Trees

## Theorem (English-S. 2025++)

If  $T$  is a tree with  $\ell \geq 2$  leaves, then every family  $\mathcal{F}$  with  $\text{ex}(n, T, \mathcal{F}) = o(n^{\ell+1})$  satisfies

$$\text{ex}(n, T, \mathcal{F}) = O\left(n^{\ell + \frac{\ell^2 - \ell}{e(T) - 1}}\right).$$

Moreover, this bound is best possible for essentially all values of  $\ell, e(T)$ .

## Theorem (English-S. 20??++)

For the path graph  $P_t$ , every graph  $F$  with  $\text{ex}(n, P_t, F) = O(n^{\alpha(P_t)})$  has  $\text{ex}(n, P_t, F) = \Theta(n^k)$  for some integer  $k$ .

# Stability for Generalized Turán Problems

# Stability for Generalized Turán Problems

## Proposition (English-Halfpap-Krueger 2024)

*For stars  $K_{1,t}$ , every family of graphs  $\mathcal{F}$  either satisfies  $\text{ex}(n, K_{1,t}, \mathcal{F}) = \Omega(n^t)$  or  $\text{ex}(n, K_{1,t}, \mathcal{F}) = O(n)$ .*

# Stability for Generalized Turán Problems

## Proposition (English-Halfpap-Krueger 2024)

*For stars  $K_{1,t}$ , every family of graphs  $\mathcal{F}$  either satisfies  $\text{ex}(n, K_{1,t}, \mathcal{F}) = \Omega(n^t)$  or  $\text{ex}(n, K_{1,t}, \mathcal{F}) = O(n)$ .*

### Proof.

If  $\mathcal{F}$  does not contain a subgraph of a star, then  $G = K_{1,n-1}$  is  $\mathcal{F}$ -free and shows  $\text{ex}(n, K_{1,t}, \mathcal{F}) = \Omega(n^t)$ .

# Stability for Generalized Turán Problems

## Proposition (English-Halfpap-Krueger 2024)

*For stars  $K_{1,t}$ , every family of graphs  $\mathcal{F}$  either satisfies  $\text{ex}(n, K_{1,t}, \mathcal{F}) = \Omega(n^t)$  or  $\text{ex}(n, K_{1,t}, \mathcal{F}) = O(n)$ .*

### Proof.

If  $\mathcal{F}$  does not contain a subgraph of a star, then  $G = K_{1,n-1}$  is  $\mathcal{F}$ -free and shows  $\text{ex}(n, K_{1,t}, \mathcal{F}) = \Omega(n^t)$ . Otherwise,  $\mathcal{F}$  must contain some subgraph of some star  $K_{1,r}$ , which means  $\text{ex}(n, K_{1,t}, \mathcal{F}) = O_r(n)$ .  $\square$

# Stability for Generalized Turán Problems

## Proposition (English-Halfpap-Krueger 2024)

*For stars  $K_{1,t}$ , every family of graphs  $\mathcal{F}$  either satisfies  $\text{ex}(n, K_{1,t}, \mathcal{F}) = \Omega(n^t)$  or  $\text{ex}(n, K_{1,t}, \mathcal{F}) = O(n)$ .*

### Proof.

If  $\mathcal{F}$  does not contain a subgraph of a star, then  $G = K_{1,n-1}$  is  $\mathcal{F}$ -free and shows  $\text{ex}(n, K_{1,t}, \mathcal{F}) = \Omega(n^t)$ . Otherwise,  $\mathcal{F}$  must contain some subgraph of some star  $K_{1,r}$ , which means  $\text{ex}(n, K_{1,t}, \mathcal{F}) = O_r(n)$ .  $\square$

We want to generalize this idea by saying that for *any* graph  $H$ , there exists some “simple” family  $\mathcal{F}_H$  such that the behavior of  $\text{ex}(n, K_{1,t}, \mathcal{F})$  depends on how  $\mathcal{F}$  “interacts” with  $\mathcal{F}_H$ .

# Stability for Generalized Turán Problems

## Definition


Given a graph  $H$ , a subset  $R \subseteq V(H)$ , and an integer  $q$ , we define the sunflower-power  $H_R^q$  to be the graph obtained by taking  $q$  copies of  $H$  which all agree on  $R$  and which are otherwise disjoint.



# Stability for Generalized Turán Problems

## Definition


Given a graph  $H$ , a subset  $R \subseteq V(H)$ , and an integer  $q$ , we define the sunflower-power  $H_R^q$  to be the graph obtained by taking  $q$  copies of  $H$  which all agree on  $R$  and which are otherwise disjoint.




# Stability for Generalized Turán Problems

## Definition

Given a graph  $H$ , a subset  $R \subseteq V(H)$ , and an integer  $q$ , we define the sunflower-power  $H_R^q$  to be the graph obtained by taking  $q$  copies of  $H$  which all agree on  $R$  and which are otherwise disjoint.




$$(P_5)_{\{x_3, x_4\}}^3$$




# Stability for Generalized Turán Problems

## Definition

Given a graph  $H$ , a subset  $R \subseteq V(H)$ , and an integer  $q$ , we define the sunflower-power  $H_R^q$  to be the graph obtained by taking  $q$  copies of  $H$  which all agree on  $R$  and which are otherwise disjoint.



$(P_5)_{\{x_3, x_4\}}^3$




We define

$$\mathcal{F}_{H,k}^q = \{H_R^q : H - R \text{ has at least } k \text{ connected components}\}.$$


# Stability for Generalized Turán Problems

## Definition

Given a graph  $H$ , a subset  $R \subseteq V(H)$ , and an integer  $q$ , we define the sunflower-power  $H_R^q$  to be the graph obtained by taking  $q$  copies of  $H$  which all agree on  $R$  and which are otherwise disjoint.



$(P_5)_{\{x_3, x_4\}}^3$



We define

$$\mathcal{F}_{H,k}^q = \{H_R^q : H - R \text{ has at least } k \text{ connected components}\}.$$

## Claim

Every graph in  $\mathcal{F}_{H,k}^q$  has at least  $q^k$  copies of  $H$ .

# Stability for Generalized Turán Problems

## Proposition (Key Observation)

*For every  $H, k, \mathcal{F}$ , either  $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$  or there exists some  $q$  such that every  $\mathcal{F}$ -free graph is  $\mathcal{F}_{H,k}^q$ -free.*

# Stability for Generalized Turán Problems

## Proposition (Key Observation)

*For every  $H, k, \mathcal{F}$ , either  $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$  or there exists some  $q$  such that every  $\mathcal{F}$ -free graph is  $\mathcal{F}_{H,k}^q$ -free.*

## Proof.

If some  $H_R^q \in \mathcal{F}_{H,k}^q$  is always  $\mathcal{F}$ -free, then the previous claim with  $q \approx n/v(H)$  shows  $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$ .

# Stability for Generalized Turán Problems

## Proposition (Key Observation)

*For every  $H, k, \mathcal{F}$ , either  $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$  or there exists some  $q$  such that every  $\mathcal{F}$ -free graph is  $\mathcal{F}_{H,k}^q$ -free.*

## Proof.

If some  $H_R^q \in \mathcal{F}_{H,k}^q$  is always  $\mathcal{F}$ -free, then the previous claim with  $q \approx n/v(H)$  shows  $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$ . Otherwise there exists some  $q$  so that every  $H_R^q \in \mathcal{F}_{H,k}^q$  contains an element of  $\mathcal{F}$  as a subgraph.  $\square$

# Stability for Generalized Turán Problems

## Proposition (Key Observation)

*For every  $H, k, \mathcal{F}$ , either  $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$  or there exists some  $q$  such that every  $\mathcal{F}$ -free graph is  $\mathcal{F}_{H,k}^q$ -free.*

## Proof.

If some  $H_R^q \in \mathcal{F}_{H,k}^q$  is always  $\mathcal{F}$ -free, then the previous claim with  $q \approx n/v(H)$  shows  $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$ . Otherwise there exists some  $q$  so that every  $H_R^q \in \mathcal{F}_{H,k}^q$  contains an element of  $\mathcal{F}$  as a subgraph.  $\square$

## Corollary (General Stability for Generalized Turán)

*If  $\text{ex}(n, H, \mathcal{F}_{H,k}^q) = O_q(n^\beta)$ , then every family  $\mathcal{F}$  either satisfies  $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$  or  $\text{ex}(n, H, \mathcal{F}) = O(n^\beta)$ .*



# Stability for Generalized Turán Problems

## Proposition

*For every graph  $H$  and family  $\mathcal{F}$ , either  $\text{ex}(n, H, \mathcal{F}) = \Omega(n)$  or  $\text{ex}(n, H, \mathcal{F}) = O(1)$ .*

# Stability for Generalized Turán Problems

## Proposition

*For every graph  $H$  and family  $\mathcal{F}$ , either  $\text{ex}(n, H, \mathcal{F}) = \Omega(n)$  or  $\text{ex}(n, H, \mathcal{F}) = O(1)$ .*


By the previous corollary, it suffices to prove  $\text{ex}(n, H, \mathcal{F}_{H,1}^q) = O_q(1)$ .

# Stability for Generalized Turán Problems

## Proposition

*For every graph  $H$  and family  $\mathcal{F}$ , either  $\text{ex}(n, H, \mathcal{F}) = \Omega(n)$  or  $\text{ex}(n, H, \mathcal{F}) = O(1)$ .*

By the previous corollary, it suffices to prove  $\text{ex}(n, H, \mathcal{F}_{H,1}^q) = O_q(1)$ .




# Stability for Generalized Turán Problems

## Proposition

*For every graph  $H$  and family  $\mathcal{F}$ , either  $\text{ex}(n, H, \mathcal{F}) = \Omega(n)$  or  $\text{ex}(n, H, \mathcal{F}) = O(1)$ .*

By the previous corollary, it suffices to prove  $\text{ex}(n, H, \mathcal{F}_{H,1}^q) = O_q(1)$ .



This is essentially equivalent to the Erdős-Rado Sunflower lemma. □

# Stability for Generalized Turán Problems

## Theorem (English-S. 2025+)

*For any tree  $T$ , integer  $k \geq 1$ , and family of graphs  $\mathcal{F}$ , either*

$$\text{ex}(n, T, \mathcal{F}) = \Omega(n^k),$$

*or*

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

# Stability for Generalized Turán Problems

## Theorem (English-S. 2025+)

*For any tree  $T$ , integer  $k \geq 1$ , and family of graphs  $\mathcal{F}$ , either*

$$\text{ex}(n, T, \mathcal{F}) = \Omega(n^k),$$

*or*

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

## Proposition (Key Observation)

*For every  $H, k, \mathcal{F}$ , either  $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$  or there exists some  $q$  such that every  $\mathcal{F}$ -free graph is  $\mathcal{F}_{H,k}^q$ -free.*

# Stability for Generalized Turán Problems

## Theorem (English-S. 2025+)

*For any tree  $T$ , integer  $k \geq 1$ , and family of graphs  $\mathcal{F}$ , either*

$$\text{ex}(n, T, \mathcal{F}) = \Omega(n^k),$$

*or*

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

## Proposition (Key Observation)

*For every  $H, k, \mathcal{F}$ , either  $\text{ex}(n, H, \mathcal{F}) = \Omega(n^k)$  or there exists some  $q$  such that every  $\mathcal{F}$ -free graph is  $\mathcal{F}_{H,k}^q$ -free.*

## Theorem (English-S. 2025+)

*If  $T$  is a tree and if  $G$  is  $\mathcal{F}_{T,k}^q$ -free, then the number of copies of  $T$  in  $G$  is at most  $O(e(G)^{k-1})$ .*

# Proof Sketch



# Proof Vibes

# Proof Vibes

## Theorem (English-S. 2025+)

*If  $T$  is a tree and if  $G$  is  $\mathcal{F}_{T,k}^q$ -free, then the number of copies of  $T$  in  $G$  is at most  $O(e(G)^{k-1})$ .*

# Proof Vibes

## Theorem (English-S. 2025+)

*If  $T$  is a tree and if  $G$  is  $\mathcal{F}_{T,k}^q$ -free, then the number of copies of  $T$  in  $G$  is at most  $O(e(G)^{k-1})$ .*


Strategy: identify each copy of  $T$  in  $G$  by a set of  $k - 1$  edges  $E$  such that each  $E$  identifies at most  $O(1)$  copies of  $T$ .

# Proof Vibes

## Theorem (English-S. 2025+)

*If  $T$  is a tree and if  $G$  is  $\mathcal{F}_{T,k}^q$ -free, then the number of copies of  $T$  in  $G$  is at most  $O(e(G)^{k-1})$ .*

Strategy: identify each copy of  $T$  in  $G$  by a set of  $k-1$  edges  $E$  such that each  $E$  identifies at most  $O(1)$  copies of  $T$ .




# Proof Vibes

## Theorem (English-S. 2025+)

If  $T$  is a tree and if  $G$  is  $\mathcal{F}_{T,k}^q$ -free, then the number of copies of  $T$  in  $G$  is at most  $O(e(G)^{k-1})$ .

Strategy: identify each copy of  $T$  in  $G$  by a set of  $k - 1$  edges  $E$  such that each  $E$  identifies at most  $O(1)$  copies of  $T$ .




For this example, to specify a given copy of  $P_5$ , we **must** identify its last edge.

# Proof Vibes

## Theorem (English-S. 2025+)

If  $T$  is a tree and if  $G$  is  $\mathcal{F}_{T,k}^q$ -free, then the number of copies of  $T$  in  $G$  is at most  $O(e(G)^{k-1})$ .

Strategy: identify each copy of  $T$  in  $G$  by a set of  $k - 1$  edges  $E$  such that each  $E$  identifies at most  $O(1)$  copies of  $T$ .



For this example, to specify a given copy of  $P_5$ , we **must** identify its last edge. More generally, whatever set of edges  $E$  we choose to identify a given copy  $K$  in a graph, the edges of  $E$  **must** intersect every subtree  $K' \subseteq K$  which has “many extensions.”

# Proof Vibes

## Question

Given a set of subtrees  $\mathcal{T}$  of a tree  $T$ , when can we guarantee that there exist a set of  $k - 1$  edges  $E$  which intersect all of these subtrees?

# Proof Vibes

## Question


Given a set of subtrees  $\mathcal{T}$  of a tree  $T$ , when can we guarantee that there exist a set of  $k - 1$  edges  $E$  which intersect all of these subtrees?

## Theorem (Helly Property for Trees)


*If  $\mathcal{T}$  is a set of subtrees of a tree  $T$  such that the vertex sets of the subtrees pairwise intersect, then there exists a vertex  $v \in V(T)$  which intersects the vertex set of every subtree of  $\mathcal{T}$ .*



# Proof Vibes



# Proof Vibes



## Definition

We call a subtree  $T' \subseteq T$  leaf-cuttable if  $e(T) \geq 1$  and if every edge in  $E(T) \setminus E(T')$  which intersects  $T'$  intersects a leaf of  $T'$ .

# Proof Vibes

## Theorem (English-S. 2025+)

*If  $\mathcal{T}$  is a set of leaf-cuttable subtrees of a tree  $T$  such that the edge sets of the subtrees pairwise intersect, then there exists an edge  $e \in E(T)$  which intersects the edge set of every subtree of  $\mathcal{T}$ .*

# Proof Vibes

## Theorem (English-S. 2025+)

*If  $\mathcal{T}$  is a set of leaf-cuttable subtrees of a tree  $T$  such that the edge sets of the subtrees pairwise intersect, then there exists an edge  $e \in E(T)$  which intersects the edge set of every subtree of  $\mathcal{T}$ .*

## Theorem (English-S. 2025+)

*If  $\mathcal{T}$  is a set of leaf-cuttable subtrees of a tree  $T$  such that every  $k$  subtrees from  $\mathcal{T}$  contains two subtrees with intersecting edge sets, then there exists a set  $E \subseteq E(T)$  of at most  $k - 1$  edges which intersects the edge set of every subtree of  $\mathcal{T}$ .*

# Proof Vibes

## Theorem (English-S. 2025+)

*If  $\mathcal{T}$  is a set of leaf-cuttable subtrees of a tree  $T$  such that the edge sets of the subtrees pairwise intersect, then there exists an edge  $e \in E(T)$  which intersects the edge set of every subtree of  $\mathcal{T}$ .*

## Theorem (English-S. 2025+)

*If  $\mathcal{T}$  is a set of leaf-cuttable subtrees of a tree  $T$  such that every  $k$  subtrees from  $\mathcal{T}$  contains two subtrees with intersecting edge sets, then there exists a set  $E \subseteq E(T)$  of at most  $k - 1$  edges which intersects the edge set of every subtree of  $\mathcal{T}$ .*

This result implies the vertex-Helly result (and also König's Theorem for trees).

# Proof Vibes

We are now at the “starting line” of the proof, the rest of which requires us to do a few more things:

# Proof Vibes

We are now at the “starting line” of the proof, the rest of which requires us to do a few more things:

- Verify that the “highly extendable” subtrees  $\mathcal{T}_K$  are leaf-cuttable (easy),

# Proof Vibes

We are now at the “starting line” of the proof, the rest of which requires us to do a few more things:

- Verify that the “highly extendable” subtrees  $\mathcal{T}_K$  are leaf-cuttable (easy),
- Verify that any  $k$  subtrees from  $\mathcal{T}_K$  have a pair with intersecting edge sets (true because  $\mathcal{F}_{T,k}^q$ -free, but not so easy to prove),



# Proof Vibes

We are now at the “starting line” of the proof, the rest of which requires us to do a few more things:

- Verify that the “highly extendable” subtrees  $\mathcal{T}_K$  are leaf-cuttable (easy),
- Verify that any  $k$  subtrees from  $\mathcal{T}_K$  have a pair with intersecting edge sets (true because  $\mathcal{F}_{T,k}^q$ -free, but not so easy to prove),
- Verify that finding this set of  $k - 1$  edges for each  $\mathcal{T}_K$  is enough to give the desired bound (annoying but doable).

# Going Further

## Theorem (English-S. 2025+)

*For any tree  $T$ , integer  $k \geq 1$ , and family of graphs  $\mathcal{F}$ , either*

$$\text{ex}(n, T, \mathcal{F}) = \Omega(n^k),$$

*or every  $\mathcal{F}$ -free graph is  $\mathcal{F}_{T,k}^q$ -free and*

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

## Going Further

### Theorem (English-S. 2025+)

*For any tree  $T$ , integer  $k \geq 1$ , and family of graphs  $\mathcal{F}$ , either*

$$\text{ex}(n, T, \mathcal{F}) = \Omega(n^k),$$

*or every  $\mathcal{F}$ -free graph is  $\mathcal{F}_{T,k}^q$ -free and*

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

### Question

Does this hold with  $T$  replaced by some more general type of graph?

## Going Further

### Theorem (English-S. 2025+)

*For any tree  $T$ , integer  $k \geq 1$ , and family of graphs  $\mathcal{F}$ , either*

$$\text{ex}(n, T, \mathcal{F}) = \Omega(n^k),$$

*or every  $\mathcal{F}$ -free graph is  $\mathcal{F}_{T,k}^q$ -free and*

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

### Question

Does this hold with  $T$  replaced by some more general type of graph?

If this result holds at  $k = \alpha(H) + 1$ , then

$$\text{ex}(n, H, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{\alpha(H)})$$

## Going Further

### Theorem (English-S. 2025+)

*For any tree  $T$ , integer  $k \geq 1$ , and family of graphs  $\mathcal{F}$ , either*

$$\text{ex}(n, T, \mathcal{F}) = \Omega(n^k),$$

*or every  $\mathcal{F}$ -free graph is  $\mathcal{F}_{T,k}^q$ -free and*

$$\text{ex}(n, T, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

### Question

Does this hold with  $T$  replaced by some more general type of graph?

If this result holds at  $k = \alpha(H) + 1$ , then

$$\text{ex}(n, H, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{\alpha(H)}),$$

and this holds for bipartite graphs without isolated edges by König's Theorem.

# Going Further

## Question

Is it the case that for every bipartite graph  $H$  without isolated vertices, integer  $k$ , and family of graphs  $\mathcal{F}$  that either

$$\text{ex}(n, H, \mathcal{F}) = \Omega(n^k),$$

or

$$\text{ex}(n, H, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

# Going Further

## Question

Is it the case that for every bipartite graph  $H$  without isolated vertices, integer  $k$ , and family of graphs  $\mathcal{F}$  that either

$$\text{ex}(n, H, \mathcal{F}) = \Omega(n^k),$$

or

$$\text{ex}(n, H, \mathcal{F}) = O(\text{ex}(n, \mathcal{F})^{k-1}).$$

## Fact

*This is false for  $C_4$  and  $k = 2$  :(*

## Going Further

We've seen a lot of ways that  $\text{ex}(n, H, \mathcal{F})$  *can't* behave like; what about the ways it *can* behave?



## Going Further

We've seen a lot of ways that  $\text{ex}(n, H, \mathcal{F})$  *can't* behave like; what about the ways it *can* behave?

### Definition

We say that a rational number  $r$  is *realizable* for a graph  $H$  if there exists a (finite) family of graphs  $\mathcal{F}$  such that  $\text{ex}(n, H, \mathcal{F}) = \Theta(n^r)$ .

# Going Further

We've seen a lot of ways that  $\text{ex}(n, H, \mathcal{F})$  *can't* behave like; what about the ways it *can* behave?

## Definition

We say that a rational number  $r$  is *realizable* for a graph  $H$  if there exists a (finite) family of graphs  $\mathcal{F}$  such that  $\text{ex}(n, H, \mathcal{F}) = \Theta(n^r)$ .

## Theorem (Bukh-Conlon 2018)

*Every rational in  $[1, 2]$  is realizable for  $H = K_2$ .*

## Going Further

Theorem (English-Halfpap-Krueger 2024)

*Every rational in  $[1, t]$  is realizable for  $H = K_t$ .*

# Going Further

## Theorem (English-Halfpap-Krueger 2024)

*Every rational in  $[1, t]$  is realizable for  $H = K_t$ .*

## Theorem (English-Halfpap-Krueger 2024)

*Every rational in  $[t, t + 1]$  is realizable for  $H = K_{1,t}$  and no rational in  $(1, t)$  is.*

# Going Further

## Theorem (English-S. 2025+)

*For every graph  $H$  of maximum degree  $\Delta$ , every rational in the range*

$$\left[ v(H) - \frac{e(H)}{2\Delta^2}, v(H) \right]$$

*is realizable.*

# Going Further

## Theorem (English-S. 2025+)

*For every graph  $H$  of maximum degree  $\Delta$ , every rational in the range*

$$\left[ v(H) - \frac{e(H)}{2\Delta^2}, v(H) \right]$$

*is realizable.*

## Theorem (English-S. 2025+)

*For every tree  $T$  of maximum degree  $\Delta$ , every rational in the range*

$$\left[ v(H) - \frac{e(H)}{\Delta}, v(H) \right]$$

*is realizable.*

This is best possible for stars.

# Open Problems

## Question

Given a graph  $H$ , what rationals  $r$  are realizable?

# Open Problems

## Question

Given a graph  $H$ , what rationals  $r$  are realizable?

From our general stability theorem, we know that if  $\text{ex}(n, H, \mathcal{F}_{H,k}^q) = o(n^\beta)$ , then no rational in  $(\beta, k)$  is realizable.



# Open Problems

## Question

Given a graph  $H$ , what rationals  $r$  are realizable?

From our general stability theorem, we know that if  $\text{ex}(n, H, \mathcal{F}_{H,k}^q) = o(n^\beta)$ , then no rational in  $(\beta, k)$  is realizable.

## Question

Is this the only way an exponent can fail to be rational?

# Open Problems

## Question

Given a graph  $H$ , what rationals  $r$  are realizable?

From our general stability theorem, we know that if  $\text{ex}(n, H, \mathcal{F}_{H,k}^q) = o(n^\beta)$ , then no rational in  $(\beta, k)$  is realizable.

## Question

Is this the only way an exponent can fail to be rational?

We believe we can verify that this is true for all graphs on at most 4 vertices through various ad hoc techniques.

# Open Problems

## Question

Is it the case that every rational in  $[\alpha(H), v(H)]$  is realizable?

This is true and best possible for cliques and stars.

## Conjecture

*For every graph  $H$ , every rational in*

$$\left[ v(H) - \frac{e(H)}{\Delta}, v(H) \right]$$

*is realizable.*

# Open Problems

Say that a rational  $r$  is strongly realizable if there exists a single graph  $F$  such that  $\text{ex}(n, H, F) = \Theta(n^r)$ .

# Open Problems

Say that a rational  $r$  is strongly realizable if there exists a single graph  $F$  such that  $\text{ex}(n, H, F) = \Theta(n^r)$ .

## Question

Is it the case that for all  $H$ , the number of strongly realizable exponents below  $\alpha(H)$  is  $O_H(1)$ ?

# Open Problems

Say that a rational  $r$  is strongly realizable if there exists a single graph  $F$  such that  $\text{ex}(n, H, F) = \Theta(n^r)$ .

## Question

Is it the case that for all  $H$ , the number of strongly realizable exponents below  $\alpha(H)$  is  $O_H(1)$ ?

The idea here is that any  $F$  of this form must be a subgraph of every member of  $\mathcal{F}_{H, \alpha(H)}^q \neq \emptyset$  for some  $q$ . For any fixed  $q$  this is a finite number of possibilities, and I don't think arbitrarily large  $q$  should give new behaviors (analogous to  $F \subseteq K_{2,t}$  implying  $\text{ex}(n, F) = \Theta(n^r)$  for some  $r \in \{0, 1, 3/2\}$ ).

# Open Problems

## Question

Can one say anything about generalized Turán numbers of trees  $T$  satisfying

$$n^{\ell+1} \ll \text{ex}(n, T, \mathcal{F}) \ll n^{\ell+2}?$$

For this it would suffice to prove upper bounds on  $\text{ex}(n, T, \mathcal{F}_{T, \ell+2}^q)$ .

# Open Problems

Our new Helly theorem for edge sets of trees turns out to imply the usual Helly theorem for vertex sets.

## Question

Does the usual vertex Helly Theorem imply the edge Helly Theorem?



# Open Problems

Our new Helly theorem for edge sets of trees turns out to imply the usual Helly theorem for vertex sets.

## Question

Does the usual vertex Helly Theorem imply the edge Helly Theorem? In particular, can you derive König's Theorem for trees?

# Open Problems

Our new Helly theorem for edge sets of trees turns out to imply the usual Helly theorem for vertex sets.

## Question

Does the usual vertex Helly Theorem imply the edge Helly Theorem? In particular, can you derive König's Theorem for trees?

I'm willing to offer up to \$1 for a proof\* or disproof of either of these questions.

# Open Problems: Supersaturation

# Open Problems: Supersaturation

## Proposition

*If  $G$  contains  $kn^{2-1/s}$  edges with  $k = \omega(1)$ , then it contains at least  $\Omega(k^{st}n^s)$  copies of  $K_{s,t}$ .*

Note that this is best possible by considering a random graph with  $kn^{2-1/s}$  edges.

# Open Problems: Supersaturation

## Proposition

*If  $G$  contains  $kn^{2-1/s}$  edges with  $k = \omega(1)$ , then it contains at least  $\Omega(k^{st}n^s)$  copies of  $K_{s,t}$ .*

Note that this is best possible by considering a random graph with  $kn^{2-1/s}$  edges.

## Question

Does the same phenomenon hold if  $G$  contains many copies of  $K_r$ ?

# Open Problems: Supersaturation

## Conjecture (Dubroff-Gunby-Narayanan-S.)

*If  $2 \leq r \leq s \leq t$  and if  $G$  contains at least  $kn^{r - \binom{r}{2}/s}$  copies of  $K_r$ , then it contains at least  $k^{st/\binom{r}{2}} n^{s - o(1)}$  copies of  $K_{s,t}$ .*

# Open Problems: Supersaturation

## Conjecture (Dubroff-Gunby-Narayanan-S.)

*If  $2 \leq r \leq s \leq t$  and if  $G$  contains at least  $kn^{r - \binom{r}{2}/s}$  copies of  $K_r$ , then it contains at least  $k^{st/\binom{r}{2}} n^{s - o(1)}$  copies of  $K_{s,t}$ .*

Things seem to behave differently if  $r > s$  and  $k$  is small.

# Open Problems: Supersaturation

## Conjecture (Dubroff-Gunby-Narayanan-S.)

*If  $2 \leq r \leq s \leq t$  and if  $G$  contains at least  $kn^{r - \binom{r}{2}/s}$  copies of  $K_r$ , then it contains at least  $k^{st/\binom{r}{2}} n^{s-o(1)}$  copies of  $K_{s,t}$ .*

Things seem to behave differently if  $r > s$  and  $k$  is small.

## Conjecture (Dubroff-Gunby-Narayanan-S.)

*For all  $1 \leq k \leq n^{1/2t}$ , there exist  $n$ -vertex graphs with  $kn^{3/2}$  triangles and at most  $k^t n^{3/2+o(1)}$  copies of  $K_{2,t}$ .*