

Card Guessing with Partial Feedback

Sam Spiro, UC San Diego.

Joint work with Persi Diaconis, Ron Graham, and Xiaoyu He

Feedback Models

Feedback Models

Start with a deck of mn cards where there are n card types each appearing with multiplicity m .

Feedback Models

Start with a deck of mn cards where there are n card types each appearing with multiplicity m . For example, $n = 13$ and $m = 4$ corresponds to a usual deck of playing cards.

Feedback Models

Start with a deck of mn cards where there are n card types each appearing with multiplicity m . For example, $n = 13$ and $m = 4$ corresponds to a usual deck of playing cards.

To begin the game, shuffle the deck uniformly at random.

Feedback Models

Start with a deck of mn cards where there are n card types each appearing with multiplicity m . For example, $n = 13$ and $m = 4$ corresponds to a usual deck of playing cards.

To begin the game, shuffle the deck uniformly at random. Each round a Guesser tries to guess what the next card in the deck is, and then the card is revealed and discarded, and we continue this way until the deck is depleted.

Feedback Models

Start with a deck of mn cards where there are n card types each appearing with multiplicity m . For example, $n = 13$ and $m = 4$ corresponds to a usual deck of playing cards.

To begin the game, shuffle the deck uniformly at random. Each round a Guesser tries to guess what the next card in the deck is, and then the card is revealed and discarded, and we continue this way until the deck is depleted. The score at the end of the game is the total number of correct guesses made during the mn rounds.

Feedback Models

If \mathbf{G} is a strategy for Guesser, we define $C_{m,n}(\mathbf{G})$ to be the expected number of points scored if Guesser follows strategy \mathbf{G} and the deck is shuffled uniformly at random.

Feedback Models

If \mathbf{G} is a strategy for Guesser, we define $C_{m,n}(\mathbf{G})$ to be the expected number of points scored if Guesser follows strategy \mathbf{G} and the deck is shuffled uniformly at random. We are particularly interested in $\mathcal{C}_{m,n}^+ = \max_{\mathbf{G}} C_{m,n}(\mathbf{G})$ and $\mathcal{C}_{m,n}^- = \min_{\mathbf{G}} C_{m,n}(\mathbf{G})$, as well as the strategies that achieve these quantities.

Feedback Models

If \mathbf{G} is a strategy for Guesser, we define $C_{m,n}(\mathbf{G})$ to be the expected number of points scored if Guesser follows strategy \mathbf{G} and the deck is shuffled uniformly at random. We are particularly interested in $C_{m,n}^+ = \max_{\mathbf{G}} C_{m,n}(\mathbf{G})$ and $C_{m,n}^- = \min_{\mathbf{G}} C_{m,n}(\mathbf{G})$, as well as the strategies that achieve these quantities.

Theorem (Diaconis-Graham, 1981)

The strategy G^\pm of guessing a most/least likely card at each stage achieves $C_{m,n}^\pm$.

Feedback Models

If \mathbf{G} is a strategy for Guesser, we define $C_{m,n}(\mathbf{G})$ to be the expected number of points scored if Guesser follows strategy \mathbf{G} and the deck is shuffled uniformly at random. We are particularly interested in $\mathcal{C}_{m,n}^+ = \max_{\mathbf{G}} C_{m,n}(\mathbf{G})$ and $\mathcal{C}_{m,n}^- = \min_{\mathbf{G}} C_{m,n}(\mathbf{G})$, as well as the strategies that achieve these quantities.

Theorem (Diaconis-Graham, 1981)

The strategy G^\pm of guessing a most/least likely card at each stage achieves $\mathcal{C}_{m,n}^\pm$. Moreover, for n fixed

$$\mathcal{C}_{m,n}^\pm = m \pm c_n \sqrt{m} + o_n(\sqrt{m}).$$

Feedback Models

Feedback Models

In the “no feedback model”, the Guesser guesses the next card which is discarded without being revealed to them.

Feedback Models

In the “no feedback model”, the Guesser guesses the next card which is discarded without being revealed to them. It is not hard to see that every strategy gives m points in expectation in this model.

Feedback Models

Feedback Models

In the “partial feedback model”, the Guesser guesses the next card and is only told whether their guess was correct or not.

Feedback Models

In the “partial feedback model”, the Guesser guesses the next card and is only told whether their guess was correct or not. The partial feedback model is significantly harder to analyze because we do not know what the optimal strategies are.

Feedback Models

Define the expected score under the partial feedback model by $P_{m,n}(\mathbf{G})$ and its optimal scores by $\mathcal{P}_{m,n}^+ = \max_{\mathbf{G}} P(\mathbf{G})$ and $\mathcal{P}_{m,n}^- = \min_{\mathbf{G}} P(\mathbf{G})$.

Feedback Models

Define the expected score under the partial feedback model by $P_{m,n}(\mathbf{G})$ and its optimal scores by $\mathcal{P}_{m,n}^+ = \max_{\mathbf{G}} P(\mathbf{G})$ and $\mathcal{P}_{m,n}^- = \min_{\mathbf{G}} P(\mathbf{G})$. We have

$$m \leq \mathcal{P}_{m,n}^+$$

Feedback Models

Define the expected score under the partial feedback model by $P_{m,n}(\mathbf{G})$ and its optimal scores by $\mathcal{P}_{m,n}^+ = \max_{\mathbf{G}} P(\mathbf{G})$ and $\mathcal{P}_{m,n}^- = \min_{\mathbf{G}} P(\mathbf{G})$. We have

$$m \leq \mathcal{P}_{m,n}^+ \leq \mathcal{C}_{m,n}^+$$

Feedback Models

Define the expected score under the partial feedback model by $P_{m,n}(\mathbf{G})$ and its optimal scores by $\mathcal{P}_{m,n}^+ = \max_{\mathbf{G}} P(\mathbf{G})$ and $\mathcal{P}_{m,n}^- = \min_{\mathbf{G}} P(\mathbf{G})$. We have

$$m \leq \mathcal{P}_{m,n}^+ \leq \mathcal{C}_{m,n}^+ = m + o_n(m)$$

Feedback Models

Define the expected score under the partial feedback model by $P_{m,n}(\mathbf{G})$ and its optimal scores by $\mathcal{P}_{m,n}^+ = \max_{\mathbf{G}} P(\mathbf{G})$ and $\mathcal{P}_{m,n}^- = \min_{\mathbf{G}} P(\mathbf{G})$. We have

$$m \leq \mathcal{P}_{m,n}^+ \leq \mathcal{C}_{m,n}^+ = m + o_n(m),$$

so for n fixed we have $\mathcal{P}_{m,n}^+ \sim m$, and similarly $\mathcal{P}_{m,n}^- \sim m$.

Feedback Models

Define the expected score under the partial feedback model by $P_{m,n}(\mathbf{G})$ and its optimal scores by $\mathcal{P}_{m,n}^+ = \max_{\mathbf{G}} P(\mathbf{G})$ and $\mathcal{P}_{m,n}^- = \min_{\mathbf{G}} P(\mathbf{G})$. We have

$$m \leq \mathcal{P}_{m,n}^+ \leq \mathcal{C}_{m,n}^+ = m + o_n(m),$$

so for n fixed we have $\mathcal{P}_{m,n}^+ \sim m$, and similarly $\mathcal{P}_{m,n}^- \sim m$. What happens when n is large?

Feedback Models

Theorem (Diaconis-Graham-He-S., 2020)

For m fixed,

$$\mathcal{C}_{m,n}^+ \sim H_m \log(n),$$

$$\mathcal{C}_{m,n}^- = \Theta(n^{-1/m}),$$

where H_m is the m th harmonic number.

Feedback Models

Theorem (Diaconis-Graham-He-S., 2020)

For m fixed,

$$\mathcal{C}_{m,n}^+ \sim H_m \log(n),$$

$$\mathcal{C}_{m,n}^- = \Theta(n^{-1/m}),$$

where H_m is the m th harmonic number.

With this we have the trivial bounds

$$m \leq \mathcal{P}_{m,n}^+ \leq \mathcal{C}_{m,n}^+ = O_m(\log n).$$

Feedback Models

$$m \leq \mathcal{P}_{m,n}^+ \leq O_m(\log n).$$

Feedback Models

$$m \leq \mathcal{P}_{m,n}^+ \leq O_m(\log n).$$

Theorem (Diaconis-Graham-He-S., 2020+)

There exist $c, C > 0$ such that if n is sufficiently large in terms of m , we have

$$m + c\sqrt{m} \leq \mathcal{P}_{m,n}^+$$

Feedback Models

$$m \leq \mathcal{P}_{m,n}^+ \leq O_m(\log n).$$

Theorem (Diaconis-Graham-He-S., 2020+)

There exist $c, C > 0$ such that if n is sufficiently large in terms of m , we have

$$m + c\sqrt{m} \leq \mathcal{P}_{m,n}^+ \leq m + Cm^{3/4} \log m.$$

Feedback Models

$$m \leq \mathcal{P}_{m,n}^+ \leq O_m(\log n).$$

Theorem (Diaconis-Graham-He-S., 2020+)

There exist $c, C > 0$ such that if n is sufficiently large in terms of m , we have

$$m + c\sqrt{m} \leq \mathcal{P}_{m,n}^+ \leq m + Cm^{3/4} \log m.$$

In particular, with partial information and fixed m , you can't get many points compared to having no feedback at all.

Feedback Models

$$m \leq \mathcal{P}_{m,n}^+ \leq O_m(\log n).$$

Theorem (Diaconis-Graham-He-S., 2020+)

There exist $c, C > 0$ such that if n is sufficiently large in terms of m , we have

$$m + c\sqrt{m} \leq \mathcal{P}_{m,n}^+ \leq m + Cm^{3/4} \log m.$$

In particular, with partial information and fixed m , you can't get many points compared to having no feedback at all. This is somewhat intuitive since saying “this card is not type i ” becomes less useful as the number of card types grow.

A Partial Proof

Lemma

Assume that we have played in the partial feedback model for $t - 1$ rounds such that we have guessed card type i a total of g_i times

A Partial Proof

Lemma

Assume that we have played in the partial feedback model for $t - 1$ rounds such that we have guessed card type i a total of g_i times, and let S be the total number of points scored.

A Partial Proof

Lemma

Assume that we have played in the partial feedback model for $t - 1$ rounds such that we have guessed card type i a total of g_i times, and let S be the total number of points scored. Given this, we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S}.$$

A Partial Proof

Lemma

Assume that we have played in the partial feedback model for $t - 1$ rounds such that we have guessed card type i a total of g_i times, and let S be the total number of points scored. Given this, we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S}.$$

That is, our upper bound is strongest when g_i and S is small.

A Partial Proof

Lemma

Assume that we have played in the partial feedback model for $t - 1$ rounds such that we have guessed card type i a total of g_i times, and let S be the total number of points scored. Given this, we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S}.$$

That is, our upper bound is strongest when g_i and S is small.
These conditions are necessary

A Partial Proof

Lemma

Assume that we have played in the partial feedback model for $t - 1$ rounds such that we have guessed card type i a total of g_i times, and let S be the total number of points scored. Given this, we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S}.$$

That is, our upper bound is strongest when g_i and S is small.

These conditions are necessary: if i has been guessed incorrectly $g_i = mn - m$ times, then we know the card must be an i

A Partial Proof

Lemma

Assume that we have played in the partial feedback model for $t - 1$ rounds such that we have guessed card type i a total of g_i times, and let S be the total number of points scored. Given this, we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S}.$$

That is, our upper bound is strongest when g_i and S is small. These conditions are necessary: if i has been guessed incorrectly $g_i = mn - m$ times, then we know the card must be an i , and similarly if the first $S = mn - m$ cards were guessed correctly and not equal to i , then the next card must be i .

A Partial Proof

Lemma

If we have guessed i a total of g_i times such that S points were scored, then

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S}.$$

A Partial Proof

Lemma

If we have guessed i a total of g_i times such that S points were scored, then

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S}.$$

Lemma

For any strategy G , with high probability the total number of points S is at most ϵmn .

A Partial Proof

Lemma

If we have guessed i a total of g_i times such that S points were scored, then

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S}.$$

Lemma

For any strategy G , with high probability the total number of points S is at most ϵmn .

Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

A Partial Proof

Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

A Partial Proof

Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all i and t , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S}$$

A Partial Proof

Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all i and t , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S} \approx \frac{m}{mn - g_i},$$

A Partial Proof

Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all i and t , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S} \approx \frac{m}{mn - g_i},$$

This bound is ineffective if g_i is large

A Partial Proof

Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all i and t , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S} \approx \frac{m}{mn - g_i},$$

This bound is ineffective if g_i is large, but at most one i is guessed more than $mn/2$ times (and this is guessed correctly at most m times).

A Partial Proof

Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all i and t , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S} \approx \frac{m}{mn - g_i},$$

This bound is ineffective if g_i is large, but at most one i is guessed more than $mn/2$ times (and this is guessed correctly at most m times). Every other j has $\Pr[\pi_t = j] \leq \frac{2}{n}$ for all t .

A Partial Proof

Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all i and t , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S} \approx \frac{m}{mn - g_i},$$

This bound is ineffective if g_i is large, but at most one i is guessed more than $mn/2$ times (and this is guessed correctly at most m times). Every other j has $\Pr[\pi_t = j] \leq \frac{2}{n}$ for all t . Thus in expectation at most $mn \cdot (2/n) = 2m$ cards are guessed correctly from this part

A Partial Proof

Corollary

$$\mathcal{P}_{m,n}^+ \leq 3m + o(m).$$

For all i and t , we have

$$\Pr[\pi_t = i] \leq \frac{m}{mn - g_i - S} \approx \frac{m}{mn - g_i},$$

This bound is ineffective if g_i is large, but at most one i is guessed more than $mn/2$ times (and this is guessed correctly at most m times). Every other j has $\Pr[\pi_t = j] \leq \frac{2}{n}$ for all t . Thus in expectation at most $mn \cdot (2/n) = 2m$ cards are guessed correctly from this part, and in total at most $3m$ are guessed correctly in expectation. □

Open Problems

Theorem (Diaconis-Graham-He-S., 2020)

$$m + \Omega(\sqrt{m}) \leq \mathcal{P}_{m,n}^+ \leq m + O(m^{3/4} \log m).$$

Open Problems

Theorem (Diaconis-Graham-He-S., 2020)

$$m + \Omega(\sqrt{m}) \leq \mathcal{P}_{m,n}^+ \leq m + O(m^{3/4} \log m).$$

Conjecture (Diaconis-Graham-He-S., 2020)

$$\mathcal{P}_{m,n}^+ = m + m^{1/2+o(1)}.$$

Open Problems

Theorem (Diaconis-Graham-He-S., 2020)

If m is a fixed constant,

$$\mathcal{P}_{m,n}^+ = \Theta_m(1),$$

$$\mathcal{C}_{m,n}^+ = \Theta_m(\log n).$$

Open Problems

Theorem (Diaconis-Graham-He-S., 2020)

If m is a fixed constant,

$$\mathcal{P}_{m,n}^+ = \Theta_m(1),$$

$$\mathcal{C}_{m,n}^+ = \Theta_m(\log n).$$

Question

Is there a reasonable choice of feedback so that the maximum expected score is, say, $\Theta_m(\log \log n)$?

Open Problems

Recall that $\mathcal{P}_{m,n}^-$ is the minimum number of points one can get in expectation with partial feedback.

Open Problems

Recall that $\mathcal{P}_{m,n}^-$ is the minimum number of points one can get in expectation with partial feedback. We trivially have

$$m \geq \mathcal{P}_{m,n}^- \geq \mathcal{C}_{m,n}^- = \Theta(n^{-1/m}).$$

Open Problems

Recall that $\mathcal{P}_{m,n}^-$ is the minimum number of points one can get in expectation with partial feedback. We trivially have

$$m \geq \mathcal{P}_{m,n}^- \geq \mathcal{C}_{m,n}^- = \Theta(n^{-1/m}).$$

Theorem (Diaconis-Graham-S., 2020)

$$\mathcal{P}_{m,n}^- \geq 1 - e^{-m} - o_m(1) \geq \frac{1}{2}.$$

Open Problems

Recall that $\mathcal{P}_{m,n}^-$ is the minimum number of points one can get in expectation with partial feedback. We trivially have

$$m \geq \mathcal{P}_{m,n}^- \geq \mathcal{C}_{m,n}^- = \Theta(n^{-1/m}).$$

Theorem (Diaconis-Graham-S., 2020)

$$\mathcal{P}_{m,n}^- \geq 1 - e^{-m} - o_m(1) \geq \frac{1}{2}.$$

This lower bound is simply the probability of guessing at least one card correctly (the best strategy for this is to just guess each card type exactly m times).

Open Problems

Theorem (Diaconis-Graham-S., 2020)

$$\mathcal{P}_{m,n}^- \geq 1 - e^{-m} - o_m(1) \geq \frac{1}{2}.$$

Open Problems

Theorem (Diaconis-Graham-S., 2020)

$$\mathcal{P}_{m,n}^- \geq 1 - e^{-m} - o_m(1) \geq \frac{1}{2}.$$

Conjecture (Diaconis-Graham-S., 2020)

If n is sufficiently large in terms of m , then

$$\mathcal{P}_{m,n}^- \sim m.$$

Increasing Subsequences

If π is a word where each letter in $\{1, 2, \dots, n\}$ exactly m times, we define $L(\pi)$ to be the largest integer p so that π contains a subsequence of the form $123 \cdots p$.

Increasing Subsequences

If π is a word where each letter in $\{1, 2, \dots, n\}$ exactly m times, we define $L(\pi)$ to be the largest integer p so that π contains a subsequence of the form $123 \cdots p$. For example,

$$\pi = 2345124351 \implies L(\pi) = 3.$$

Increasing Subsequences

If π is a word where each letter in $\{1, 2, \dots, n\}$ exactly m times, we define $L(\pi)$ to be the largest integer p so that π contains a subsequence of the form $123 \cdots p$. For example,

$$\pi = 2345124351 \implies L(\pi) = 3.$$

Let $\mathcal{L}_{m,n} = \mathbb{E}[L(\pi)]$ where π is chosen uniformly at random.

Increasing Subsequences

If π is a word where each letter in $\{1, 2, \dots, n\}$ exactly m times, we define $L(\pi)$ to be the largest integer p so that π contains a subsequence of the form $123 \cdots p$. For example,

$$\pi = 2345124351 \implies L(\pi) = 3.$$

Let $\mathcal{L}_{m,n} = \mathbb{E}[L(\pi)]$ where π is chosen uniformly at random.

Proposition

$$\mathcal{L}_{m,n} \leq \mathcal{P}_{m,n}^+.$$

Increasing Subsequences

Proposition

$$\mathcal{L}_{m,n} = \mathbb{E}[L(\pi)] \leq \mathcal{P}_{m,n}^+.$$

Increasing Subsequences

Proposition

$$\mathcal{L}_{m,n} = \mathbb{E}[L(\pi)] \leq \mathcal{P}_{m,n}^+.$$

Consider the following strategy in the partial feedback model: guess 1 until you guess one correct, then 2 until you guess one correct, then 3, and so on. After guessing n play arbitrarily.

Increasing Subsequences

Proposition

$$\mathcal{L}_{m,n} = \mathbb{E}[L(\pi)] \leq \mathcal{P}_{m,n}^+.$$

Consider the following strategy in the partial feedback model: guess 1 until you guess one correct, then 2 until you guess one correct, then 3, and so on. After guessing n play arbitrarily. If the deck is shuffled according to π , then the Guesser's score is at least $L(\pi)$.

Increasing Subsequences

Proposition

$$\mathcal{L}_{m,n} = \mathbb{E}[L(\pi)] \leq \mathcal{P}_{m,n}^+.$$

Consider the following strategy in the partial feedback model: guess 1 until you guess one correct, then 2 until you guess one correct, then 3, and so on. After guessing n play arbitrarily. If the deck is shuffled according to π , then the Guesser's score is at least $L(\pi)$. Thus,

$$\mathbb{E}[L(\pi)] \leq \mathcal{P}_{m,n}^+.$$

Increasing Subsequences

Corollary

If n is sufficiently large in terms of m ,

$$\mathcal{L}_{m,n} \leq m + O(m^{3/4} \log m).$$

Increasing Subsequences

Corollary

If n is sufficiently large in terms of m ,

$$\mathcal{L}_{m,n} \leq m + O(m^{3/4} \log m).$$

Theorem (Clifton-Deb-Huang-S.-Yoo)

We have

$$\left| \lim_{n \rightarrow \infty} \mathcal{L}_{m,n} - \left(m + 1 - \frac{1}{m+2} \right) \right| \leq O(e^{-\beta m})$$

for some $\beta > 0$.

Increasing Subsequences

Corollary

If n is sufficiently large in terms of m ,

$$\mathcal{L}_{m,n} \leq m + O(m^{3/4} \log m).$$

Theorem (Clifton-Deb-Huang-S.-Yoo)

We have

$$\left| \lim_{n \rightarrow \infty} \mathcal{L}_{m,n} - \left(m + 1 - \frac{1}{m+2} \right) \right| \leq O(e^{-\beta m})$$

for some $\beta > 0$.

In particular, the strategy “guess 1 until you get one correct, then 2,...” does not give much more than the trivial bound.

Increasing Subsequences

More precisely: if $\alpha_1, \dots, \alpha_m$ are the zeroes of $\sum_{k=0}^m \frac{x^k}{k!}$, then

$$\lim_{n \rightarrow \infty} \mathcal{L}_{m,n} = -1 - \sum \alpha_i^{-1} e^{-\alpha_i}.$$

Increasing Subsequences

More precisely: if $\alpha_1, \dots, \alpha_m$ are the zeroes of $\sum_{k=0}^m \frac{x^k}{k!}$, then

$$\lim_{n \rightarrow \infty} \mathcal{L}_{m,n} = -1 - \sum \alpha_i^{-1} e^{-\alpha_i}.$$

This implies $\mathcal{L}_{1,n} \rightarrow e - 1$

Increasing Subsequences

More precisely: if $\alpha_1, \dots, \alpha_m$ are the zeroes of $\sum_{k=0}^m \frac{x^k}{k!}$, then

$$\lim_{n \rightarrow \infty} \mathcal{L}_{m,n} = -1 - \sum \alpha_i^{-1} e^{-\alpha_i}.$$

This implies $\mathcal{L}_{1,n} \rightarrow e - 1$ and that

$$\mathcal{L}_{2,n} \rightarrow e(\cos(1) + \sin(1)) - 1.$$

Card Guessing

Card Guessing

The probability of drawing four aces in a row with a deck shuffled uniformly at random is $1/270725$.

Card Guessing

The probability of drawing four aces in a row with a deck shuffled uniformly at random is $1/270725$. We can thus conclude that either you were really unlucky

Card Guessing

The probability of drawing four aces in a row with a deck shuffled uniformly at random is $1/270725$. We can thus conclude that either you were really unlucky, or the deck was not shuffled uniformly at random.

Online Card Guessing

The probability of drawing four aces in a row with a deck shuffled uniformly at random is $1/270725$. We can thus conclude that either you were really unlucky, or the deck was not shuffled uniformly at random.

Online Card Guessing

More precisely, we are now considering a two player game played by Shuffler and Guesser.

Online Card Guessing

More precisely, we are now considering a two player game played by Shuffler and Guesser. First Shuffler chooses some ordering π for the deck according to a distribution/strategy **S**

Online Card Guessing

More precisely, we are now considering a two player game played by Shuffler and Guesser. First Shuffler chooses some ordering π for the deck according to a distribution/strategy **S**, and then Guesser plays the complete feedback model with π according to some strategy **G**.

Online Card Guessing

More precisely, we are now considering a two player game played by Shuffler and Guesser. First Shuffler chooses some ordering π for the deck according to a distribution/strategy **S**, and then Guesser plays the complete feedback model with π according to some strategy **G**.

Mathematically it's equivalent to have Shuffler iteratively choose each subsequent card in an online fashion

Online Card Guessing

More precisely, we are now considering a two player game played by Shuffler and Guesser. First Shuffler chooses some ordering π for the deck according to a distribution/strategy **S**, and then Guesser plays the complete feedback model with π according to some strategy **G**.

Mathematically it's equivalent to have Shuffler iteratively choose each subsequent card in an online fashion (but your friends may disagree if you try this in real life).

Online Card Guessing

In any case, we let $\mathcal{C}_{m,n}(\mathbf{G}, \mathbf{S})$ be the expected number of points Guesser scores when the two players follow strategies \mathbf{G}, \mathbf{S} .

Online Card Guessing

In any case, we let $\mathcal{C}_{m,n}(\mathbf{G}, \mathbf{S})$ be the expected number of points Guesser scores when the two players follow strategies \mathbf{G}, \mathbf{S} .

$$\Theta_m(n^{-1/m}) \leq \mathcal{C}_{m,n}(\mathbf{G}, \text{Uniform}) \leq H_m \log n + o_m(\log n).$$

Online Card Guessing

In any case, we let $\mathcal{C}_{m,n}(\mathbf{G}, \mathbf{S})$ be the expected number of points Guesser scores when the two players follow strategies \mathbf{G}, \mathbf{S} .

$$\Theta_m(n^{-1/m}) \leq \mathcal{C}_{m,n}(\mathbf{G}, \text{Uniform}) \leq H_m \log n + o_m(\log n).$$

Theorem (S., 2021+)

If Shuffler wants to minimize the number of correct guesses and Guesser wants to maximize this, then under their optimal strategies G', S' we have

$$\mathcal{C}_{m,n}(G', S') = \log n + o_m(\log n).$$

Online Card Guessing

In any case, we let $\mathcal{C}_{m,n}(\mathbf{G}, \mathbf{S})$ be the expected number of points Guesser scores when the two players follow strategies \mathbf{G}, \mathbf{S} .

$$\Theta_m(n^{-1/m}) \leq \mathcal{C}_{m,n}(\mathbf{G}, \text{Uniform}) \leq H_m \log n + o_m(\log n).$$

Theorem (S., 2021+)

If Shuffler wants to minimize the number of correct guesses and Guesser wants to maximize this, then under their optimal strategies G', S' we have

$$\mathcal{C}_{m,n}(G', S') = \log n + o_m(\log n).$$

This theorem is a first for me, since normally I prove a result, then makes jokes about it during my talk.

Online Card Guessing

Theorem

There exists a strategy S' for Shuffler so that
$$\mathcal{C}_{m,n}(G, S') \leq \log n + o_m(\log n).$$

Online Card Guessing

Theorem

There exists a strategy S' for Shuffler so that
$$\mathcal{C}_{m,n}(G, S') \leq \log n + o_m(\log n).$$

A strategy that gives this is the “greedy strategy”, which is such that if there are r types of cards remaining in the deck, then Shuffler draws each of these card types with probability r^{-1} (regardless of how many copies are left in the deck of each type).

Online Card Guessing

Theorem

There exists a strategy S' for Shuffler so that
$$C_{m,n}(G, S') \leq \log n + o_m(\log n).$$

A strategy that gives this is the “greedy strategy”, which is such that if there are r types of cards remaining in the deck, then Shuffler draws each of these card types with probability r^{-1} (regardless of how many copies are left in the deck of each type). E.g. if the deck has a hundred 1's and one 2, we draw a 1 or 2 with probability $\frac{1}{2}$.

Online Card Guessing

Theorem

There exists a strategy S' for Shuffler so that
$$C_{m,n}(G, S') \leq \log n + o_m(\log n).$$

A strategy that gives this is the “greedy strategy”, which is such that if there are r types of cards remaining in the deck, then Shuffler draws each of these card types with probability r^{-1} (regardless of how many copies are left in the deck of each type). E.g. if the deck has a hundred 1's and one 2, we draw a 1 or 2 with probability $\frac{1}{2}$. This gives the desired bound due to a variant of the coupon collector problem. □

Online Card Guessing

Theorem (S., 2021+)

The greedy strategy is the unique strategy that minimizes the number of correct guesses if Guesser tries to maximize their score.

Online Card Guessing

Theorem (S., 2021+)

The greedy strategy is the unique strategy that minimizes the number of correct guesses if Guesser tries to maximize their score.

Interestingly, the greedy strategy is also the “unique” strategy which maximizes the number of correct guesses if Guesser tries to minimize their score.

Online Card Guessing

There is a classical game called “Matching Pennies” where two players simultaneously choose one of n numbers, and if the two match player A gets a point and otherwise player B gets a point.

Online Card Guessing

There is a classical game called “Matching Pennies” where two players simultaneously choose one of n numbers, and if the two match player A gets a point and otherwise player B gets a point.

The online card guessing game can be viewed as a “semi-restricted” version of this game where mn rounds of Matching Pennies is played and player B must use each number exactly m times.

Online Card Guessing

There is a classical game called “Matching Pennies” where two players simultaneously choose one of n numbers, and if the two match player A gets a point and otherwise player B gets a point.

The online card guessing game can be viewed as a “semi-restricted” version of this game where mn rounds of Matching Pennies is played and player B must use each number exactly m times.

Question

Consider a “semi-restricted” version of Rock Paper Scissors: $3m$ rounds of the game is played but one of the players must use each move exactly m times. What are the optimal strategies/scores in this game?

Online Card Guessing

One can also consider a “balanced” version of this game where Guesser must guess each card type at most m times.

Online Card Guessing

One can also consider a “balanced” version of this game where Guesser must guess each card type at most m times. Equivalently one can give both players a deck and have each of them simultaneously draw from their respective decks.

Online Card Guessing

One can also consider a “balanced” version of this game where Guesser must guess each card type at most m times. Equivalently one can give both players a deck and have each of them simultaneously draw from their respective decks.

Theorem (S., 2021+)

In this situation, an optimal strategy for both players is to draw cards uniformly at random, which gives m matches in expectation.

Online Card Guessing

One can also consider a “balanced” version of this game where Guesser must guess each card type at most m times. Equivalently one can give both players a deck and have each of them simultaneously draw from their respective decks.

Theorem (S., 2021+)

In this situation, an optimal strategy for both players is to draw cards uniformly at random, which gives m matches in expectation.

If Guesser can guess each card type at most mn times, then this is equivalent to having no restrictions and they can score as many as $\log n + o_m(\log n)$ points.

Online Card Guessing

One can also consider a “balanced” version of this game where Guesser must guess each card type at most m times. Equivalently one can give both players a deck and have each of them simultaneously draw from their respective decks.

Theorem (S., 2021+)

In this situation, an optimal strategy for both players is to draw cards uniformly at random, which gives m matches in expectation.

If Guesser can guess each card type at most mn times, then this is equivalent to having no restrictions and they can score as many as $\log n + o_m(\log n)$ points.

Question

What happens if Guesser can guess each card type at most k times?

The End

Thank You!