# automatically removing high missing patients or features

### wei zou

#### 2022-02-10 09:55:01

### using Jaccard distance

```
c1 <- clusterMiss(ds1, dis_measure = list('row'='jaccard', 'col'='jaccard'), numCluster = c('row'=2, 'c

## raw input dimension: 126 520

## dimension after removing full missing cols/rows: 126 484

## matrix used for clustering: 115 438

## using jaccard, jaccard distance

## Metric: 'jaccard'; comparing: 115 vectors.

## Metric: 'jaccard'; comparing: 438 vectors.

hclust of missingness (>0: not missing)


0.4

0.2

0.2
```

-0.2

-0.4



```
1
## 113
## [1] "column groups:"
## ft_groups
        2
            3
## 357
        7 56 18
## [1] "summary of percent non-missing values among excluded items in the original matrix"
## G028915-264866-401004 G028915-266618-423025
##
              0.6198347
                                    0.5061983
## summary for 81 items
     Min. 1st Qu. Median
                             Mean 3rd Qu.
                                             Max.
## 0.4683 0.5159 0.5794 0.5684 0.5873 0.7619
## [1] "summary of percent non-missing values after removing patients and samples with high missing"
## summary for 124 items
     Min. 1st Qu. Median
                             Mean 3rd Qu.
                                             Max.
  0.8065 0.9404 0.9851 0.9648 1.0000 1.0000
## summary for 403 items
     Min. 1st Qu. Median
                             Mean 3rd Qu.
## 0.7823 0.9516 0.9839 0.9648 0.9919 1.0000
```

## using eucledian distance

```
this is an option; but not the default setting
```


```
c1 <- clusterMiss(ds1, dis_measure = list('row'='euclidean', 'col'='euclidean'), numCluster = c('row'=2)

## raw input dimension: 126 520

## dimension after removing full missing cols/rows: 126 484

## matrix used for clustering: 115 438

## using euclidean, euclidean distance
```



```
## [1] "row groups:"
## pt_groups
##
   1
## 113
## [1] "column groups:"
## ft_groups
   1
        2
        7 56 18
## 357
## [1] "summary of percent non-missing values among excluded items in the original matrix"
## G028915-264866-401004 G028915-266618-423025
##
              0.6198347
                                    0.5061983
## summary for 81 items
##
     Min. 1st Qu. Median
                             Mean 3rd Qu.
  0.4683 0.5159 0.5794 0.5684 0.5873 0.7619
## [1] "summary of percent non-missing values after removing patients and samples with high missing"
## summary for 124 items
     Min. 1st Qu. Median
                             Mean 3rd Qu.
##
                                             Max.
  0.8065 0.9404 0.9851 0.9648 1.0000 1.0000
## summary for 403 items
     Min. 1st Qu. Median
                             Mean 3rd Qu.
##
## 0.7823 0.9516 0.9839 0.9648 0.9919 1.0000
```