

Selecting Genetic Variants Using Knockoff Under Collider Bias

Qian Zhao

Postdoctoral Scholar, Biomedical Data Science



Stanford | Biomedical Data Science
SCHOOL OF MEDICINE

A motivating question

- Severe mental illnesses overlap in symptoms and share some genetic risks

A motivating question

- Severe mental illnesses overlap in symptoms and share some genetic risks

What are common and distinct pathways of severe mental illnesses?

A motivating question

- Severe mental illnesses overlap in symptoms and share some genetic risks
- Leveraging genetic variants can identify biological pathways and thus help psychiatrists define subtypes in a more biologically sound way

What are common and distinct pathways of severe mental illnesses?

A motivating question

- Severe mental illnesses overlap in symptoms and share some genetic risks
- Leveraging genetic variants can identify biological pathways and thus help psychiatrists define subtypes in a more biologically sound way

What are common and distinct pathways of severe mental illnesses?



Which genetic variants are associated with an **endophenotype**?

A motivating question

What are common and distinct pathways of severe mental illnesses?



Which genetic variants are associated with an **endophenotype**?

A motivating question




What are common and distinct pathways of severe mental illnesses?



Which genetic variants are associated with an **endophenotype**?

A motivating question

Genotype




What are common and distinct pathways of severe mental illnesses?



Which genetic variants are associated with an **endophenotype**?

A motivating question

Genotype



Symptoms
& Diagnosis




What are common and distinct pathways of severe mental illnesses?



Which genetic variants are associated with an **endophenotype**?

A motivating question

Genotype



Symptoms
& Diagnosis



Neurocognitive
function




What are common and distinct pathways of severe mental illnesses?



Which genetic variants are associated with an **endophenotype**?

A motivating question

Genotype



Symptoms
& Diagnosis



Neurocognitive
function



- Demographics
- Medication
- Substance use
- ...

What are common and distinct pathways of severe mental illnesses?




Which genetic variants are associated with an **endophenotype**?

Case-control study design

Case-control study design


Cases

Controls




Population (P)


Case-control study design




Case-control study design



Case-control study design



Case-control study design




Example: Is academic ability associated with athletic ability?



Academic Ability
(X)

Athletic Ability
(Y)


Population $P(X, Y)$
(Red + Black points)




Example: Is academic ability associated with athletic ability?




Collider Bias



Case-control study




Collider Bias




Can we test if X affects Y in the population (P) using case case-control sample (S)?

Case-control study




Collider Bias




Can we test if X affects Y in the population (P) using case case-control sample (S)?

Case-control study



Current approaches usually do not apply when the number of variables is large!

A knockoff approach to select variables in the high-dimensional setting



- Single nucleotide polymorphisms (SNP) are alterations in a single nucleotide (A, T, G, C) in the DNA sequence.

- Denote the SNPs as

$$(X_1, X_2, \dots, X_p)$$

- Denote the endophenotype as Y .
- Identify important SNPs by testing

$$\mathcal{H}_j : X_j \perp\!\!\!\perp Y \mid X_{-j}$$

The knockoff method

The knockoff method

- The knockoff method is a variable selection procedure that controls the false discovery rate

The knockoff method

- The knockoff method is a variable selection procedure that controls the false discovery rate
- For every observation (X_1, X_2, \dots, X_p)

We construct **knockoff variables** $(\tilde{X}_1, \dots, \tilde{X}_p)$

Such that

$$(X_1, \dots, X_p, \tilde{X}_1, \dots, \tilde{X}_p)_{\text{Swap } S} \stackrel{d}{=} (X_1, \dots, X_p, \tilde{X}_1, \dots, \tilde{X}_p)$$

For every set S containing only **null** variables

The knockoff method

	$(X_1, X_2, X_3, X_4, X_5)$	$(\tilde{X}_1, \tilde{X}_2, \tilde{X}_3, \tilde{X}_4, \tilde{X}_5)$										
Sample 1	<table><tr><td>A</td><td>T</td><td>G</td><td>G</td><td>C</td></tr></table>	A	T	G	G	C	<table><tr><td>T</td><td>T</td><td>A</td><td>G</td><td>C</td></tr></table>	T	T	A	G	C
A	T	G	G	C								
T	T	A	G	C								
Sample 2	<table><tr><td>A</td><td>T</td><td>A</td><td>G</td><td>C</td></tr></table>	A	T	A	G	C	<table><tr><td>A</td><td>T</td><td>G</td><td>G</td><td>A</td></tr></table>	A	T	G	G	A
A	T	A	G	C								
A	T	G	G	A								
Sample 3	<table><tr><td>T</td><td>T</td><td>G</td><td>G</td><td>A</td></tr></table>	T	T	G	G	A	<table><tr><td>T</td><td>T</td><td>A</td><td>G</td><td>A</td></tr></table>	T	T	A	G	A
T	T	G	G	A								
T	T	A	G	A								
Sample 4	<table><tr><td>A</td><td>T</td><td>A</td><td>T</td><td>C</td></tr></table>	A	T	A	T	C	<table><tr><td>A</td><td>C</td><td>A</td><td>T</td><td>C</td></tr></table>	A	C	A	T	C
A	T	A	T	C								
A	C	A	T	C								

The knockoff method

	$(X_1, X_2, X_3, X_4, X_5)$	$(\tilde{X}_1, \tilde{X}_2, \tilde{X}_3, \tilde{X}_4, \tilde{X}_5)$
Sample 1	<div>A</div> <div>T</div> <div>G</div> <div>G</div> <div>C</div>	<div>T</div> <div>T</div> <div>A</div> <div>G</div> <div>C</div>
Sample 2	<div>A</div> <div>T</div> <div>A</div> <div>G</div> <div>C</div>	<div>A</div> <div>T</div> <div>G</div> <div>G</div> <div>A</div>
Sample 3	<div>T</div> <div>T</div> <div>G</div> <div>G</div> <div>A</div>	<div>T</div> <div>T</div> <div>A</div> <div>G</div> <div>A</div>
Sample 4	<div>A</div> <div>T</div> <div>A</div> <div>T</div> <div>C</div>	<div>A</div> <div>C</div> <div>A</div> <div>T</div> <div>C</div>



We should not be able to tell
between a **null** variable and
its knockoff!

The knockoff method

	$(X_1, X_2, X_3, X_4, X_5)$					$(\tilde{X}_1, \tilde{X}_2, \tilde{X}_3, \tilde{X}_4, \tilde{X}_5)$				
Sample 1	A	T	G	G	C	T	T	A	G	C
Sample 2	A	T	A	G	C	A	T	G	G	A
Sample 3	T	T	G	G	A	T	T	A	G	A
Sample 4	A	T	A	T	C	A	C	A	T	C

Feature importance statistics

$(Z_1, Z_2, \dots, Z_5, \tilde{Z}_1, \tilde{Z}_2, \dots, \tilde{Z}_5)$

E.g. LASSO regression coefficients



We should not be able to tell
between a **null** variable and
its knockoff!

The knockoff method

	$(X_1, X_2, X_3, X_4, X_5)$	$(\tilde{X}_1, \tilde{X}_2, \tilde{X}_3, \tilde{X}_4, \tilde{X}_5)$										
Sample 1	<table><tr><td>A</td><td>T</td><td>G</td><td>G</td><td>C</td></tr></table>	A	T	G	G	C	<table><tr><td>T</td><td>T</td><td>A</td><td>G</td><td>C</td></tr></table>	T	T	A	G	C
A	T	G	G	C								
T	T	A	G	C								
Sample 2	<table><tr><td>A</td><td>T</td><td>A</td><td>G</td><td>C</td></tr></table>	A	T	A	G	C	<table><tr><td>A</td><td>T</td><td>G</td><td>G</td><td>A</td></tr></table>	A	T	G	G	A
A	T	A	G	C								
A	T	G	G	A								
Sample 3	<table><tr><td>T</td><td>T</td><td>G</td><td>G</td><td>A</td></tr></table>	T	T	G	G	A	<table><tr><td>T</td><td>T</td><td>A</td><td>G</td><td>A</td></tr></table>	T	T	A	G	A
T	T	G	G	A								
T	T	A	G	A								
Sample 4	<table><tr><td>A</td><td>T</td><td>A</td><td>T</td><td>C</td></tr></table>	A	T	A	T	C	<table><tr><td>A</td><td>C</td><td>A</td><td>T</td><td>C</td></tr></table>	A	C	A	T	C
A	T	A	T	C								
A	C	A	T	C								



We should not be able to tell
between a **null** variable and
its knockoff!

Feature importance statistics

$(Z_1, Z_2, \dots, Z_5, \tilde{Z}_1, \tilde{Z}_2, \dots, \tilde{Z}_5)$

E.g. LASSO regression coefficients

Scores

$(W_1, W_2, W_3, W_4, W_5)$

$$W_j = f(Z_j, \tilde{Z}_j) = -f(\tilde{Z}_j, Z_j)$$

The knockoff method

	$(X_1, X_2, X_3, X_4, X_5)$					$(\tilde{X}_1, \tilde{X}_2, \tilde{X}_3, \tilde{X}_4, \tilde{X}_5)$				
Sample 1	A	T	G	G	C	T	T	A	G	C
Sample 2	A	T	A	G	C	A	T	G	G	A
Sample 3	T	T	G	G	A	T	T	A	G	A
Sample 4	A	T	A	T	C	A	C	A	T	C



We should not be able to tell between a variable and its knockoff!

Feature importance statistics

$(Z_1, Z_2, \dots, Z_5, \tilde{Z}_1, \tilde{Z}_2, \dots, \tilde{Z}_5)$

E.g. LASSO regression coefficients

Scores

$(W_1, W_2, W_3, W_4, W_5)$

$$W_j = f(Z_j, \tilde{Z}_j) = -f(\tilde{Z}_j, Z_j)$$

A list of selected variables

How do we construct knockoff variables?

$$(X_j, X_{-j}, \tilde{X}_j, \tilde{X}_{-j}, Y) \stackrel{d}{=} (\tilde{X}_j, X_{-j}, X_j, \tilde{X}_{-j}, Y)$$

How do we construct knockoff variables?

$$(X_j, X_{-j}, \tilde{X}_j, \tilde{X}_{-j}, Y) \stackrel{d}{=} (\tilde{X}_j, X_{-j}, X_j, \tilde{X}_{-j}, Y)$$



$$\mathcal{L}(X_j, X_{-j}, \tilde{X}_j, \tilde{X}_{-j}, Y) = P(X_j, X_{-j}, Y | S = 1) Q(\tilde{X}_j, \tilde{X}_{-j} | X_j, X_{-j}, Y)$$

How do we construct knockoff variables?

$$(X_j, X_{-j}, \tilde{X}_j, \tilde{X}_{-j}, Y) \stackrel{d}{=} (\tilde{X}_j, X_{-j}, X_j, \tilde{X}_{-j}, Y)$$







$$\mathcal{L}(X_j, X_{-j}, \tilde{X}_j, \tilde{X}_{-j}, Y) = \mathbb{P}(X_j, X_{-j}, Y | S = 1) Q(\tilde{X}_j, \tilde{X}_{-j} | X_j, X_{-j}, Y)$$



User defined!

How do we construct knockoff variables?

$$(X_j, X_{-j}, \tilde{X}_j, \tilde{X}_{-j}, Y) \stackrel{d}{=} (\tilde{X}_j, X_{-j}, X_j, \tilde{X}_{-j}, Y)$$




$$\mathcal{L}(X_j, X_{-j}, \tilde{X}_j, \tilde{X}_{-j}, Y) = P(X_j, X_{-j}, Y | S = 1) Q(\tilde{X}_j, \tilde{X}_{-j} | X_j, X_{-j}, Y)$$


We don't know what is
 $P(X, Y)$!

User defined!

How do we construct knockoff variables?

$$(X_j, X_{-j}, \tilde{X}_j, \tilde{X}_{-j}, Y) \stackrel{d}{=} (\tilde{X}_j, X_{-j}, X_j, \tilde{X}_{-j}, Y)$$


$$\mathcal{L}(X_j, X_{-j}, \tilde{X}_j, \tilde{X}_{-j}, Y) = \boxed{P(X_j, X_{-j}, Y | S = 1)} Q(\tilde{X}_j, \tilde{X}_{-j} | X_j, X_{-j}, Y)$$


We don't know what is
 $P(X, Y)$!

How do we construct knockoff variables?

$$(X_j, X_{-j}, \tilde{X}_j, \tilde{X}_{-j}, Y) \stackrel{d}{=} (\tilde{X}_j, X_{-j}, X_j, \tilde{X}_{-j}, Y)$$



$$\mathcal{L}(X_j, X_{-j}, \tilde{X}_j, \tilde{X}_{-j}, Y) = \boxed{P(X_j, X_{-j}, Y | S = 1)} Q(\tilde{X}_j, \tilde{X}_{-j} | X_j, X_{-j}, Y)$$



$$P(X_j, X_{-j}) P(S = 1 | X_j, X_{-j}, Y)$$

We can use this distribution instead!

(Selection probability)

How do we construct knockoff variables?

$$(X_j, X_{-j}, \tilde{X}_j, \tilde{X}_{-j}, Y) \stackrel{d}{=} (\tilde{X}_j, X_{-j}, X_j, \tilde{X}_{-j}, Y)$$



$$\mathcal{L}(X_j, X_{-j}, \tilde{X}_j, \tilde{X}_{-j}, Y) = P(X_j, X_{-j}, Y | S = 1) Q(\tilde{X}_j, \tilde{X}_{-j} | X_j, X_{-j}, Y)$$



$$P(X_j, X_{-j}) P(S = 1 | X_j, X_{-j}, Y)$$


We can use this distribution instead!

(Selection probability)



$$\propto P(X_j | X_{-j}, Y, S = 1)$$

How do we construct knockoff variables?



If

$$P(X_j, X_{-j}) P(S = 1 | X_j, X_{-j}, Y)$$

Is multivariate Gaussian, then we can sample knockoffs from another multivariate Gaussian.

Number of cases = Number of controls = 1,000; Number of variables = 200; X is from a multivariate Gaussian distribution where the covariance matrix is block diagonal (block size = 10); Y is from a linear model where 10% of the variables are non-nulls;


$$P(D = 1 | X, Y) = e^{-v^2/2}, \quad v = \gamma_0 Y + X^\top \gamma.$$

Conclusion and remaining challenges

- Collider bias can occur in many studies!
- We describe one way to adjust the knockoff sampling procedure to select variables that controls the false discovery rate (1) in high-dimensions (2) under collider bias


Conclusion and remaining challenges

- Collider bias can occur in many studies!
- We describe one way to adjust the knockoff sampling procedure to select variables that controls the false discovery rate (1) in high-dimensions (2) under collider bias
- It's unclear how to define $Q(\tilde{X}_j, \tilde{X}_{-j} | X_j, X_{-j}, Y)$



Conclusion and remaining challenges

- Collider bias can occur in many studies!
- We describe one way to adjust the knockoff sampling procedure to select variables that controls the false discovery rate (1) in high-dimensions (2) under collider bias
- It's unclear how to define $Q(\tilde{X}_j, \tilde{X}_{-j} | X_j, X_{-j}, Y)$
- One idea: approximate $P(X_j, X_{-j}) P(S = 1 | X_j, X_{-j}, Y)$ by another distribution for which we know Q



Thank you! Questions?