

Discrete Mathematics: Lecture 27

Shortest Paths and Dijkstra's Algorithm, Traveling Salesperson Problem, Planar Graph, Euler's Formula, Kuratowski's Theorem

Xuming He

Associate Professor

School of Information Science and Technology
ShanghaiTech University

Spring Semester, 2022


Notes by Prof. Liangfeng Zhang

Shortest Path Problem

Definition

A **weighted graph** is a graph $G = (V, E)$ such that each edge is assigned with a strictly positive number.


The **length** of a path in weighted graph is the sum of the weights of the edges of this path.



a, b, c is a path of length 7 and b, e, d, c is a path of length 6


Remark: Observe that in a non-weighted graph the length of a path is the number of edges in the path!

Dijkstra's Algorithm



- 1** Find the closest vertex to $a \rightsquigarrow$ analyse all the edges starting from a :
 a, b of length 4
 a, f of length 2
 $\Rightarrow f$ is the closest vertex to a . The shortest path from a to f has length 2.
- 2** Find the second closest vertex to $a \rightsquigarrow$ shortest paths from a to a vertex in $\{a, f\}$ followed by an edge from a vertex in $\{a, f\}$ to a vertex not in this set:
 a, b of length 4
 a, f, e of length 5
 $\Rightarrow b$ is the second closest vertex to a . The shortest path from a to b has length 4.

Dijkstra's Algorithm



- 3** Find the third closest vertex to $a \rightsquigarrow$ shortest path from a to a vertex in $\{a, f, b\}$ followed by an edge from a vertex in $\{a, f, b\}$ to a vertex not in this set:

a, b, c of length 7

a, b, e of length 7

a, f, e of length 5

$\Rightarrow e$ is the third closest vertex to a . The shortest path from a to e has length 5.

- 4** Find the fourth closest vertex to $a \rightsquigarrow$ shortest path from a to a vertex in $\{a, f, b, e\}$ followed by an edge from a vertex in $\{a, f, b, e\}$ to a vertex not in this set:

a, b, c of length 7

a, f, e, d of length 6

$\Rightarrow d$ is the fourth closest vertex to a . The shortest path from a to d has length 6.

Dijkstra's Algorithm

Goal: find the length of a shortest path from a to z with a series of iterations.

- A distinguished set of vertices is constructed by adding one vertex at each iteration.
- A labeling procedure is carried out at each iteration: a vertex w is labeled with the length of a shortest path from a to w that contains only vertices in the distinguished set.
- The vertex added to the distinguished set is one with minimal label among those vertices not already in the set.


Notations: $S_k :=$ distinguished set after k iterations, $L_k(v) :=$ length of a shortest path from a to v containing only vertices in S_k ("label" of v).

Initialization: $L_0(a) = 0,$
 $L_0(v) = \infty$ for every vertex $v \neq a,$
 $S_0 = \emptyset.$

k th iteration:


- S_k is formed from S_{k-1} by adding a vertex u not in S_{k-1} with smallest label,
- Update the labels of all vertices not in S_k so that $L_k(v)$ is the length of a shortest path from a to v containing only vertices in S_k , i.e.
 $L_k(v) = \min\{L_{k-1}(v), L_{k-1}(u) + w(u, v)\}$ (with $w(u, v)$ length of the edge (u, v))

Dijkstra's Algorithm



- **k=0 (initialization):** $S_0 = \emptyset$,
 $L_0(a) = 0$, $L_0(b) = L_0(c) =$
 $L_0(d) = L_0(e) = L_0(z) = \infty$


Dijkstra's Algorithm



- **k=0 (initialization):** $S_0 = \emptyset$,
 $L_0(a) = 0$, $L_0(b) = L_0(c) =$
 $L_0(d) = L_0(e) = L_0(z) = \infty$


- **k=1:** $u := a \rightsquigarrow S_1 = \{a\}$,
 $L_0(a) + w(a, b) = 4 < L_0(b) \rightsquigarrow L_1(b) = 4$
 $L_0(a) + w(a, c) = 2 < L_0(c) \rightsquigarrow L_1(c) = 2$
- **k=2:** $u := c \rightsquigarrow S_1 = \{a, c\}$,
 $L_1(c) + w(c, b) = 3 < L_1(b) \rightsquigarrow L_2(b) = 3$
 $L_1(c) + w(c, d) = 10 < L_1(d) \rightsquigarrow L_2(d) = 10$
 $L_1(c) + w(c, e) = 12 < L_1(e) \rightsquigarrow L_2(e) = 12$
- **k=3:** $u := b \rightsquigarrow S_1 = \{a, c, b\}$,
 $L_2(b) + w(b, d) = 8 < L_2(d) \rightsquigarrow L_3(d) = 8$
- **k=4:** $u := d \rightsquigarrow S_1 = \{a, c, b, d\}$,
 $L_3(d) + w(d, e) = 10 < L_3(e) \rightsquigarrow L_4(e) = 10$
 $L_3(d) + w(d, z) = 14 < L_3(z) \rightsquigarrow L_4(z) = 14$
- **k=5:** $u := e \rightsquigarrow S_1 = \{a, c, b, d, e\}$,
 $L_4(e) + w(e, z) = 13 < L_4(z) \rightsquigarrow L_5(z) = 13$
- **k=6:** $u := z \rightsquigarrow S_1 = \{a, c, b, d, z\}$,
- **return:** $L(z) = 13$

Traveling Salesperson Problem



Traveling salesperson problem: a traveling salesperson wants to visit each of the cities once and return to his starting point. In which order should he visit these cities to travel the minimum total distance?
⇒ **Hamiltonian circuit with minimum total weight in the complete graph.**

Traveling Salesperson Problem



Route	Tot. dist.
A, B, C, D, E, A	610
A, B, C, E, D, A	516
A, B, E, D, C, A	588
A, B, E, C, D, A	458
A, B, D, E, C, A	540
A, B, D, C, E, A	504
A, D, B, C, E, A	598
A, D, B, E, C, A	576
A, D, E, B, C, A	682
A, D, C, B, E, A	646
A, C, D, B, E, A	670
A, C, B, D, E, A	728



Traveling salesperson problem: a traveling salesperson wants to visit each of the cities once and return to his starting point. In which order should he visit these cities to travel the minimum total distance?

⇒ **Hamiltonian circuit with minimum total weight in the complete graph.**



Planar Graph

DEFINITION: Let $G = (V, E)$ be an undirected graph. G is called a **planar graph** 平面图 if it can be drawn in the plane without any edges crossing.

- Crossing of edges: an intersection other than endpoints (vertices)
- **planar representation** 平面表示: a drawing w/o edge crossing; **nonplanar** 非平面的




- K_1, K_2, K_3, K_4 are planar graphs
- $K_{1,n}, K_{2,n}$ are planar graphs
- C_n ($n \geq 3$), W_n ($n \geq 3$) are planar graphs
- Q_1, Q_2, Q_3 are planar graphs



Planar Graph

Examples



A graph may be planar even if it is usually drawn with crossings, because it may be possible to draw it in a different way without crossings.

Nonplanar Graph




约旦曲线

Jordan Curve Theorem: Every simple closed planar curve Γ separates the plane into a bounded interior region and an unbounded exterior region. Any planar curve connecting the two regions must intersect Γ .

有界内部区域

无界外部区域


EXAMPLE: The bipartite graph $K_{3,3}$ is not planar.



- choose a simple circuit v_1, v_5, v_2, v_4, v_1 in $K_{3,3}$
- If $K_{3,3}$ is a planar, then the circuit forms a simple closed planar curve
- Add v_3, v_6 and the edges incident with them.
 - Intersection occurs (due to the Jordan curve Theorem).

Nonplanar Graph

EXAMPLE: The complete graph K_5 is not planar.




- $v_1, v_2, v_3, v_4, v_5, v_1$ is a simple closed curve in the planar representation of K_5
- Every remaining edge is in the interior region or in the exterior region
 - at least one is in the interior region
- No matter how you draw the remaining edges, crossing occurs.

Regions

DEFINITION: Let $G = (V, E)$ be a planar graph. Then the plane is divided into several **regions**面 by the edges of G .

- The infinite region is **exterior region** ^{R_0} 外部面. The others are **interior regions**内部面.
- The **boundary**边界 of a region is a subset of E .
- The **degree**度数 of a region is the number of edges on its boundary.
 - If an edge is shared by R_i, R_j , then it contributes 1 to $\deg(R_i), \deg(R_j)$
 - If an edge is on the boundary of a single region R_i , then it contributes 2 to $\deg(R_i)$



- The plane is divided into 5 regions R_0, R_1, R_2, R_3, R_4
 - R_0 is the exterior region
 - R_1, R_2, R_3, R_4 are interior regions
- The boundary of R_1 ; $\deg(R_1) = 4$
- There are 4 edges on the boundary of R_4
 - $\deg(R_4) = 1 + 1 + 1 + 2 = 5$ because one of the edges contribute 2 to $\deg(R_4)$
- $\deg(R_0) = 11, \deg(R_1) = 4, \deg(R_2) = 3, \deg(R_3) = 3, \deg(R_4) = 5$

Euler's Formula

THEOREM: Let $G = (V, E)$ be a connected planar simple graph with e edges and v vertices. Let r be the number of regions in a planar representation of G . Then $r = e - v + 2$.

THEOREM: Let G be a planar simple graph with p connected components. Then $|V(G)| - |E(G)| + |R(G)| = p + 1$.

- Let G_1, G_2, \dots, G_p be the connected components of G .
 - By Euler's formula, $|R(G_i)| = |E(G_i)| - |V(G_i)| + 2$ for all $i \in [p]$
- $|V(G)| = |V(G_1)| + |V(G_2)| + \dots + |V(G_p)|$
- $|E(G)| = |E(G_1)| + |E(G_2)| + \dots + |E(G_p)|$
- $|R(G)| = |R(G_1)| + |R(G_2)| + \dots + |R(G_p)| - p + 1$
- $|V(G)| - |E(G)| + |R(G)| = \sum_{i=1}^p (|V(G_i)| - |E(G_i)| + |R(G_i)|) - p + 1$
 $= 2p - p + 1 = p + 1$

Euler's Formula: Proof*

Proof of Euler's formula by induction on the number e of edges

- A simple connected planar graph with 0 edges has only one vertex and one face (unbounded). The relation $f = e - v + 2$ is satisfied.
- Suppose the relation is satisfied for all simple connected planar graphs with k edges.

Consider a simple connected planar graph G with $k + 1$ edges, $k \geq 0$. This graph can be seen as a simple connected planar graph G' with k edges (satisfying the relation by induction hypothesis) to which we add one edge. There are two ways to add an edge to G' to get G :

- either the two endpoints of the edge are already in G' : in this case, adding the edge adds also one face,
- either only one of the endpoint is already in G' : in this case, adding the edge adds also one vertex but no other face.

In both cases, the relation $f = e - v + 2$ is satisfied by G .

Application

THEOREM: Let G be a connected planar simple graph. If every region has degree $\geq l$ in a planar representation of G , then

then $|E(G)| \leq \frac{l}{l-2} (|V(G)| - 2)$.

- Let R_1, \dots, R_t be the regions given by a planar representation of G // $t = |R(G)|$
 - $\deg(R_i) \geq l$ for every $i = 1, 2, \dots, t$
- Let $r = \deg(R_1) + \deg(R_2) + \dots + \deg(R_t)$. Then $r = 2|E(G)|$.
 - Every edge contributes 2 to r
 - If $e \in E$ is on the boundary of a single region R_i , then e contributes 2 to $\deg(R_i)$;
 - If $e \in E$ is shared by R_i and R_j , then e contributes 1 to $\deg(R_i)$ and 1 to $\deg(R_j)$;
- $2|E(G)| = r = \deg(R_1) + \deg(R_2) + \dots + \deg(R_t) \geq lt = l|R(G)|$
- $|R(G)| = |E(G)| - |V(G)| + 2$
- Hence, $|E(G)| \leq \frac{l}{l-2} (|V(G)| - 2)$

Application

推论

COROLLARY: Let G be a connected planar simple graph. If $|V(G)| \geq 3$, then $|E(G)| \leq 3|V(G)| - 6$.

- Every region has degree ≥ 3 in a planar representation of G
- Let $l = 3$ in the previous theorem
 - $|E(G)| \leq \frac{3}{3-2} (|V(G)| - 2) = 3|V(G)| - 6$.

EXAMPLE: The complete graph K_5 is not planar.

- $|E(K_5)| = \binom{5}{2} = 10, |V(K_5)| = 5, K_5$ is connected simple and of order ≥ 3
- $|E(K_5)| > 3|V(K_5)| - 6$
 - Hence, K_5 cannot be planar

COROLLARY: Let G be a connected planar simple graph. Then G has a vertex of degree ≤ 5 .

- $|V(G)| < 3$: the statement is true.
- $|V(G)| \geq 3: \forall u \in V(G), \deg(u) \geq 6 \Rightarrow 2|E(G)| = \sum_u \deg(u) \geq 6|V(G)|$
 - G cannot be planar


Application

COROLLARY: Let G be a connected planar simple graph. If $|V(G)| \geq 3$ and there is no circuits of length 3 in G , then $|E(G)| \leq 2|V(G)| - 4$.

- Let R_1, \dots, R_t be the regions given by a planar representation of G // $t = |R(G)|$
 - $\deg(R_i) \geq 4$ for every $i = 1, 2, \dots, t$
- Hence, $|E(G)| \leq \frac{4}{4-2} (|V(G)| - 2) = 2|V(G)| - 4$

EXAMPLE: The complete bipartite graph $K_{3,3}$ is not planar.

- $|E(K_{3,3})| = 3 \times 3 = 9, |V(K_{3,3})| = 3 + 3 = 6 \geq 3$
- $K_{3,3}$ is connected, simple and of order ≥ 3 .
- There is no circuits of length 3 in $K_{3,3}$
- $|E(K_{3,3})| = 9 > 8 = 2|V(K_{3,3})| - 4$
- Hence, $K_{3,3}$ cannot be planar




REMARKS: K_5 and $K_{3,3}$ are fundamental nonplanar graphs.

Homeomorphic

DEFINITION: Let $G = (V, E)$ be a graph and $\{u, v\} \in E$.

- **elementary subdivision** 初等细分: $G' = (V \cup \{w\}, E - \{u, v\} + \{u, w\} + \{v, w\})$
- Two graphs are **homeomorphic** 同胚的 if they can be obtained from the same graph via elementary subdivisions






G_2 and G_3 are homeomorphic

Kuratowski's Theorem




库拉托夫斯基定理

THEOREM: A graph G is nonplanar if and only if it has a subgraph homeomorphic to $K_{3,3}$ or K_5 .

EXAMPLE: The following graph is nonplanar.




There is a subgraph
homeomorphic to K_5



There is a subgraph
homeomorphic to $K_{3,3}$


Dual Graph 对偶图

Let G be a planar graph and assume we take a planar representation of G that we denote also G . The **dual of G** is the graph G^* that has a vertex for each face of G and an edge connecting two vertices if the corresponding faces in G have a common edge in their boundary.



Remark: The dual of a planar simple graph is not necessarily simple.

Coloring a Map





Coloring regions of the map \Leftrightarrow Coloring vertices of the dual graph

Graph Coloring

DEFINITION: Let $G = (V, E)$ be a simple graph. A **k -coloring** _{k -着色} of G is a map $f: V \rightarrow [k]$ such that $f(u) \neq f(v)$ whenever $\{u, v\} \in E$.

- **chromatic number** ($\chi(G)$)_{色数}: the least k s.t. G has a k -coloring.





$$\chi(G) = 3$$

The chromatic number is at least 3 because $a; b; c$ is a circuit of length 3

Graph Coloring

DEFINITION: Let $G = (V, E)$ be a simple graph. A **k -coloring** _{k -着色} of G is a map $f: V \rightarrow [k]$ such that $f(u) \neq f(v)$ whenever $\{u, v\} \in E$.

- **chromatic number** $(\chi(G))$ _{色数}: the least k s.t. G has a k -coloring.





$$\chi(G) = 4$$

Graph Coloring



THEOREM: Let $G = (V, E)$ be a simple graph.

- $1 \leq \chi(G) \leq |V|$
- $\chi(G) = 1$ iff $E = \emptyset$
- $\chi(G) = 2$ iff G is bipartite and $|E| \geq 1$.
- $\chi(K_n) = n$ for every integer $n \geq 1$.
 - $\chi(G) \geq n$ if G has a subgraph isomorphic to K_n
- $\chi(C_n) = 2$ if $2|n$; $\chi(C_n) = 3$ if $2 \nmid (n - 1)$; ($n \geq 3$)
- $\chi(G) \leq \Delta(G) + 1$, where $\Delta(G) = \max\{\deg(v) : v \in V\}$.



Application

PROBLEM: How can the final exams at a university be scheduled so that no student has two exams at the same time?



- There are 7 different courses, they are vertices of a graph.
- Two courses are adjacent if there is a student registered both courses.
- Choose time slots for the courses such that no two adjacent courses take place at the same time.

$$1 \leq \chi(G) \leq 7$$

- $\chi(G)$ time slots is needed. $1 \leq \chi(G) \leq \Delta(G) + 1 = 6$

$$\chi(G) \geq 4: G \text{ has a subgraph isomorphic to } K_4$$

4-coloring Theorem

Theorem (Four coloring Theorem)

The chromatic number of a simple planar graph is no greater than 4.

Remarks: The proof of the 4-coloring Theorem depends on a computer. The two previous theorems are true for planar graphs only. A non planar graph can have an arbitrarily large chromatic number.