

SI251 - Convex Optimization, 2024 Spring  
Homework 2

Name: **Zhou Shouchen**

Student ID: 2021533042

Due 23:59 (CST), Apr. 10, 2024

1. **(50 pts) Robust quadratic programming.** In the lecture, we have learned about robust linear programming as an application of second-order cone programming. Now we will consider a similar robust variation of the convex quadratic program

$$\begin{aligned} & \text{minimize} && (1/2)x^T Px + q^T x + r \\ & \text{subject to} && Ax \preceq b. \end{aligned}$$

For simplicity, we assume that only the matrix  $P$  is subject to errors, and the other parameters  $(q, r, A, b)$  are exactly known. The robust quadratic program is defined as

$$\begin{aligned} & \text{minimize} && \sup_{P \in \mathcal{E}} ((1/2)x^T Px + q^T x + r) \\ & \text{subject to} && Ax \preceq b \end{aligned}$$

where  $\mathcal{E}$  is the set of possible matrices  $P$ .

For each of the following sets  $\mathcal{E}$ , express the robust QP as a convex problem in a standard form (e.g., QP, QCQP, SOCP, SDP).

- (a) A finite set of matrices:  $\mathcal{E} = \{P_1, \dots, P_K\}$ , where  $P_i \in S_+^n, i = 1, \dots, K$ .
- (b) A set specified by a nominal value  $P_0 \in S_+^n$  plus a bound on the eigenvalues of the deviation  $P - P_0$ :

$$\mathcal{E} = \{P \in \mathbf{S}^n \mid -\gamma I \preceq P - P_0 \preceq \gamma I\}$$

where  $\gamma \in \mathbf{R}$  and  $P_0 \in \mathbf{S}_+^n$ .

- (c) An ellipsoid of matrices:

$$\mathcal{E} = \left\{ P_0 + \sum_{i=1}^K P_i u_i \mid \|u\|_2 \leq 1 \right\}.$$

You can assume  $P_i \in \mathbf{S}_+^n, i = 0, \dots, K$ .

**Solution:**

2. **(50 pts) Water-filling.** Please consider the convex optimization problem and calculate its solution

$$\begin{aligned} & \text{minimize} && - \sum_{i=1}^n \log(\alpha_i + x_i) \\ & \text{subject to} && x \succeq 0, \quad \mathbf{1}^T x = 1 \end{aligned}$$

**Solution:**