

# Math 500 - Topology and Geometry

Notes by Zach Schutzman

University of Pennsylvania, Fall 2017

<b>Lecture 1 (2017-08-30)</b>	<b>1</b>	<b>Lecture 4 (09-08-2017)</b>	<b>8</b>
<b>Lecture 2 (2017-09-01)</b>	<b>3</b>	<b>Lecture 5 (09-11-2017)</b>	<b>10</b>
<b>Lecture 3 (09-06-2017)</b>	<b>5</b>	<b>1 Topological Bases</b>	<b>10</b>

## Introduction

Math 500 is a Masters-level first-course in Topology and Geometry. The course follows James Munkres' *Topology, 2ed.* and this set of notes is based on the Fall 2017 offering.

These notes are being live-Texed, though I edit for typos and add diagrams requiring the *TikZ* package separately. I am using the editor TeXstudio. The template for these notes was created by Zev Chonoles and is made available (and being used here) under a Creative Commons License.

I am responsible for all faults in this document, mathematical or otherwise; any merits of the material here should be credited to the lecturer, not to me.

Please email any corrections or suggestions to [ianzach+notes@seas.upenn.edu](mailto:ianzach+notes@seas.upenn.edu).

## Lecture 1 (2017-08-30)

### What is Topology?

**Definition.** A **topology** is a set  $X$  with a collection of subsets  $\mathcal{A} \subset \mathcal{P}(X)$  such that:

- 1  $\emptyset, X \in \mathcal{A}$
- 2  $\mathcal{A}$  is closed under finite intersection (the intersection of a finite subset of  $\mathcal{A}$  is in  $\mathcal{A}$ )
- 3  $\mathcal{A}$  is closed under arbitrary union (the union of any (possibly infinite) subset of  $\mathcal{A}$  is in  $\mathcal{A}$ )

The phrases “ $\mathcal{A}$  is a topology on  $X$ ”, “ $X$  is a topological space with topology  $\mathcal{A}$ ”, and the notation  $(X, \mathcal{A})$  all refer to the same concept.

**Definition.** The **standard topology** (also called the euclidean topology or metric topology) on  $\mathbb{R}^n$  is the set of subsets  $U \subset \mathbb{R}^n$  such that for every  $U$ , every point  $x \in U$  is interior, meaning that there exists some radius  $r > 0$  such that the ball of radius  $r$  centered at  $x$  is entirely contained in  $U$ .

**Definition.** A set is **open** in a topological space  $X$  if it belongs to the topology on  $X$ .

**Example.** The standard topology is a topology over  $\mathbb{R}^n$ :

- 1 Every point in the empty set is vacuously interior, and every point of  $\mathbb{R}^n$  is trivially interior
- 2 If we take two open sets and intersect them, any point in the intersection must be an interior point in both constituent sets. The smaller of the two balls witnessing this must lie entirely within both constituent sets, and therefore entirely within the intersection. By induction, we have the finite intersection of open sets being open.
- 3 Intuitively, taking any union of open sets only creates a bigger set. The ball witnessing any point as interior to some open set clearly lies in any union including that open set.

We can see from this example why it's important to specify closure under *finite* intersection. Singleton sets are not open in the standard topology on  $\mathbb{R}^n$ , but the Nested Interval Theorem gives us a way to construct a singleton set from the countable intersection of open intervals.

**Example.** If  $X$  is our topological space,  $\{\emptyset, X\}$  is a topology, called the **trivial topology**.

**Example.** Similarly, all of  $\mathcal{P}(X)$  is a topology, called the **discrete topology**.

**Example.** The **Zariski topology** on  $\mathbb{R}^n$  is a little more interesting. A set is open in the Zariski topology if it is the complement of the root set of some polynomial. Open sets in the one-dimensional case look like the real line minus a finite number of points. It gets a little more complicated in higher dimensions, as we can have zeroes along entire dimensions of a euclidean space. Let's verify that this is a topology:

- 1 The empty set is the complement of the root set of the zero function, and the entire space  $\mathbb{R}^n$  is the complement of the root set of a polynomial which has no real roots, such as  $f(\vec{x}) = 6$ .
- 2 The intersection of two open sets, corresponding to polynomials  $P$  and  $Q$  is, by DeMorgan's Laws,  $\mathbb{R}^n \setminus \{x \mid x \text{ is a root of } P \text{ or } Q\}$ . Something is a root of  $P$  or  $Q$ , it must be a root of the product  $PQ$ . Since the finite product of polynomials is a polynomial, this set is still the complement of the root set of some polynomial, and is therefore open, and we have closure under finite intersection.

3 Again by DeMorgan's Laws, the union of two open sets corresponding to polynomials  $P$  and  $Q$  is the set  $\mathbb{R}^n \setminus \{x \mid x \text{ is a root of } P \text{ and } Q\}$ . The set of points which are roots of  $P$  and  $Q$  are the roots of the greatest common polynomial divisor of  $P$  and  $Q$ . Since this is also a polynomial, our set is the complement of the root set of a polynomial and is therefore open. Since the greatest common polynomial divisor of any set of polynomials has root set no greater than any of the constituent polynomials, we properly have closure under arbitrary union.

The Zariski topology is an object of importance in the area of algebraic geometry.

**Definition.** If  $X$  is a topological space with topology  $\mathcal{A}$  and  $Y \subset X$ , then  $\mathcal{B}$  is a topology on  $Y$  where a subset  $V \subset Y$  is open in  $\mathcal{B}$  if and only if there is a  $U$  open in  $\mathcal{A}$  such that  $V = U \cap Y$ . This is called the **subset** or **subspace topology**.

**Example.** Let  $H^2$  denote the closed upper-half plane in  $\mathbb{R}^2$ . That is, the set of points  $(x, y) \in \mathbb{R}^2$  such that  $y \geq 0$ . Any set which was open in  $\mathbb{R}^2$  and does not intersect the  $x$ -axis is still open in  $H^2$ . However, a set like an open half-disk against the  $x$ -axis together with the line segment where it rests up against the  $x$ -axis was not an open set in  $\mathbb{R}^2$ , as the boundary points are not interior, but it is open in  $H^2$  with the subspace topology, as it is the intersection of an open disk in  $\mathbb{R}^2$  with the upper half-plane.

## Lecture 2 (2017-09-01)

### Continuous Maps

Continuous maps are the standard morphisms in topology.

In Analysis, we have a definition of continuity which looks like:

**Definition.** A function  $f : X \rightarrow Y$  is **continuous** at  $x \in X$  if for any  $\delta > 0$  there exists an  $\epsilon > 0$  such that  $\|x - y\| < \epsilon$  implies  $\|f(x) - f(y)\| < \delta$ .

The issue with this definition is that we have no natural notion of distance in topology. Instead, we use the definition:

**Definition.** A function  $f : X \rightarrow Y$  is **continuous** if the inverse image of an open set in  $Y$  is open in  $X$ . Equivalently, the inverse image of closed sets are closed.

It turns out that in metric spaces like  $\mathbb{R}^n$  with the standard topology, these definitions are equivalent.

**Example.** Let's consider two topological spaces:  $(\mathbb{R}, std)$ , the real numbers with the standard topology, and  $(\mathbb{R}, \mathcal{P}(\mathbb{R}))$ , the real numbers with the discrete topology. The map  $f : (\mathbb{R}, \mathcal{P}(\mathbb{R})) \rightarrow (\mathbb{R}, std)$ , where  $f(x) = x$  is continuous. Since every set is open in the discrete topology, the inverse image of any set, in particular any open set, is open. The map  $g : (\mathbb{R}, std) \rightarrow (\mathbb{R}, \mathcal{P}(\mathbb{R}))$ , where  $g(x) = x$  is not continuous. To see this, take any set that is closed with respect to the standard topology. This set is open in the discrete topology, but its inverse image is closed.

This raises the question: is there any  $g : (\mathbb{R}, std) \rightarrow (\mathbb{R}, \mathcal{P}(\mathbb{R}))$  which is continuous?

**Theorem.** *The only continuous functions  $g : (\mathbb{R}, std) \rightarrow (\mathbb{R}, \mathcal{P}(\mathbb{R}))$  are the constant maps.*

*Proof.* First, it is easy to see that a constant map is continuous. Without loss of generality, we'll assume that  $g(x) = 0$ . Let  $V$  be open in the discrete topology. If  $V$  contains 0, then the inverse image of  $V$  is all of  $\mathbb{R}$ . If  $V$  does not contain zero, then the inverse image of  $V$  is the empty set. Since both of these are open in the standard topology, the inverse image of any open set is open, and the map is continuous.

To see that such a continuous map must be constant, first observe that  $\mathbb{R}$  and  $\emptyset$  are the only sets which are both closed and open with respect to the standard topology. Let  $g : (\mathbb{R}, std) \rightarrow (\mathbb{R}, \mathcal{P}(\mathbb{R}))$  be a continuous map and pick some  $x \in \mathbb{R}$ . The set  $\{g(x)\}$  is both closed and open in the discrete topology (as every set is closed and open), so its inverse image must be, in particular, open. But the inverse image cannot be empty, as we know for sure it contains  $x$ , and the only non-empty closed and open set in the standard topology is the entire space. Therefore, for any  $x, y \in \mathbb{R}$ , we have  $g(x) = g(y)$ , which is only true for constant maps.

□

**Definition.** A **homeomorphism** is a continuous bijection between two topological spaces such that the inverse is also continuous.

Under a homeomorphism, we also have the property that the image of open sets is open. This induces a bijection between the open sets of the two topological spaces. In a sense, the existence of a homeomorphism means that two topological spaces are the same.

**Example.** The two spaces  $(-1, 1)$  and  $(-2, 2)$  with the standard topology are homeomorphic under the map  $f(x) = 2x$ .

**Example.** The two spaces  $(-1, 1)$  and  $\mathbb{R}$  with the standard topology on each are homeomorphic under the map  $f(x) = \tan(\frac{\pi}{2}x)$ .

**Example.** Let  $S^n = \mathbb{R}^n \cup \{\infty\}$ . A set  $U \subset S^n$  is open if:

$$U = \emptyset \text{ or } U = S^n$$

$U \subset \mathbb{R}^n$  and  $U$  is open with respect to the standard topology.

$\infty \in U$  and  $U \cap \mathbb{R}^n$  is the complement of a compact subset of  $\mathbb{R}^n$ . That is,  $U$  looks like all of  $\mathbb{R}^n$  with a closed and bounded chunk removed, and an additional point  $\infty$ .

This forms a topology, and the set  $S^n$  is the surface of the  $n$ -dimensional sphere. If we think about  $S^2$ , there's a natural embedding in  $\mathbb{R}^3$ , but it turns out that  $S^2 \setminus \{(0, 0, 1)\}$  is homeomorphic to  $\mathbb{R}^2$ . If we use the (north polar) stereographic projection, which maps points in  $S^2$  to the point in the  $\mathbb{R}^2$  plane according to the straight line passing through the north pole and that point, we get a nice homeomorphism, and this is easy to see from the subspace topology that  $S^2$  inherits from  $\mathbb{R}^3$ . If we then include that the north pole maps to our added point  $\infty$ , we get a map from all of  $S^2$  to the set  $\mathbb{R}^2 \cup \{\infty\}$  which is a homeomorphism.

## The Quotient Topology

Let  $(X, \mathcal{A})$  be a topological space and  $\sim$  an equivalence relation on  $X$ . Then  $X/\sim$  inherits a topology, which is that  $U' \subset X/\sim$  is open if and only if there is some open set  $U \subset X$  such that  $U' = U/\sim$ .

**Definition.** This topology is called the **quotient topology**, or the **identification map**.

**Example.** Take  $\mathbb{R}^2$  with the standard topology and define an equivalence relation  $(x, y) \sim (x, -y)$ . The quotient space looks like the closed (upper or lower) half-plane.

**Example.**  $\mathbb{R}^2$  with the standard topology quotiented by the equivalence relation  $(x, y) \sim (-x, -y)$  looks like a cone, and is actually homeomorphic to  $\mathbb{R}^2$ .

**Example.**  $\mathbb{R}^2$  with the standard topology and the equivalence relation  $(x, y) \sim (x+1, y) \sim (x, y+1)$  has a quotient space that looks like the unit square with opposite sides glued together. This is homeomorphic to a (genus 1) torus.

## Lecture 3 (09-06-2017)

### The Pullback Topology

Let  $(X, \mathcal{A})$  be a topological space and  $Y$  some set. Given a map  $f : X \rightarrow Y$ ,  $Y$  inherits a topology from  $X$  where  $V \subset Y$  is open if and only if  $f^{-1}(V) \subset X$  is open.

**Definition.** This topology on  $Y$  is called the **pullback topology**.

The pullback topology is the finest topology on  $Y$  which makes  $f$  a continuous map.

**Example.** Take  $f : (-1, 1) \rightarrow \mathbb{R}$  with  $f(x) = x$  and the standard topology on each. The pullback topology on  $\mathbb{R}$  has open sets  $\emptyset$  and  $\mathbb{R}$ , whose inverse images are themselves. Also, any set in  $\mathbb{R}$  which does not intersect the open interval  $(-1, 1)$ , as all of these sets have empty inverse image. Finally, any set which is open in  $(-1, 1)$  or whose intersection with  $(-1, 1)$  is open is also open in the pullback topology.

### Group Actions and Fundamental Regions

Lets think about  $\mathbb{Z}^2$  as a group action on  $\mathbb{R}^2$ , where applying  $(a, b) \in \mathbb{Z}^2$  to  $(x, y) \in \mathbb{R}^2$  means shifting  $(x, y)$  right by  $a$  and up by  $b$  (left, down if  $a$  or  $b$  is negative, of course). We write this as  $\mathbb{Z}^2$  acts on  $\mathbb{R}^2$  by  $(a, b).(x, y) = (x + a, y + b)$ . This establishes an equivalence relation on  $\mathbb{R}^2$  :  $(x_0, y_0) \sim (x_1, y_1)$  if there exists  $(a, b) \in \mathbb{Z}^2$  such that  $(a, b).(x_0, y_0) = (x_1, y_1)$ . This divides  $\mathbb{R}^2$  into  $1 \times 1$  squares, where each square is equivalent to any other, and we identify the left and right edges and the top and bottom edges, but no two points in the interior of any given square are equivalent. We call the squares fundamental regions.

**Definition.** A **fundamental region** of a group action and is the (closure of) largest region such that no two interior points are identified with respect to the induced equivalence relation.

**Example.** The fundamental region described above, the square with opposite edges identified, defines a torus.




Figure 1:

Open sets in the torus  $\mathbb{T}$  are  $\emptyset$  and  $\mathbb{T}$ , and the intersection of any standard open set with the fundamental region. This is the same as the inherited subspace topology from  $\mathbb{R}^2$ .

**Example.** If we consider  $\mathbb{R}^2/\mathbb{Z}$ , where  $a \in \mathbb{Z}$  acts on  $\mathbb{R}^2$  by  $a.(x, y) = (x + a, y)$ . A fundamental domain of this action is a vertical strip of unit width. Again this inherits a subspace topology from  $\mathbb{R}^2$ .



Figure 2:

**Example.** Consider again  $\mathbb{R}^2/\mathbb{Z}$  but this time with the group action  $a.(x, y) = (x + a, (-1)^a y)$ . The fundamental region is still a strip of unit width, but this time instead of identifying points on the boundary with their horizontal translation, we identify them with their horizontal translation composed with reflection about the  $x$ -axis. This space is homeomorphic to an infinite Moebius strip, which is difficult to draw.

**Example.** Consider the equivalence relation on  $\mathbb{R}^2$  described by  $(x, y) \sim (x, y)$ ,  $(0, y) \sim (1, y)$ , and  $(x, 0) \sim (1 - x, 1)$ . The fundamental region again is a square with the left and right edges identified by simple translation, but the top and bottom edges are now identified by translation plus a flip across the square's vertical axis of symmetry. This is homeomorphic to the Klein bottle, which is, again, hard to draw.

**Example.** The previous example where we also identify the left and right edges by translation and a flip is called the real projective plane, denoted  $\mathbb{R}P^2$ . Both vertical and horizontal strips of this space look like Moebius strips. This is, once again, not easy to draw.

**Example.** This one we can draw! Take the unit square as the fundamental region, but identify the top and left edge with each other by symmetry about the corner where they intersect, and do the same for the bottom and right edge. This space is homeomorphic to the 2-sphere  $\mathbb{S}^2$ .





Figure 3:

## Lecture 4 (09-08-2017)

### Mapping Cylinders and Tori

**Definition.** Let  $I$  denote the closed unit interval  $[0, 1]$ . The **mapping cylinder** of a continuous map  $f : X \rightarrow Y$  is the quotient space defined by  $X \times I / \sim$ , where  $(x, 0) \sim (f(x), 1)$ .

**Definition.** The **mapping torus** of a map  $f : X \rightarrow X$  is similar, except that we require that the map be from a space to a copy of itself and we define the equivalence relation as  $(x, 0) \sim (f(x), 0)$ .

**Example.** The mapping cylinder of  $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$  where  $f(x) = x$  is a regular old cylinder. The mapping torus is a regular old torus.

**Example.** We can equivalently think of  $\mathbb{S}^1$  as  $\{(x, y) | x^2 + y^2 = 1\}$  in Euclidean space or as  $\{(r, \theta) | r = 1\}$  in polar coordinates. Using this second formulation, consider the map  $f : \mathbb{S}^1 \rightarrow \mathbb{S}^1$  where  $f(\theta) = 2\theta$ . This is a two-to-one map which maps antipodal points to each other. The mapping cylinder of  $f$  is a Moebius strip.

**Example.** What about the three-to-one map  $f(\theta) = 3\theta$ ? The mapping cylinder of this looks like a three-bladed wing with a one-third twist and the ends glued together.

### Boundaries and Exteriors

Let  $(X, \mathcal{A})$  be a topological space and let  $K \subseteq X$ .

**Definition.** The **interior** of  $K$  is the largest open subset contained in  $K$ . That is, it is the union of all  $U \subset K$  such that  $U \in \mathcal{A}$ .

**Definition.** The **closure** of  $K$  is the smallest closed subset containing  $K$ . That is, it is the intersection of all  $V \supset K$  such that  $(X - V) \in \mathcal{A}$ .

**Definition.** The **boundary** of  $K$  is the intersection of the closure of  $K$  with the closure of the complement of  $K$ , that is  $Bd(K) = \overline{K} \cap \overline{X - K}$ . If  $K$  is open, then  $Bd(K) = \overline{K} - K$ . If  $K$  is closed, then  $Bd(K) = \partial K$ .

**Example.** Take  $K = \mathbb{Q} \cap [0, 1] \subset \mathbb{R}$  with the standard topology on  $\mathbb{R}$ . The interior of this set is empty, as there is no open interval which doesn't contain an irrational number, so  $\emptyset$  is the largest open subset in  $K$ . The closure of  $K$  is the entire interval  $[0, 1]$ , as there is no smaller closed set which contains all of the rationals in that interval. We also have that the boundary  $Bd(K) = [0, 1]$ .

**Example.** Take  $K = \mathbb{R} - \{0\}$  with the Zariski topology on  $\mathbb{R}$ . The interior of  $K$  is  $K$ , as  $K$  is open. The closure of  $K$  is all of  $\mathbb{R}$ , and the boundary is  $\{0\}$ .

**Definition.** A point  $x$  is a **limit point** of  $K \subset X$  if every open set containing  $x$  has non-empty intersection with  $K$ . Equivalently,  $x$  is a limit point of  $K$  if  $x \in \overline{K - \{x\}}$ .

**Example.** Take  $\mathbb{R}$  with the Zariski topology. If  $U$  is an open set, then every  $x \in \mathbb{R}$  is a limit point of  $U$ . In fact, for any infinite subset of  $\mathbb{R}$ , every point in  $\mathbb{R}$  is a limit point.

**Definition.** A **neighborhood** of a point  $x \in X$  is an open set containing  $x$ .

**Example.** In the either-or topology on  $[-1, 1] \subset \mathbb{R}$  has open sets  $\emptyset$ ,  $[-1, 1]$ , a set is open if and only if it does not contain 0 or it contains  $(-1, 1)$ . If  $U$  is an open set and  $0 \in U$ , then

$U = (-1, 1)$ ,  $[-1, 1)$ ,  $(-1, 1]$ , or  $[-1, 1]$ . Closed sets are subsets of  $\{-1, 1\}$ , all of  $[-1, 1]$ ,  $\emptyset$ , and any set that contains 0.

What are the continuous functions? From  $EO$  to  $std$ , constant functions are continuous. Anything else? From  $std$  to  $EO$ , continuous functions look like  $f(x) = \frac{1}{2}\text{sgn}(x)$ .

**Example.** Take  $\frac{1}{2} \in [-1, 1]$  with the either-or topology. Is  $\frac{1}{2}$  a limit point of  $[-1, 1] - \{\frac{1}{2}\}$ ? No! That set is already closed, so it contains all of its limit points.

Is 0 a limit point of  $(\frac{1}{2}, \frac{3}{4})$ ? Yes! Every open set containing 0 contains  $(-1, 1)$ , so it obviously also contains  $(\frac{1}{2}, \frac{3}{4})$ .

## Lecture 5 (09-11-2017)

### Topological Bases

**Definition.** Let  $X$  be a set. A collection  $\mathcal{B}$  of subsets of  $X$  is called a **base** (or **basis**) of  $X$  if:

- 1 If  $x \in X$  then there is a  $B \in \mathcal{B}$  such that  $x \in B$ . Equivalently,  $\mathcal{B}$  covers  $X$ .
- 2 If  $B_1, B_2 \in \mathcal{B}$  and  $x \in B_1 \cap B_2$ , then there is a  $B_3 \in \mathcal{B}$  such that  $B_3 \subset B_1 \cap B_2$  and  $x \in B_3$ .

This is a weaker concept than a topology; we don't require that the union of base elements is a base element and we only require that the intersection of base elements contains another base element.

**Example.** Consider  $\mathbb{R}^2$  with the standard topology. Define  $\mathcal{B} = \{B_x(r) | x \in \mathbb{R}^2, r > 0\}$  as the set of open balls in  $\mathbb{R}^2$ . This is a base. It is easy to see the first criterion is satisfied. To see the second, consider two balls which both contain some point  $x$ . Then there is a small ball centered at  $x$  which is fully contained in the intersection of the two balls. This smaller ball is also a base element, so we are done.

**Lemma.** If  $\mathcal{B}$  is a base and  $B_1, B_2, \dots, B_n \in \mathcal{B}$ , and  $x \in B_1 \cap B_2 \cap \dots \cap B_n$  then there exists a base element  $B' \subset B_1 \cap \dots \cap B_n$  which contains  $x$ .

*Proof.* We proceed by induction. Since  $x \in B_1 \cap B_2$ , there exists some  $D_1 \in \mathcal{B}$  such that  $x \in D_1 \subset B_1 \cap B_2$ . Then  $x \in D_1 \cap B_3$ , so there exists some  $D_2 \in \mathcal{B}$  with  $x \in D_2$ . We proceed iteratively like this to find there is some  $D_{n-1} \in \mathcal{B}$  with  $x \in D_{n-1}$ , and we set  $B' = D_{n-1}$ . □

**Definition.** A **topology generated by a base** is the collection of sets which are unions of base elements.

If  $X$  is a set with a base  $\mathcal{B}$ , then there is a smallest (coarsest) topology on  $X$  containing  $\mathcal{B}$ , which is the topology generated by  $\mathcal{B}$ . Open sets are the base elements, arbitrary unions of base elements, and  $\emptyset$  and  $X$  by definition. Do we get the intersection property as well?

**Claim.** Yes.

*Proof.* If  $B_1, \dots, B_n$  are base elements, then we can write the intersection  $\bigcap_{i \in [n]} B_i$  as the union of base elements just by taking neighborhoods of each point in the intersection. If we have  $U_1, \dots, U_n$  open in  $X$  and  $x \in \bigcap_{i \in [n]} U_i$ , then there is some base element in the intersection containing  $x$ . If we do this for all points in the intersection, we can write the intersection as an arbitrary union of base elements, and we are done. □