

# Braid Presentations for Concurrent Programs

Zachary Stone

September 23, 2026

## Abstract

Ziemiański's cube chain category  $\text{Ch}(K)$  is a combinatorial model for the space of directed paths on a precubical set  $K$ . Paliga and Ziemiański built a braid invariant for precubical sets by observing a relation between  $\pi_1|\text{Ch}(Z)|$  and braids, where  $Z$  is the terminal precubical set. However, that invariant trivializes for Euclidean  $K$  because  $|\text{Ch}(I^n)|$  is contractible [PZ21, Prop. 6.1]. This paper constructs a localization of  $\text{Ch}(K)$  at morphisms that preserve the order of coordinates, called  $\text{Br}(K)$ . And we will show that this structure is the source of the braid structure of  $\pi_1|\text{Ch}(Z)|$ . But, there is no nerve or geometric realization in this technique. Instead we build a 2-polygraph giving an artin braid-like presentation of  $\text{Br}(K)$ . The  $\{0, 1, 2\}$ -cells of the polygraph are the dimension  $\{0, 1, 2\}$  chains of  $K$ . Intuitively, 3-cells of  $K$  contribute braid relations, and pairs of non-intersecting 2-cells of  $K$  contribute commuting relations. This is not just a metaphor: at  $K = Z$ , we get the artin relations and generators exactly. So  $\text{Br}(Z)$  is equivalent to the positive braid category. The presentation is natural in  $K$ , providing a generalization of the existing work. As an application, we establish a link to the hyperplane arrangement literature. The way braids are constructed via Salvetti's theorem for the braid arrangement exactly matches the polygraph we build here for the cube. All this work is formalized in Lean [Sto26].

## 1 Introduction

Ziemiański's cube chain category  $\text{Ch}(K)$  is a combinatorial model for the executions of a bipointed precubical set  $K$  [Zie20, Thm. 7.6]. Paliga and Ziemiański established a homotopy equivalence between  $|\text{Ch}(Z)|$  and configuration spaces of points in the plane, where  $Z$  is the terminal precubical set. It extends, by naturality, to an invariant of any precubical set  $K$ . However, it trivializes on any Euclidean complex, see [PZ21, Prop. 6.1]. The proof is one line: the invariant factors through  $|\text{Ch}(I^n)|$ , which is contractible.

This paper explains that the braid structure is present in  $\text{Ch}(K)$  without passing through any homotopy equivalences. We will construct a set of morphisms,  $W$ , in  $\text{Ch}(K)$ . And define the localization  $\text{Br}(K) := \text{Ch}(K)^{\text{op}}[(W^{\text{op}})^{-1}]$ . Then we will build a 2-polygraph,  $\text{Art}(K)$ , giving an artin-like presentation for  $\text{Br}(K)$  in geometric terms. This approach gives us several advantages. Firstly, we preserve more structure of  $\text{Ch}(K)$ , so  $\text{Br}(I^n)$  does not collapse. Secondly, it permits us to stay with positive braids which yields a simpler theory.

The cells of our polygraph are straightforward. A chain of  $K$  has a list of positive integers for the dimension of the cubes. Let the degree of a chain be the sum of the list minus the length. Then the polygraph's  $n$ -cells are the degree  $n$  chains, for  $n \in \{0, 1, 2\}$ . The attachments of the 2 cells to the 1 cells will take some work to formalize, but the following pictures gives a clear intuition about the attachments.




Figure 1: The four runs over the degree-two chain  $e = \vee[2, 2]$ . Each arrow is decorated with the square that witnesses the reordering. And each node is decorated with the degree 0 refinement's permutation. So the 2 cell attaches  $\sigma_1\sigma_3 \rightarrow \sigma_3\sigma_1$ .




Figure 2: The six runs over the degree-two chain  $e = \vee[3]$ . Again each arrow is decorated with the square that witnesses the reordering. And each node is decorated with the degree 0 refinement's permutation. So we get the attachment  $\sigma_1\sigma_2\sigma_1 \rightarrow \sigma_2\sigma_1\sigma_2$ .

The main results of this paper will be

**Theorem 1.1** (Theorem 5.3, Theorem 6.1, Theorem 7.5). *Let  $K$  be a bipointed precubical set*

1.  $\text{Art}(K)$  presents  $\text{Br}(K)$ , in the sense that there is an equivalence  $E_K : \langle \text{Art}(K) \rangle \rightarrow \text{Br}(K)$ .
2. Both constructions are functorial in  $K$ , and  $E$  is natural: for every  $f : K \rightarrow L$ ,

$$E_L \circ \langle \text{Art}(f) \rangle = \text{Br}(f) \circ E_K.$$

3.  $\text{Art}(Z)$  is isomorphic to positive Artin's polygraph for the braid category. Consequently  $\text{Br}(Z)$  is equivalent to the positive braid category.

We are begging the question by naming our polygraph  $\text{Art}$ , but it is well justified in the spirit of [GGM15]. At  $K = Z$ , we see above that we will prove it is indeed the literal artin presentation of the positive braids. So in general, we get a subset of the artin relations in this presentation depending the precense of 3-cells of  $K$ .

**Formalisation.** All of the below is formalised in Lean 4 on mathlib [Sto26], sorry-free. The objects  $\text{Br}(K)$ ,  $W$  and  $\text{Art}(K)$  are `ChainCat.Br`, `ChainCat.W` and `Paper.poly`; the three parts of Theorem 1.1 are `Paper.paperPresents`, `Paper.paperSquare` and `Paper.paperArtinIso`. Each statement below appears in `CubeChains/Paper.lean` as an `example` discharged by one citation.

**Conventions.** We assume the reader is familiar with [PZ21; Zie20] and use its notation:  $\square_*$  is the category of bipointed precubical sets, with  $Z$  final and  $I^n$  the  $n$ -cube. Put  $[n] := \{1, \dots, n\}$ . Also, note that the ops are there because we have chosen to use the convention for  $\text{Ch}(K)$  as in [Zie20]. But that won't translate across to braids conveniently. So we will need the op in the definition of  $\text{Br}$ .

## 2 Background

► *One paragraph placing Proposition 2.1: everything downstream is bookkeeping about the events of a wedge, and this is the only place the precubical structure is touched.*

**Cube chains.** A **shape** is a list  $a = [a_1, \dots, a_r]$  of positive integers; it names the serial wedge  $\bigvee a := I^{a_1} \vee \dots \vee I^{a_r}$ , whose **blocks** are the  $I^{a_i}$ , and we put  $|a| := \sum_i a_i$  and  $\deg a := |a| - r$ . Objects of  $\text{Ch}(K)$  are the bipointed maps  $\bigvee a \rightarrow K$  and its morphisms, the refinements, the maps of wedges over  $K$ . We write  $|d|$  and  $\deg d$  for the event count and degree of a chain's shape, and  $\text{Ch}(u)$  for the pushforward along  $u : K \rightarrow L$ , which postcomposes the classifying map and leaves the shape alone. A chain of degree 0 is a **run**, and the subcategory of runs is discrete, a refinement losing no block being an identity.

We import three facts from [PZ21, §2]: serial wedges are non-self-linked and admit an altitude, for which the final vertex of  $\bigvee a$  has altitude  $|a|$ ; a bipointed map between connected objects admitting altitude preserves it [PZ21, Prop. 2.7]; and concatenation  $\text{Ch}(P) \times \text{Ch}(Q) \rightarrow \text{Ch}(P \vee Q)$  is an isomorphism when  $P \vee Q$  is non-self-linked and admits altitude [PZ21, Lem. 2.11].

**The events of a shape.** Send  $I^n$  to its set of directions  $[n]$ , and a face embedding to the monotone injection naming the directions it leaves free. The left Kan extension of that functor along Yoneda is cocontinuous and vanishes at  $I^0$ , so the pushout defining a wedge becomes a coproduct: the **events**

of  $\vee a$  are the pairs  $(i, c)$  with  $i \in [r]$  and  $c \in [a_i]$ , one for each direction of each block. Order them lexicographically and put  $\text{blk}_a(i, c) := i$ , so the blocks of  $a$  are the fibres of  $\text{blk}_a$  and are intervals for that order; we drop the subscript when the shape is clear. Numbering the events  $1, \dots, N$  in order identifies them with  $[N]$ ,  $N = |a|$ , and we use whichever description is convenient. Write  $f_*$  for the map on events induced by a map  $f$  of serial wedges.

**Proposition 2.1.** *Let  $f : \vee a \rightarrow \vee b$  in  $\square_*$ , with  $a$  and  $b$  shapes on  $N$  and  $M$  events. Then  $N = M$ ; the map  $f_*$  is a permutation of  $[N]$ , and  $f \mapsto f_*$  is injective; and there is a monotone  $\text{blk}(f)$  with*

$$\text{blk}_b \circ f_* = \text{blk}(f) \circ \text{blk}_a, \quad (1)$$

*the restriction of  $f_*$  to each block of  $a$  being increasing.*

*Proof.* By the pushout,  $f$  is determined by its restrictions to the blocks of  $a$ , each of which lands in a single block of  $b$  because  $a_i > 0$ ; that gives  $\text{blk}(f)$ , equation (1) and the last clause, and a face embedding is determined by the directions it leaves free, so  $f_*$  determines  $f$ . That  $f_*$  is injective and  $\text{blk}(f)$  monotone is an induction on the length of  $b$ : by [PZ21, Lem. 2.11] a map into  $I^m \vee \vee b'$  splits over a splitting of  $a$ , and at each glued vertex the two halves have disjoint images, since the coordinates a face embedding leaves free before the vertex are already 1 there. Finally  $f$  preserves altitude [PZ21, Prop. 2.7], so  $N = M$  and the injection  $f_*$  is a permutation.  $\square$

**Presented categories.** A **2-polygraph**  $\mathcal{P}$  consists of a set  $V$  of 0-cells, families  $\mathcal{P}_1(x, y)$  and  $\mathcal{P}_2(x, y)$  for  $x, y \in V$ , and two maps  $\text{src}, \text{tgt} : \mathcal{P}_2(x, y) \rightarrow \{\text{paths } x \rightarrow y\}$  into the free category on the 1-cells;  $\text{src}(\alpha)$  and  $\text{tgt}(\alpha)$  are the **words** of  $\alpha$ . A morphism of polygraphs is a map in each dimension commuting with  $\text{src}$  and  $\text{tgt}$  on the nose, and **Poly** is the resulting category. The category  $\langle \mathcal{P} \rangle$  **presented by**  $\mathcal{P}$  is the free category on the 1-cells modulo the congruence generated by  $\text{src}(\alpha) = \text{tgt}(\alpha)$ , with quotient functor  $q$ .

**Proposition 2.2** ([Ara+23, Ch. 2]). *Let  $\mathcal{P}$  be a 2-polygraph and  $C$  a category.*

1. *To give a functor  $\langle \mathcal{P} \rangle \rightarrow C$  is to give an object of  $C$  for each 0-cell and an arrow of  $C$  for each 1-cell, subject to the two words of every 2-cell being sent to the same arrow.*
2. *Two functors out of  $\langle \mathcal{P} \rangle$  that agree on 0-cells and 1-cells are equal.*
3. *A morphism of polygraphs bijective in every dimension is an isomorphism.*

**Proposition 2.3** ([GZ67, Ch. I]). *Let  $W$  be a class of morphisms of a category  $C$ . There is a functor  $Q : C \rightarrow C[W^{-1}]$  inverting  $W$  through which every functor inverting  $W$  factors uniquely, and two functors out of  $C[W^{-1}]$  that agree after  $Q$  are equal.*

► *One sentence: every arrow of  $\text{Br}(K)$  written down below is a composite  $Q(g)Q(m)^{-1}$ , and every identity used about them is an instance of  $Q$  being a functor.*

**Newman's lemma.** Write  $\text{Hasse}(P)$  for the quiver of covers of a poset  $P$ . A map of quivers  $\text{Hasse}(P) \rightarrow C$  into a category extends to paths by composition, and we read it on paths without further comment.

**Theorem 2.4** (coherent Newman's lemma; [Hue80], [Ara+23, Ch. 4]). *Let  $P$  be a finite poset and  $\varphi : \text{Hasse}(P) \rightarrow C$  a map of quivers. Suppose that for every pair of distinct covers  $x \triangleleft y_1$ ,  $x \triangleleft y_2$  and every common upper bound  $z$  of  $y_1$  and  $y_2$  there are  $j \leq z$  and paths  $p_i$  from  $y_i$  to  $j$  with  $\varphi(x \triangleleft y_1) \varphi(p_1) = \varphi(x \triangleleft y_2) \varphi(p_2)$ . Then  $\varphi$  sends parallel paths to equal arrows, so it extends uniquely to a functor  $P \rightarrow C$ ; and a functor out of  $P$  is determined by its values on covers.*

### 3 Squares, crossings, and the braiding

► *One paragraph: a square is a pair of events of one block, and the whole construction runs on the question of which squares a map reverses.*

Throughout,  $a$  and  $b$  are shapes on  $N$  events.

**Definition 3.1.** A **square** of  $a$  is a pair  $s = (p, q)$  of events with  $p < q$  and  $\text{blk}(p) = \text{blk}(q)$ ; write  $\text{Sq}(a)$  for the set of them, so  $|\text{Sq}(a)| = \sum_i \binom{a_i}{2}$ . We write  $\text{Sq}(d)$  for the squares of a chain's shape, and  $f_*$  also for the induced injection  $\text{Sq}(a) \rightarrow \text{Sq}(b)$ ,  $(p, q) \mapsto (f_*p, f_*q)$ , well defined because  $f_*$  is increasing on blocks (Proposition 2.1).

**Definition 3.2.** The **crossings** of a map of serial wedges  $f : \bigvee a \rightarrow \bigvee b$  are the inversions of the relabelling it performs:

$$\text{Cr}(f) := \{ (p, q) : p < q \text{ and } f_*^{-1}(q) < f_*^{-1}(p) \}.$$

**Lemma 3.3.** *The map  $f_*^{-1}$  preserves the order of any two events lying in distinct blocks of  $b$ . Hence  $\text{Cr}(f) \subseteq \text{Sq}(b)$ . A square  $(p, q)$  of  $b$  lies in the image of  $f_*$  if and only if  $f_*^{-1}p$  and  $f_*^{-1}q$  lie in one block of  $a$ .*

*Proof.* Let  $u < v$  lie in distinct blocks of  $b$ , so  $\text{blk}_b(u) < \text{blk}_b(v)$ , and put  $w = f_*^{-1}u$ ,  $w' = f_*^{-1}v$ . Then (1) gives  $\text{blk}(f)(\text{blk}_a w) < \text{blk}(f)(\text{blk}_a w')$ , so  $\text{blk}_a(w) < \text{blk}_a(w')$  as  $\text{blk}(f)$  is monotone, so  $w < w'$  as blocks are intervals. An inversion of  $f_*^{-1}$  therefore has both events in one block of  $b$ . For the last claim, a square in the image pulls back to a square, whose events share a block; conversely if the two preimages share a block then,  $f_*$  being increasing there, they form a square mapping to  $(p, q)$ .  $\square$

**Lemma 3.4.** *For  $f : \bigvee a \rightarrow \bigvee b$  and  $g : \bigvee b \rightarrow \bigvee c$ ,*

$$\text{Cr}(gf) = g_*(\text{Cr } f) \sqcup \text{Cr}(g),$$

*and  $\text{Cr}(\text{id}) = \emptyset$ .*

*Proof.* Let  $s$  be a square of  $c$ . If  $s = g_*(t)$  then  $s \notin \text{Cr}(g)$ , the two events of  $t$  sharing a block (Lemma 3.3) — which also gives disjointness — and  $(gf)_*^{-1}$  agrees with  $f_*^{-1}$  on the events of  $t$ , so  $s \in \text{Cr}(gf)$  if and only if  $t \in \text{Cr}(f)$ . If  $s$  is not in the image of  $g_*$  then it is not in  $g_*(\text{Cr } f)$ , and its two events pull back along  $g_*$  into distinct blocks of  $b$ , so the second part of Lemma 3.3 applied to  $f$  gives  $s \in \text{Cr}(gf)$  if and only if  $s \in \text{Cr}(g)$ .  $\square$

**Definition 3.5.** A refinement  $f$  is **order preserving** when  $\text{Cr}(f) = \emptyset$ . Write  $W = W_K$  for the class of order-preserving refinements of  $\text{Ch}(K)$ , and

$$\text{Br}(K) := \text{Ch}(K)^{\text{op}}[(W^{\text{op}})^{-1}].$$

Write  $Q : \text{Ch}(K)^{\text{op}} \rightarrow \text{Br}(K)$  for the localization functor of Proposition 2.3. We leave the  $^{\text{op}}$  on morphisms implicit: for a refinement  $u : c \rightarrow d$  we write  $Q(u) : Q(d) \rightarrow Q(c)$ , so  $Q$  reverses composition.

**Proposition 3.6.**  $\text{Br}(K) \simeq (\text{Ch}(K)[W^{-1}])^{\text{op}}$ .

*Proof.* A functor  $\text{Ch}(K)^{\text{op}} \rightarrow D$  inverting  $W^{\text{op}}$  is a functor  $\text{Ch}(K) \rightarrow D^{\text{op}}$  inverting  $W$ , and conversely. So  $(\text{Ch}(K)[W^{-1}])^{\text{op}}$  has the universal property Proposition 2.3 gives  $\text{Br}(K)$ .  $\square$

**Proposition 3.7.** *A refinement  $f$  is order preserving if and only if  $f_* = \text{id}_{[N]}$ . Hence  $W$  contains the identities;  $gf \in W$  if and only if  $f \in W$  and  $g \in W$ ; any two order-preserving refinements with the same source and target are equal; and  $\text{Ch}(u)$  carries  $W_K$  into  $W_L$ , so it descends to  $\text{Br}(u) : \text{Br}(K) \rightarrow \text{Br}(L)$ , strictly functorially in  $u$ .*

*Proof.* A permutation with no inversions is the identity, and  $\text{Cr}(f)$  is the inversion set of  $f_*^{-1}$ , so  $\text{Cr}(f) = \emptyset$  if and only if  $f_*^{-1} = \text{id}$ , that is  $f_* = \text{id}$ . The consequences follow: the second from Lemma 3.4 with  $g_*$  injective, the third because  $f_*$  determines  $f$  (Proposition 2.1), and the last because  $\text{Ch}(u)$  changes neither shape nor wedge map, with the descent given by Proposition 2.3.  $\square$

► *One or two sentences reading Proposition 3.7 back: an order-preserving refinement is one that only glues adjacent blocks, the elementary ones being  $\bigvee[\dots, p, q, \dots] \rightarrow \bigvee[\dots, p + q, \dots]$ .*

## 4 The Artin polygraph of $K$

► *Frame the section: the runs over a chain form a poset; its covers are the degree-one chains over that chain, and the rank-two intervals are the degree-two ones. That is the whole polygraph.*

Fix  $K$ , and let  $d$  be a chain of  $K$  on  $N$  events.

### 4.1 Runs over a chain

**Definition 4.1.** A **run over  $d$**  is a refinement  $x : X \rightarrow d$  whose source is a run; write  $\mathcal{R}(d)$  for the set of these.

Such a refinement is determined by its map of wedges, the classifying map of its source being forced, so  $\mathcal{R}(d)$  is the set of runs of  $\bigvee a$ . That source has shape  $[1, \dots, 1]$ , so  $x_*$  enumerates the events of  $d$ ; write  $p <_x q$  when  $x_*$  reaches  $p$  before  $q$ . Then  $\text{Cr}(x)$  is the inversion set of  $<_x$ .

**Lemma 4.2.** *If  $\text{blk}(p) < \text{blk}(q)$  then  $p <_x q$ ; hence  $\text{blk}$  is monotone for  $<_x$ , with interval fibres, and  $x$  is exactly a choice of linear order on each block. In particular  $\text{Cr}$  is injective on  $\mathcal{R}(d)$ .*

*Proof.* The blocks of the source are singletons, so  $\text{blk} \circ x_* = \text{blk}(x)$ , monotone by Proposition 2.1. A monotone surjection out of a chain has interval fibres, here the blocks of  $d$ ; the only freedom left is the order inside each. A linear order is determined by its inversion set, and  $\text{Cr}(x)$  restricts on each block to the inversion set there.  $\square$

► *Note that for a single block  $\text{Cr}$  is the classical inversion set and inclusion of them is the right weak order [BB05, §3.1]; the order theory below is that statement, block by block.*

We order  $\mathcal{R}(d)$  by  $x \leq y : \iff \text{Cr}(x) \subseteq \text{Cr}(y)$ , and write  $\perp_d$  and  $\top_d$  for the runs taking each block in event order and in reverse, its least and greatest elements.

**Proposition 4.3.** *Let  $q : o \rightarrow d$  be a refinement. Composition with  $q$  is an order embedding  $\mathcal{R}(o) \rightarrow \mathcal{R}(d)$ , and*

$$\text{Cr}(qx) = q_*(\text{Cr}(x)) \sqcup \text{Cr}(q).$$

Moreover  $q \perp_o = \perp_d$  if and only if  $q \in W$ ; for  $x \in \mathcal{R}(d)$  the conditions  $x \in W$ ,  $\text{Cr}(x) = \emptyset$  and  $x = \perp_d$  agree;  $\perp_X = \text{id}_X$  for a run  $X$ ; and  $\text{dom } \perp_d = \text{dom } \perp_c$  whenever  $c \rightarrow d$  lies in  $W$ .

*Proof.* The formula is Lemma 3.4. Since  $q_*$  is injective with image disjoint from  $\text{Cr}(q)$  (Lemma 3.3),  $qx \leq qy$  if and only if  $x \leq y$ , and  $\text{Cr}(q \perp_o) = \text{Cr}(q)$ . The remaining equivalences are Definition 3.5 and the injectivity of  $\text{Cr}$  (Lemma 4.2), and the last clause is the formula applied to  $c \rightarrow d$ .  $\square$

## 4.2 Steps, covers, and cells

A square  $s$  is a **step** of  $x \in \mathcal{R}(d)$  when its two events are consecutive for  $<_x$ ; the **flip**  $x^s$  is  $x$  with those two swapped.

**Lemma 4.4.**  $\text{Cr}(x^s) = \text{Cr}(x) \triangle \{s\}$ , and  $x$  has exactly  $\deg d$  steps.

*Proof.* Swapping a consecutive pair changes the order of that pair and of no other. The steps are the covers of  $<_x$  lying inside a block;  $<_x$  is a chain on  $N$  elements and  $\text{blk}$  is a monotone surjection from it onto the  $r$  blocks (Lemma 4.2), and such a map collapses exactly  $N - r = \deg d$  covers.  $\square$

**Lemma 4.5.** If  $x < z$  in  $\mathcal{R}(d)$  then  $\text{Cr}(z)$  contains a step of  $x$  that  $\text{Cr}(x)$  does not. Consequently  $x < y$  exactly when  $\text{Cr}(y) = \text{Cr}(x) \cup \{s\}$  for such a step  $s$ , and then  $y = x^s$ .

*Proof.* If  $z$  agreed with  $x$  on every pair covering for  $<_x$ , then  $<_x$  would be contained in  $<_z$  by transitivity, and two linear orders so related are equal, so  $z = x$ . So some covering pair  $p <_x q$  has  $q <_z p$ . Its two events share a block, since otherwise Lemma 4.2 would make every run fire  $p$  first, so the pair is a step  $s$  of  $x$ . If  $s = (p, q)$  then  $s \in \text{Cr}(z) \setminus \text{Cr}(x)$ , as wanted; if  $s = (q, p)$  then  $s \in \text{Cr}(x) \subseteq \text{Cr}(z)$ , which says  $p <_z q$ , a contradiction.

Given such an  $s$  we have  $x \leq x^s \leq y$ , forcing  $y = x^s$ ; conversely if  $\text{Cr}(y) = \text{Cr}(x) \cup \{s\}$  then any  $x \leq w \leq y$  has  $\text{Cr}(x) \subseteq \text{Cr}(w) \subseteq \text{Cr}(y)$ , so  $w$  is  $x$  or  $y$  by injectivity (Lemma 4.2).  $\square$

Say that a set  $S$  of steps of  $x$  **ris**es from  $x$  when  $S \cap \text{Cr}(x) = \emptyset$ .

**Proposition 4.6.** Let  $x \in \mathcal{R}(d)$ . Sending a factorisation  $X \rightarrow m \rightarrow d$  of  $x$  to the set of steps of  $x$  whose two events share a block of  $m$  is a bijection onto the sets of steps of  $x$ . Write  $m_S$  for the middle it sends to  $S$ , and  $\iota_S : X \rightarrow m_S$ ,  $\pi_S : m_S \rightarrow d$  for the two legs. Then  $\deg m_S = |S|$ , and  $\iota_S$  is order preserving whenever  $S$  rises from  $x$ .

*Proof.* A middle  $m$  makes the  $N$  events into its blocks, which by Lemma 4.2 applied to  $\iota$  are intervals of  $<_x$ , and  $\pi$  requires each such interval to lie inside one block of  $d$ ; conversely such an interval partition arises from the chain whose blocks are its parts, in order. The factorisation is determined by that shape: a map of wedges is determined by its action on events and injective there (Proposition 2.1), hence a monomorphism, while the first leg is forced by which events fall in which part. An interval partition of a chain is recorded by the set of covers it does not cut; those covers lie inside a block, so they are steps; and  $\deg m$  is  $N$  minus the number of parts, which is the number of uncut covers.

Let  $S$  rise from  $x$ . Since  $\iota_S$  is a run over  $m_S$ , its crossings are the squares of  $m_S$  that  $x$  reverses, that is the pairs inside one part that  $x$  reverses. The covers of  $<_x$  inside a part all lie in  $S$ , hence none is reversed, so  $<_x$  agrees with the event order on consecutive pairs of the part and hence, by transitivity, on every pair of it. So  $\text{Cr}(\iota_S) = \emptyset$ .  $\square$

**Definition 4.7.** For runs  $X, Y$  of  $K$ , an  $n$ -cell from  $X$  to  $Y$  is a chain  $e$  with  $\deg e = n$ ,  $\text{dom } \perp_e = X$  and  $\text{dom } \top_e = Y$ .

► *One sentence: so the  $n$ -cells are the chains of degree  $n$ , sorted by the two runs each spans, and a 0-cell is a run.*

**Lemma 4.8.** Let  $S$  rise from  $x \in \mathcal{R}(d)$ . Then

$$\text{Cr}(\pi_S \top_{m_S}) = \text{Cr}(x) \cup \pi_{S*}(\text{Sq}(m_S)),$$

and every  $z \geq x$  with  $S \subseteq \text{Cr}(z)$  satisfies  $\pi_{S*}(\text{Sq}(m_S)) \subseteq \text{Cr}(z)$ . So  $\pi_S \top_{m_S}$  is the least element of  $\mathcal{R}(d)$  above  $x$  reversing all of  $S$ .

*Proof.* Proposition 4.3 gives  $\text{Cr}(\pi_S \top_{m_S}) = \pi_{S*} \text{Sq}(m_S) \cup \text{Cr}(\pi_S)$ , and Lemma 3.4 applied to  $\pi_S \iota_S = x$  with  $\text{Cr}(\iota_S) = \emptyset$  gives  $\text{Cr}(\pi_S) = \text{Cr}(x)$ .

For the second claim list a part as  $p_1 < \dots < p_k$ ; as in the previous proof this is also the restriction of  $<_z$  to it, so each  $(p_i, p_{i+1})$  lies in  $S$  and is reversed by  $z$ . Then  $p_k <_z \dots <_z p_1$ , so  $z$  reverses every pair of the part.  $\square$

**Corollary 4.9.** Let  $x \leq y$  in  $\mathcal{R}(d)$ , with step  $s$ . Then  $\gamma(x \leq y) := m_{\{s\}}$  is a 1-cell from  $\text{dom } x$  to  $\text{dom } y$ , with  $\pi_{\{s\}} \perp_{m_{\{s\}}} = x$  and  $\pi_{\{s\}} \top_{m_{\{s\}}} = y$ ; and every 1-cell  $\alpha$  is  $\gamma$  of the unique cover  $\perp_\alpha \leq \top_\alpha$  of  $\mathcal{R}(\alpha)$ . So  $\gamma$ , together with  $x \mapsto \text{dom } x$  on vertices, is a map of quivers from  $\text{Hasse}(\mathcal{R}(d))$  to the quiver of 1-cells, and  $\gamma(qe) = \gamma(e)$  for every refinement  $q : o \rightarrow d$  and every cover  $e$  of  $\mathcal{R}(o)$ .

*Proof.*  $m_{\{s\}}$  has degree 1 and  $\text{dom } \perp_{m_{\{s\}}} = \text{dom } x$  by Proposition 4.6; by Lemma 4.8 the image of its top is the least element above  $x$  that reverses  $s$ , which is  $x^s = y$ . Conversely a chain  $\alpha$  of degree 1 has one square, so  $\mathcal{R}(\alpha)$  is the two-element poset  $\perp_\alpha \leq \top_\alpha$ , and the factorisation through its one step is  $\alpha$ . For the last claim the step of  $qe$  is  $q_*$  of the step of  $e$  (Proposition 4.3), and  $q_*$  carries  $<_x$  to  $<_{qx}$ , so the two factorisations cut the same partition and have the same middle.  $\square$

### 4.3 The attachment of a 2-cell

**Proposition 4.10.** Let  $\deg e = 2$ . Then either  $e$  has one block of dimension 3 and  $|\text{Sq}(e)| = 3$ , or  $e$  has two blocks of dimension 2 and  $|\text{Sq}(e)| = 2$ . In both cases  $\mathcal{R}(e)$  has exactly two maximal chains from  $\perp_e$  to  $\top_e$ , each of length  $|\text{Sq}(e)|$ , and they leave  $\perp_e$  by different covers. Write  $\text{src}(e)$  and  $\text{tgt}(e)$  for the words of 1-cells that  $\gamma$  reads off them, labelled so that  $\text{src}(e)$  begins with the least square for the lexicographic order on  $\text{Sq}(e)$ , by the smaller event and then the larger.

*Proof.*  $\deg e = 2$  forces the block dimensions above 1 to be  $\{3\}$  or  $\{2, 2\}$ , and Definition 3.1 counts  $\binom{3}{2} = 3$  squares in the first case and  $2\binom{2}{2} = 2$  in the second. A run over  $e$  is a linear order on each block (Lemma 4.2) and each cover swaps a consecutive pair (Lemma 4.5).

Two blocks of dimension 2 therefore give four runs, a Boolean square on the two squares. One block of dimension 3, on events  $p < q < r$ , gives six, and the covers are

$$pqr \leq qpr \leq qrp \leq rqp, \quad pqr \leq prq \leq rpq \leq rqp,$$

a hexagon. Either way the two displayed chains are the only maximal ones and leave  $\perp_e$  by different covers.  $\square$

In both cases  $\text{src}(e)$  reverses the squares of  $e$  in lexicographic order and  $\text{tgt}(e)$  in the reverse order.<sup>1</sup>

The two species are drawn in Figure 1 and Figure 2.

**Definition 4.11.** The **Artin polygraph**  $\text{Art}(K)$  has the runs of  $K$  as 0-cells. Its 1-cells and 2-cells from  $X$  to  $Y$  are the 1-cells and 2-cells from  $X$  to  $Y$  of Definition 4.7, that is the chains of degree 1 and 2 spanning those two runs, and each 2-cell  $e$  is attached along the pair of words  $\text{src}(e)$ ,  $\text{tgt}(e)$  of Proposition 4.10.

► *One paragraph on the name:  $\text{Art}(K)$  is built from  $K$  alone, with no braid theory in it, and Proposition 4.10 already exhibits its 2-cells as braid and commutation relations; Theorem 7.5 is what makes the name literal.*

**Proposition 4.12.** Read in  $\langle \text{Art}(K) \rangle$ , the map of quivers  $\gamma$  satisfies the hypothesis of Theorem 2.4 on  $\mathcal{R}(d)$ . So it extends uniquely to a functor  $\Gamma_d : \mathcal{R}(d) \rightarrow \langle \text{Art}(K) \rangle$ , and  $\Gamma_d(qx) = \Gamma_o(x)$  for every refinement  $q : o \rightarrow d$  and every  $x \in \mathcal{R}(o)$ .

*Proof.* Let  $x \leq y_1$ ,  $x \leq y_2$  be distinct covers with steps  $s_1 \neq s_2$ , and  $z$  a common upper bound. Then  $S = \{s_1, s_2\}$  rises from  $x$ , so  $m_S$  is a chain of degree 2, that is a 2-cell (Proposition 4.6), and  $j := \pi_S \top_{m_S}$  satisfies  $j \leq z$  (Lemma 4.8). By Proposition 4.10 the two maximal chains of  $\mathcal{R}(m_S)$  are  $\text{src}(m_S)$  and  $\text{tgt}(m_S)$ , and they leave  $\perp_{m_S}$  by its two covers. Composition with  $\pi_S$  carries  $\perp_{m_S}$  to  $x$  and  $\top_{m_S}$  to  $j$ , carries covers to covers (Proposition 4.3, Lemma 4.5) and keeps the 1-cell each cover reads as (Corollary 4.9). So the two chains become two paths from  $x$  to  $j$ , leaving by  $y_1$  and by  $y_2$  and reading as  $\text{src}(m_S)$  and  $\text{tgt}(m_S)$ , which are equal in  $\langle \text{Art}(K) \rangle$ .

For the last claim, composition with  $q$  is monotone, so  $x \mapsto \Gamma_d(qx)$  and  $\Gamma_o$  are functors out of  $\mathcal{R}(o)$ ; they agree on covers by Corollary 4.9, hence agree, by the last clause of Theorem 2.4.  $\square$

## 5 The presentation

► *Frame the proof as a comparison of two universal properties: a functor out of  $\langle \text{Art}(K) \rangle$  is an arrow for each degree-one chain identifying the two words of every degree-two chain (Proposition 2.2), and a functor out of  $\text{Br}(K)$  is a functor on  $\text{Ch}(K)^{\text{op}}$  inverting  $W$  (Proposition 2.3). Each is built from the other.*

Fix  $K$  throughout this section, and write  $E$ ,  $\Theta$ ,  $\Phi$  for the functors called  $E_K$ ,  $\Theta_K$ ,  $\Phi_K$  elsewhere.

An  $n$ -cell  $\alpha$  from  $X$  to  $Y$  comes with the two refinements  $\perp_\alpha : X \rightarrow \alpha$  and  $\top_\alpha : Y \rightarrow \alpha$ , of which the first is order preserving; since  $Q$  inverts  $\perp_\alpha$ , we may set

$$E(X) := Q(X), \quad E(\alpha) := Q(\top_\alpha) \circ Q(\perp_\alpha)^{-1} : Q(X) \rightarrow Q(Y).$$

**Lemma 5.1.** Let  $d$  be a chain and  $z \in \mathcal{R}(d)$ . Every path from  $\perp_d$  to  $z$  in  $\text{Hasse}(\mathcal{R}(d))$  reads, under  $E \circ \gamma$ , as  $Q(z) \circ Q(\perp_d)^{-1}$ . Consequently  $E$  extends to a functor  $\langle \text{Art}(K) \rangle \rightarrow \text{Br}(K)$ .

<sup>1</sup>The formalisation [Sto26] works with the sets  $\text{Cr}(x)$  in place of the orders  $<_x$ : `reversedIso` identifies  $\mathcal{R}(d)$  with the subsets of  $\text{Sq}(d)$  that are *biclosed*, meaning that no block's induced tournament has a directed 3-cycle, and it takes the last sentence above as the definition of the two words. The four and the six runs are `card_biclosed_of_isEmpty_triangles` and `card_biclosed_of_triangle`, and this proposition is derived twice there: as `eq_srcPath_or_tgtPath_of_triangles`, from `subsingleton_triangles` and `biclosed_iff_of_triangle`, and as `eq_srcPath_or_tgtPath`, from 2-regularity of  $\text{Hasse}(\mathcal{R}(e))$ .

*Proof.* Induct on the length of the path, the empty path reading as the identity. For a cover  $x \leq y$  with step  $s$ , put  $\beta := \gamma(x \leq y)$  and  $q := \pi_{\{s\}}$ , so  $x = q \perp_\beta$  and  $y = q \top_\beta$  (Corollary 4.9), so  $Q(x) = Q(\perp_\beta)Q(q)$ . The middle two factors of

$$E(\beta) \circ Q(x) \circ Q(\perp_d)^{-1} = Q(\top_\beta) \circ Q(\perp_\beta)^{-1} \circ Q(\perp_\beta) \circ Q(q) \circ Q(\perp_d)^{-1}$$

cancel, leaving  $Q(\top_\beta)Q(q)Q(\perp_d)^{-1} = Q(y)Q(\perp_d)^{-1}$ .

The two words of a 2-cell  $e$  are both paths from  $\perp_e$  to  $\top_e$  (Proposition 4.10), hence have equal image, so Proposition 2.2(i) gives the extension.  $\square$

**Lemma 5.2.** *Setting  $\Theta(d) := \text{dom } \perp_d$  and, for a refinement  $u : c \rightarrow d$ ,*

$$\Theta(u) := \Gamma_d(\perp_d \leq u \perp_c) : \text{dom } \perp_d \rightarrow \text{dom } \perp_c,$$

*defines a functor  $\Theta : \text{Ch}(K)^{\text{op}} \rightarrow \langle \text{Art}(K) \rangle$ ; it sends  $u$  to an identity whenever  $u$  is order preserving, so it descends uniquely to  $\Phi : \text{Br}(K) \rightarrow \langle \text{Art}(K) \rangle$  with  $\Phi Q = \Theta$ .*

*Proof.* The stated target is the right one:  $u \perp_c$  lies in  $\mathcal{R}(d)$  with source  $\text{dom } \perp_c$ . Let  $v : b \rightarrow c$  and  $u : c \rightarrow d$ . Composition with  $u$  is monotone (Proposition 4.3), so  $\perp_d \leq u \perp_c \leq uv \perp_b$ , and  $\Gamma_d$  being a functor,  $\Theta(uv) = \Gamma_d(u \perp_c \leq uv \perp_b) \circ \Theta(u)$ ; the left factor is  $\Gamma_c(\perp_c \leq v \perp_b) = \Theta(v)$  by the last clause of Proposition 4.12. If  $u$  is order preserving then  $u \perp_c = \perp_d$  (Proposition 4.3), so the comparison is an identity. The descent is Proposition 2.3.  $\square$

**Theorem 5.3.** *For every bipointed precubical set  $K$ , the functor  $E$  is an equivalence  $\langle \text{Art}(K) \rangle \rightarrow \text{Br}(K)$ , with  $\Phi$  inverse to it.*

*Proof.* On a 0-cell,  $\Phi(E(X)) = \Theta(X) = \text{dom } \perp_X = X$  (Proposition 4.3). On a 1-cell  $\alpha$ , applying  $\Phi$  to the defining composite gives  $\Phi(E(\alpha)) = \Theta(\top_\alpha) \circ \Theta(\perp_\alpha)^{-1}$ , whose second factor is an identity because  $\perp_\alpha$  is order preserving (Lemma 5.2). Since  $\top_\alpha \perp_Y = \top_\alpha$  and  $\mathcal{R}(\alpha)$  is the two-element poset  $\perp_\alpha \leq \top_\alpha$ ,

$$\Theta(\top_\alpha) = \Gamma_\alpha(\perp_\alpha \leq \top_\alpha) = q(\gamma(\perp_\alpha \leq \top_\alpha)) = q(\alpha)$$

by Corollary 4.9. So  $\Phi E$  agrees with the identity on 0- and 1-cells, hence is the identity (Proposition 2.2(ii)).

For the other composite, Lemma 5.1 gives  $E(\Theta(u)) = Q(u \perp_c)Q(\perp_d)^{-1} = Q(\perp_c) \circ Q(u) \circ Q(\perp_d)^{-1}$ , which is  $Q(u)$  conjugated by the isomorphisms  $Q(c) \cong Q(\text{dom } \perp_c)$  that  $Q$  makes of the order-preserving refinements  $\perp_c$ . So  $E\Theta \cong Q$ , and  $E\Phi \cong \text{id}$  by uniqueness of descent (Proposition 2.3).  $\square$

► *One sentence: the comparison is an equivalence and not an isomorphism because  $\text{Br}(K)$  is not skeletal — a chain and the run below it become isomorphic there without being equal — so the runs are a skeleton of  $\text{Br}(K)$ .*

## 6 Naturality

**Theorem 6.1.** *Sending  $K$  to  $\text{Art}(K)$ , and  $f : K \rightarrow L$  to  $\text{Ch}(f)$  on cells in every dimension, is a functor  $\square_* \rightarrow \mathbf{Poly}$ ; and for every  $f : K \rightarrow L$ ,*

$$E_L \circ \langle \text{Art}(f) \rangle = \text{Br}(f) \circ E_K : \langle \text{Art}(K) \rangle \longrightarrow \text{Br}(L).$$

*Proof.* A run over a chain  $e$  is determined by its map of wedges (Definition 4.1), and  $\text{Ch}(f)e$  has the same shape and the same maps of wedges into it, so  $\mathcal{R}(e) = \mathcal{R}(\text{Ch}(f)e)$  as ordered sets. Crossings, steps, flips and  $\gamma$  (Definition 3.2, Corollary 4.9) are read off the map of wedges alone, so  $\text{Ch}(f)$  preserves  $\perp$ ,  $\top$ , cells, steps, covers and  $\gamma$ ; and it carries the lexicographic order on  $\text{Sq}(e)$  to the lexicographic order on  $\text{Sq}(\text{Ch}(f)e) = \text{Sq}(e)$ , so it preserves  $\text{src}$  and  $\text{tgt}$  letter by letter. Hence  $\text{Art}(f)$  is a morphism of polygraphs, functorially in  $f$  because  $\text{Ch}$  is a functor.

Both composites send a run  $X$  to  $Q(\text{Ch}(f)X)$ . On a 1-cell  $\alpha$ ,  $\text{Br}(f)$  is the descent of the functor induced by  $\text{Ch}(f)$ , so it sends  $E_K(\alpha)$  to  $Q(\text{Ch}(f)\top_\alpha) \circ Q(\text{Ch}(f)\perp_\alpha)^{-1}$ . Here  $\text{Ch}(f)\perp_\alpha$  is an order-preserving refinement into  $\text{Ch}(f)\alpha$  out of a run, hence is  $\perp_{\text{Ch}(f)\alpha}$  (Proposition 4.3), and likewise  $\text{Ch}(f)\top_\alpha = \top_{\text{Ch}(f)\alpha}$ . So both composites send  $\alpha$  to  $E_L(\text{Ch}(f)\alpha)$ , and Proposition 2.2(ii) applies.  $\square$

► *One sentence: the square is an equality, not an isomorphism, because an order-preserving refinement is determined by its source and target (Proposition 3.7), so transporting the two legs of a cell involves no choice. The tree states it as an isomorphism pinned at the identity, which is this equality read through eqToIso.*

## 7 The point

Since  $Z$  is terminal, every serial wedge has exactly one bipointed map to it and every map of wedges commutes over it, so  $\text{Ch}(Z)$  is the category of shapes and maps of wedges. Write  $Z_N$  for the run of  $Z$  on  $N$  events. Both  $\perp_e$  and  $\top_e$  have source a run on  $|e|$  events, so every cell of  $Z$  is a loop at  $Z_{|e|}$ .

For a shape  $a$  on  $N$  events put  $\text{cut}(a) := \{k \in [N-1] : \text{blk}_a(k) = \text{blk}_a(k+1)\}$ , and write  $\text{cut}(d)$  for the cuts of a chain's shape; for  $i \neq k$  in  $[N-1]$  let  $\text{cox}(i, k) := 3$  when  $|i - k| = 1$  and  $\text{cox}(i, k) := 2$  otherwise.

**Lemma 7.1.**  *$a \mapsto \text{cut}(a)$  is a bijection from the shapes on  $N$  events onto the subsets of  $[N-1]$ , with  $|\text{cut}(a)| = \deg a$ . Hence the  $n$ -cells at  $Z_N$  are the  $n$ -element subsets of  $[N-1]$ . A chain  $e$  of  $Z$  with  $\deg e = 2$  and  $\text{cut}(e) = \{i < k\}$  has  $|\text{Sq}(e)| = \text{cox}(i, k)$ .*

*Proof.* A shape on  $N$  events is a composition of  $N$  and  $\text{cut}(a)$  is the complement of its set of block boundaries in  $[N-1]$ , so the first claim is the classical correspondence [Sta12, §1.2] and  $|\text{cut}(a)| = N - r = \deg a$ . For the last claim,  $e$  has a block of dimension 3 exactly when three consecutive events share a block, that is exactly when  $k = i + 1$ ; then  $|\text{Sq}(e)| = 3$ , and otherwise  $|\text{Sq}(e)| = 2$ , by Proposition 4.10.  $\square$

Write  $\sigma_k$  for the 1-cell at  $Z_N$  with  $\text{cut} = \{k\}$ . For a step  $s$  of  $x$ , write  $\text{pos}_x(s)$  for the earlier of the two positions  $<_x$  gives the events of  $s$ ; so  $x$  passes  $s$  between positions  $\text{pos}_x(s)$  and  $\text{pos}_x(s) + 1$ .

**Lemma 7.2.** *Let  $d$  be a chain on  $N$  events and  $x \in \mathcal{R}(d)$ . Then:*

1.  $\text{pos}_x$  is a bijection from the steps of  $x$  onto  $\text{cut}(d)$ ;
2.  $\text{pos}_{x^s}(s) = \text{pos}_x(s)$ ;
3.  $\text{pos}_{\perp_d}(s) = p$ , where  $s = (p, q)$ .

*If  $d$  is a chain of  $Z$ , then a cover  $x \triangleleft y$  with step  $s$  has  $\gamma(x \triangleleft y) = \sigma_{\text{pos}_x(s)}$ .*

*Proof.* (i) The fibres of  $\text{blk}$  for  $<_x$  are consecutive intervals of lengths  $a_1, \dots, a_r$  (Lemma 4.2), independent of  $x$ , so the set of positions  $t$  with  $t$  and  $t+1$  in one block is  $\text{cut}(d)$  for every  $x$ . A step is a cover of  $<_x$  inside a block, so  $\text{pos}_x$  lands in  $\text{cut}(d)$ , and it is injective because distinct covers of a chain have distinct lower ends; both sides have  $\deg d$  elements (Lemma 4.4, Lemma 7.1). (ii) Flipping  $s$  exchanges its two times and changes no other. (iii)  $\perp_d$  fires the events in their own order.

For the last claim,  $\gamma(x < y) = m_{\{s\}}$ , whose blocks are the parts of  $<_x$  cut at every cover but  $s$  (Proposition 4.6); so  $m_{\{s\}}$  has one block of dimension 2, at positions  $\text{pos}_x(s)$  and  $\text{pos}_x(s) + 1$ , and edges elsewhere, so its cut set is  $\{\text{pos}_x(s)\}$ .  $\square$

**Proposition 7.3.** *Let  $e$  be a chain of  $Z$  with  $\deg e = 2$  and  $\text{cut}(e) = \{i < k\}$ . Then*

$$\text{src}(e) = \sigma_i \sigma_k \sigma_i \cdots, \quad \text{tgt}(e) = \sigma_k \sigma_i \sigma_k \cdots,$$

*both of length  $\text{cox}(i, k) = |\text{Sq}(e)|$ .*

*Proof.* A cover with step  $s$  reads as  $\sigma_{\text{pos}_x(s)}$  (Lemma 7.2), so it is enough to read the two chains of Proposition 4.10, and  $\perp_e$  passes  $(p, q)$  at  $p$  (Lemma 7.2(iii)).

If  $k > i + 1$  then  $e$  has blocks of dimension 2 at positions  $\{i, i + 1\}$  and  $\{k, k + 1\}$  (Lemma 7.1), with squares  $s_1$  and  $s_2$ . Flipping one leaves the positions of the other alone (Lemma 7.2(ii)), so the two chains read  $\sigma_i \sigma_k$  and  $\sigma_k \sigma_i$ .

If  $k = i + 1$  then  $e$  has one block of dimension 3, on the events at positions  $i, i + 1, i + 2$ . Along the two chains of Proposition 4.10 the swapped pair sits at positions  $i, i + 1, i$  and at  $i + 1, i, i + 1$ .

The lengths are  $|\text{Sq}(e)| = \text{cox}(i, k)$  (Lemma 7.1).  $\square$

**Definition 7.4.** **Artin's braid polygraph**  $\mathcal{A}$  has one 0-cell for each  $N \in \mathbb{N}$ ; at  $N$ , the loops  $\sigma_1, \dots, \sigma_{N-1}$ ; and, for each pair  $i < k$  in  $[N - 1]$ , one 2-cell with words  $\sigma_i \sigma_k \sigma_i \cdots$  and  $\sigma_k \sigma_i \sigma_k \cdots$ , both of length  $\text{cox}(i, k)$ .

► *One or two sentences: unwind Definition 7.4 to Artin's relations [Art47], so  $\langle \mathcal{A} \rangle$  has endomorphism monoid  $(B_N^+)^{\text{op}}$  at  $N$  and no arrows between distinct objects; then invoke Corollary 7.6's reversal to read it as **Braid**<sup>+</sup>.*

**Theorem 7.5.**  $\text{Art}(Z) \cong \mathcal{A}$  in **Poly**.

*Proof.* The runs of  $Z$  are the  $Z_N$  and every cell of  $Z$  is a loop, so both polygraphs are disjoint unions over  $N$  and it suffices to compare at each  $N$ . There the 1-cells are indexed by  $[N - 1]$  and the 2-cells by the pairs  $i < k$  (Lemma 7.1), and by Proposition 7.3 the words of the 2-cell at  $\{i < k\}$  are those of Definition 7.4. The resulting morphism of polygraphs is bijective in every dimension, hence an isomorphism (Proposition 2.2(iii)).  $\square$

**Corollary 7.6.**  $\text{Br}(Z) \simeq \text{Braid}^+$ .

*Proof.* Theorem 5.3 at  $K = Z$  and Theorem 7.5 give  $\text{Br}(Z) \simeq \langle \mathcal{A} \rangle$ , whose endomorphism monoid at  $N$  is  $B_N^+$  read with words composing left to right, that is  $(B_N^+)^{\text{op}}$ . Reversing words is an anti-automorphism of  $B_N^+$ , both Artin relations being invariant under reversal, so  $(B_N^+)^{\text{op}} \cong B_N^+$ .  $\square$

► *Closing paragraph: combining with Theorem 6.1, the terminal map gives  $\text{Br}(!_K) : \text{Br}(K) \rightarrow \text{Br}(Z)$ , computed cell by cell — a degree-one chain goes to  $\sigma_{\text{pos}}$  for the position at which its run passes its one step, a degree-two chain to the relation of its two positions. Say what this sharpens in [PZ21]. Note that Matsumoto’s theorem plays no role,  $B_N^+$  being taken here as Artin’s presentation.*

## References

- [Ara+23] Dimitri Ara, Albert Burroni, Yves Guiraud, Philippe Malbos, François Métayer, and Samuel Mimram. *Polygraphs: From Rewriting to Higher Categories*. 2023. arXiv: [2312.00429](#) [math.CT].
- [Art47] Emil Artin. “Theory of Braids”. In: *Annals of Mathematics* 48.1 (1947), pp. 101–126. DOI: [10.2307/1969218](#).
- [BB05] Anders Björner and Francesco Brenti. *Combinatorics of Coxeter Groups*. Vol. 231. Graduate Texts in Mathematics. Springer, 2005.
- [GGM15] Stéphane Gaussent, Yves Guiraud, and Philippe Malbos. “Coherent presentations of Artin monoids”. In: *Compositio Mathematica* 151.5 (2015), pp. 957–998. DOI: [10.1112/S0010437X14007842](#).
- [GZ67] Peter Gabriel and Michel Zisman. *Calculus of Fractions and Homotopy Theory*. Vol. 35. Ergebnisse der Mathematik und ihrer Grenzgebiete. Springer, 1967.
- [Hue80] Gérard Huet. “Confluent Reductions: Abstract Properties and Applications to Term Rewriting Systems”. In: *Journal of the ACM* 27.4 (1980), pp. 797–821. DOI: [10.1145/322217.322230](#).
- [PZ21] Jakub Paliga and Krzysztof Ziemiański. *Configuration spaces and directed paths on the final precubical set*. 2021. arXiv: [2103.05336](#) [math.AT].
- [Sta12] Richard P. Stanley. *Enumerative Combinatorics, Volume 1*. 2nd ed. Vol. 49. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2012.
- [Sto26] Zachary Stone. *Cubical: a Lean 4 formalisation of cube chains and their complexification*. 2026. URL: <https://github.com/zstone1/Cubical>.
- [Zie20] Krzysztof Ziemiański. “Spaces of directed paths on pre-cubical sets II”. In: *Journal of Applied and Computational Topology* 4 (2020), pp. 45–78. DOI: [10.1007/s41468-019-00040-z](#). arXiv: [1901.05206](#).