

↳

Some Basic Statistical Theorems

Review



Review

Given a **probability space** (Ω, \mathcal{F}, P) , and a **measurable space** (E, \mathcal{E}) , an E -valued **random variable** is a **measurable function** $X : \Omega \rightarrow E$.

⋮

Review


Given a **probability space** (Ω, \mathcal{F}, P) , and a **measurable space** (E, \mathcal{E}) , an E -valued **random variable** is a **measurable function** $X : \Omega \rightarrow E$.

$$\omega \in \Omega$$



Review


Given a **probability space** (Ω, \mathcal{F}, P) , and a **measurable space** (E, \mathcal{E}) , an E -valued **random variable** is a **measurable function** $X : \Omega \rightarrow E$.



⋮


Review

Given a **probability space** (Ω, \mathcal{F}, P) , and a **measurable space** (E, \mathcal{E}) , an E -valued **random variable** is a **measurable function** $X : \Omega \rightarrow E$.



Review


Given a **probability space** (Ω, \mathcal{F}, P) , and a **measurable space** (E, \mathcal{E}) , an E -valued **random variable** is a **measurable function** $X : \Omega \rightarrow E$.




$$\mathcal{F} = \sigma(\{\text{red}\}) = \{\Omega, \text{red}, \text{blue}, \emptyset\}$$

Review

Given a **probability space** (Ω, \mathcal{F}, P) , and a **measurable space** (E, \mathcal{E}) , an E -valued **random variable** is a **measurable function** $X : \Omega \rightarrow E$.



$$\mathcal{F} = \sigma(\{\text{green region}\}) = \{\Omega, \text{green region}, \text{blue region}, \emptyset\}$$



Review

Convergence:

- Sure ("pointwise")
- Almost Sure
- In Probability
- Weak ("in distribution"/"in law")

Review


Convergence:

- Sure ("pointwise")

$$X_n(\omega) \rightarrow X(\omega) \quad \forall \omega \in \Omega$$

- Almost Sure
- In Probability
- Weak ("in distribution"/"in law")

Review



Convergence:

- Sure ("pointwise")

$$X_n(\omega) \rightarrow X(\omega) \quad \forall \omega \in \Omega$$

- Almost Sure

$$P(\{\omega : X_n(\omega) \rightarrow X\}) = 1$$

- In Probability

- Weak ("in distribution"/"in law")

Review

Convergence:

- Sure ("pointwise") $X_n(\omega) \rightarrow X(\omega) \quad \forall \omega \in \Omega$
- Almost Sure $P(\{\omega : X_n(\omega) \rightarrow X\}) = 1$
- In Probability $P(\{\omega : |X_n(\omega) - X(\omega)| > \epsilon\}) \rightarrow 0 \quad \forall \epsilon > 0$
- Weak ("in distribution"/"in law")

Review

Convergence:

- Sure ("pointwise") $X_n(\omega) \rightarrow X(\omega) \quad \forall \omega \in \Omega$
- Almost Sure $P(\{\omega : X_n(\omega) \rightarrow X\}) = 1$
- In Probability $P(\{\omega : |X_n(\omega) - X(\omega)| > \epsilon\}) \rightarrow 0 \quad \forall \epsilon > 0$
- Weak ("in distribution"/"in law")

$F_{X_n}(\alpha) \rightarrow F_X(\alpha)$ for each continuity point.

Outline for Today

Outline for Today

1. Concentration Inequalities

Outline for Today

1. Concentration Inequalities
2. Proof of the weak law of large numbers and $\frac{1}{\sqrt{N}}$ convergence rates

Outline for Today

1. Concentration Inequalities

Break

2. Proof of the weak law of large numbers and $\frac{1}{\sqrt{N}}$ convergence rates

Outline for Today

1. Concentration Inequalities

Break

2. Proof of the weak law of large numbers and $\frac{1}{\sqrt{N}}$ convergence rates

3. Central limit theorem

4. Importance sampling


Concentration Inequalities

Concentration Inequalities

Intuition: If an r.v. has a finite variance, the probability that a random variable takes a value far from its mean should be small


Concentration Inequalities

Intuition: If an r.v. has a finite variance, the probability that a random variable takes a value far from its mean should be small



Concentration Inequalities

Intuition: If an r.v. has a finite variance, the probability that a random variable takes a value far from its mean should be small



Concentration inequalities take the form

$$P(X \geq t) \leq \phi(t)$$

where ϕ goes to zero (quickly) as $t \rightarrow \infty$

Concentration Inequalities

Concentration Inequalities

Markov's Inequality:

If $X \geq 0$, then

$$P(X \geq t) \leq \frac{E[X]}{t} \quad \forall t \geq 0$$

Proof:

$$E[X] = \int_{-\infty}^{\infty} x p(x) dx$$


$$= \int_0^{\infty} x p(x) dx$$

$$= \underbrace{\int_0^a x p(x) dx}_{\geq 0} + \int_a^{\infty} x p(x) dx$$

$$\geq \int_a^{\infty} x p(x) dx \geq \int_a^{\infty} a p(x) dx = a \underbrace{\int_a^{\infty} p(x) dx}_{= P(X \geq a)}$$

$$= a P(X \geq a)$$

$$\therefore \frac{E[X]}{a} \geq P(X \geq a)$$



Concentration Inequalities

Concentration Inequalities

Chebyshev's Inequality:

Let X be any real-valued random variable with $\text{Var}(X) < \infty$. Then

Concentration Inequalities

Chebyshev's Inequality:

Let X be *any* real-valued random variable with $\text{Var}(X) < \infty$. Then

$$P(|X - E[X]| \geq t) \leq \frac{\text{Var}(X)}{t^2}.$$


Proof: Recall $\text{Var}(X) = E[(X - E[X])^2]$

Use Markov's inequality on $(X - E[X])^2$

$$P((X - E[X])^2 \geq t^2) \leq \frac{E[(X - E[X])^2]}{t^2}$$

$$P(|X - E[X]| \geq t) \leq \frac{\text{Var}(X)}{t^2}$$

$$0.05 = \frac{\text{Var}(X)}{t^2}$$



Concentration Inequalities

Chebyshev's Inequality:


Let X be *any* real-valued random variable with $\text{Var}(X) < \infty$. Then

$$P(|X - E[X]| \geq t) \leq \frac{\text{Var}(X)}{t^2}.$$

Very general, but still loose

Concentration Inequalities

k	Min. % within k standard deviations of mean	Max. % beyond k standard deviations from mean
1	0%	100%
$\sqrt{2}$	50%	50%
1.5	55.56%	44.44%
2	75%	25%
$2\sqrt{2}$	87.5%	12.5%
3	88.8889%	11.1111%
4	93.75%	6.25%
5	96%	4%
6	97.2222%	2.7778%
7	97.9592%	2.0408%
8	98.4375%	1.5625%
9	98.7654%	1.2346%
10	99%	1%



Concentration Inequalities

k	Min. % within k standard deviations of mean	Max. % beyond k standard deviations from mean
1	0%	100%
$\sqrt{2}$	50%	50%
1.5	55.56%	44.44%
2	75%	25%
$2\sqrt{2}$	87.5%	12.5%
3	88.8889%	11.1111%
4	93.75%	6.25%
5	96%	4%
6	97.2222%	2.7778%
7	97.9592%	2.0408%
8	98.4375%	1.5625%
9	98.7654%	1.2346%
10	99%	1%

For Normal Distribution

Concentration Inequalities

k	Min. % within k standard deviations of mean	Max. % beyond k standard deviations from mean
1	0%	100%
$\sqrt{2}$	50%	50%
1.5	55.56%	44.44%
2	75%	25%
$2\sqrt{2}$	87.5%	12.5%
3	88.8889%	11.1111%
4	93.75%	6.25%
5	96%	4%
6	97.2222%	2.7778%
7	97.9592%	2.0408%
8	98.4375%	1.5625%
9	98.7654%	1.2346%
10	99%	1%

68% For Normal Distribution

95%

99%

Concentration Inequalities

Concentration Inequalities

Moment generating function: $M_X(t) \equiv E[e^{tX}]$

Concentration Inequalities

Moment generating function: $M_X(t) \equiv E[e^{tX}]$

Chernoff Bound: If the moment-generating function M_X exists, then

$$P(X \geq \underline{a}) \leq \frac{E[e^{tX}]}{e^{ta}} \quad \forall t > 0$$

$\forall a$

*infinite number
of bounds, one
for each t*

and

$$P(X \leq a) \leq \frac{E[e^{tX}]}{e^{ta}} \quad \forall t < 0$$

Proof: Apply Markov's inequality to e^{tX}

$$P(X \geq a) = P(e^{tX} \geq e^{ta}) \leq \frac{E[e^{tX}]}{e^{ta}}$$

Concentration Inequalities

Moment generating function: $M_X(t) \equiv E[e^{tX}]$

Chernoff Bound: If the moment-generating function M_X exists, then

$$P(X \geq a) \leq \frac{E[e^{tX}]}{e^{ta}} \quad \forall t > 0$$

and

$$P(X \leq a) \leq \frac{E[e^{tX}]}{e^{ta}} \quad \forall t < 0$$

Tighter than Markov and Chebyshev

Concentration Inequalities

Concentration Inequalities

Name	Requirements	Bound

Concentration Inequalities

Name	Requirements	Bound
Markov	$X \geq 0$, $E[X]$ exists	$P(X \geq t) \leq \frac{E[X]}{t} \quad \forall t \geq 0$

Concentration Inequalities

Name	Requirements	Bound
Markov	$X \geq 0, E[X]$ exists	$P(X \geq t) \leq \frac{E[X]}{t} \quad \forall t \geq 0$
Chebyshev	$\text{Var}(X) < \infty$	$P(X - E[X] \geq t) \leq \frac{\text{Var}(X)}{t^2}$


Concentration Inequalities

Name	Requirements	Bound
Markov	$X \geq 0, E[X]$ exists	$P(X \geq t) \leq \frac{E[X]}{t} \quad \forall t \geq 0$
Chebyshev	$\text{Var}(X) < \infty$	$P(X - E[X] \geq t) \leq \frac{\text{Var}(X)}{t^2}$
Chernoff	M_X exists	$P(X \geq a) \leq \frac{E[e^{tX}]}{e^{ta}} \quad \forall t > 0$ $P(X \leq a) \leq \frac{E[e^{tX}]}{e^{ta}} \quad \forall t < 0$

Concentration Inequalities

Name	Requirements	Bound
Markov	$X \geq 0, E[X]$ exists	$P(X \geq t) \leq \frac{E[X]}{t} \quad \forall t \geq 0$
Chebyshev	$\text{Var}(X) < \infty$	$P(X - E[X] \geq t) \leq \frac{\text{Var}(X)}{t^2}$
Chernoff	M_X exists	$P(X \geq a) \leq \frac{E[e^{tX}]}{e^{ta}} \quad \forall t > 0$ $P(X \leq a) \leq \frac{E[e^{tX}]}{e^{ta}} \quad \forall t < 0$

More restrictive



Concentration Inequalities

Name	Requirements	Bound
Markov	$X \geq 0, E[X]$ exists	$P(X \geq t) \leq \frac{E[X]}{t} \quad \forall t \geq 0$
Chebyshev	$\text{Var}(X) < \infty$	$P(X - E[X] \geq t) \leq \frac{\text{Var}(X)}{t^2}$
Chernoff	M_X exists <i>More restrictive</i>	$P(X \geq a) \leq \frac{E[e^{tX}]}{e^{ta}} \quad \forall t > 0$ $P(X \leq a) \leq \frac{E[e^{tX}]}{e^{ta}} \quad \forall t < 0$ <i>Tighter</i>

Break

Name	Requirements	Bound
Markov	$X \geq 0, E[X]$ exists	$P(X \geq t) \leq \frac{E[X]}{t} \quad \forall t \geq 0$
Chebyshev	$\text{Var}(X) < \infty$	$P(X - E[X] \geq t) \leq \frac{\text{Var}(X)}{t^2}$
Chernoff	M_X exists	$P(X \geq a) \leq \frac{E[e^{tX}]}{e^{ta}} \quad \forall t > 0$ $P(X \leq a) \leq \frac{E[e^{tX}]}{e^{ta}} \quad \forall t < 0$

Let Y be a r.v. that takes values in $[-1, 1]$ with mean -0.5 . Give an upper bound on the probability that $Y \geq 0.5$.

$$\text{Let } Z \equiv Y + 1 \quad P(Y \geq 0.5) = P(Z \geq 1.5) \leq \frac{0.5}{1.5} = \frac{1}{3}$$



(Weak) Law of large numbers

(Weak) Law of large numbers

Let X_i be independent identically distributed r.v.s with mean μ and variance σ^2 . If $Q_N \equiv \frac{1}{N} \sum_{i=1}^N X_i$, then $Q_N \rightarrow_p \mu$.

(Weak) Law of large numbers

Let X_i be independent identically distributed r.v.s with mean μ and variance σ^2 . If $Q_N \equiv \frac{1}{N} \sum_{i=1}^N X_i$, then $Q_N \rightarrow_p \mu$.

Proof:

$$E[Q_N] = \mu$$

$$\begin{aligned} \text{Var}(Q_N) &= \text{Var}\left(\frac{1}{N} \sum_i X_i\right) \\ &= \frac{1}{N^2} \text{Var}\left(\sum_i X_i\right) \\ &= \frac{1}{N^2} \sum_i \text{Var}(X_i) \quad (\text{Bienaymé}) \\ &= \frac{1}{N^2} N \sigma^2 = \frac{\sigma^2}{N} \end{aligned}$$

$$\begin{aligned} P(|Q_N - \mu| \geq \varepsilon) &\leq \frac{\text{Var}(Q_N)}{\varepsilon^2} \quad (\text{Chebyshev}) \\ &= \frac{\sigma^2}{N \varepsilon^2} \end{aligned}$$

$$P(|Q_N - \mu| \geq \varepsilon) \rightarrow 0 \text{ as } N \rightarrow \infty \quad \forall \varepsilon > 0 \quad \therefore Q_N \rightarrow_p \mu \quad \square$$

Law of Large Numbers

Two somewhat astounding takeaways:

Law of Large Numbers


Two somewhat astounding takeaways:



Law of Large Numbers

Two somewhat astounding takeaways:


1. Standard deviation decays at $\frac{1}{\sqrt{N}}$ *regardless of dimension.*



Law of Large Numbers

Two somewhat astounding takeaways:

1. Standard deviation decays at $\frac{1}{\sqrt{N}}$ regardless of dimension.



2. You can estimate the "standard error" with

$$SE = \frac{s}{\sqrt{N}}$$

← empirical estimate of standard deviation of X_i

where s is the sample standard deviation.

Confidence Intervals

Confidence Intervals

How do you estimate $|Q_N - \mu|$?

Confidence Intervals

How do you estimate $|Q_N - \mu|$?

Given a random variable Q , a γ **Confidence Interval**, $[u(Q), v(Q)]$, is a random interval that contains μ with probability γ , i.e.

$$P(u(Q) \leq \mu \leq v(Q)) = \gamma$$

Confidence Intervals

How do you estimate $|Q_N - \mu|$?

Given a random variable Q , a γ **Confidence Interval**, $[u(Q), v(Q)]$, is a random interval that contains μ with probability γ , i.e.

$$P(u(Q) \leq \mu \leq v(Q)) = \gamma$$

Example: $Q_N \equiv \frac{1}{N} \sum_{i=1}^N X_i$

Confidence Intervals

How do you estimate $|Q_N - \mu|$?

Given a random variable Q , a γ **Confidence Interval**, $[u(Q), v(Q)]$, is a random interval that contains μ with probability γ , i.e.

$$P(u(Q) \leq \mu \leq v(Q)) = \gamma$$

Example: $Q_N \equiv \frac{1}{N} \sum_{i=1}^N X_i$

Idea for approximate confidence interval: estimate $\text{Var}(Q_N)$ with $SE^2 = \frac{s^2}{N}$ and use Chebyshev.

Confidence Intervals

Idea for approximate confidence interval: estimate $\text{Var}(Q_N)$ with $SE^2 = \frac{s^2}{N}$ and use Chebyshev.

Confidence Intervals

Idea for approximate confidence interval: estimate $\text{Var}(Q_N)$ with $SE^2 = \frac{s^2}{N}$ and use Chebyshev.

Use $\gamma = 0.95$

Confidence Intervals

Idea for approximate confidence interval: estimate $\text{Var}(Q_N)$ with $SE^2 = \frac{s^2}{N}$ and use Chebyshev.

Use $\gamma = 0.95$

$$P(|X - E[X]| \geq t) \leq \frac{\text{Var}(X)}{t^2} = 1 - \gamma = 0.05$$

Confidence Intervals

Idea for approximate confidence interval: estimate $\text{Var}(Q_N)$ with $SE^2 = \frac{s^2}{N}$ and use Chebyshev.

Use $\gamma = 0.95$

$$P(|X - E[X]| \geq t) \leq \frac{\text{Var}(X)}{t^2} = 1 - \gamma = 0.05$$

$$t = \sqrt{\frac{\text{Var}(X)}{0.05}}$$

Confidence Intervals

Idea for approximate confidence interval: estimate $\text{Var}(Q_N)$ with $SE^2 = \frac{s^2}{N}$ and use Chebyshev.

Use $\gamma = 0.95$

$$P(|X - E[X]| \geq t) \leq \frac{\text{Var}(X)}{t^2} = 1 - \gamma = 0.05$$

$$t = \sqrt{\frac{\text{Var}(X)}{0.05}}$$

$$t \approx \frac{SE}{\sqrt{0.05}} \approx 4.47SE$$

Confidence Intervals

Idea for approximate confidence interval: estimate $\text{Var}(Q_N)$ with $SE^2 = \frac{s^2}{N}$ and use Chebyshev.

Use $\gamma = 0.95$

$$P(|X - E[X]| \geq t) \leq \frac{\text{Var}(X)}{t^2} = 1 - \gamma = 0.05$$

$$t = \sqrt{\frac{\text{Var}(X)}{0.05}}$$

$$t \approx \frac{SE}{\sqrt{0.05}} \approx 4.47 SE$$

Approximate 95% CI: $[Q_N - 4.47 SE, Q_N + 4.47 SE]$

Confidence Intervals

Idea for approximate confidence interval: estimate $\text{Var}(Q_N)$ with $SE^2 = \frac{s^2}{N}$ and use Chebyshev.

Use $\gamma = 0.95$

$$P(|X - E[X]| \geq t) \leq \frac{\text{Var}(X)}{t^2} = 1 - \gamma = 0.05$$

$$t = \sqrt{\frac{\text{Var}(X)}{0.05}}$$

$$t \approx \frac{SE}{\sqrt{0.05}} \approx 4.47 SE$$

Approximate 95% CI: $[Q_N - 4.47 SE, Q_N + 4.47 SE]$

We can do much better if we know something about the distribution of Q_N !

Central Limit Theorem

Central Limit Theorem

Lindeberg-Levy CLT: If $\text{Var}[X_i] = \sigma^2 < \infty$, then

$$\sqrt{N}(Q_N - \mu) \xrightarrow{D} \mathcal{N}(0, \sigma)$$

Central Limit Theorem

Lindeberg-Levy CLT: If $\text{Var}[X_i] = \sigma^2 < \infty$, then

$$\sqrt{N}(Q_N - \mu) \xrightarrow{D} \mathcal{N}(0, \sigma)$$

After many samples Q_N starts to look
distributed like $\mathcal{N}(\mu, \frac{\sigma}{\sqrt{N}})$


Central Limit Theorem

Lindeberg-Levy CLT: If $\text{Var}[X_i] = \sigma^2 < \infty$, then

$$\sqrt{N}(Q_N - \mu) \xrightarrow{D} \mathcal{N}(0, \sigma)$$

After many samples Q_N starts to look distributed like $\mathcal{N}(\mu, \frac{\sigma}{\sqrt{N}})$

$$Q_1 \stackrel{D}{=} X_i$$




Central Limit Theorem

Lindeberg-Levy CLT: If $\text{Var}[X_i] = \sigma^2 < \infty$, then



$$\sqrt{N}(Q_N - \mu) \xrightarrow{D} \mathcal{N}(0, \sigma)$$

After many samples Q_N starts to look distributed like $\mathcal{N}(\mu, \frac{\sigma}{\sqrt{N}})$

$Q_1 \stackrel{D}{=} X_i$



Q_{10}



Central Limit Theorem


Lindeberg-Levy CLT: If $\text{Var}[X_i] = \sigma^2 < \infty$, then

$$\sqrt{N}(Q_N - \mu) \xrightarrow{D} \mathcal{N}(0, \sigma)$$



After many samples Q_N starts to look distributed like $\mathcal{N}(\mu, \frac{\sigma}{\sqrt{N}})$



$Q_1 \stackrel{D}{=} X_i$




Q_{10}



Q_{100}

Q_{1000}



Confidence Intervals

Idea for approximate confidence interval: estimate $\text{Var}(Q_N)$ with $SE^2 = \frac{s^2}{N}$ and use ~~Chebyshev~~ the central limit theorem.

Confidence Intervals

Idea for approximate confidence interval: estimate $\text{Var}(Q_N)$ with $SE^2 = \frac{s^2}{N}$ and use ~~Chebyshev~~ the central limit theorem. Use $\gamma = 0.95$

Confidence Intervals

Idea for approximate confidence interval: estimate $\text{Var}(Q_N)$ with $SE^2 = \frac{s^2}{N}$ and use ~~Chebyshev~~ the central limit theorem.

Use $\gamma = 0.95$

For a normal distribution,

$$P(|X - \mu| \geq t) = 1 - \text{erf}\left(\frac{t - \mu}{\sqrt{2}\sigma}\right)$$


Confidence Intervals

Idea for approximate confidence interval: estimate $\text{Var}(Q_N)$ with $SE^2 = \frac{s^2}{N}$ and use ~~Chebyshev~~ the central limit theorem.

Use $\gamma = 0.95$

For a normal distribution,

$$P(|X - \mu| \geq t) = 1 + \text{erf}\left(\frac{t - \mu}{\sqrt{2}\sigma}\right)$$




Confidence Intervals

Idea for approximate confidence interval: estimate $\text{Var}(Q_N)$ with $SE^2 = \frac{s^2}{N}$ and use ~~Chebyshev~~ the central limit theorem.

Use $\gamma = 0.95$

For a normal distribution,

$$P(|X - \mu| \geq t) = 1 + \text{erf}\left(\frac{t - \mu}{\sqrt{2}\sigma}\right)$$



$$t \approx 1.96SE$$


Confidence Intervals

Idea for approximate confidence interval: estimate $\text{Var}(Q_N)$ with $SE^2 = \frac{s^2}{N}$ and use ~~Chebyshev~~ the central limit theorem.

Use $\gamma = 0.95$

For a normal distribution,

$$P(|X - \mu| \geq t) = 1 + \text{erf}\left(\frac{t - \mu}{\sqrt{2}\sigma}\right)$$



$$t \approx 1.96 SE$$

Approximate 95% CI: $[Q_N - 1.96 SE, Q_N + 1.96 SE]$


Confidence Intervals

Idea for approximate confidence interval: estimate $\text{Var}(Q_N)$ with $SE^2 = \frac{s^2}{N}$ and use ~~Chebyshev~~ the central limit theorem.

Use $\gamma = 0.95$

For a normal distribution,

$$P(|X - \mu| \geq t) = 1 + \text{erf}\left(\frac{t - \mu}{\sqrt{2}\sigma}\right)$$



$$t \approx 1.96 SE$$

Approximate 95% CI: $[Q_N - 1.96 SE, Q_N + 1.96 SE]$

(Chebyshev gave 4.47)

Importance Sampling

Want to estimate $X \sim p$ with samples from $Y_i \sim q$.

Importance Sampling

Want to estimate $X \sim p$ with samples from $Y_i \sim q$.

$$E[X] = \int x p(x) dx$$

Importance Sampling

Want to estimate $X \sim p$ with samples from $Y_i \sim q$.

$$\begin{aligned} E[X] &= \int x p(x) dx \\ &= \int x \frac{p(x)}{q(x)} q(x) dx \end{aligned}$$

Importance Sampling


Want to estimate $X \sim p$ with samples from $Y_i \sim q$.

$$\begin{aligned} E[X] &= \int x p(x) dx \\ &= \int x \left(\frac{p(x)}{q(x)} \right) q(x) dx \\ &\approx \frac{1}{N} \sum Y_i \frac{p(Y_i)}{q(Y_i)} \end{aligned}$$

Importance Sampling

Want to estimate $X \sim p$ with samples from $Y_i \sim q$.

$$\begin{aligned} E[X] &= \int x p(x) dx \\ &= \int x \frac{p(x)}{q(x)} q(x) dx \\ &\approx \frac{1}{N} \sum Y_i \frac{p(Y_i)}{q(Y_i)} \quad \swarrow \\ &\approx \frac{1}{N} \sum Y_i w_i \quad \swarrow \\ \text{where } w_i &= \frac{p(Y_i)}{q(Y_i)} \end{aligned}$$



Summary

1. Concentration Inequalities

$$P(X \geq t) \leq \phi(t)$$

2. Law of large numbers

$$Q_N \rightarrow_p \mu$$

3. Central Limit Theorem

$$Q_N \xrightarrow{D} \mathcal{N}(\mu, \frac{\sigma}{\sqrt{N}})$$

4. Importance Sampling

$$E[X] \approx \frac{1}{N} \sum Y_i w_i$$

where $w_i = \frac{p(Y_i)}{q(Y_i)}$