

## The American Mathematical Monthly



ISSN: 0002-9890 (Print) 1930-0972 (Online) Journal homepage: https://www.tandfonline.com/loi/uamm20

# A One-Sentence Proof That Every Prime $p \equiv 1 \pmod{4}$ Is a Sum of Two Squares

## D. Zagier

To cite this article: D. Zagier (1990) A One-Sentence Proof That Every Prime  $p \equiv 1 \pmod{4}$  Is a Sum of Two Squares, The American Mathematical Monthly, 97:2, 144-144, DOI: 10.1080/00029890.1990.11995565

To link to this article: https://doi.org/10.1080/00029890.1990.11995565



## THE TEACHING OF MATHEMATICS

EDITED BY MELVIN HENRIKSEN AND STAN WAGON

## A One-Sentence Proof That Every Prime $p \equiv 1 \pmod{4}$ Is a Sum of Two Squares

#### D. ZAGIER

Department of Mathematics, University of Maryland, College Park, MD 20742

The involution on the finite set  $S = \{(x, y, z) \in \mathbb{N}^3 : x^2 + 4yz = p\}$  defined by

$$(x,y,z) \mapsto \begin{cases} (x+2z, z, y-x-z) & \text{if } x < y-z \\ (2y-x, y, x-y+z) & \text{if } y-z < x < 2y \\ (x-2y, x-y+z, y) & \text{if } x > 2y \end{cases}$$

has exactly one fixed point, so |S| is odd and the involution defined by  $(x, y, z) \mapsto (x, z, y)$  also has a fixed point.  $\square$ 

This proof is a simplification of one due to Heath-Brown [1] (inspired, in turn, by a proof given by Liouville). The verifications of the implicitly made assertions—that S is finite and that the map is well-defined and involutory (i.e., equal to its own inverse) and has exactly one fixed point—are immediate and have been left to the reader. Only the last requires that p be a prime of the form 4k + 1, the fixed point then being (1,1,k).

Note that the proof is not constructive: it does not give a method to actually find the representation of p as a sum of two squares. A similar phenomenon occurs with results in topology and analysis that are proved using fixed-point theorems. Indeed, the basic principle we used: "The cardinalities of a finite set and of its fixed-point set under any involution have the same parity," is a combinatorial analogue and special case of the corresponding topological result: "The Euler characteristics of a topological space and of its fixed-point set under any continuous involution have the same parity."

For a discussion of constructive proofs of the two-squares theorem, see the Editor's Corner elsewhere in this issue.

#### REFERENCE

1. D. R. Heath-Brown, Fermat's two-squares theorem, *Invariant* (1984) 3-5.

## **Inverse Functions and their Derivatives**

### ERNST SNAPPER

Department of Mathematics and Computer Science, Dartmouth College, Hanover, NH 03755

If the concept of inverse function is introduced correctly, the usual rule for its derivative is visually so obvious, it barely needs a proof. The reason why the standard, somewhat tedious proofs are given is that the inverse of a function f(x) is