

Figure 1 | Islands and ports as windows onto butterfly resource geography



No historical host shipment, larval feeding, butterfly movement or breeding inferred.


Distinct grains: 239 resource butterflies; 98 botanical island regions; 225 candidates/51 landmass groups. Diagrams do not establish a port-to-plant-to-butterfly colonisation path.


Figure 2 | Islands amplify the potential larval-host resources of butterflies




Source: 239 butterflies / frozen HOSTS–WCVP; 126 insular, 237 mainland WGSRPD3 regions. Overall +54.9% potential resource units (206 species gained); NOT butterfly occurrence.

Figure 3 | Port-rich islands contain introduced floras that include butterfly hosts









Source: 1,596 complete port queries / 142 country codes / 687 candidate host plants / 25,742 plant×port records; 98 island regions, 38 early port buffers. Not plant carriage or larval use.


Figure 4 | Butterfly records and compatible introduced-host geography are associated





Distinct sources: A 225 butterfly species; B 51 landmasses/1,275 dependent pairs; C 1,163 butterfly×pair records; D 155 species/208,482 scores (contemporary hosts). No feeding, shipment or colonisation inferred.

Figure 5 | Butterfly indexing supports geographic association, not a demonstrated island-to-island mechanism









Different data supports: A 113 first-index cells/17 landmasses; B 83 indexed butterflies×landmass cells; C/D 225 candidate taxa×51 sampled landmasses, 199 matched trait taxa. All first INDEX records, not true colonisation.